

Abundant Stable Gauge Field Hair for Black Holes in Anti-de Sitter Space

BAXTER, Erik <<http://orcid.org/0000-0002-7524-7353>>, HELBLING, Marc and WINSTANLEY, Elizabeth

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/11461/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

BAXTER, Erik, HELBLING, Marc and WINSTANLEY, Elizabeth (2008). Abundant Stable Gauge Field Hair for Black Holes in Anti-de Sitter Space. *Physical Review Letters*, 100 (1).

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

Abundant stable gauge field hair for black holes in anti-de Sitter space

J. E. Baxter,¹ Marc Helbling,² and Elizabeth Winstanley^{1,*}

¹*Department of Applied Mathematics, The University of Sheffield, Hicks Building, Hounsfield Road, Sheffield, S3 7RH, United Kingdom.*

²*INSA de Rouen, Laboratoire de Mathématiques (LMI), Place Emile Blondel BP 08, 76131 Mont Saint Aignan Cedex, France.*

(Dated: February 22, 2013)

We present new hairy black hole solutions of $\mathfrak{su}(N)$ Einstein-Yang-Mills theory (EYM) in asymptotically anti-de Sitter (adS) space. These black holes are described by $N + 1$ independent parameters, and have $N - 1$ independent gauge field degrees of freedom. Solutions in which all gauge field functions have no zeros exist for all N , and for sufficiently large (and negative) cosmological constant. At least some of these solutions are shown to be stable under classical, linear, spherically symmetric perturbations. Therefore there is no upper bound on the amount of stable gauge field hair with which a black hole in adS can be endowed.

PACS numbers: 04.20.Jb, 04.40.Nr, 04.70.Bw

The “no-hair” conjecture [1] states that black hole equilibrium states possess extremely simple geometries, determined completely by the mass, angular momentum and charge of the black hole. While hairy black hole solutions of the Einstein equations have been discovered, particularly in Einstein-Yang-Mills (EYM) theory and its variants (see [2] for a review), many of the plethora of new black hole solutions found in the literature are classically unstable. Those hairy black holes which are stable (such as the $\mathfrak{su}(2)$ EYM black holes in anti-de Sitter space (adS) [3, 4]) have, at least to date, been described by only a small number of parameters additional to the mass, angular momentum and charge of the black hole. This means that the “spirit” if not the “letter” of the no-hair conjecture is maintained.

In recent years there has been an explosion of interest in hairy black holes in adS, partly because at least some of these configurations are stable, but also because of the importance of the adS/CFT correspondence [5] in string theory. In particular, it has been suggested [6] that there should be observables in the dual (deformed) CFT which are sensitive to the presence of black hole hair (see also [7] for an adS/CFT interpretation of some stable seven-dimensional black holes with $\mathfrak{so}(5)$ gauge fields). Our purpose in this letter is to present new stable, asymptotically adS, hairy black hole solutions of $\mathfrak{su}(N)$ EYM for sufficiently large $|\Lambda|$ which are described by an unbounded number of parameters. The existence of these solutions casts the status of the “no-hair” conjecture in a completely new light: equilibrium black holes in adS are no longer simple objects, but rather require an infinite number of parameters in order to fully determine their geometry.

We consider static, spherically symmetric, four-dimensional black holes with metric

$$ds^2 = -\mu S^2 dt^2 + \mu^{-1} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2, \quad (1)$$

where the metric functions μ and S depend on the ra-

dial co-ordinate r only. Here, and throughout this letter, the metric has signature $(-, +, +, +)$ and we use units in which $4\pi G = c = 1$. In the presence of a negative cosmological constant Λ , we write the metric function μ as

$$\mu(r) = 1 - \frac{2m(r)}{r} - \frac{\Lambda r^2}{3}. \quad (2)$$

The most general, spherically symmetric, ansatz for the $\mathfrak{su}(N)$ gauge potential has been given in [8]. Here, we assume that the gauge potential is purely magnetic and has the gauge-fixed form:

$$\mathcal{A} = \frac{1}{2} (C - C^H) d\theta - \frac{i}{2} [(C + C^H) \sin \theta + D \cos \theta] d\phi, \quad (3)$$

where C is an $(N \times N)$ upper-triangular matrix with non-zero entries immediately above the diagonal:

$$C_{j,j+1} = \omega_j(r), \quad (4)$$

for $j = 1, \dots, N - 1$, with C^H the Hermitian conjugate of C , and D is a constant diagonal matrix:

$$D = \text{Diag}(N - 1, N - 3, \dots, -N + 3, -N + 1). \quad (5)$$

The $(N - 1)$ Yang-Mills equations take the form

$$r^2 \mu \omega_j'' + \left(2m - 2r^3 p_\theta - \frac{2\Lambda r^3}{3} \right) \omega_j' + W_j \omega_j = 0 \quad (6)$$

for $j = 1, \dots, N - 1$, where a prime $'$ denotes d/dr , and

$$p_\theta = \frac{1}{4r^4} \sum_{j=1}^N [(\omega_j^2 - \omega_{j-1}^2 - N - 1 + 2j)^2], \quad (7)$$

$$W_j = 1 - \omega_j^2 + \frac{1}{2} (\omega_{j-1}^2 + \omega_{j+1}^2), \quad (8)$$

with $\omega_0 = \omega_N = 0$. The Einstein equations take the form

$$m' = \mu G + r^2 p_\theta, \quad \frac{S'}{S} = \frac{2G}{r}, \quad (9)$$

where

$$G = \sum_{j=1}^{N-1} \omega_j'^2. \quad (10)$$

The field equations (6,9) have the following trivial solutions. Setting $\omega_j(r) \equiv \pm\sqrt{j(N-j)}$ for all j gives the Schwarzschild-adS black hole with $m(r) = M = \text{constant}$ (which can be set to zero to give pure adS space). Setting $\omega_j(r) \equiv 0$ for all j gives the Reissner-Nordström-adS black hole with magnetic charge. There is an additional special class of solutions, given by setting

$$\omega_j(r) = \pm\sqrt{j(N-j)}\omega(r) \quad \forall j = 1, \dots, N-1. \quad (11)$$

In this case, it is possible to show, using a rescaling method along the lines of that in [9], that the field variables $\omega(r)$, $m(r)$ and $S(r)$ satisfy the $\mathfrak{su}(2)$ EYM field equations with a negative cosmological constant. Furthermore, the boundary conditions (as discussed below) are also preserved. Therefore any $\mathfrak{su}(2)$, asymptotically adS, EYM black hole solution can be embedded into $\mathfrak{su}(N)$ EYM to give another asymptotically adS black hole.

In this letter we study black hole solutions of the field equations (6,9), returning to soliton solutions elsewhere [10]. We assume there is a regular, non-extremal, black hole event horizon at $r = r_h$. The field variables $\omega_j(r)$, $m(r)$ and $S(r)$ will have regular Taylor series expansions about $r = r_h$. These expansions are determined by the $N+1$ quantities $\omega_j(r_h)$, r_h , $S(r_h)$ for fixed cosmological constant Λ . Since the field equations (6,9) are invariant under the transformation $\omega_j(r) \rightarrow -\omega_j(r)$ (for any j independently), we may consider $\omega_j(r_h) > 0$ without loss of generality. For the event horizon to be non-extremal, it must be the case that

$$2m'(r_h) = 2r_h^2 p_\theta(r_h) < 1 - \Lambda r_h^2, \quad (12)$$

which constrains the possible values of the gauge field functions $\omega_j(r_h)$ at the event horizon. At infinity, the boundary conditions are considerably less stringent than in the asymptotically flat case. In order for the metric (1) to be asymptotically adS, we simply require that the field variables $\omega_j(r)$, $m(r)$ and $S(r)$ converge to constant values as $r \rightarrow \infty$, and have regular Taylor series expansions in r^{-1} near infinity. Since $\Lambda < 0$, there is no cosmological horizon.

The field equations (6,9) are integrated numerically using standard ‘shooting’ techniques [11]. The equation for $S(r)$ decouples from the other Einstein equation and the Yang-Mills equations so can be integrated separately if required. We start integrating just outside the event horizon, using as our shooting parameters the N variables $\omega_j(r_h)$ and r_h , subject to the weak constraint (12). The field equations are then integrated outwards in the radial co-ordinate r until either the field variables start

to diverge or they have converged to the asymptotic form at infinity.

As in the $\mathfrak{su}(2)$ case [3], we find black hole solutions in open subsets of the N -dimensional parameter space $(\omega_j(r_h), r_h)$ for fixed Λ . For sufficiently large $|\Lambda|$ (where how large ‘sufficiently large’ is depends on the radius of the event horizon r_h), we find that the gauge field functions $\omega_j(r)$ all have no zeros. In figure 1 we show a typical nodeless solution, for $\mathfrak{su}(4)$ EYM. It can be seen




FIG. 1: A typical black hole solution of $\mathfrak{su}(4)$ EYM in which all the gauge field functions $\omega_j(r)$ are nodeless. For this solution, $\Lambda = -10$ and $r_h = 1$. The values of the gauge field functions on the event horizon are: $\omega_1(r_h) = 2.3$, $\omega_2(r_h) = 2.6$ and $\omega_3(r_h) = 2.2$.

that the metric functions $m(r)$ and $S(r)$ have very similar behaviour to the $\mathfrak{su}(2)$ case, and that, since $|\Lambda|$ is so large, the gauge field functions do not vary significantly from their values at the event horizon.

The phase space of black hole solutions in the $\mathfrak{su}(3)$ case, with $\Lambda = -10$ and $r_h = 1$ is shown in figure 2, and is typical of the phase space for large values of $|\Lambda|$. In figure 2 we have examined, for $\Lambda = -10$ and $r_h = 1$, all values of the $\omega_1(r_h)$ and $\omega_2(r_h)$ which satisfy the constraint (12). The inequality in (12) is saturated on the outer-most curve in figure 2. It can be seen from figure 2 that not all values of $(\omega_1(r_h), \omega_2(r_h))$ give black hole solutions; those values for which no regular black hole solution satisfying the boundary conditions at infinity could be found lie in the narrow band on the outside of the plot. The region between this narrow band and the coordinate axes contains black hole solutions in which both gauge field functions $\omega_1(r)$ and $\omega_2(r)$ have no zeros. We have also plotted in figure 2 the line $\omega_1(r_h) = \omega_2(r_h)$, on which lie embedded $\mathfrak{su}(2)$ solutions given by (11). The significance of the shaded region in figure 2 will be described shortly. More detailed properties of the phase space of black hole solutions will be discussed elsewhere [10].




FIG. 2: Phase space of black hole solutions in $\mathfrak{su}(3)$ EYM with $\Lambda = -10$ and $r_h = 1$. The shaded region shows where solutions exist which satisfy the inequalities (16) at the event horizon.

In [3], the existence of black hole solutions for which the gauge function $\omega(r)$ had no zeros was proven analytically in the $\mathfrak{su}(2)$ case. Since $\mathfrak{su}(2)$ solutions can be embedded as $\mathfrak{su}(N)$ solutions via (11), we have automatically an analytic proof of the existence of nodeless $\mathfrak{su}(N)$ EYM black holes in adS. However, these embedded solutions are ‘trivial’ in the sense that they are described by just three parameters: r_h , Λ and $\omega(r_h)$. An important question is whether the existence of ‘non-trivial’ (that is, genuinely $\mathfrak{su}(N)$) solutions in which all the gauge field functions $\omega_j(r)$ have no zeros can be proven analytically. The answer to this question is affirmative, and involves a generalization to $\mathfrak{su}(N)$ of the continuity-type argument used in [3]. The details are lengthy and will be presented elsewhere. However, the main thrust of the argument can be simply stated. We firstly prove (generalizing the analysis of [9] to include Λ) that the field equations (6,9) and initial conditions at the event horizon possess, locally in a neighborhood of the horizon, solutions which are analytic in r , r_h , Λ and the parameters $\omega_j(r_h)$. This enables us to prove that, in a sufficiently small neighborhood of any embedded $\mathfrak{su}(2)$ solution in which $\omega(r)$ has no nodes, there exists (at least in a neighborhood of the event horizon) an $\mathfrak{su}(N)$ solution in which all the $\omega_j(r)$ have no nodes. The key part of the proof lies in then showing that these $\mathfrak{su}(N)$ solutions can be extended out to $r \rightarrow \infty$ and that they satisfy the boundary conditions at infinity. This gives genuinely $\mathfrak{su}(N)$ black hole solutions in which all the gauge field functions have no zeros, and which are characterized by the $N + 1$ parameters r_h , Λ and $\omega_j(r_h)$.

The other outstanding question is whether these new black holes, with potentially unbounded amounts of

gauge field hair, are stable. We consider linear, spherically symmetric perturbations only for simplicity. Even for spherically symmetric perturbations, the analysis is highly involved in the $\mathfrak{su}(N)$ case and the details will be presented elsewhere. Here we briefly outline just the key features.

Firstly we consider spherically symmetric perturbations of the gauge potential (3), fixing the gauge so that the perturbed potential is purely magnetic and has the form [8]

$$A = \mathcal{B} dr + \frac{1}{2} (C - C^H) d\theta - \frac{i}{2} [(C + C^H) \sin \theta + D \cos \theta] d\phi. \quad (13)$$

Here, the matrices \mathcal{B} and C depend on both t and r , and matrix D is still constant and given by (5). The matrix $\mathcal{B}(t, r)$ is traceless, diagonal and has purely imaginary entries. The only non-zero entries of the matrix $C(t, r)$ are:

$$C_{j,j+1}(t, r) = \omega_j(t, r) \exp(i\gamma_j(t, r)). \quad (14)$$

As usual, the metric retains the form (1) but now the functions m and S depend on both t and r . With this choice of gauge potential (13), the perturbation equations decouple into two sectors:

- the *sphaleronic sector* consisting of entries of \mathcal{B} and the functions γ_j ;
- the *gravitational sector* which consists of the perturbations of the metric functions δm and δS and the perturbations of the gauge field functions $\delta \omega_j$.

The form of the perturbation equations in the sphaleronic sector is little changed from the asymptotically flat case [12]. It consists of $2N - 1$ coupled equations for the $2N - 1$ variables (N diagonal entries of the matrix \mathcal{B} and $N - 1$ functions γ_j). In addition, there is the *Gauss constraint*, which gives N coupled consistency conditions. After much algebra (along the lines of [12]), the sphaleronic sector perturbation equations can be cast in the form

$$-\ddot{\Psi} = \mathcal{U}\Psi, \quad (15)$$

where a dot denotes $\partial/\partial t$, the $(2N - 1)$ -dimensional vector Ψ consists of combinations of perturbations and \mathcal{U} is a self-adjoint, second order, differential operator (involving derivatives with respect to r but not t), depending on the equilibrium functions $\omega_j(r)$, $m(r)$ and $S(r)$. It can be shown that the operator \mathcal{U} is regular and positive provided the unperturbed gauge functions $\omega_j(r)$ have no zeros and satisfy the $N - 1$ inequalities

$$\omega_j^2 > 1 + \frac{1}{2} (\omega_{j+1}^2 + \omega_{j-1}^2) \quad (16)$$

for all $j = 1, \dots, N - 1$. These inequalities define a non-empty subset of the parameter space, which is shown in the $\mathfrak{su}(3)$ case in figure 2.

The shaded region in figure 2 shows where the inequalities (16) are satisfied for the gauge field functions at the event horizon. However, the requirements of (16) are considerably stronger, as the inequalities have to be satisfied for *all* $r \geq r_h$. Our analytic work shows that, in fact, for sufficiently large $|\Lambda|$, there do exist solutions to the field equations for which the inequalities (16) are indeed satisfied for all r (an example of such a solution is shown in figure 3). This involves proving that for at least some




FIG. 3: An example of an $\mathfrak{su}(3)$ solution for which the inequalities (16) are satisfied for all $r \geq r_h$. In this example, $\Lambda = -10$, $r_h = 1$ and the values of the gauge field functions at the event horizon are $\omega_1(r_h) = 2$, $\omega_2(r_h) = 1.95$.

solutions for which the gauge field function values at the event horizon lie within the region where the inequalities (16) are satisfied, the gauge field functions remain within this open region.

For the gravitational sector, the metric perturbations can be eliminated to yield a set of $N - 1$, coupled perturbation equations of the form

$$-\delta\ddot{\omega} = \mathcal{M}\delta\omega, \quad (17)$$

where $\delta\omega = (\delta\omega_1, \dots, \delta\omega_{N-1})^T$, and \mathcal{M} is a self-adjoint, second order, differential operator (involving derivatives with respect to r but not t), depending on the equilibrium functions $\omega_j(r)$, $m(r)$ and $S(r)$. The operator \mathcal{M} is more difficult to analyze than the operator \mathcal{U} . For sufficiently large $|\Lambda|$, it can be shown that \mathcal{M} is a positive operator for embedded $\mathfrak{su}(2)$ solutions, provided that $\omega^2(r) > 1$ for all r (the existence of such $\mathfrak{su}(2)$ solutions is proved, for sufficiently large $|\Lambda|$, in [3]). As described

above, our analytic work ensures the existence of genuinely $\mathfrak{su}(N)$ solutions in a sufficiently small neighborhood of these embedded $\mathfrak{su}(2)$ solutions. These $\mathfrak{su}(N)$ solutions are such that the inequalities (16) are satisfied for all $r \geq r_h$ (and therefore the solutions are stable under sphaleronic perturbations). The positivity of \mathcal{M} can then be extended to these genuinely $\mathfrak{su}(N)$ solutions using an analyticity argument, based on the nodal theorem of [13]. The technical details of this argument will be presented elsewhere. Therefore at least some of our solutions are linearly stable in both the gravitational and sphaleronic perturbation sectors.

For sufficiently large $|\Lambda|$ (for each fixed r_h), we have shown the existence of $\mathfrak{su}(N)$ EYM black holes in adS, which are described by $N + 1$ parameters and are stable under linear, spherically symmetric perturbations. If the cosmological constant is very large and negative, there are potentially a very large number of possible gauge field configurations giving the same mass and magnetic charge at infinity. As explained in the introduction, we anticipate that these solutions may well have interesting consequences for the adS/CFT correspondence [5]. We hope to return to these questions in the near future.

We thank Eugen Radu for many informative discussions. The work of JEB is supported by UK EPSRC, and the work of EW is supported by UK PPARC, grant reference number PPA/G/S/2003/00082.

* Electronic address: E.Winstanley@sheffield.ac.uk

- [1] R. Ruffini and J. A. Wheeler, Phys. Today **24** 30 (1971).
- [2] M. S. Volkov and D. V. Gal'tsov, Phys. Rept. **319** 1 (1999).
- [3] E. Winstanley, Class. Quant. Grav. **16** 1963 (1999).
- [4] J. Bjoraker and Y. Hosotani, Phys. Rev. Lett. **84**, 1853 (2000); Phys. Rev. **D62** 043513 (2000).
- [5] J. Maldacena, Adv. Theor. Math. Phys. **2** 231 (1998); E. Witten, Adv. Theor. Math. Phys. **2** 253 (1998); *ibid* **2** 505 (1998).
- [6] T. Hertog and K. Maeda, JHEP **07** (2004) 051.
- [7] J. P. Gauntlett, N. Kim and D. Waldram, Phys. Rev. **D63** 126001 (2001).
- [8] H. P. Kunzle, Class. Quant. Grav. **8** 2283 (1991).
- [9] H. P. Kunzle, Comm. Math. Phys. **162** 371 (1994).
- [10] J. E. Baxter, M. Helbling and E. Winstanley, Phys. Rev. **D76** 104017 (2007).
- [11] W. Vetterling, W. Press, S. Teukolsky and B. Flannery, *Numerical Recipes in FORTRAN* (Cambridge University Press, 1992).
- [12] O. Brodbeck and N. Straumann, J. Math. Phys. **37** 1414 (1996).
- [13] H. Amann and P. Quittner, J. Math. Phys. **36** 4553 (1995).