

**The decision-making process for senior cancer patients:
treatment allocation of older women with operable breast
cancer in the UK.**

MORGAN, Jenna L, RICHARDS, Paul, ZAMAN, Osama, WARD, Sue,
COLLINS, Karen <<http://orcid.org/0000-0002-4317-142X>>, ROBINSON,
Thompson, CHEUNG, Kwok-Leung, AUDISIO, Riccardo A., REED, Malcolm
W and WYLD, Lynda

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/11477/>

This document is the author deposited version. You are advised to consult the
publisher's version if you wish to cite from it.

Published version

MORGAN, Jenna L, RICHARDS, Paul, ZAMAN, Osama, WARD, Sue, COLLINS,
Karen, ROBINSON, Thompson, CHEUNG, Kwok-Leung, AUDISIO, Riccardo A.,
REED, Malcolm W and WYLD, Lynda (2015). The decision-making process for
senior cancer patients: treatment allocation of older women with operable breast
cancer in the UK. *Cancer Biology & Medicine*, 12 (4), 308-315.

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

Article Type: Original Article

Title: The decision-making process for senior cancer patients: treatment allocation of older women with operable breast cancer in the UK.

Authors: Jenna L. Morgan¹, Paul Richards², Osama Zaman¹, Sue Ward², Karen Collins³, Thompson Robinson⁴, Kwok-Leung Cheung⁵, Riccardo Audisio⁶, Malcolm W. Reed⁷, Lynda Wyld¹, on behalf of the Bridging the Age Gap in Breast Cancer Trial Management Team.

1. Academic Unit of Surgical Oncology, University of Sheffield Medical School, Beech Hill Road, Sheffield, UK.
2. Health Economics and Decision Science, School for Health and Related Research, University of Sheffield, Sheffield, UK.
3. Centre for Health and Social Care Research, Sheffield Hallam University, Collegiate Crescent, Sheffield, UK.
4. Department of Cardiovascular Sciences, University of Leicester, Leicester, LE2 7LX, UK.
5. School of Medicine, University of Nottingham, Royal Derby Hospital Centre, Uttoxeter Road, Derby DE22 3DT, UK.
6. Department of Surgery, University of Liverpool, St Helens Teaching Hospital, Marshalls Cross Road, St Helens, WA9 3DA, UK.
7. Brighton and Sussex Medical School, University of Sussex, Falmer, Brighton, UK.

Corresponding author: Miss Jenna Morgan, Academic Department of Surgical Oncology, University of Sheffield Medical School, Beech Hill Road, Sheffield, S10 2RX, UK.

Email: j.morgan@sheffield.ac.uk. Tel: +44 (0)114 271 3611. Fax: +44 (0)114 271 3314.

Author Contributions:

Jenna Morgan – drafted the manuscript, performed data analysis and interpretation.

Paul Richards – drafted the manuscript, performed data analysis and interpretation.

Osama Zaman – collected data, performed data analysis and interpretation, commented on manuscript.

Sue Ward – commented on manuscript, performed data interpretation.

Karen Collins – commented on manuscript, involved in study design.

Thompson Robinson - commented on manuscript, involved in study design.

Kwok-Leung Cheung - commented on manuscript, involved in study design.

Riccardo Audisio - commented on manuscript, involved in study design.

Malcolm Reed – commented on manuscript, involved in study design, set up, management and deputy study lead.

Lynda Wyld – commented on manuscript, involved in study design, set up, management and overall study lead.

Source of Support: This paper presents independent research funded by the National Institute for Health Research (NIHR) under its Programme Grants for Applied Research Programme (Grant Reference Number RP-PG-1209-10071). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

Running Title: Treatment allocation of older women with operable breast cancer.

Abstract.

Objectives: Up to 40% of women over 70 years with primary operable breast cancer in the UK are treated with Primary Endocrine Therapy (PET) as an alternative to surgery. A variety of factors are important in determining treatment for older breast cancer patients. This study aimed to identify the patient and tumor factors associated with treatment allocation in this population.

Methods: Prospectively collected data on treatment received (surgery vs. PET) were analysed with multivariable logistic regression using the variables age, modified Charlson Comorbidity Index (CCI), Activities of Daily Living (ADL) score, Mini-Mental State Examination (MMSE) score, HER2 status, tumour size, grade and nodal status.

Results: Data were available for 1122 cancers in 1098 patients recruited between April 2013 and May 2015 from 51 UK hospitals. Seventy-eight per cent of the population were treated surgically, with the remainder being treated with PET. Increasing patient age at diagnosis, increasing CCI score, large tumor size (five cm or more) and dependence in one or more ADL categories were all strongly associated with non-surgical treatment ($p < 0.05$).

Conclusions: Increasing comorbidity, large tumor size and reduced functional ability are associated with reduced likelihood of surgical treatment of breast cancer in older patients. However, age itself remains a significant factor for non-surgical treatment; reinforcing the need for evidence-based guidelines.

Keywords.

Frail elderly, breast neoplasms, decision-making.

Introduction.

A third of new breast cancer diagnoses occur in women age over 70 years in the UK¹. The number of women affected will increase over time due to the ageing population². The prevalence of comorbidity and frailty increases with age, resulting in deaths from other causes exceeding breast cancer mortality in older women with breast cancer^{3,4}. Additionally, tolerance of some breast cancer therapies is reduced in this age group^{5,6}. The majority (about 90%) of older women are diagnosed with oestrogen receptor (ER) positive disease³. Older patients with operable breast cancer may be offered alternative treatment modalities, such as primary endocrine therapy (PET)^{7,8}, where ER positive disease is treated with endocrine therapy alone without surgery. Non-operative management of older breast cancer patients accounts for up to 40% of treatment in the UK⁹.

Primary endocrine therapy gained popularity in the 1980s for the management of older women after tamoxifen was shown to be effective in this setting¹⁰ and a succession of randomised controlled trials comparing its efficacy with surgery followed. A subsequent Cochrane review comparing PET with surgery in the over 70s demonstrated superior rates of local control with surgery but no difference in survival rates¹¹. However the studies included in the review were small and flawed by modern standards because tumor ER status was not always tested and the age range of the trials included relatively young older women with no documentation of health, fitness or frailty, in fact all patients were deemed fit for surgery under general anaesthesia and therefore frail patients were excluded. A recent review of case series indicated that older frailer women tend to be treated with PET in routine practice and have inferior survival rates as would be expected due to higher other-cause mortality¹². Since the trials included in the Cochrane review were conducted there have been significant increases in population life expectancy, so it is not clear that results of these trials are applicable in contemporary clinical practice.

Recent reports have advocated the use of PET only in the very old or frail¹³. Current national guidelines state that patients with operable breast cancer should be treated with surgery, and not PET, "irrespective of age" unless this is precluded by comorbidities¹⁴; whilst the International Society of Geriatric Oncology (SIOG) and European Society of Breast Cancer Specialists (EUSOMA) recommend that PET should only be offered to patients with a "short estimated life expectancy (less than 2 to three years), who are considered unfit for surgery... or who refuse surgery"¹⁵. However, as a large number of older women are treated with PET in UK and other countries it is not clear that this guidance is being followed consistently.

Population cohort studies have been carried out to investigate the factors associated with non-surgical treatment of breast cancer in older women and found that increasing age, comorbidities and

worsening scores on Activities of Daily Living (ADL) were all associated with lower rates of surgery¹⁶. An analysis of English cancer registry data showed that after adjusting for disease characteristics, age and comorbidity, there remained significant variation in the rate of PET use at the hospital level¹⁷. This may reflect genuine variation in practice, which would contradict current recommendations. However, this analysis could not account for other potential confounding factors such as frailty and cognitive function, which would be expected to influence treatment decision making. There are also limitations concerning data quality such as missing data due to the retrospective and routine nature of data collection. A further confounding factor is the influence of the patient's preference for one or other option if they are given a choice.

The study presented here aims to identify patient and tumor factors associated with treatment allocation in the older UK breast cancer population, including data on functional status, cognitive function and comorbidities, as well as disease characteristics. A planned interim analysis was undertaken using prospectively collected patient data from a large multicentre UK cohort study.

Materials and Methods.

Study Design.

This study is part of a prospective observational cohort study of women aged over 70 years diagnosed with operable primary breast cancer in 51 UK breast cancer units between February 2013 and June 2015. Recruitment for this trial is ongoing, and results presented here represent an interim analysis of the first 1451 patients. Data were collected on patient and tumor characteristics, and treatment type. Data collection time points for the study are at baseline (prior to any treatment being administered), six weeks post-treatment, six months post treatment, 12 months post-treatment, 18 months post-treatment and 24 months post-treatment. Analyses are restricted to data collected up to six months post-treatment as the outcome of interest was primary treatment allocation.

Analyses are also restricted to the subgroup of patients with operable ER positive disease at diagnosis. Variables used within the analysis are shown in table i. Patients with missing treatment data or with known ER negative disease were excluded from the analysis. Patients with missing ER status who did not receive endocrine therapy were assumed to be ER negative and also excluded from the analysis. Bilateral cancers are counted individually. Figure i shows the baseline features of the study population.

Statistical Analysis.

Primary treatment was dichotomised as “surgery” or “PET” according to whether or not the patient received surgery to the primary tumor. Simple logistic regression was used to investigate associations between complete covariates and treatment. The joint effect of patient level factors on the probability of surgical treatment was assessed using multivariable logistic regression.

Variables with missing data cannot be included in standard regression models, but excluding cases with missing values (complete case analysis) can produce biased results where data is not missing completely at random, as well as reducing statistical precision. Missing data on disease characteristics and comorbidity was handled using the method of multiple imputation by chained equations (MICE)¹⁸. Missing covariate values were imputed using predictive mean matching¹⁹, and all variables included in the main analysis were incorporated into the imputation models. Twenty-five completed datasets were used for subsequent analysis. Results of the regression models for each dataset were combined using the standard methods described by Rubin²⁰. Covariates with over 50% missing data were not included in the regression models.

The performance of the regression model was assessed by considering the Receiver-Operating Characteristic (ROC) curve of the model, along with its corresponding area under the curve (AUC) statistic. This is a measure of the ability of the model to discriminate between individuals treated with surgery and PET²¹. The AUC can range from 0.5 to 1, where a value of 1 indicates perfect discrimination, and a value of 0.5 less indicates a model which performs no better than random chance.

All analyses were conducted using the open source statistical programming language R (version 3.0.1). Multiple imputation was implemented using the “mi” package (version 1.0).

Ethics and Research Governance.

The study protocols were approved by the UK National Research Ethics Committee (12/LO/1808) and institutional approvals were granted at each site. Written informed consent was obtained from the patient, or patient consultee in the case of patients with cognitive impairment.

Results.

Data were available on 1122 ER positive breast cancers in 1098 patients with a treatment record of either primary surgery with adjuvant therapy as deemed appropriate by the treating clinician (“surgery”) or endocrine therapy without surgery (“PET”). Characteristics of the study population can be seen in table i.

Seventy-eight per cent of the study population were treated surgically, with the remainder being treated with PET. The patients treated with surgery were younger overall than those treated with PET (see table i). The imputation algorithm converged successfully, and exploratory analysis of the distributions of the imputed values did not raise concerns about their validity or plausibility.

The multivariable logistic regression model is shown in Table ii. Increased age at diagnosis, increasing levels of comorbidity, large tumor size (five cm or more) and dependence in one or more ADL categories are all strongly associated with non-surgical treatment (Table ii). The ROC curve shows that the model discriminates well between individuals treated with PET and surgery, with the AUC statistic equal to 0.824 (95% CI, 0.792-0.855) (see Figure ii).

Discussion.

Of the 1122 ER positive operable breast cancers enrolled into the study to date, 78% were treated with surgery. Age was confirmed as one of the most important factors in determining treatment in older women with operable ER positive breast cancer, with increasing age associated with higher rates of PET, even accounting for other patient and disease characteristics. This is consistent with other studies^{7,22,23} despite current UK guidelines, which state that patients should be treated with surgery and appropriate adjuvant therapy “irrespective of age”¹⁴.

Increasing age was associated with increasing rates of comorbidity and multi-morbidity²⁴, this in turn may reduce the survival benefit of more aggressive breast cancer therapies, such as surgery⁴. It was therefore unsurprising that higher levels of co-morbidity in the study population are strongly associated with lower levels of surgical treatment, however this does not account for the effect seen with age.

Functional dependence in one or more of the ADL categories was strongly associated with non-surgical treatment in this study. Rates of functional dependence also increase with age and have been shown to affect treatment for in a small population of older breast cancer patients²².

Functional dependence has been associated with increased rates of post-operative complications and longer hospital stays in elderly cancer patients²⁵, possibly explaining why it has an impact on treatment allocation here.

Additionally, large tumor size (five cm or more) was also associated with higher rates of PET. Tumors of this size usually mandate a mastectomy²⁶, and so these findings may represent patients and clinicians trying to avoid more major surgery.

The rate of surgical treatment was higher in this study than some of the published figures for the UK, which range from 55-83%^{8,16,27-30}. This may result from a recruitment bias as patients within this study had to be recruited prior to commencing treatment, which was logistically more difficult with patients treated with PET as there is a shorter time period between diagnosis and treatment compared with surgery. Another possible bias could be that clinicians may be more likely to ‘follow the guidelines’ knowing that their data were being captured. This will ultimately result in an under-representation of PET patients, and may explain why Lavelle and colleagues found a similar rate of 83% having surgical treatment in their prospective cohort study¹⁶.

This study is limited by missing data, particularly with relation to ADL and MMSE, which is due, in part to patients opting out of completing the questionnaires. We have used multiple imputation to account for these missing data as it is less prone to bias than other commonly used methods to

account for missing data. However, whilst exploratory analysis of the imputed data suggested that the values were plausible, it is not possible to verify the extent to which the distribution of the imputed data accurately represents that of the missing values. By using 25 imputations, uncertainty around the missing data is incorporated into the probabilities used to adjust for case mix which mitigates against any small biases due to problems with the imputation model.

The findings of this study confirm existing research showing that age remains a significant predictor of non-surgical treatment after adjusting for disease characteristics and the underlying health status of the patient. However, it remains unclear as to whether or not current practice resulted in sub-optimal outcomes. An attempt to compare outcomes under PET and surgery with adjuvant endocrine therapy using a randomised clinical trial failed due to lack of recruitment, which was explained by an unwillingness on behalf of both patients and clinicians to choose treatment at random³¹. As a result, observational research is needed to investigate the effects of non-surgical treatment in this population. As recruitment and follow up of this cohort matures it will be possible to investigate the relationship between treatment, individual characteristics and outcomes such as survival and treatment failure. These data will be crucial in the development of evidence based guidance for patients and clinicians to help target treatments effectively.

Acknowledgements.

The authors would like to thank all the members of the Bridging the Age Gap in Breast Cancer Trial Management Group: Charlene Martin, Jenny Newhouse, Rosie Cooper, Alistair Ring, Stephen Walters, Tim Stephenson, Kirsty Pemberton, Tim Chater, Jaqui Gath, Lisa Caldon, Kate Lifford, Kate Brain. In addition, the authors would like to acknowledge the Principal Investigators and staff working at each of the recruiting sites: Ms J Dicks (Barnsley), Ms C Rogers (Doncaster and Bassetlaw), Ms S Edwards (Milton Keynes), Mr R Vijn (North Lincolnshire and Goole), Miss M Kaushik (Leicester), Ms S Ashworth (Burnley), Ms A Norton (Harrogate), Ms R Nasr (York), Mr C Holcolme (Royal Liverpool), Mr A Nejum (Airedale), Ms S Hartup (Leeds), Mr R Linford (Bradford), Ms C Morris (Cardiff and Vale), Ms H Sweetland (Cardiff), Ms A Williams (Nevill Hall Hospital), Mr S Parker (Coventry) Mr A Modi (Grantham), Ms E Gullaksen (Hull), Ms L Whisker (Nottingham), Mr A Haq (Southport), Miss V Pope (Leighton), Ms Karen Brooks (Royal Marsden), Ms J Forkes (Cheltenham), Mr M Douek (Guy's and St Thomas), Ms C Osborne (Dorset County), Ms S Miles-Dua (Broomfield), Ms S Buckley (Pinderfields), Ms K Kirby (Southmead, Bristol), Miss I Azmy (Chesterfield), Mr S Nicol (Rotherham), Miss S Seetharam (Dartford), Ms C Osborne (Yeovil), Mr S Joshi (Croyden), Ms A Chilvers (North

Tees), Mr Cheema (James Cook South Tees), Ms L Rivett (Luton and Dunstable), Miss R Ainsworth (Great Weston).

Conflict of Interest.

The authors declare they have no conflict of interest.

References

1. Office of National Statistics O. 2010 9 october 2014. New cases of cancer diagnosed in England, 2010: selected sites by age group and sex. In Cancer Registrations in England 2010. <<http://www.ons.gov.uk/ons/rel/vsob1/cancer-registrations-in-england/2010/rft-cancer-registrations-in-england--2010.xls>>. 9 october 2014.
2. Alberg AJ, Singh S. Epidemiology of breast cancer in older women: implications for future healthcare. *Drugs Aging* 2001;18(10):761-72.
3. Diab S, Elledge R, Clark G. Tumor characteristics and clinical outcome of elderly women with breast cancer. *Journal of the National Cancer Institute* 2000;92(7):550-556.
4. Satariano W, Ragland D. The effect of co-morbidity on 3-year survival of women with primary breast cancer. *Ann Intern Med* 1994;120:104–110.
5. Ring A, Reed M, Leonard R, Kunkler I, Muss H, Wildiers H, Fallowfield L, Jones A, Coleman R. The treatment of early breast cancer in women over the age of 70. *Br J Cancer* 2011;105(2):189-93.
6. Muss H, Berry D, Cirincione C, Budman D, Henderson I, Citron M, Norton L, Winer E, Hudis C. Toxicity of older and younger patients treated with adjuvant chemotherapy for node-positive breast cancer: the Cancer and Leukaemia Group B experience. Muss, HB, Berry, DA, Cirincione, C, Budman, DR, Henderson IC, Citron ML, Norton L, Winer EP, Hudis CA. *Journal Clinical Oncology* 2007;25(24):3699-3704.
7. Lavelle K, Todd C, Moran A, Howell A, Bundred N, Campbell M. Non-standard management of breast cancer increases with age in the UK: a population based cohort of women \geq 65 years. *British Journal of Cancer* 2007;96(8):1197-1203.
8. Wyld L, Garg DK, Kumar ID, Brown H, Reed MW. Stage and treatment variation with age in postmenopausal women with breast cancer: compliance with guidelines. *Br J Cancer* 2004;90(8):1486-91.
9. BCCOM. Breast Cancer Clinical Outcome Measures (BCCOM) Project: Analysis of the management of symptomatic breast cancers diagnosed in 2004. 3rd Year Report December 2007. 2007.
10. Preece P, Wood R, Mackie C, Cischieri A. Tamoxifen as initial sole treatment of localized breast-cancer in elderly women - a pilot-study. *British Medical Journal* 1982;284(6319):869-870.
11. Morgan J, Wyld L, Collins K, Reed M. Surgery versus primary endocrine therapy for operable primary breast cancer in elderly women (70 years plus). *Cochrane database systematic reviews* 2014;5:CD004272.
12. Morgan J, Reed M, Wyld L. Primary endocrine therapy as a treatment for older women with operable breast cancer - a comparison of randomised controlled trial and cohort study findings. *European Journal Surgical Oncology* 2014;40(6):676-684.
13. Chakrabarti J, Kenny F, Syed B, Robertson J, Blamey R, Cheung K. A randomised trial of mastectomy only versus tamoxifen for treating elderly patients with operable primary breast cancer-Final results at 20-year follow-up. *Critical Reviews in Oncology Hematology* 2011;78(3):260-264.
14. NICE. CG80 Early and locally advanced breast cancer: full guideline. London: National Institute for Clinical Excellence; 2009.
15. Biganzoli L, Wildiers H, Oakman C, Marotti L, Loibl S, Kunkler I, Reed M, Ciatto S, Voogd AC, Brain E and others. Management of elderly patients with breast cancer: updated recommendations of the International Society of Geriatric Oncology (SIOG) and European Society of Breast Cancer Specialists (EUSOMA). *Lancet Oncology* 2012;13(4):e148-60.
16. Lavelle K, Sowerbutts A, Bundred N, Pilling M, Degner L, Stockton C, Todd C. Is lack of surgery for older breast cancer patients in the UK explained by patient choice or poor health? A prospective cohort study. *British Journal of Cancer* 2014;110:573-583.

17. Morgan J, Richards P, Ward S, Francis M, Lawrence G, Collins K, Reed M, Wyld L. Case-mix analysis and variation in rates of non-surgical treatment of older women with operable breast cancer. *Br J Surg* 2015;102(9):1056-63.
18. Su Y-S, Yajima M, Gelman AE, Hill J. Multiple Imputation with Diagnostics (mi) in R: Opening Windows into the Black Box. *Journal of Statistical Software* 2011;45(2):1-31.
19. Horton NJ, Kleinman KP. Much ado about nothing: A comparison of missing data methods and software to fit incomplete data regression models. *Am Stat* 2007;61(1):79-90.
20. Rubin D. *Multiple Imputation for Nonresponse in Surveys*. New York: J Wiley & Sons; 1987.
21. Hanley J, McNeil B. The meaning and use of the area under a receiver operating characteristic (ROC) curve. *Radiology* 1982;143(1):29-36.
22. Lavelle K, Moran A, Howell A, Bundred N, Campbell M, Todd C. Older women with operable breast cancer are less likely to have surgery. *British Journal of Surgery* 2007;94(10):1209-1215.
23. Ali A, Greenberg D, Wishart G, Pharoah P. Patient and tumour characteristics, management, and age-specific survival in women with breast cancer in the East of England. *British Journal of Cancer* 2011;104(4):564-570.
24. Barnett K, Mercer SW, Norbury M, Watt G, Wyke S, Guthrie B. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet* 2012;380(9836):37-43.
25. Audisio R, Pope D, Ramesh H, Gennari R, van Leeuwen B, West Cea. Shall we operate? Preoperative assessment in elderly cancer patients (PACE) can help. A SIOG surgical task force prospective study. *Crit Rev Oncol Hematol* 2008;65(2):156-63.
26. NHSIC. National Mastectomy and Breast Reconstruction Audit 2011. Leeds: National Health Service Information Centre; 2011.
27. Lavelle K, Downing A, Thomas J, Lawrence G, Forman D, Oliver S. Are lower rates of surgery amongst older women with breast cancer in the UK explained by comorbidity? *British Journal of Cancer* 2012;170(7):1175-1180.
28. Syed BM, Johnston SJ, Wong DW, Green AR, Winterbottom L, Kennedy H, Simpson N, Morgan DA, Ellis IO, Cheung KL. Long-term (37 years) clinical outcome of older women with early operable primary breast cancer managed in a dedicated clinic. *Ann Oncol* 2012;23(6):1465-71.
29. Dordea M, Jones R, Nicolas A, Sudeshna S, Solomon J, Truran P, Fetherson A, Iwuchukwu O. Surgery for breast cancer in the elderly - How relevant? *Breast* 2011;20(3):212-214.
30. Morgan J, Richards P, Ward S, Francis M, Lawrence G, Collins K, Reed M, Wyld L. Case-mix analysis and variation in rates of non-surgical treatment of older women with operable breast cancer. *Br J Surg* 2015.
31. Reed MW, Wyld L, Ellis P, Bliss J, Leonard R, Groups AaETM. Breast cancer in older women: trials and tribulations. *Clin Oncol (R Coll Radiol)* 2009;21(2):99-102.