

## **Analysis of finger motion coordination during packaging interactions**

YOXALL, Alaster <<http://orcid.org/0000-0002-0954-2725>>, GONZALEZ, V. and ROWSON, J.

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/15730/>

---

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

### **Published version**

YOXALL, Alaster, GONZALEZ, V. and ROWSON, J. (2018). Analysis of finger motion coordination during packaging interactions. *Packaging Technology and Science*, 31 (6), 389-400.

---

### **Copyright and re-use policy**

See <http://shura.shu.ac.uk/information.html>

# ANALYSIS OF FINGER MOTION COORDINATION DURING PACKAGING INTERACTIONS

Yoxall A.<sup>1</sup>, Gonzalez, V.<sup>2</sup>, Rowson, J.<sup>2</sup>

1 Art and Design Research Centre, Sheffield Hallam University, Sheffield, UK

2 Department of Mechanical Engineering, University of Sheffield, Sheffield, UK

## ABSTRACT

Packaging accessibility is a significant problem for many older people. Whilst the majority of studies have focused on issues surrounding strength, work has shown that dexterity required to open a pack is also a major issue for many older people.

Hence, the work undertaken here, reports a quantitative study that aimed to analyse motion coordination patterns across digits 2–5 (index to little finger) during interactions with three of the most common types of packaging: plastic bottles, jars, and crisps packets, and comparing those interactions to a common measure of dexterity, the Perdue Pegboard. Ten subjects (6 males and 4 females) were examined while reaching forward to grasp and open a 300ml plastic bottle and a 500g jar. A ten-camera opto-electronic motion capture system measured trajectories of 25 miniature reflective markers placed on the dorsal surface landmarks of the hand. Joint angular profiles for 12 involved flexion–extension movements were derived from the measured coordinates of surface markers.

The results showed that finger correlations vary widely across the differing pack formats with the crisps having the lowest finger movement correlation and the jar having the highest. Speed and jerk metrics were also seen to vary across the various pack formats. However, finger correlations were seen to be more relevant to perceived dexterity of pack opening than finger speeds and jerk motions.

## INTRODUCTION

The frustration of accessing packaging has been termed 'Wrap rage' [1] and a significant number of studies have questioned, observed and measured people's interaction with packaging in order to gain insights into what makes packaging difficult to open [2]–[5].

Issues around packaging accessibility and capability have largely been split into three areas, lack of strength to access a pack, lack of dexterity and an inability to read and/or understand the instructions necessary to open the pack [6]. The majority of work examining the openability of packaging has concentrated on the strength and grip used by consumers, from the study by Rholes, Moldurp and Laviana [7], The Department of Trade and Industry [8], Voorbij and Steenbekkers [9], Yoxall et al [6] and more recently by Yoxall and Janson [6], Su et al [10], Kuo et al [11], Chihara and Leitkam and Bix [12].

Less work has been undertaken on issues surrounding dexterity and visual acuity or cognition of packaging. Leitch et al. [13], have undertaken a series of studies examining font sizes and labelling for consumers, whilst Yoxall et al. [2], [14], has undertaken several studies comparing the time taken to open packaging with dexterity as measured using a Purdue pegboard.

The Purdue Pegboard (PBT) is one of the most widely used tests of hand function for therapy, rehabilitation, and treatment assessment purposes. It was developed by Dr. Joseph Tiffin, an Industrial Psychologist at Purdue University, in 1948 [15], and originally intended for assessing the dexterity of assembly line workers.

The PBT tests the quality and the speed of performance of the hand as the person accomplishes a task. More precisely, it assesses proficiency of one particular grasping pattern, the precision grip [16]. It has been shown, however, that there are several factors that account for manipulative tasks [17]–[21], and the degree with which the Purdue Pegboard Test assesses individual factors has yet to be investigated.

Whilst the work by Yoxall [2], [5] attempted to understand the effect of dexterity related to different forms of packaging using the dexterity measure as a way of ranking the packaging, no attempt was made to understand the actual nature of the dexterity required to open the pack, i.e. what the fingers needed to do or the movements required. Hence, this study aims to understand the motion, trajectory, speed and effort required to open several forms of packaging and relate that to a normative dexterity method.

Previous works have shown that finger movements during manipulative tasks rarely involve motion or rotation at a single joint. Anatomical factors, such as interdigit webbings, connections between various tendons, insertions of extrinsic finger muscles, and neuronal connections result in mechanical and neural couplings between various joints. The sum of mechanical and neural coupling generates coordinated movements between various joints [22]–[25]. Thus the proficient grasping of an object entails simultaneous motion at multiple joints, with correlated rotations [23]. Simultaneous correlated motion at multiple joints has been studied during more dexterous uses of the hand, such as typing [16], playing the piano [26], and haptic interactions [27], but a standard procedure to assess such movement synergies has not been developed. Moreover, a study involving packaging interactions and hand postures has yet to be developed.

Movement smoothness measures are kinematic variables that have been used as measures of motor performance of both healthy subjects and persons with motor control and musculoskeletal impairments [28]–[30]. Although smoothness metrics have often been based on minimizing jerk, the rate of change of acceleration, [28], many other measures are possible, including three-dimensional curvature, and counting peaks in speed [31]–[34]. Smoothness has been used to assess individuals with arm ataxia [33], Parkinson's disease [35], children with cerebral palsy [36], and, more generally, it has been shown to account for the two-thirds power law, widely considered an invariant in human movement [37], [38].

Previous works in patients recovering from stroke and other motor related impairments revealed a reduction in trajectory smoothness and segmentation of continuous movements [31], [39]. However, evidence of discrete sub- movements has also been found in the movements of healthy subjects [40], and decomposition of complex movements into sub-movements has been implemented as analysis tool as they account for many patterns in human movement [41]. However, studies investigating the degree to which speed and jerk metrics can reflect dexterity when interacting with packaging and, particularly the relation of such metrics with finger correlated movement and perception of dexterity are still lacking.

## METHODOLOGY

### EXPERIMENTAL PROTOCOL

This study examined 10 healthy participants (6 male, 4 female, all right-handed, age 22-38 years,  $26 \pm 6.2$  years) performing four tasks: grasping and opening a 300ml plastic bottle, grasping and opening a 500g jar, grasping and opening a 25g crisps packet, and the Purdue Pegboard Test.

Following the tasks, the participants were asked to rate the perceived dexterity and strength required to perform each task from low (1) to high (5) in an ordinal scale.

All movements began in a consistent seated posture with the torso upright, the right upper arm approximately vertical and forearm horizontal, the fingers in natural full extension (abduction/adduction not specified), and the palm resting on a specified area on the table.

The participants carried out three repetitions of each task with a 10-second pause between each trial.

### ETHICAL APPROVAL FOR THE STUDY

The University of Sheffield's Department of Mechanical Engineering Ethics Committee approved the experimental protocol.

### DATA ACQUISITION

The acquisition technique consisted of the placement of 25 reflective markers (diameter 4mm) on different anatomical hand landmarks.

From the index to little fingers, five markers were placed as follows: first marker on the metacarpal base, second marker on the knuckle, third on the proximal interphalangeal (PIP) joint, fourth on the distal interphalangeal (DIP) joint and, finally, the fifth marker on the nail.

For the thumb, first marker was placed on the metacarpal base, second marker on the MCP joint, fourth on the IP joint and the fifth marker on the nail. One marker was placed on the wrist, aligned with the middle finger, on the wrist dorsum.

A ten-camera Vicon T-160 opto-electronic motion capture system (Oxford Metrics Ltd., UK) recorded the reflective marker movements at a sampling frequency of 120 Hz, and produced the time-varying marker coordinates in a three-dimensional laboratory coordinate system (X–Y–Z) established through calibration. The experimental set-up used for the analysis is shown in Figure 1 below.

A local coordinate system  $X_0$ – $Y_0$ – $Z_0$  was established to facilitate kinematic descriptions and definitions (Figure 2). The origin of this local coordinate system was the marker adhered to the dorsal landmark of wrist again as shown in Figure 2. The coordinates of the markers measured in the global (laboratory) coordinate system (X –Y –Z) were transformed and expressed in the local coordinate system ( $X_0$ – $Y_0$ – $Z_0$ ). From the local coordinates, the time-varying angles for all the involved joints were derived through a computational procedure.

The flexion portions of angular profiles for the metacarpophalangeal (MCP), proximal interphalangeal (PIP), and distal interphalangeal (DIP) joints of digits 2–5, a total of 12 movements, were analyzed in this study.

The starting time was defined as the frame during which the first increase in the moving of joint angles occurred, and the final time was defined as the frame during which the last decrease in the moving of joint angles occurred. The angular profiles were later normalised and a total movement time period was determined.

## DATA ANALYSIS

The analysis consisted of the computation of a cross-correlation coefficient matrix for the joint angles of interest. A matrix  $X$ , whose rows are observations (instantaneous joint angles) and whose columns are variables (degree of freedom), was defined from data from the last trial of each task for each subject in order to reduce error due to learning effect.

The matrix  $R$  of correlation coefficients was calculated from the matrix  $X$ . The matrix  $R$  is related to the covariance matrix  $C$  by:

$$R(i,j) = \frac{C(i,j)}{\sqrt{(C(i,i)C(j,j))}}$$

Where  $R$ , is the zeroth lag of the normalized covariance function.

Significance of the correlation values was examined for  $p < 0.05$ , and for all correlation coefficients  $n = 10$ ,  $df = 8$ .

Data from each matrix  $R$  was represented as a colour map, with red indicating low correlation (coefficients between 0 - 0.5) and yellow indicating high correlation (coefficients between 0.6 and 1.0) (Figure 4).

A second analysis consisted of the inspection of the joint angular profiles' sigmoidal shape. The proximal-to-distal flexion sequences across digits 2–5 for each joint type were then examined and compared with data from the correlation matrices to identify their possible relation.

The analysis consisted of the computation of the magnitude of the velocities, accelerations, and jerk by two-point numerical differentiation of positional data.

Jerk at each time point was computed according to the following equation,

$$J = \left( \ddot{x}^2 + \ddot{y}^2 + \ddot{z}^2 \right)^{1/2}$$

The jerk metric ( $Jm$ ) used for this study was calculated by dividing the negative mean jerk magnitude by the peak speed:

$$Jm = -\frac{1}{v_{peak}(t_2 - t_1)} \int_{t_1}^{t_2} |\ddot{x}(t)| dt$$

Taking the negative of the mean jerk makes the jerk metric directly proportional with smoothness, transforming this metric into a measure of smoothness. Dividing the jerk magnitude by peak speed normalizes the metric, making it less sensitive to changes in overall movement speed. The jerk metric has units of  $1/s^2$ .

The speed metric ( $Sm$ ) was calculated as the mean of the speed divided by the peak speed. The resulting speed profile from a non-smooth movement has a series of peaks with deep valleys in between, representing sudden stops between sub-movements. An example of a speed and jerk plot for opening a jar is shown in Figure 5 below.

The mean speed of such a movement is much less than its peak, making the normalized mean speed relatively low. A smooth movement tends to have fewer sub-movements and thus, fewer sudden stops, resulting in significantly higher normalized mean speeds.

## Results

### Perception Analysis

All the of the 10 users were asked to open the packaging and perform the Purdue Pegboard Test and asked to rate the interactions on a scale of 1 to 5 on their perception of the strength and dexterity. Participants were provided with a definition of strength and dexterity before undertaking the test. A score of 1 represented that the perception of the dexterity needed was low and a score of 5 rated that the perception of the dexterity needed was high (Figure 6).

The 10 participants were also asked to rate the manipulative task based on their perception of the strength needed to perform the task. A score of 1 represented that the perception the strength needed was low and a score of 5 rated that the perception of the strength needed was high (Table 1).

Scores from the Purdue Pegboard across participants were averaged and are presented along the perception results in Table 1.

Item	Mean Strength Score (SD)	Mean Dexterity Score (SD)	Mean Score (Purdue Pegboard Test)
<b>Purdue Pegboard</b>	1.13 (0.35)	3.75 (0.71)	16.4
<b>Bottle</b>	3.25 (0.71)	3.13 (0.83)	
<b>Jar</b>	4.25 (0.46)	2.13 (1.36)	
<b>Crisps</b>	1.38 (0.52)	3.25 (1.04)	

Table 1: Perceived strength and dexterity rating scores for all tasks

### Correlation Analysis

Correlation plots are shown for the Purdue pegboard and the jar, bottle and crisps opening events for two subjects (Figures 7a-f). Table 2 also shows the percentage of

high finger correlation, i.e. over 0.85 or low finger correlation, i.e. less than 0.5 average (and standard deviation) for all 10 subjects tested.

The plots show that the degree of finger correlation, i.e. fingers moving together in a coordinated way, is highest for the jar (58.9%, with a standard deviation of 10.6%) than the bottle (54.7%, with a standard deviation of 11.5%) compared to the Purdue pegboard (48.3%, with a standard deviation of 11.0%) with the crisp opening event having the lowest finger correlation (43.1%, with a standard deviation of 11.5%).

Item	Average (SD) % of movements with correlation coefficients > 0.85	Average (SD) % of movements with correlation coefficients < 0.50
Purdue Pegboard	48.3 (11.0)	23.9 (5.3)
Bottle	54.7 (11.5)	22.8 (6.9)
Jar	58.9 (10.6)	16.6 (7.7)
Crisps	43.1 (11.5)	46.2 (13.1)

Table 2: Average Finger Correlation % with correlation coefficients > 0.85 and correlation coefficients < 0.50

## Speed and jerk Analysis

Plots of speed and jerk smoothness metrics for the Purdue pegboard, bottle jar and crisp packet are shown in the following Figures 8a-h. The figures are chosen as representative examples of the measured outputs for various tasks and differing participants and are presented to show trends for each pack type. From the figures it can be readily seen that peak finger speed and jerk are similar for all tasks measured. Examining the 'peakiness' of each event, i.e. the number of peaks over 500 m/s the bottle that last for over 2.5 m/s here again the events are very similar with typically three peaks measured for the bottle, jar and crisps. However, the Purdue pegboard has a slightly lower average of 2.55 peaks.

Examining the packaging events in more detail, Table 3 below shows the average speed and jerk smoothness metrics for the bottle, jar and crisp packet. The larger the number, the smoother the task hence the jar and the crisps are seen to be smoother than the bottle. For the jerk analysis the larger (more negative) the jerk metric the less smooth the task is. Here again, the bottle is seen to be less smooth than the jar or crisps packet opening.



Item	Average Speed metric (Standard Deviation) m/s	Average Jerk Metric (Standard Deviation) m/s <sup>3</sup>
<b>Perdue Pegboard</b>	0.24 (0.03)	-0.009 (0.0025)
<b>Bottle</b>	0.28 (0.07)	-0.016 (0.008)
<b>Jar</b>	0.33 (0.03)	-0.013 (0.006)
<b>Crisps</b>	0.33 (0.04)	-0.012 (0.004)

Table3: Speed and jerk metrics for all measured tasks

## Discussion

This research has used several metrics to understand finger movements and relate them to packaging opening and dexterity. To date little work has been undertaken looking at dexterity and packaging accessibility and the understanding of finger movements and generating smoothness metrics is also a relatively new field.

Moreover, although the complex correlated movements of the hand have been investigated in previous studies, an in-depth investigation into the effects of correlated movement and velocity in packaging accessibility and their relation with perceived dexterity had yet to be made [44]–[47].

To that end this work has taken three packaging formats that are generally different in their opening properties, a jar (known to be largely an issue of strength), a bottle (that has elements of fine finger movements) and crisp pack which has elements of precision grip and fine fingers and studied them in detail using optical methods to measure finger correlations, finger speed and movement smoothness.

The opening events were compared to a known measure of dexterity, the Perdue pegboard and a participant perception test, and scores from the dexterity test were used to assess the average manual ability of the participants.

This study presents evidence of identifiable finger correlation patterns during packaging interactions, with clear differences based on the grasping pattern, strength, and dexterity required. Moreover, the current study discovered that finger correlation patterns are consistent with the complexity of the task, the number of sub-movements involved and the degree of independent finger movement required.

Furthermore, it has been shown that the Purdue Pegboard Test does not accurately assess the true differences in movement coordination occurring during the packaging

tasks under analysis. It has been shown how the wide range of strategies and grasping patterns used to interact with packaging limits the accuracy and robustness of a single time-based test.

Normative data for the Purdue Pegboard Test indicated participants were in line with average healthy adults performing the right hand (RH) Purdue Pegboard Test [42], [43].

Correlation coefficients from the Purdue Pegboard Test were the lowest when compared with the packaging tasks, indicating a lower degree of finger coordination and less coordinated fingers flexion was required to perform this task. In addition, participants coincided in rating the Purdue Pegboard Test as the most dexterous task (3.75/5). These results may suggest that finger correlated movements are strongly associated with independent finger movement and, particularly, perceived dexterity. Normative data for the Purdue Pegboard Test indicates participants are in line with average healthy adults performing the right hand (RH) Purdue Pegboard Test [42], [43].

From the selection of packaging tasks, bottle opening produced the lowest correlation coefficients, with participants rating perceived dexterity for this task higher when compared to the jar opening task, further suggesting the relation between finger correlated movements and perception of dexterity.

Speed and jerk metrics proved to be consistent with the number of movements required to perform the task, with bottle opening having the lowest movement smoothness as measured from speed and jerk. These results, however, were not consistent with perceived dexterity, with participants considering the Purdue Pegboard Test as the more dexterous task. Furthermore, results from the analysis of trajectory suggest a strong relation between object size and movement smoothness, with tasks requiring manipulation of larger objects resulting in smoother trajectories (larger speed and jerk metrics).

## Conclusions

A series of metrics have been successfully used to examine finger dexterity and packaging accessibility, namely; finger correlations analysis, speed and jerk measurements, and a dexterity perception test.

It can be seen that finger correlations vary widely across the differing pack formats with the crisps having the lowest finger movement correlation across participants and the jar having the highest. Empirically we might expect this since the use of the Purdue pegboard precludes the formation of a pinch grip leaving the remaining fingers to move freely as the task progresses. Work by Yoxall et al., [43] showed that for jars the most common grip is typically a highly correlated spherical grip and so we would expect this packaging type to have the highest measured finger correlation.

However, in examining speed and jerk, whilst the jar and crisps have very different finger correlation measurements, their speed and jerk metrics are almost identical whilst the Perdue pegboard and the bottle have the lowest speed and jerk metrics.

Examining perception scores the Perdue pegboard scores highest on perceived dexterity and the jar the lowest. Given that the Perdue pegboard has the lowest speed and jerk scores but the highest perceived dexterity score, we can assume that perceived dexterity of the packaging accessibility task is related to finger correlation and not finger speed. Similarly the jar has the lowest perceived dexterity score and the highest correlation, again indicating that there is a likely link to finger coordination and perceived dexterity.

Hence packaging that forces users to use highly uncoordinated grip patterns is likely to result in users perceiving the packaging to need high amounts of dexterity. Packaging that needs high amounts of dexterity to access is often perceived as 'fiddly' by consumers and has a higher chance of being unopenable by older consumers. Therefore 'non-fiddly' easy open packaging should be designed to facilitate the use of coordinated grip patterns, whilst minimizing the strength needed to also access the pack.

## References

- [1] American Dialect Society, "American Dialect Society," *Word of the Year 2007*, p. 81, 2008.
- [2] B. A. Yoxall, J. Luxmoore, M. Austin, L. Canty, K. J. Margrave, C. J. Richardson, J. Wearn, I. C. Howard, and R. Lewis, "Getting to Grips with Packaging : Using Ethnography and Computer Simulation to Understand Hand – Pack Interaction and Science," *Packag. Technol. Sci.*, no. October 2006, pp. 217–229, 2007.
- [3] B. Carse, A. Thomson, and B. Stansfield, "Use of biomechanical data in the Inclusive Design process : packaging design and the older adult," *J. Eng. Des.*, vol. 21, no. June, pp. 289–303, 2010.
- [4] M. Galotto and P. Ulloa, "Effect of high-pressure food processing on the mass transfer properties of selected packaging materials," *Packag. Technol. Sci.*, vol. 23, no. May, pp. 253–266, 2010.
- [5] J. Rowson and a Yoxall, "Hold, grasp, clutch or grab: consumer grip choices during food container opening.," *Appl. Ergon.*, vol. 42, no. 5, pp. 627–33, Jul. 2011.
- [6] A. Yoxall and R. Janson, "Fact or friction: A model for understanding the openability of wide mouth closures," *Packag. Technol. Sci.*, vol. 21, no. 3, pp. 137–147, 2008.
- [7] F. Rholes, K. L. Moldrup, and J. E. Laviana, "Opening jars: an anthropometric study of the wrist twisting strength in elderly," in *Proceedings of the Human Factors Society 27th Annual Meeting*, 1983, pp. 112–116.
- [8] Department of Trade and Industry, "Tha Handbook of Measurements and capabilities of the Older Adult. Strength Data for Design Safety.," 2000.
- [9] A. I. M. Voorbij and L. P. A. Steenbekkers, "The twisting force of aged consumers when opening a jar," *Appl. Ergon.*, vol. 33, no. 1, pp. 105–109, 2002.

- [10] F.-C. Su, H.-Y. Chiu, J.-H. Chang, C.-F. Lin, R.-F. Hong, and L.-C. Kuo, "Jar-opening challenges. Part 1: an apparatus for assessing hand and finger torques and forces in a jar-opening activity.," *Proc. Inst. Mech. Eng. H.*, vol. 223, no. 1, pp. 121–129, 2009.
- [11] L. C. Kuo, J. H. Chang, C. F. Lin, H. Y. Hsu, K. Y. Ho, and F. C. Su, "Jar-opening challenges. Part 2: estimating the force-generating capacity of thumb muscles in healthy young adults during jar-opening tasks.," *Proc. Inst. Mech. Eng. H.*, vol. 223, no. 5, pp. 577–588, 2009.
- [12] S. T. Leitch, T. R. Bush, and L. Bix, "Determining functional finger capabilities of healthy adults: comparing experimental data to a biomechanical model.," *J. Biomech. Eng.*, vol. 136, no. 2, p. 021022, 2014.
- [13] S. T. Leitch, L. Bix, J. de la Fuente, and T. Reid Bush, "Mapping kinematic functional abilities of the hand to three dimensional shapes for inclusive design," *J. Biomech.*, vol. 48, no. 11, pp. 2903–2910, 2015.
- [14] A. Yoxall, E. M. Rodriguez-Falcon, and J. Luxmoore, "Carpe diem, Carpe ampulla: a numerical model as an aid to the design of child-resistant closures.," *Appl. Ergon.*, vol. 44, no. 1, pp. 18–26, Jan. 2013.
- [15] J. Tiffin and E. Asher, "The Purdue pegboard test; norms and studies of reliability and validity," *J. Appl. Psychol.*, vol. 32, no. 3, pp. 234–47, 1948.
- [16] J. F. Soechting and M. Flanders, "Flexibility and repeatability of finger movements during typing: Analysis of multiple degrees of freedom," *J. Comput. Neurosci.*, vol. 4, no. 1, pp. 29–46, 1997.
- [17] E. a. Fleishman and W. E. Hempel, "A Factor Analysis of Dexterity Tests," *Pers. Psychol.*, vol. 7, no. 1, pp. 15–32, Mar. 1954.
- [18] A. Pennathur, L. R. Contreras, K. Arcaute, and W. Dowling, "Manual dexterity of older Mexican American adults: a cross-sectional pilot experimental investigation," *Int. J. Ind. Ergon.*, vol. 32, no. 6, pp. 419–431, Dec. 2003.
- [19] T. Vanbellingen, B. Kersten, M. Bellion, P. Temperli, F. Baronti, R.

Müri, and S. Bohlhalter, "Impaired finger dexterity in Parkinson's disease is associated with praxis function.," *Brain Cogn.*, vol. 77, no. 1, pp. 48–52, Oct. 2011.

- [20] F. J. Valero-Cuevas, N. Smaby, M. Venkadesan, M. Peterson, and T. Wright, "The strength-dexterity test as a measure of dynamic pinch performance.," *J. Biomech.*, vol. 36, no. 2, pp. 265–70, Feb. 2003.
- [21] M. L. Latash and M. T. Turvey, *Dexterity and its development*, . Mahwah, New Jersey: Lawrence Erlbaum Associates, Inc, 1996.
- [22] F.-C. Su, Y. L. Chou, C. S. Yang, G. T. Lin, and K. N. An, "Movement of finger joints induced by synergistic wrist motion.," *Clin. Biomech. (Bristol, Avon)*, vol. 20, no. 5, pp. 491–7, Jun. 2005.
- [23] M. Santello and J. F. Soechting, "Force synergies for multifingered grasping," *Exp. Brain Res.*, vol. 133, no. 4, pp. 457–467, Aug. 2000.
- [24] I. V Grinyagin, E. V Biryukova, and M. a Maier, "Kinematic and dynamic synergies of human precision-grip movements.," *J. Neurophysiol.*, vol. 94, no. 4, pp. 2284–94, Oct. 2005.
- [25] M. Tagliabue, A. L. Ciancio, T. Brochier, S. Eskiizmirli, and M. a. Maier, "Differences between kinematic synergies and muscle synergies during two-digit grasping," *Front. Hum. Neurosci.*, vol. 9, no. March, pp. 1–17, 2015.
- [26] K. C. Engel, M. Flanders, and J. F. Soechting, "Anticipatory and sequential motor control in piano playing.," *Exp. brain Res.*, vol. 113, no. 2, pp. 189–199, 1997.
- [27] P. H. Thakur, A. J. Bastian, and S. Hsiao, "Multidigit movement synergies of the human hand in an unconstrained haptic exploration task.," *J Neurosci*, vol. 28, no. 6, pp. 1271 – 1281, 2008.
- [28] N. Hogan and T. Flash, "The Coordination of Arm Movements: Mathematical Model," *J. Neurosci.*, vol. 5, no. 7, pp. 1688–1703, 1985.
- [29] Q. C. Pham, H. Hicheur, G. Arechavaleta, J. P. Laumond, and A. Berthoz, "The formation of trajectories during goal-oriented locomotion in humans. II. A maximum smoothness model," *Eur. J. Neurosci.*, vol. 26, pp. 2391–2403, 2007.

- [30] B. Poston, A. W. a Van Gemmert, S. Sharma, S. Chakrabarti, S. H. Zavaremi, and G. Stelmach, "Movement trajectory smoothness is not associated with the endpoint accuracy of rapid multi-joint arm movements in young and older adults.," *Acta Psychol. (Amst).*, vol. 143, no. 2, pp. 157–67, Jun. 2013.
- [31] H. I. Krebs, M. L. Aisen, B. T. Volpe, and N. Hogan, "Quantization of continuous arm movements in humans with brain injury.," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 96, no. 8, pp. 4645–9, Apr. 1999.
- [32] R. C. Miall and P. N. Haggard, "The curvature of human arm movements in the absence of visual experience.," *Exp. brain Res.*, vol. 103, no. 3, pp. 421–8, Jan. 1995.
- [33] D. Goldvasser, C. a. McGibbon, and D. E. Krebs, "High curvature and jerk analyses of arm ataxia," *Biol. Cybern.*, vol. 84, pp. 85–90, 2001.
- [34] J. J. Boessenkool, E. J. Nijhof, and C. J. Erkelens, "A comparison of curvatures of left and right hand movements in a simple pointing task.," *Exp. brain Res.*, vol. 120, no. 3, pp. 369–76, Jun. 1998.
- [35] T. Flash, R. Inzelberg, E. Schechtman, and a. D. Korczyn, "Kinematic analysis of upper limb trajectories in Parkinson's disease," *Exp. Neurol.*, vol. 118, no. 2, pp. 215–226, 1992.
- [36] S. Schneiberg, P. Mckinley, E. Gisell, H. Sveistrup, and M. F. Levin, "Reliability of kinematic measures of functional reaching in children with cerebral palsy," *Dev. Med. Child Neurol.*, vol. 52, no. 7, pp. 167–173, 2010.
- [37] E. Todorov and M. I. Jordan, "Smoothness maximization along a predefined path accurately predicts the speed profiles of complex arm movements.," *J. Neurophysiol.*, vol. 80, pp. 696–714, 1998.
- [38] S. Schaal and D. Sternad, "Origins and violations of the 2/3 power law in rhythmic three-dimensional arm movements," *Exp. Brain Res.*, vol. 136, no. 1, pp. 60–72, 2001.
- [39] B. Rohrer, S. Fasoli, H. I. Krebs, R. Hughes, B. Volpe, W. R. Frontera, J. Stein, and N. Hogan, "Movement smoothness changes during stroke recovery.," *J. Neurosci.*, vol. 22, no. 18, pp. 8297–304, Sep. 2002.

- [40] A. B. Vallbo and J. Wessberg, "Organization of motor output in slow finger movements in man," *J. Physiol.*, pp. 673–691, 1993.
- [41] J. a. Doeringer and N. Hogan, "Serial processing in human movement production," *Neural Networks*, vol. 11, no. 7–8, pp. 1345–1356, 1998.
- [42] N. Amirjani, N. L. Ashworth, J. L. Olson, M. Morhart, and K. M. Chan, "Validity and reliability of the Purdue Pegboard Test in carpal tunnel syndrome," *Muscle Nerve*, vol. 43, no. 2, pp. 171–7, 2011.
- [43] D. K. Lindstrom-Hazel and N. VanderVlies Veenstra, "Examining the Purdue Pegboard Test for Occupational Therapy Practice," *Open J. Occup. Ther.*, vol. 3, no. 3, 2015.
- [44] M. L. Latash, J. F. Scholz, F. Danion, and G. Schöner, "Finger coordination during discrete and oscillatory force production tasks.," *Exp. brain Res.*, vol. 146, no. 4, pp. 419–32, Oct. 2002.
- [45] C. Castellini and P. Van Der Smagt, "Evidence of muscle synergies during human grasping," *Biol. Cybern.*, vol. 107, no. 2, pp. 233–245, 2013.
- [46] P. H. Thakur, A. J. Bastian, and S. S. Hsiao, "Multidigit movement synergies of the human hand in an unconstrained haptic exploration task.," *J. Neurosci.*, vol. 28, no. 6, pp. 1271–1281, 2008.
- [47] D. J. Berger and A. d'Avella, "Effective force control by muscle synergies.," *Front. Comput. Neurosci.*, vol. 8, no. April, p. 46, Jan. 2014.





Figure 1: Ten-camera Vicon T-160 motion capture system



Figure 2: Marker positions for hand motion analysis




Figure 3: Participant undergoing testing opening a bottle




Figure 4: Example of a correlation plot from participant 6




Figure 5: Speed and jerk plot for participant #3 opening a jar



Figure 6: Participant interacting with the crisps packet during the perception test




Figure 7a: Correlation plot for participant #3 undertaking the Perdue Pegboard test




Figure 7b: Correlation plot for participant #12 undertaking the Perdue Pegboard test




Figure 7c: Correlation plot for participant #3 opening a jar




Figure 7d: Correlation plot for participant #10 opening a jar




Figure 7e: Correlation plot for participant #10 opening a bottle




Figure 7f: Correlation plot for participant #8 opening a bottle




Figure 7g: Correlation plot for participant #1 opening a crisp packet




Figure 7h: Correlation plot for participant #8 opening a crisp packet




Figure 8a: Speed and jerk plot for participant #2 undertaking the Perdue pegboard test




Figure 8b: Speed and jerk plot for participant #10 undertaking the Perdue pegboard test




Figure 8c: Speed and jerk plot for participant #8 opening a bottle




Figure 8d: Speed and jerk plot for participant #13 opening a bottle



