

Fibromyalgia pathogenesis provides drug target clues

LAWSON, Kim <<http://orcid.org/0000-0002-5458-1897>>

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/15785/>


This document is the Published Version [VoR]

Citation:

LAWSON, Kim (2016). Fibromyalgia pathogenesis provides drug target clues. Drug Target Review, 3, 45-49. [Article]

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>



Fibromyalgia pathogenesis provides drug target clues

Kim Lawson

Sheffield Hallam University

Fibromyalgia (FM) has been described as a condition of heightened generalised sensitisation to sensory input presenting as a complex of symptoms dominated by chronic widespread pain characterised by hyperalgesia and allodynia. A range of co-morbidities of variable intensity, such as fatigue, sleep disturbance, cognitive impairment, anxiety and depression, are often present (Figure 1; page 46). The prevalence of this condition, which is more common in females than males, is reported to be 2-8% of the population and presents a major financial and social burden to patients and healthcare systems. Neuronal excitability associated with amplified responses of the central nervous system (CNS) to peripheral input leading to central sensitisation is believed to underlie the pathophysiology^{1,2}. Peripheral nociceptive generators, such as nerve pathologies, neuro-inflammation, skeletal muscle abnormalities and ischaemia, play a role in the enhancement of the central components and the pain experienced by FM patients^{3,4}.

People with chronic pain experience balance shifts to an enhanced excitation and reduced inhibition within the CNS. The enhanced excitability of the neurophysiology reflects altered neurotransmitter functionality and neuroplasticity augmenting sensory processing⁵. Pharmacological treatments aimed at increasing antinociceptive neurotransmission in the CNS or lower levels of pronociceptive excitatory neurotransmitters are required. In the cerebrospinal fluid (CSF) of FM patients increased concentrations of substance P (two to

three-fold), endogenous opioids (three to four-fold), glutamine (two-fold), nerve growth factor (four-fold) and brain-derived neurotrophic factor (two to four-fold) have been observed^{4,6}.

In contrast, the CSF levels of 5-hydroxy indoleacetic acid (5-HIAA) – the main metabolite of serotonin, and 3-methoxy-4-hydroxyphenethylene (MPHG) – the main metabolite of noradrenaline, as well as blood levels of L-tryptophan and serotonin are lower compared with healthy controls^{4,6}. Thus, the changed biochemistry observed in FM

patients is consistent with the proposed central neuronal excitability. The raised glutamate-stimulating N-methyl-D-aspartic acid (NMDA) receptors in the dorsal horn of the spinal cord may be responsible for a wind-up phenomenon observed in FM patients.

Diffuse noxious inhibitory control

In healthy subjects application of intense painful stimuli activates a whole-body analgesia referred to as diffuse noxious inhibitory control (DNIC), which involves descending opioidergic and serotonergic-noradrenergic efferent pathways from the brain to the spinal cord that downregulate the pain signal. In FM, compared with healthy controls, DNIC is consistently reduced or absent⁷. The altered biochemistry of serotonin and noradrenaline in the CSF and serum is consistent with a decreased endogenous serotonergic and noradrenergic activity responsible for the reduced DNIC in FM patients. In contrast, evidence from FM patients indicates normal or increased endogenous opioid activity with high baseline occupancy of the receptors, rather than a deficiency of endogenous opioid release⁸. Thus, low-dose naltrexone (LDN), an opioid receptor antagonist, has been proposed as an effective treatment strategy in some FM patients⁸. Microglia antagonism, rather than blocking endogenous opioid release, has been suggested as the mechanism by which LDN reduces the severity of the FM symptoms. Pro-inflammatory factors released from microglia, which in FM may be abnormally sensitised, interact with CNS neurons leading to central facilitation of pain processing⁹.

Pharmacological treatments that simultaneously raise both serotonin and noradrenaline, for example, tricyclic antidepressants and serotonin noradrenaline reuptake inhibitors have gained interest and demonstrated efficacy in treating FM, further supporting the involvement of the low levels of these neurotransmitters in



“The changed biochemistry observed in FM patients is consistent with the proposed central neuronal excitability”

the pathogenesis of the disease². Treatment with the tricyclic antidepressant amitriptyline also correlated the clinical response with normalisation of function within the bilateral thalamus and basal ganglia, demonstrating involvement of serotonin and noradrenaline in several central aspects of the pathophysiology of FM. In contrast, drug treatments (fluoxetine, paroxetine, trazodone, esreboxetine, droxidopa) that selectively raise either of the neurotransmitter levels, e.g., selective serotonin reuptake inhibitors and noradrenaline reuptake inhibitors, have not demonstrated the success of the broader spectrum interventions². Interestingly, when amitriptyline was administered in combination with melatonin a superior improvement in symptoms was observed compared with amitriptyline alone¹⁰. Thus,




Figure 1: Interrelationship of centrally- and peripherally-derived symptoms of fibromyalgia

stimulation of melatonergic receptors improved the serotonergic-noradrenergic components of the descending endogenous pain-modulating system and, thereby, abnormality of the melatonergic system may also play a role in the pathogenesis of FM.

NMDA receptor

The enhanced wind-up and central sensitisation in FM has been associated with elevated excitatory CNS neurotransmitters. Glutamate levels in FM patients are elevated in key pain-processing areas of the brain and these levels change in response to successful

treatment that attenuates the pain^{11,12}. For example, glutamatergic activity in the insula was decreased by pregabalin, a voltage-gated calcium channel $\alpha_2\delta$ subunit ligand, with those FM patients with the highest levels of glutamate before treatment being the most likely to respond to the drug¹³. Calcium ions enable neurotransmitters to bind to vesicles at the presynaptic membrane terminal, thus inhibiting the $\alpha_2\delta$ subunit, and results in decreased calcium influx and a reduction in the release of neurotransmitters that transmit nociceptive signals between neurons.

In addition, a subset of FM patients responded to NMDA receptor antagonists (ketamine, dextromethorphan and memantine) which would be consistent with suppression of an increased glutamatergic activity, however such treatments are often not well tolerated and have limited use in the clinic¹⁴. Improved understanding of the contribution of the activity of NMDA receptor subtypes in pain – and the other symptoms of FM – and recent development of subtype-specific modulators should address those limitations of this therapeutic approach¹⁴. In addition to reducing pain, the $\alpha_2\delta$ subunit ligands pregabalin and gabapentin have been reported to improve other symptoms characteristic of FM, such as sleep, anxiety and fatigue, supporting the involvement of the neurotransmitter imbalances (e.g., glutamate) targeted by these drugs in the co-morbid symptoms¹⁵.

Identification of subtypes of the $\alpha_2\delta$ subunits, $\alpha_2\delta$ -1 and $\alpha_2\delta$ -2, have shown that analgaesic effects are primarily due to ligands binding





Figure 2: Central and peripheral mechanisms related to the pathophysiology of fibromyalgia

“Pharmacological treatments that simultaneously raise both serotonin have gained interest and demonstrated efficacy”

to the former, whereas binding to the latter subtype appears to contribute to CNS side effects¹⁶. Evaluation of the preferentially $\alpha_2\delta$ -1 subunit selective ligand mirogabalin (DS5565) demonstrated a preferable efficacy profile in a treatment study of diabetic peripheral neuropathic pain¹⁷. These findings support testing of mirogabalin in clinical trials with FM patients, where clinically meaningful differences in efficacy and safety may provide a more successful therapeutic option.

Although efficacy is observed with modulation of bioamines or the $\alpha_2\delta$ ligands in the management of FM symptoms, the outcomes are often limited with a partial reduction of symptoms only in a proportion of patients. Thus, the potential of greater benefit or more preferable outcomes from combined therapies targeting multiple pharmacological mechanisms has been considered. In a single-blind randomised trial, the combined use of pregabalin plus paroxetine, amitriptyline or venlafaxine resulted in significantly lower somatic symptom and depression scale scores; better life satisfaction; mood and sleep quality; higher medication tolerability and less frequent adverse effects in patients with FM¹⁸. Suppression of neuronal excitability and neurotransmitter release appears to be a key pharmacological property to the management of FM as observed with the $\alpha_2\delta$ ligands and the bioamine modulators. Thus, anticonvulsant drugs such as levetiracetam, topiramate and lacosamide, which suppress neuronal hyperexcitability by a mechanism other than $\alpha_2\delta$ subunits have been evaluated as a potential treatment approach, but have failed to provide sufficient evidence of efficacy against the symptoms of FM¹⁹. The mechanism(s) of action of the gabapentanoids, pregabalin and



The enhanced wind-up and central sensitisation in FM has been associated with raised excitatory neurotransmitters



Cannabinoids such as nabilone and dronabinol improved quality of life in FM patients in clinical trials

gabapentin appear to provide a unique, albeit limited, control of the neuronal hyperexcitability and central sensitisation associated with FM.

Although raised substance P levels have been reported in FM patients, substance P antagonists have provided inconsistent outcomes or failed in clinical trials in FM and other chronic pain states²⁶, casting questions about how critical this neurotransmitter is as a drug target in pain transmission. Interestingly, topically applied capsaicin significantly improved the symptoms of FM⁹. Capsaicin binds to the transient receptor potential vanilloid 1 subunits (TRPV1) located in peripheral nociceptors and desensitises nociceptive processes, possibly due to depletion of substance P21. These findings infer, at least, a peripheral role of substance P in the regulation of peripheral generators and

sensory factors activating central neuronal mechanisms responsible for the symptoms of FM.

Dopamine deficiency and endocannabinoid deficiency in the CNS, where there is an involvement in regulation of pain processing and chronic stress, have also been implicated in the pathophysiology of FM^{22,23}. In patients with FM dopamine release into the basal ganglia in response to painful stimuli is attenuated or absent²². Attempts to rectify this dysfunction

with dopamine receptor agonists have been inconsistent with pramipexole improving symptoms but similar outcomes have not been observed with ropinirole and terguride^{24,26}. Evaluation of cannabinoids, such as nabilone and dronabinol, as a treatment approach demonstrated in clinical trials an improvement of quality of life in patients with FM, but with limiting adverse effects². Psychotropic effects due to hepatic metabolites from first-pass metabolism could limit the clinical utility of cannabinoids, which has led to the evaluation of a transdermal application of the D-(-)-glyceric acid ester of delta-9-tetrahydrocannabinol, ZYN001 in FM patients²⁷.

Systemic stress-related effects associated with alterations in the hypothalamic pituitary adrenal axis, and autonomic and cardiovascular system have also been suggested to enhance or underlie the symptoms, particularly the pain, of FM^{2,6}. Although studies in FM generally have shown alterations of these stress systems, the findings of the studies are often inconsistent with abnormal HPA or autonomic function in only a small percentage of patients, with significant overlap between patients and

“The diversity of pharmacological targets gaining interest emphasises the complexity of FM”




Figure 3: Current and emerging drug targets for treatments of fibromyalgia

healthy subjects. Further data also suggests that HPA and autonomic abnormalities occur as a consequence of the pain in FM patients²⁸.

Although inflammation has also been suggested to underlie the pathology in FM, literature on cytokines has been variable and the studies have several limitations that could influence the findings^{12,6}. Pro-inflammatory cytokines might play a role in the generation and enhancement of the chronic pain characteristic of FM, however evidence does not as yet allow conclusions to be drawn as to whether the inflammatory response is the cause of the symptoms or due to the changed physiology initiating the pain. It is important to note that anti-inflammatory drugs such as non-steroidal anti-inflammatory drugs and steroids have been shown in clinical trials to have no or very limited efficacy in the treatment of FM^{2,6}.

Conclusion

Advances in the understanding of the pathophysiology of FM are providing clues as to underlying mechanisms as targets for new medications (Figure 2; page 47). The alteration of multiple systems and biological mechanisms, both at central and peripheral levels, has often introduced confusion as to which processes are a cause or consequence of FM. Bioamine modulation and voltage-gated Ca²⁺ channel $\alpha 2\delta$ subunits in addition to dopamine receptors, NMDA receptors, cannabinoid receptors and melatonin receptors are emerging as drug targets (Figure 3; page 48). However, the inconsistency

of clinical data for drugs that target single mechanisms support the requirement of therapies that modulate more than one dysfunctional neurotransmitter.

The diversity of pharmacological targets gaining interest emphasises the complexity of FM, but insight into potential drug treatment profiles offers important clues for condition-focused therapies and improved diagnosis. Further evidence of the symptoms indicates heterogeneity with multiple potential aetiologies. Such an interpretation is consistent with an heterogeneity of drug treatment outcomes often leading to the requirement of individualised management. Finally, greater understanding of the central and peripheral contributions to the pathophysiology is required so that therapies can be targeted toward both components to manage the characteristic symptoms, but also to consider accompanying co-morbid conditions. 



Dr Kim Lawson obtained a PhD in pharmacology at the University of Sunderland, UK. After industrial engagements at Rhone Poulenc Sante, Sanofi-Labaz, Recherche Syntex France and British Biotechnology, in 1995 he became Senior Lecturer in Pharmacology in the Department of Biosciences and Chemistry at Sheffield Hallam University, UK. His research interests are focused on the identification of treatments of fibromyalgia. Kim holds the honorary positions of Chair of the Medical Advisory Board to Fibromyalgia Action UK and is a Patron to Folly Pogs Fibromyalgia Research UK (2012). Kim can be contacted at: k.lawson@shu.ac.uk.

References

- Borchers AT & Gershwin ME. Fibromyalgia: A critical and comprehensive review. *Clinic Rev. Allerg. Immunol.* 2016; 49, 100-151
- Lawson K. Potential drug therapies for the treatment of fibromyalgia. *Exp. Opin. Invest. Drugs.* 2016; 16, 1-11
- Üçeyler N, Zeller D, Kahn AK, Kewenig S, Kittel-Schneider S, Schmid A, Casanova-Molla J, Reiners K & Sommer C. Small fibre pathology in patients with fibromyalgia syndrome. *Brain.* 2013; 136, 1857-67
- Albrecht PJ, Hou Q, Argoff CE, Storey JR, Wymer JP & Rice FL. Excessive peptidergic sensory innervation of cutaneous arteriole-venule shunts (AVS) in the palmar glabrous skin of fibromyalgia patients: implications for widespread deep tissue pain and fatigue. *Pain Med.* 2013; 14(6), 895-915
- Clauw DJ. Fibromyalgia: a clinical review. *JAMA* 2014; 311, 1547-1555
- Sluka KA & Clauw DJ. Neurobiology of fibromyalgia and chronic widespread pain. *Neurosci.* 2016
- Julien N, Goffaux P, Arseneault P & Marchand S. Widespread pain in fibromyalgia is related to a deficit of endogenous pain inhibition. *Pain* 2005; 114, 295-302
- Younger J, Noor N, McCue R, & Mackey S. Low-dose naltrexone for the treatment of fibromyalgia: findings of a small, randomized, double-blind, placebo-controlled, counterbalanced, crossover trial assessing daily pain levels. *Arthritis Rheum.* 2013; 65(2), 529-38
- Tsuda M, Masuda T, Tozaki-Saitoh H, & Inoue K. Microglial regulation of neuropathic pain. *J Pharmacol. Sci.* 2013; 121(2), 89-94
- De Zanette SA, Vercellino R, Laste G, Rozisky JR, Schwertner A, Machado CB, Xavier F, de Souza IC, Deitos A, Torres IL & Caumo W. Melatonin analgesia is associated with improvement of the descending endogenous pain-modulating system in fibromyalgia: a phase II, randomized, double-dummy, controlled trial. *BMC Pharmacol. Toxicol.* 2014; 15, 40
- Foerster BR, Nascimento TD, DeBoer M, Bender MA, Rice IC, Truong DQ, Bikson M, Clauw DJ, Zubieta JK, Harris RE & DaSilva AF. Excitatory and inhibitory brain metabolites as targets of motor cortex transcranial direct current stimulation therapy and predictors of its efficacy in fibromyalgia. *Arthritis Rheumatol.* 2015; 67, 576-581
- Harte SE, Clauw DJ, Napadow V & Harris RE. Pressure pain sensitivity and insular combined glutamate and glutamine (Glx) are associated with subsequent clinical response to sham but not traditional acupuncture in patients who have chronic pain. *Med. Acupunct.* 2013; 25, 154-160
- Harris RE, Napadow V, Huggins JP, Pauer L, Kim J, Hampson J, Sundgren PC, Foerster B, Petrou M, Schmidt-Wilcke T & Clauw DJ. Pregabalin rectifies aberrant brain chemistry, connectivity, and functional response in chronic pain patients. *Anesthesiology* 2013; 119, 1453-1464
- Zhou HY, Chen SR & Pan HL. Targeting N-methyl-D-aspartate receptors for treatment of neuropathic pain. *Expert Rev. Clin. Pharmacol.* 2011; 4(3), 379-388
- Baidya DK, Agarwal A, Khanna P & Arora MK. Pregabalin in acute and chronic pain. *J. Anaesthesiol. Clin. Pharmacol.* 2011; 27, 307-314
- Field MJ, Cox PJ, Stott E, Melrose H, Offord J, Su TZ, Bramwell S, Corradini L, England S, Winks J, Kinloch RA, Hendrich J, Dolphin AC, Webb T & Williams D. Identification of the alpha2-delta-1 subunit of voltage-dependent calcium channels as a molecular target for pain mediating the analgesic actions of pregabalin. *Proc Natl Acad Sci U S A* 2006; 103:17537-42
- Vinik A, Rosenstock J, Sharma U, Feins K, Hsu C & Merante D. Efficacy and safety of mirogabalin (DS-5565) for the treatment of diabetic peripheral neuropathic pain: a randomized, double-blind, placebo- and active comparator-controlled, adaptive proof-of-concept phase 2 study. *Diabetes Care.* 2014; 37(12), 3253-61
- Ramzy EA. Comparative efficacy of newer antidepressants in combination with pregabalin for fibromyalgia syndrome: A Controlled, Randomized Study. *Pain Pract.* 2016; doi: 10.1111/papr.12409
- Üçeyler N, Sommer C, Walitt B & Häuser W. Anticonvulsants for fibromyalgia. *Cochrane Database Syst. Rev.* 2013; 10, CD010782
- Casanueva B, Rodero B, Quintal C, Llorca J & González-Gay MA. Short-term efficacy of topical capsaicin therapy in severely affected fibromyalgia patients. *Rheumatol. Int.* 2013; 33(10), 2665-70
- Schumacher M & Pasvankas G. Topical capsaicin formulations in the management of neuropathic pain. *Prog. Drug. Res.* 2014; 68, 105-28
- Wood PB & Holman AJ. An elephant among us: the role of dopamine in the pathophysiology of fibromyalgia. *J. Rheumatol.* 2009; 36(2), 221-4
- Russo EB. Clinical endocannabinoid deficiency (CECD): can this concept explain therapeutic benefits of cannabis in migraine, fibromyalgia, irritable bowel syndrome and other treatment-resistant conditions? *Neuro. Endocrinol. Lett.* 2008; 29(2), 192-200
- Holman AJ & Myers RR. A randomized, double-blind, placebo-controlled trial of pramipexole, a dopamine agonist, in patients with fibromyalgia receiving concomitant medications. *Arthritis Rheum.* 2005; 52(8), 2495-505
- Holman AJ. Ropinirole, open preliminary observations of a dopamine agonist for refractory fibromyalgia. *J. Clin. Rheumatol.* 2003; 9(4), 277-9
- Distler O, Eich W, Dokoupilova E, Dvorak Z, Fleck M, Gaubitz M, Hechler M, Jansen JP, Krause A, Bendzus M, Pache L, Reiter R & Müller-Ladner U. Evaluation of the efficacy and safety of tergicide in patients with fibromyalgia syndrome: results of a twelve-week, multicenter, randomized, double-blind, placebo-controlled, parallel-group study. *Arthritis Rheum.* 2010; 62(1), 291-300
- THC Pro-Drug Patch – ZYN001. Being Studied in Fibromyalgia and Peripheral Neuropathic Pain. <http://zynerba.com/in-development/thc-pro-drug-patch-zyn001/> [Last accessed 4 July 2016]
- McLean SA, Williams DA, Stein PK, Harris RE, Lyden AK, Whalen G, Park KM, Liberzon I, Sen A, Gracely RH, Baraniuk JN & Clauw DJ. Cerebrospinal fluid corticotropin-releasing factor concentration is associated with pain but not fatigue symptoms in patients with fibromyalgia. *Neuropsychopharmacol.* 2006; 31, 2776-2782