

Prevalence of laceration injuries in professional and amateur rugby union: a systematic review and meta-analysis

OUDSHOORN, Bodil <<http://orcid.org/0000-0001-5081-4078>>, DRISCOLL, Heather, KILNER, Karen <<http://orcid.org/0000-0003-0196-8518>>, DUNN, Marcus <<http://orcid.org/0000-0003-3368-8131>> and JAMES, David <<http://orcid.org/0000-0002-1135-626X>>

Available from Sheffield Hallam University Research Archive (SHURA) at:
<https://shura.shu.ac.uk/16093/>

This document is the Accepted Version [AM]

Citation:

OUDSHOORN, Bodil, DRISCOLL, Heather, KILNER, Karen, DUNN, Marcus and JAMES, David (2017). Prevalence of laceration injuries in professional and amateur rugby union: a systematic review and meta-analysis. *BMJ Open Sport & Exercise Medicine*, 3 (1), e000239. [Article]

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

Prevalence of laceration injuries in professional and amateur rugby union: a systematic review and meta-analysis.

Bodil Yucki Oudshoorn^{1*}, Heather Frances Driscoll², Karen Kilner³, Marcus Dunn¹ and David James¹

¹Centre for Sports Engineering Research, Faculty of Health and Wellbeing, Sheffield Hallam University, Sheffield, United Kingdom

²School of Mechanical Engineering, Manchester Metropolitan University, Manchester, United Kingdom

³Centre for Health and Social Care Research, Faculty of Health and Wellbeing, Sheffield Hallam University, Sheffield, United Kingdom

*Corresponding author.

Postal address: Centre for Sports Engineering Research, 11 Broomgrove Road, Sheffield, S10 2LX.

E-mail: b.oudshoorn@shu.ac.uk

Telephone number: +44 (0) 114 225 2355

Word count: 4208 words

ABSTRACT

Background: Studded footwear can cause severe lacerations in rugby union; the prevalence of these injuries is currently unknown.

Objective: To summarise the skin and laceration injury prevalence in published epidemiological studies and to investigate any differences in skin injury risk between amateur and professional players.

Design: Systematic literature review and meta-analysis of epidemiological studies.

Data sources: Pubmed, Web of Science, Scopus and Ovid.

Eligibility criteria for selecting studies: Prospective, epidemiological studies published in English after 1995, measuring a minimum of 400 match or 900 training exposure hours. Participants should be adult rugby union players (amateur or professional). The study should report a separate skin or laceration injury category and provide sufficient detail to calculate injury prevalence within this category.

Results: Twelve studies were included. Mean skin injury prevalence during matches was 2.4 injuries per 1000 exposure hours; during training sessions the prevalence was 0.06 injuries per 1000 exposure hours. Skin injuries accounted for 5.3% of match injuries and 1.7% of training injuries. Skin injury risk was similar for amateur compared to professional players during matches (odds ratio: 0.63, $p = 0.46$), but higher during training sessions (odds ratio: 9.24, $p = 0.02$).

Summary/conclusions: The skin injury prevalence of 2.4 injuries per 1000 exposure hours is equivalent to one time-loss injury sustained during matches per team, per season. Amateur players are more likely to sustain skin injuries during training sessions than professional players. There is a need for more studies observing injuries among amateur players.

Systematic review registration: PROSPERO CRD42015024027.

Keywords: Epidemiology, Injury, Rugby, Dermatology, Review

What is already known?

- Injury prevalence in professional rugby union is 81 injuries per 1000 match exposure hours.
- Laceration and skin injury prevalence in professional rugby union is 1 - 3 injuries per 1000 match exposure hours.
- Injury definitions influence the reporting of overall injury prevalence; this makes it difficult to compare different studies.

INTRODUCTION

Rugby union is a popular sport with approximately 7.23 million players worldwide.[1] Rugby union has a high injury incidence with a self-reported injury rate of 96 injuries per 1000 occasions of participation. This is higher than other popular team sports such as association football (64 injuries per 1000 occasions) and field hockey (62 injuries per 1000 occasions).[2] The full-contact nature of rugby union has previously been identified as a major risk factor for injury; an estimated 72% of match injuries resulted from player-to-player contact.[3]

Studded footwear is worn in rugby union and other field sports played on grass, as a means to increase the traction between the shoe and the playing surface. Studs have been shown to cause severe laceration injuries to players.[4] In 2008, a studded footwear manufacturer was sued by a player who sustained a laceration to the head.[5] Since the introduction of the bladed stud in 1994, manufacturers have made innovations in the design and material of studded footwear. Traditionally, rugby boots have been equipped with screw-in studs with a round profile, made out of aluminium. Bladed studs are moulded out of plastic, and their elongated profile gives rise to their name. Nowadays, rugby players wear both shoes with moulded studs and with aluminium rounded studs.[6] World Rugby, the international governing body of rugby, has specified a test method for assessing the laceration injury risk of studded footwear; however, these regulations currently lack validation and are optional for manufacturers to follow. Identifying the proportion and prevalence of skin injuries in rugby union will aid the understanding of skin injuries and is the first step in injury prevention models such as van Mechelen *et al.*[7] Previously, there has been no specific published data available on the skin injury prevalence caused by stud-skin interactions in rugby union.

In rugby union, a survey study[6] investigating the cause of stud laceration injuries during the game identified that 97% of respondents had experienced *minor* (i.e. did not hinder the player from continuing to play) skin injuries caused by studs and 71% of respondents had experienced *major* (i.e. lost playing time) skin injuries caused by studs at least once during their rugby career. No stud laceration injury incidence (i.e. injuries per thousand exposure hours (ITEH)) was reported in this study.

The injury definition used in sports injury epidemiological studies influences their results and has been a topic of debate.[8–10] Four commonly used injury definitions are:

- **Time-loss (one-day):** Any physical complaint sustained by a player during a match or training session that prevents the player from taking full part in a match or training session one day after the injury.[10]

- **Time-loss (immediate):** Any physical complaint sustained by a player during a match or training session that forces the player to retire and / or prevents the player from taking full part in a training session or match the following day.
- **Medical attention:** The player receives medical treatment for his / her injury.
- **Player reported:** The player reports his / her physical complaint as an injury.
- **Trainer reported:** The coach or (head) trainer reports a player injured.

A consensus statement on injury definitions and data collection procedures for epidemiological studies in rugby union was published in 2007.[10] This document recommends using the one day time-loss injury definition and laceration injuries are recommended to have their own injury type category. If a time-loss injury definition is used, minor injuries and injuries which fall under the blood injury rule[11] could be treated on the side of the field (stitched or glued) and therefore be ignored in the injury count, leading to a likely underestimation of the skin injury prevalence.[12, 13] An example of such a situation is given in the consensus statement,[10] p330:

"A loose-head prop forward sustained a laceration to his head during a match; the player left the field of play to enable the team doctor to suture and protect the injury. The player returned to the field of play. The player continued to train and play with his head bandaged for the next three weeks. This episode should not be recorded as an injury"

In a systematic review on skin injuries in football, van den Eijnde *et al*[12] warned that skin injuries were underreported in current literature and suggested this resulted from restrictive injury definitions. The use of a medical attention injury definition rather than a time-loss injury definition led to a two to fourfold increase in the injury prevalence of abrasions and lacerations compared to the use of a time-loss injury definition.[12]

Level of play has previously been identified as a risk factor for injury.[14] In a meta-analysis of professional rugby union injuries by Williams *et al*,[15] international match injury risk was found to be higher than level 1 and level 2 clubs match injury risk, suggesting that more elite players sustain a higher risk of injury during matches. This effect was not found for training injuries. Overall, mean match injury prevalence during senior men's professional rugby was 81 ITEH and mean training injury prevalence was 3 ITEH.[15] This same systematic review also investigated injury patterns in professional rugby union; they found a laceration and skin injury prevalence during match play of 1 ITEH (95% confidence interval (CI): 1 - 3 ITEH).[15] The review did not report on the proportion of laceration and skin injuries, the laceration injury prevalence during training sessions, and solely focussed on injuries sustained by professional players. Therefore no comparisons could be made between injury risks for amateur versus professional players. Differences between professional and amateur rugby could influence the skin and laceration injury prevalence between playing levels.

These differences include; playing styles, the quality and potential wear of the footwear worn, intensity of movements and impact forces. This study systematically reviews published epidemiological literature reporting on laceration or skin injury prevalence in rugby union during match play and training sessions and identifies potential differences in skin injury risk for amateur and professional players.

METHODS

This review was registered in the PROSPERO International Prospective Register of Systematic Reviews (registration number CRD42015024027). The 2009 PRISMA-P guidelines for preferred reporting items for systematic review and meta-analysis protocols[16] were implemented in preparing and conducting this review.

Literature search strategy

The literature search was conducted in July 2016 using the search engines Pubmed, Ovid, Scopus and Web of Science. The search strategy was designed with the purpose of finding epidemiological studies on rugby union injuries. Search terms for all search engines were:

[rugby] AND [epidemiology OR epidemiological OR epidemiologic OR injury OR injured OR injuries].

Search engine limits were set to find only articles published after January 1995, due to an innovative change in stud design in the previous year (1994) and the introduction of professional rugby union in 1995. The title, author, date of publication and abstract of each record were imported to a reference manager (RefWorks, ProQuest® LLC, 2016).

Selection criteria

After removing duplicates, the title and abstract of the remaining records were screened for eligibility by the first author. From resulting records the full-text article was obtained and again screened for eligibility. The following inclusion criteria were used throughout:

1. The study must report on prospective, epidemiological findings in a rugby union cohort of players.
2. The study must observe injuries in players over 18 years.
3. The study must not report only a sub-category of injuries, e.g. tackling injuries or head injuries.

4. The study must report a skin or laceration injury category and provide enough information to calculate injury prevalence within this category.
5. The athlete exposure in the study should be over 400 match hours and / or 900 training hours.
6. The full-text article must be written in English and published in a peer-reviewed journal after 1995.

Only studies observing injuries in an adult cohort were included in this review since in junior (under 18 years) age categories, the rules of rugby union are different. The minimum athlete exposure criteria was defined such that at least one team (15 players) was followed for a full season (defined as 20 matches and / or 20 weeks of 2 training sessions a week), which results in a minimum of 400 match exposure hours or 900 training exposure hours. Studies published after 1995 were included because a change in stud designs was introduced in 1994 and professionalism was introduced in rugby union in 1995. No restriction was placed on level of play (amateur or professional), sex, or injury definition used. Athletes were defined as 'professional' if they were receiving monetary reimbursement for their participation in the team to the degree that maintaining full-time employment elsewhere was not necessary. Review papers, conference abstracts and studies using data which was already published in a different study were excluded. A study was also excluded if it did not report injury frequency and exposure hours for match and training sessions separately.

Quality and bias assessment

A four-item checklist was used to assess the risk of bias of each of the included studies. The checklist was adapted from Waldén *et al*[17] and based on the “Strengthening the reporting of observational studies in epidemiology” (STROBE) statement.[18] Studies were assessed on each item for low risk (0) or high risk (1) of bias. A cumulative score for each study was calculated. Studies with a cumulative score of 0 were defined as at low risk of bias, studies with a score of 1 or 2 as at medium risk and studies with a score >3 as at high risk. Only studies with a low or medium risk of bias were included in the meta-analysis. The full checklist can be found in supplementary materials 1.

Data extraction

For each study that met the inclusion criteria, the injury definition, skin injury category and athlete level - professional or amateur - were recorded. Where available, the observed number of total injuries, the observed number of skin or laceration injuries, and their corresponding match and training exposure hours were extracted from each study. Injury prevalence was defined as the number of observed injuries per 1000 exposure hours. For each study, separate injury prevalence (\pm 95% CI) was calculated for injuries sustained during match play and during training sessions. The percentage of skin injuries in proportion to total injuries observed in a study was also calculated for both matches and training sessions. If stated in a separate category, abrasion injuries were not included when

calculating skin injury prevalence. A weighting factor based on study size (defined as exposure hours) was implemented when calculating the mean of each group of studies.

Statistical analysis

In this meta-analysis the influence of playing level (amateur or professional) on skin injury prevalence during match and training sessions was of interest. Influence of sex on injury risk could not be investigated due to a shortage of studies publishing female injury numbers. Skin injury prevalence data was modelled using a negative binomial regression with log link. This generalised linear model is similar to the Poisson regression model previously described by Lystad *et al*[19], but also accounts for overdispersion in the data set. In our model, the response variable was the number of observed skin injuries, offset by the log of the exposure hours. A weight scaling variable was assigned to each study depending on its relative size. The odds ratio (and 95% CI) that an amateur player sustains a skin injury - as compared to a professional player - was calculated. Separate odds ratios were determined for training sessions and during match play. Alpha was set at 0.05.

RESULTS

A total of 1351 records were identified through the online search engines (Stage 1, Figure 1). After duplicates were removed, the titles and abstracts of 464 remaining records were screened (Stage 2). The screening excluded 424 articles; therefore 40 full-text articles were reviewed for eligibility (Stage 3). Twenty-eight articles were excluded based on the selection criteria, leaving twelve studies which were included in this review[3, 20–30] (Stage 4). Out of these selected studies, ten studies measured professional athletes[3, 20, 22–28, 30] and two studies observed injuries in an amateur cohort.[21, 29] Eleven out of twelve studies included a completely male cohort;[3, 20, 22–30] one study observed injuries of both males and female rugby players.[21]

*** Figure 1 near here ***

Quality and bias assessment

In the risk of bias assessment seven articles were categorised as a low risk of bias and five articles were categorised as a medium risk of bias (supplementary materials 1). No articles were excluded from the meta-analysis based on the bias assessment.

Match injuries

Eleven out of twelve studies included in this review monitored and reported the match injuries of their athletes[3, 20, 21, 23–30] (Table 1). The majority of the studies[3, 22–27, 29, 30] used a *time-loss one-day* injury definition. The other studies used *time-loss immediate*, sometimes in combination with *medical attention*. [20, 21, 28] No included study used a *player or coach reported* definition. Mean overall match injury prevalence in professional and amateur players combined was 53 ITEH (95% CI: 51 - 54).

Table 1: Total and skin injury prevalence of rugby union players during matches. Prevalence is reported as injuries per 1000 match hours. TL-1D = time-loss one-day; TL-IM = time-loss immediately; MA = medical attention.

Study	Athlete level	Injury definition	Injury category	Match exposure hours	Match injury prevalence	Skin injury prevalence	No of skin injuries [proportion]
Best <i>et al</i> , 2005[20]	Professional	TL-IM	Open wound / Laceration	1930	97.9	21.2	41 [21.7%]
Brooks <i>et al</i> , 2005[23]	Professional	TL-1D	Laceration & skin	445	218.0	4.5	2 [2.1%]
Brooks <i>et al</i> , 2005[3]	Professional	TL-1D	Laceration & skin	16782	91.4	1.1	18 [1.2%]
Fuller <i>et al</i> , 2008[24]	Professional	TL-1D	Laceration	1920	83.9	0	0 [0%]
Fuller <i>et al</i> , 2009[25]	Professional	TL-1D	Skin	5600	74.8	2.5	14 [3.3%]
Fuller & Molloy 2011[26]	Professional	TL-1D	Skin	3320	52.4	1.5	5 [2.9%]
Fuller <i>et al</i> , 2013[27]	Professional	TL-1D	Laceration	1920	89.1	1.6	3 [1.8%]
Jakoet & Noakes 1998[28]	Professional	TL-IM	Laceration	2194	31.9	8.7	19 [27.0%]
Schwellnus <i>et al</i> , 2014[30]	Professional	TL-1D	Skin	1512	83.3	4.0	6 [4.8%]
Bird <i>et al</i> , 1998[21]	Amateur	TL-IM & MA	Laceration	5871	83.3	7.3	43 [8.8%]
Kerr <i>et al</i> , 2008[29]	Amateur	TL-1D	Laceration	34324	17.0	0.8	29 [5.0%]

The skin injury prevalence during match play in amateur and professional players combined was 2.4 ITEH (95% CI: 2.0 - 2.7) (Figure 2). Mean skin injury prevalence during professional match play was 3.0 ITEH (95% CI 2.5 - 3.6) and during amateur match play 1.8 ITEH (95% CI 1.4 - 2.2). The skin injury risk for amateur players compared to professional players during matches was not significantly different (odds ratio: 0.60; 95% CI: 0.17 - 2.08; $p = 0.42$). The lowest skin injury prevalence observed was 0 laceration injuries after 1920 hours of match exposure.[24] The highest skin injury prevalence was 21.2 ITEH, where 41 injuries occurred during 1930 hours of match play.[20] The proportion of skin and laceration injuries in the included studies during match play ranged from 0 to 27%, with a weighted mean of 5.1%.

*** Figure 2 near here ***

Training injuries

Seven out of twelve studies included in this review monitored and reported the injury data of their athletes during training sessions.[21–24, 27, 29, 30] Five studies monitored a professional population,[22–24, 27, 30] and two studies an amateur population.[21, 29] One study used the *time-loss immediate* in combination with *medical attention* injury definition,[21] whilst the other six studies used the *time-loss, one day* injury definition[23, 24, 27, 29–31] (Table 2). Mean overall training injury prevalence for amateur and professional players combined was 2.8 ITEH (95% CI: 2.6 - 2.9).

Table 2: Total and skin injury prevalence of rugby union players during training sessions. Prevalence is reported as injuries per 1000 training hours. TL-1D = time-loss, one day; TL-IM = time-loss immediately; MA = medical attention.

Study	Athlete level	Injury definition	Injury category	Training exposure hours	Training injury prevalence	Skin injury prevalence	No of skin injuries [proportion]
Brooks <i>et al</i> , 2005[22]	Professional	TL-1D	Laceration & skin	196409	2.0	0.02	4 [1.0%]
Brooks <i>et al</i> , 2005[23]	Professional	TL-1D	Laceration & skin	7928	6.1	0	0 [0%]
Fuller <i>et al</i> , 2008[24]	Professional	TL-1D	Laceration	17046	3.5	0	0 [0%]
Fuller <i>et al</i> , 2013[27]	Professional	TL-1D	Laceration	15628	2.2	0	0 [0%]
Schwellnus <i>et al</i> , 2014[30]	Professional	TL-1D	Skin	15828	2.1	0	0 [0%]
Bird <i>et al</i> , 1998[21]	Amateur	TL-IM & MA	Laceration	12980	8.7	0.23	3 [2.7%]
Kerr <i>et al</i> , 2008[29]	Amateur	TL-1D	Laceration	72039	3.7	0.14	10 [3.8%]

Mean skin injury prevalence during training sessions of amateur and professional players combined was 0.05 ITEH (95% CI: 0.02 - 0.07) (Figure 3). Skin injury prevalence during training sessions in professional players was 0.02 ITEH (95% CI: 0.00 - 0.03) and in amateur players 0.15 ITEH (95% CI: 0.07 - 0.24). There was a significantly higher risk of skin injuries for amateur players compared to professional players during training sessions (odds ratio: 7.92; 95% CI: 1.19 - 52.78; $p = 0.03$). Four out of seven studies[23,24,27,30] reported that no laceration injuries were observed during training sessions. The proportion of skin and laceration injuries during training sessions ranged from 0 - 4.5%, with a weighted mean of 1.5%.

*** Figure 3 near here ***

DISCUSSION

The objective of this review was to identify the prevalence of skin injuries in rugby union during match play and training sessions and to investigate if the risk of injury differs between amateur and professional players. The skin injury prevalence during match play from the combined data of eleven studies observing rugby union players was 2.4 ITEH, accounting for 5.1% of all match injuries. This number could be interpreted as one time-loss or medical attention laceration injury per team (15 players) per year (20 matches, 80 minutes per match). Seven of the included studies in this review provided sufficient information to calculate skin and laceration injury prevalence during training sessions. Mean skin injury prevalence during training sessions was 0.05 ITEH, accounting for 1.5% of all training injuries. Injury prevalence has previously been found to be lower during training sessions than in match play,[3, 22, 27] and in this review a similar trend is observed when measuring skin injury prevalence (training: 0.05 ITEH; matches: 2.4 ITEH). In comparison, the incidence of abrasion and laceration injuries in association football matches - systematically summarised by van den Eijnde *et al*[12] - varied from 0.8 to 6.1 ITEH.

Ten out of twelve studies included in this review reported on professional rugby union injuries. No difference was found in skin injury risk during match play for amateur players compared to professional players (odds ratio: 0.60; 95% CI 0.17 - 2.08; $p = 0.42$). During training sessions the risk of skin injuries was higher for amateur players than for professional players (odds ratio: 7.9; 95% CI: 1.2 - 52.8; $p = 0.03$). Only two studies included in this review reported on amateur injuries,[21,29] from which one used a time-loss immediate in combination with a medical attention injury definition.[21] The injury definition could therefore have influenced this result. Williams *et al*[15] found no clear influence of playing level on injury risk during training sessions, though more elite levels of play were associated with a higher injury likelihood during matches (87% to 100% likelihood). The conclusions of Williams *et al*[15] were based on various levels of professional rugby (club level 2 to international), which makes it unclear if this trend could be extrapolated to injuries sustained in amateur rugby as well. The impact of professionalism on injuries has previously been investigated by Garraway *et al*[14] who concluded that professionalisation in rugby union had increased the prevalence of injuries in both amateur and professional cohorts. Financial rewards were suggested to raise the overall injury rate and increase the pressure on athletes to return to play as soon as possible, making recurrent injuries more likely. The lower risk of skin injuries found in professional players could indicate that this type of injury is frequently ignored in professional rugby injury counts, therefore underestimating the extent of the injury problem.[12, 13] Approximately one-third of amateur rugby union players have found their own studs sharpened due to wear.[6] In professional rugby union, players are more likely to have thorough stud checks, and are more likely to renew their footwear or studs frequently, therefore minimising the usage of damaged studs resulting from wear.

All studies included in this review used a 'time-loss' injury definition. Four studies used the 'time-loss immediate' definition and these studies had the highest percentage of laceration injuries of the twelve studies, suggesting that this type of injury often forces a player off the field, but not necessarily prevents them from playing subsequent matches. A consensus statement on the injury definitions used in epidemiological studies for rugby union recommends the use of time-loss, one day injury definitions.[10] Before the publication of this consensus statement, Ekstrand *et al*[32] warned that the use of time-loss injury definitions can lead to underestimation of the true prevalence of skin injuries. An example is given by Gibbs[13] who observed 62 lacerations which needed suturing during three rugby league seasons. One of these lacerations was officially marked as an injury because the wound became infected which resulted in missed training sessions and / or matches for the injured player. This situation reiterates the importance of appropriate injury definitions when estimating the skin injury risk of players; the medical attention injury definition is less likely to underestimate skin injury prevalence compared to one-day time-loss definition.[12]

Limitations

The results of this study should be interpreted in the light of its limitations. Since skin and laceration injuries are rarely the focus of epidemiological studies of sports injuries, a more accurate estimate of skin injury prevalence could have been obtained if more studies were included; specifically, 8 studies were excluded from this review because no separate skin or laceration injury category was available in the published article. Excluding articles that did not state a separate skin or laceration injury category could have biased the results if authors decided not to report this injury category because the injury frequency was too low. Any observed skin injuries would therefore be masked in the 'other' injury category. However, Fuller *et al*[24] still included a laceration injury category even though no lacerations were observed in the study. Four out of the seven studies reporting on training injuries also found no lacerations or skin injuries during training sessions in the observed time period, but still included the injury type as a separate category.[23, 24, 27, 30] According to the consensus statement on injury definitions and data collection procedures for studies of injuries in rugby union,[10] it is recommended that studies should report both separate laceration and abrasion injury categories. This is reiterated by van den Eijnde *et al*[12] who recommends reporting of contusions, lacerations and abrasions as separate injury categories since the causes of the injuries and the consequences for the athlete differ between these categories. The consensus statement on the collecting and reporting of epidemiological data had not been adopted by all studies in this review, often because studies were carried out before the statement was published. If future studies adopt these guidelines on reporting of injury categories, comparability between studies should improve. This increased standard in reporting can aid the validity of collated injury information across a number of studies and should encourage the reporting of 'null-results', where no injury in a specific category was observed.

A generalised linear model with negative binomial distribution with log link was used for the statistical analysis. Since the data showed overdispersion, it was not possible to apply a Poisson distribution as previously described.[15, 19] Using a negative binomial distribution has lowered the statistical power, therefore making a conservative estimate of the effect of the predictor on the outcome variable. For that reason, it could be that there is an effect of playing level on skin injury risk during matches (Figure 2) but more studies are needed to confirm or negate this.

Future research

There are a disproportionately small number of studies observing injuries in amateur cohorts. With less than 1% of all rugby union players receiving enough monetary reimbursement to focus on rugby full-time, professional players are over represented in published epidemiological studies. Since injury trends are not necessarily transferable between playing levels, an increase in studies observing amateur injury trends is needed. Furthermore, in this meta-analysis, the risk of bias in the two studies observing injuries in an amateur cohort[21, 29] was also higher than in the studies observing professional players[3, 20, 22–28, 30] (supplementary material 1). Therefore, future research should focus on establishing relevant skin injury frequencies in amateur cohorts, through equally robust methods as those applied to professional cohorts.

The female rugby population was underrepresented in this review. One previous study found differences in injury patterns between male and female intercollegiate rugby players,[33] but skin injuries were not investigated. The studies identified in this meta-analysis did not include enough female participants to determine if gender had an influence on skin injury risk. Due to increasing women's participation in the game,[34] focus should be given to document female rugby injuries.

In future, incidence and severity of laceration injuries could be assessed with a modified version of the Skin Damage and Severity index which was originally developed for assessing abrasive injuries in football.[35] This would allow for a more accurate estimation of the laceration injury prevalence in rugby union, minimising the non-reporting of these injuries caused by current injury definitions.

CONCLUSION

This systematic review summarised the published epidemiological literature on laceration or skin injury prevalence in rugby union. Following the inclusion criteria, twelve studies were selected for meta-analysis. Overall, skin and laceration injuries accounted for 5.1% of match injuries and 1.5% of training injuries in rugby union. A mean injury prevalence of 2.4 skin or laceration injuries per 1000 match exposure hours was found in this study, which could be interpreted as one time-loss injury per team, per season. During match play, no difference in skin and laceration injury risk was found

between amateur and professional players. In training sessions, amateur players had a higher risk of sustaining skin and laceration injuries than professional players. Future research should focus on high quality studies observing amateur players, female athletes, and potentially developing a tool for assessing acute skin injury damage in rugby union.

What are the new findings?

- **Skin injuries account for 5.1% of match injuries and 1.5% of training injuries in rugby union.**
- **Amateur players are more likely to sustain a laceration injury during training sessions than professional players**
- **Current injury definitions can lead to underestimation of skin and laceration injury prevalence.**
- **Future studies should focus on investigating amateur and female rugby skin injuries**

Competing interests

No potential conflict of interest was reported by the authors.

Funding

This project was internally funded.

6. REFERENCES

- 1 World Rugby. Player numbers. <http://www.worldrugby.org/development/player-numbers?lang=en> (accessed 13 Dec2016).
- 2 Nicholl JP, Coleman P, Williamst BT. The epidemiology of sports and exercise related injury in the United Kingdom. *Br J Sports Med* 1995;**29**:232–8. doi:10.1136/bjsm.29.4.232
- 3 Brooks JHM, Fuller CW, Kemp SPT, *et al.* Epidemiology of injuries in English professional rugby union: part 1 match injuries. *Br J Sports Med* 2005;**39**:757–66. doi:10.1136/bjsm.2005.018135
- 4 Hall MJ, Riou P. Football blades: a cause for concern. *Br J Sports Med* 2004;**38**:642–4. doi:10.1136/bjsm.2004.012245
- 5 Dennehy G. Dennehy -v- Adidas [Ireland] ltd. 2008;**134 P.**

- 6 Oudshoorn BY, Driscoll HF, Dunn M, *et al.* Causation Events of Stud Laceration Injuries in Rugby Union. *Procedia Eng* 2016;**147**:496–500. doi:10.1016/j.proeng.2016.06.227
- 7 van Mechelen W. Sports Injury Surveillance Systems. *Sport Med* 1997;**24**:164–8. doi:10.2165/00007256-199724030-00003
- 8 Ekstrand J, Karlsson J. Editorial: The risk for injury in football. There is a need for a consensus about definition of the injury and the design of studies. *Scand J Med Sci Sports* 2003;**13**:147–9.
- 9 Häggglund M, Waldén M, Bahr R, *et al.* Methods for epidemiological study of injuries to professional football players: developing the UEFA model. *Br J Sports Med* 2005;**39**:340–6. doi:10.1136/bjsm.2005.018267
- 10 Fuller CW, Molloy MG, Bagate C, *et al.* Consensus statement on injury definitions and data collection procedures for studies of injuries in rugby union. *Br J Sports Med* 2007;**41**:328–31. doi:10.1136/bjsm.2006.033282
- 11 World Rugby. Laws of the Game Rugby Union. 2016;:38.
- 12 van den Eijnde WAJ, Peppelman M, Lamers EAD, *et al.* Understanding the Acute Skin Injury Mechanism Caused by Player-Surface Contact During Soccer: A Survey and Systematic Review. *Orthop J Sport Med* 2014;**2**:1–8. doi:10.1177/2325967114533482
- 13 Gibbs N. Injuries in professional rugby league. A three-year prospective study of the South Sydney Professional Rugby League Football Club. *Am J Sports Med* 1993;**21**:696–700.
- 14 Garraway WM, Lee AJ, Hutton SJ, *et al.* Impact of professionalism on injuries in rugby union. *Br J Sports Med* 2000;**34**:348–51. doi:10.1136/bjsm.34.5.348
- 15 Williams S, Trewartha G, Kemp S, *et al.* A meta-analysis of injuries in senior men's professional Rugby Union. *Sport Med* 2013;**43**:1043–55. doi:10.1007/s40279-013-0078-1
- 16 Moher D, Liberati A, Tetzlaff J, *et al.* Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;**6**:1–7. doi:10.1371/journal.pmed.1000097
- 17 Waldén M, Häggglund M, Ekstrand J. The epidemiology of groin injury in senior football: a systematic review of prospective studies. *Br J Sports Med* 2015;**0**:1–7. doi:10.1136/bjsports-2015-094705
- 18 von Elm E, Altman DG, Egger M, *et al.* The Strengthening the Reporting of Observational

- Studies in Epidemiology (STROBE) Statement: guidelines for reporting observational studies. *Bull World Health Organ* 2007;**85**:867–72. doi:10.1136/bmj.39335.541782.AD
- 19 Lystad RP, Pollard H, Graham PL. Epidemiology of injuries in competition taekwondo: A meta-analysis of observational studies. *J Sci Med Sport* 2009;**12**:614–21. doi:10.1016/j.jsams.2008.09.013
 - 20 Best JP, McIntosh AS, Savage TN. Rugby World Cup 2003 injury surveillance project. *Br J Sports Med* 2005;**39**:812–7. doi:10.1136/bjism.2004.016402
 - 21 Bird YN, Waller AE, Marshall SW, *et al.* The New Zealand Rugby Injury and Performance Project: V. Epidemiology of a season of rugby injury. *Br J Sports Med* 1998;**32**:319–25. doi:10.1136/bjism.32.4.319
 - 22 Brooks JHM, Fuller CW, Kemp SPT, *et al.* Epidemiology of injuries in English professional rugby union: part 2 training injuries. *Br J Sports Med* 2005;**39**:767–75. doi:10.1136/bjism.2005.018408
 - 23 Brooks JHM, Fuller CW, Kemp SPT, *et al.* A prospective study of injuries and training amongst the England 2003 Rugby World Cup squad. *Br J Sports Med* 2005;**39**:288–93. doi:10.1136/bjism.2004.013391
 - 24 Fuller CW, Laborde F, Leather RJ, *et al.* International Rugby Board Rugby World Cup 2007 injury surveillance study. *Br J Sports Med* 2008;**42**:452–9. doi:10.1136/bjism.2008.047035
 - 25 Fuller CW, Raftery M, Readhead C, *et al.* Impact of the International Rugby Board's experimental law variations on the incidence and nature of match injuries in southern hemisphere professional rugby union. *South African Med J* 2009;**99**:232–7.
 - 26 Fuller CW, Molloy MG. Epidemiological study of injuries in men's international under-20 rugby union tournaments. *Clin J Sport Med* 2011;**21**:356–8. doi:10.1097/JSM.0b013e31821f5085
 - 27 Fuller CW, Sheerin K, Targett S. Rugby World Cup 2011: International Rugby Board injury surveillance study. *Br J Sports Med* 2013;**47**:1184–91. doi:10.1136/bjsports-2012-091155
 - 28 Jakoet I, Noakes TD. A high rate of injury during the 1995 Rugby World Cup. *South African Med J* 1998;**87**:45–7.
 - 29 Kerr HA, Curtis C, Micheli LJ, *et al.* Collegiate rugby union injury patterns in New England: a prospective cohort study. *Br J Sports Med* 2008;**42**:595–603. doi:10.1136/bjism.2007.035881

- 30 Schwellnus MP, Thomson A, Derman W, *et al.* More than 50% of players sustained a time-loss injury (>1 day of lost training or playing time) during the 2012 Super Rugby Union Tournament: a prospective cohort study of 17 340 player-hours. *Br J Sports Med* 2014;**48**:1306–15. doi:10.1136/bjsports-2014-093745
- 31 Fuller CW, Brooks JHM, Cancea RJ, *et al.* Contact events in rugby union and their propensity to cause injury. *Br J Sports Med* 2007;**41**:862–7. doi:10.1136/bjism.2007.037499
- 32 Ekstrand J, Timpka T, Häggglund M. Risk of injury in elite football played on artificial turf versus natural grass: a prospective two-cohort study. *Br J Sports Med* 2006;**40**:975–80. doi:10.1136/bjism.2006.027623
- 33 Peck KY, Johnston DA, Owens BD, *et al.* The incidence of injury among male and female intercollegiate rugby players. *Sports Health* 2013;**5**:327–33. doi:10.1177/1941738113487165
- 34 World Rugby. More women are playing rugby than ever before. <http://www.worldrugby.org/news/60275?lang=en> (accessed 13 Dec2016).
- 35 Van den Eijnde W, Peppelman M, Weghuis MO, *et al.* Psychosensorial assessment of skin damage caused by a sliding on artificial turf: The development and validation of a skin damage area and severity index. *J Sci Med Sport* 2014;**17**:18–22. doi:10.1016/j.jsams.2013.03.017

Figure legends

Figure 1: Flow chart of the literature search and selection process. Chart design adopted from Moher et al.[16]

Figure 2: Forest plot of skin injury prevalence during match play in studies observing amateur and professional injuries.

Figure 3: Forest plot of skin injury prevalence during training sessions in studies observing amateur and professional injuries.