

Submucosal diclofenac for acute postoperative pain in third molar surgery: A randomized, controlled clinical trial

GOECKI, P. <<http://orcid.org/0000-0002-5657-5001>>, RAINSFORD, Kim, TANEJA, P., BULSARA, Y., PEARSON, D., SAUND, D., AHMED, B. and DIETRICH, T. <<http://orcid.org/0000-0002-2557-7645>>

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/17592/>


This document is the Other

Citation:

GOECKI, P., RAINSFORD, Kim, TANEJA, P., BULSARA, Y., PEARSON, D., SAUND, D., AHMED, B. and DIETRICH, T. (2018). Submucosal diclofenac for acute postoperative pain in third molar surgery: A randomized, controlled clinical trial. *Journal of Dental Research*, 97 (4), 381-387. [Article]

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>



Supplementary Figure 1: Study Consort Flow Diagram

Eighty patients were screened for eligibility, out of which 75 were randomised and completed the randomised controlled trial.



Supplementary Figure 2: Partial flap necrosis in the 25mg/1mL HP β CD¹ diclofenac group, evident at 2-day review.

The patient was completely asymptomatic. The necrosis was present around the margins of the mucoperiosteal flap. This resolved without further intervention by the 7-day post-operative review appointment.

¹ HP β CD - hydroxypropyl- β -cyclodextrin



Supplementary Figure 3: Partial flap necrosis in the 50mg/1mL HP β CD diclofenac group evident at 7-day review.

The patient was asymptomatic and the necrosis recovered without further intervention (similar to the necrosis seen in the 25mg/1mL group – see Appendix Figure 2).

¹ HP β CD - hydroxypropyl- β -cyclodextrin