

Submucosal diclofenac for acute postoperative pain in third molar surgery: A randomized, controlled clinical trial

GORECKI, P. <<http://orcid.org/0000-0002-5657-5001>>, RAINSFORD, Kim, TANEJA, P., BULSARA, Y., PEARSON, D., SAUND, D., AHMED, B. and DIETRICH, T. <<http://orcid.org/0000-0002-2557-7645>>

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/17592/>


This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

GORECKI, P., RAINSFORD, Kim, TANEJA, P., BULSARA, Y., PEARSON, D., SAUND, D., AHMED, B. and DIETRICH, T. (2018). Submucosal diclofenac for acute postoperative pain in third molar surgery: A randomized, controlled clinical trial. *Journal of Dental Research*, 97 (4), 381-387.

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>



Supplementary Figure 1: Study Consort Flow Diagram

Eighty patients were screened for eligibility, out of which 75 were randomised and completed the randomised controlled trial.



Supplementary Figure 2: Partial flap necrosis in the 25mg/1mL HP β CD¹ diclofenac group, evident at 2-day review.

The patient was completely asymptomatic. The necrosis was present around the margins of the mucoperiosteal flap. This resolved without further intervention by the 7-day post-operative review appointment.

¹ HP β CD - hydroxypropyl- β -cyclodextrin



Supplementary Figure 3: Partial flap necrosis in the 50mg/1mL HP β CD diclofenac group evident at 7-day review.

The patient was asymptomatic and the necrosis recovered without further intervention (similar to the necrosis seen in the 25mg/1mL group – see Appendix Figure 2).

¹ HP β CD - hydroxypropyl- β -cyclodextrin