

Examining Lower Limb Injuries among Male Amateur Soccer Players

LAM, M.S.H, TUNG, K, MAN, D, LEE, K Y, FLINT, S and PEAKE, Rebecca

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/18233/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

LAM, M.S.H, TUNG, K, MAN, D, LEE, K Y, FLINT, S and PEAKE, Rebecca (2017). Examining Lower Limb Injuries among Male Amateur Soccer Players. *Journal of Yoga and Physical Therapy*, 7 (1), p. 264.

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/317322302>

Examining Lower Limb Injuries among Male Amateur Soccer Player

Article · January 2017

DOI: 10.4172/2157-7595.1000264

CITATIONS

6

READS

174

6 authors, including:



Michael Huen Sum Lam

Sheffield Hallam University

45 PUBLICATIONS 142 CITATIONS

SEE PROFILE



Ka-Yiu Lee

Sheffield Hallam University

27 PUBLICATIONS 198 CITATIONS

SEE PROFILE



Stuart Flint

Leeds Beckett University

61 PUBLICATIONS 89 CITATIONS

SEE PROFILE



Rebecca Peake

Sheffield Hallam University

10 PUBLICATIONS 11 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Unscheduled service utilisation [View project](#)



Students' National Identity (China, Hong Kong, Taiwan, Macau) and Beijing Olympics [View project](#)

All content following this page was uploaded by [Michael Huen Sum Lam](#) on 07 June 2017.

The user has requested enhancement of the downloaded file.

Examining Lower Limb Injuries among Male Amateur Soccer Player

Lam MHS^{1,2,4*}, Tung K¹, Man DP³, Lee KY^{1,2}, Lei Y⁴, Flint SW⁵ and Peake R¹

¹Academy of Sport and Physical Activity, Sheffield Hallam University, UK

²International Industry and Professional Accreditation Association, UK

³Vocational Training Council, Hong Kong

⁴School of Physical Education, University of Jinan, Shandong, China

⁵Leeds Beckett University, UK

*Corresponding Author:

Lam MHS

Academy of Sport and Physical Activity

Sheffield Hallam University, UK

Tel: 852 6542 3388

E-mail: michaelhslam@vtc.edu.hk

Journal of Yoga & Physical Therapy

Received date: March 31, 2017; **Accepted date:** May 22, 2017; **Published date:** May 29, 2017

Citation: Lam MHS, Tung K, Man DP, Lee KY, Lei Y, et al. (2017) Examining Lower Limb Injuries among Male Amateur Soccer Player. Journal of Yoga and Physical Therapy 7:264. doi:10.4172/2157-7595.1000264

Copyright: © 2017 Lam MHS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Amateur players far outnumbered professional athletes, but they are often under shadowed in literatures. Unlike professional athletes, amateur players may not possess the resources, knowledge or guidance in treating **sports injuries**. Soccer is one of the popular sports with fewer literatures addressing amateur players. Quantitative method was used for data collection and analysis. The practical contribution of this study lies in terms of the exploring of risks factors leading to lower limb injuries, most common type and area of injuries faced by amateur soccer players in contrast with professional players. Preventive measures were also discussed. The implications of these findings are valuable to coaches, **physical therapists** and governments for understanding amateur players and arouse public awareness on health and safety.

Keywords

Athletes; [Sports injuries](#); Lower limb; Injuries

<https://www.omicsonline.org/open-access/examining-lower-limb-injuries-among-male-amateur-soccer-player-2157-7595-1000264.php?aid=89483>

Introduction

Soccer is a popular sport; there are 240 million soccer players in the world. It is also one of the most popular [sports](#) among Hong Kong male citizens [1]. There are 145,780 unregistered soccer players and 4,176 registered soccer players in Hong Kong [2].

Many studies found that soccer has a high injury rate; it was reported to have the highest injury rate amongst all sports in Hong Kong [3]. Most of the injuries involved lower limbs [4-6]. Those injuries were mainly caused by tackling, running, being tackled, and shooting, twisting, turning, jumping and

landing. Lack of training, inadequate [rehabilitation](#); joint instability and muscle tightness contribute to most injuries in soccer, and the occurrence of injury for outdoor soccer was almost 2 times higher than that of indoor soccer [7,8].

Unlike professional players, amateur athletes in Hong Kong seldom play on grass pitches, and competitions are mainly held on the 180 outdoor 7-a-side hard surfaced soccer pitches provided by the Leisure and Culture Services Department. Besides, amateur athletes tended to seek self-treatment rather than approaching a doctor when they got injured [3], it is believed that most of the amateur soccer players do not know how to prevent and deal with the injuries since they do not have designated [physical therapist](#).

To reduce sports injury and better educate the public, understanding the risk factors is essential for prevention. There is lack of literature focus on amateurs; the current study is to examine the factors that lead lower limb injury and the most common types of injuries among amateur soccer players, which consists of a larger population than professional players. The results will also be compared with previous studies that focused mainly on professional players, preventive measures targeting the most common risks factors and injuries areas will then be discussed.

Literature Review

Lower limb injuries

Radelet et al. [9] defined injury as whenever a player needed any type of first aid during the event. Fuller et al. [10] refers injury as any physical complaint of a player due to soccer match or training, irrespective of the need for [medical](#) attention or time loss from football activities. An injury that results in a player receiving medical attention is referred to a 'medical attention' injury, and an injury that results in a player being unable to take part in future football training or match play as a 'time loss' injury [7-12].

Cluett [13] stated that lower limb refers to 'The part of body from the hip to toes' which includes hip, groin, thigh, knee, calf, ankle and foot [12]. Lower limb [injuries](#) are common in sports, Dick et al. [14] suggested that ankle, and knee and lower leg were the most common injury area in soccer, field

hockey, basketball, and lacrosse athletes. Chan et al. [15] also explained that lower limb injuries are easier to be found in weight-bearing sports such as soccer, basketball and distance running, etc. It is because those sports required a lot of footwork such as running, turning and jumping.

Lower limb injuries in soccer

The majority of soccer injuries are acute injuries which caused by [trauma](#) [11]. There are a lot of studies found that soccer has a high injury rate and most of the injuries in soccer involve lower limbs [3,5,6]. Elias [4] suggested ankle sprains were the most frequent injury. Chan et al. [15] indicated that lower limb injuries made up 83.07% of all injuries in Hong Kong, knee and ankle were the most common areas.

Injury types can be divided into sprain, strain, contusion, tendinitis and fracture. Wong and Hong [12] concluded that sprains, strains, contusion and tendinitis were the commonest injury types in soccer. Junge et al. [11] indicated that most of the soccer injuries occurred during matches. Keller et al. [7] shown further information that ligament sprains are one of the commonest injury types in any age or intensity of competition. Besides, muscles strains were only popular in professional and senior players and not common in youth players. Alternatively, contusion was more common in youth players than in professional and senior players.

Amateur soccer player

Federation Internationale de Football Association (FIFA) [16] explained that 'Amateurs are players who play without receiving any remuneration'. Carter et al. [17] stated that 'they engage in soccer for pleasure rather than for financial benefit or professional reasons.'

Materials and Method

Quantitative method was used to conduct this research. Questionnaire was selected for collecting data from the male amateur soccer players in Hong Kong. It is a reliable and valid method that obtains the data directly from the respondents. Apart from that, using questionnaire can collect a large number of data in short period and increase the representation of this research [18]. Besides, it is a more scientific and objective than other research method when the data has been quantified [19].

A questionnaire was constructed based on the objectives of the study. It is divided into 2 parts and there are 23 items in total. There are 7 items in Part A which collect the demographic information of the respondents such as age, weight, height, the information can also act as a tool for screening purposes. There are 16 items in Part B measuring the practices when the respondents play soccer and their injury experiences. For example, respondents were asked the duration of warm up, frequency of playing soccer and years of experience in playing soccer. The list of items were reduced to avoid redundancy after the pilot study, the questionnaire was also translated into Cantonese to prevent language barrier.

According to The Chinese University of Hong Kong [1], people who are 20-29 years old love to play sports in the evening and people who are 40 years old or above love to play sports in the morning. Respondents were recruited in three rounds from 3 7-a-side hard-surfaced soccer pitches in Hong Kong. The first round was on 11th-13th March 2016 at Fa Hui Park. The second round was on 18th-20th March 2016 at Kwai Chung Football Field. The third round was on 25th-27th March 2016 at Victoria Park. Survey was conducted from 9:00-10:00 and 17:00-19:00.

Male soccer players, aged 18 or above, reside in Hong Kong, playing at amateur or recreational level are eligible for inclusion. They generally had 1 year or more experiences in playing soccer, and had playing at least 1 practice session every month. A total of 150 subjects were successfully recruited for this study.

The statistical procedures were performed with SPSS (Version 21). Pearson Correlation and Chi-Square test were used for analyzing the data. The independent variables of this research are those factors that associated with lower limb injury prevalence of HK male amateur soccer player. The dependent variables are the injury prevalence of those factors.

Results

There are 150 subjects in total and all the subjects are male (weight 66.4 ± 6.5 kg, height 174.5 ± 4.4 cm). There are 88 subjects aged 18-29 year old, 38 subjects aged 30-54 year old and 24 subjects aged 55-64 year old. 4 subjects have played soccer for 1-2 years, 21 subjects have played soccer for 3-5 years, 29 subjects have played soccer for 6-9 years and 96 subjects have played soccer for 10 years or above.

The correlation between lower limb injury and other variables are shown in [Table 1](#).

	Not having lower limb injury Pearson Correlation	p-value
Having Warm up	-0.32	0.696
Time of Warm up	0.398**	0.000
Frequency of Playing Soccer (1 month)	0.12	0.144
Having Play Other Sport	0.88	0.286
Having worn Shin guards	0.052	0.574
Still Play Soccer when Exhausted	0.73	0.374
Age	-0.064	0.434

	Not having lower limb injury Pearson Correlation	p-value
Income	0.048	0.564
Experience in player soccer	-0.107	0.192

**Correlation is significant at the 0.01 level (2-tailed).

Table 1: Correlation between lower limb injury and other variables (N=115).

Referring to **Table 1**, there are positive correlation between lower limb injury and time of warm up ($r=0.398$) since the $p\text{-value}=0.00<0.01$ (2-tailed). It means that player is when the time of warm up increases, the tendency to suffer lower limb injury decreases.

The result of Chi-square test between lower limb injury and other variables are shown in **Table 2**.

	Chi-square Value	p-value
Education Attainment	0.56	0.456
Occupation	0.66	0.882
Type of Warm up	3.245	0.197
Type of Game	0.327	0.849

	Chi-square Value	p-value
Weather	1.898	0.387
Type of Pitch	0.671	0.715

Table 2: Results of Chi-square test between lower limb injury and other variables (N=115).

Referring to **Table 2**, those variables have no relation with lower limb injury since all the p-values>0.05.

The lower limb injury rate and common lower limb injuries in soccer are shown in **Tables 3 and 4**.

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Yes	115	76.7	76.7	76.7
	No	35	23.3	23.3	100.0
	Total	150	100.0	100.0	

Table 3: Lower limb injury rate in soccer (N=150).

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Ankle sprain	58	38.7	49.6	49.6

		Frequency	Percent	Valid Percent	Cumulative Percent
	Thigh muscle strain	27	8.0	23.1	72.6
	ACL Rupture	5	3.3	4.3	76.9
	Calf strain	7	4.7	6.0	82.9
	Displaced meniscus	2	.3	.7	84.6
	Hamstring strain	8	2.0	5.4	00.0
	Total	17	78.0	00.0	
Missing	System	33	22.0		
Total		50	00.0		

Table 4: Common lower limb injuries in soccer (N=150).

Referring to **Tables 3 and 4**, there are 76.7% (115 subjects) of the respondents reported that they have suffered from lower limb injury in 2015. 58 of them have suffered from ankle sprain, 27 subjects suffered from thigh muscle strain, 18 subjects suffered from hamstring strain, 7 subjects suffered from calf strain, 5 subjects suffered from ACL rupture and 2 subjects suffered from displaced meniscus.

Discussion

Determinants of physical activity [20-22] and sport participation [23-27] have been extensively examined as it can influence both psychological [28,29] and physical well-being [30-32]. In particular, sport participation has been growing in recent years due to increased health awareness of people. Consequently, prevention of sport injuries has been attracting attention in the community.

Time of warm up was the only factor found in this study that is associated with lower limb injuries among male amateur soccer player in HK. There was positive correlation between lower limb injury and time of warm up. It means that player is less likely to suffer lower limb injury when they have a longer duration for warm up.

The results give new insights to Keller et al.'s [7] findings, just like professional players, warm up could reduce the risk of suffering lower limb injury among amateur players, since it can improve the flexibility of the players. Player who has poor flexibility would easily cause muscle strain and tendinitis. Thus, having a proper warm up in appropriate length of time could reduce the rate of lower limb injury.

Besides, the result of the study also supported Keller et al. [7] that soccer injuries are equally distributed between practice and match situation that the type of game would not affect the injury rate, both practice and match could have the probability to cause injury. It might be because soccer players in the same level are facing similar aggression and pressure in match or training. However, the severity of the injuries between amateur and professional players might be different.

The lower limb injury rate for soccer players in this study was 76.7% and the most common lower limb injury was ankle sprain (49.6%), followed by thigh muscle strain (23.1%), hamstring strain, which has similar ranking as National Collegiate Athletic Association's findings in 1993. But the result contradicts with Chan et al [3], who suggested that the most common injured area was knee (50.47%) followed, by ankle. The possible reason of having different result is that the subjects were recruiting from different places. Subjects were recruited from football pitches in this study and subjects were recruited from the Sports Injury Clinic in the study conducted by Chan et al [3]. Recruiting subjects from the Sports Injury Clinic could have higher possibility of excluding players who suffered injury such as the grade 1 ankle sprain. Also, the amateur players, who would choose to attend bone-setter rather than doctor if they suffered minor injury [15].

In contrast to Keller et al. [7], the result in this study could not find significant correlation between weather or pitch type and lower limb injuries. It might be because most of the amateur soccer players in Hong Kong are playing on hard-surfaced pitch but not turf-surfaced pitch. Besides, most of the amateur soccer players can avoid playing soccer during bad weather.

To prevent injury in playing soccer Keller et al. [7] suggested warmup and cool-down should be implemented before and after the game. Proprioceptive neuromuscular facilitation (PNF) stretching should be introduced to the warm-up and cool-down session, which is more effective to prevent injury than static stretching. It was found that ankle sprains occur more often in players who suffered ankle injury before, therefore, for these players, it was highly recommended to tape their ankles before playing soccer. Arnason et al. [33] found that inadequate muscle strength, imbalance in the hamstring to quadriceps strength ratio and side-to-side imbalances could cause hamstring strains. Hence, the eccentric strength training can improve the strength of hamstring for preventing injury. Alentorn-Geli et al. [34] suggested modifying the landing technique of the player could help to prevent ACL injuries. Training should include drills that familiarize players with making unanticipated changes of direction.

This study has several limitations. The frequency and types of injuries were assessed using questionnaire alone in which the validity and reliability are not confirmed since this is a pilot study. The body of evidence suggests that objective measures are important in a study design [20,35-41]. Besides, the data collected was limited and may not be able to generalize amateur soccer players in other places with other local habits.

Conclusion

The findings of the study could contribute to the society by arousing public's awareness and identifying possibly lower limb injuries of amateur soccer players; effective preventive measures targeting specific injuries are also examined as apart from understanding the injury mechanism, understanding what factors associated with lower limb injuries is also important for preventing lower limb injuries [42].

References

1. The Chinese University of Hong Kong (2009) [Consultancy study on sport for all: Participation patterns of Hong Kong People in physical activities.](#)
2. Federation International de Football Association (2015) [Big count.](#)
3. Chan KM, Yuan Y, Li CK, Chien P, Tsang G (1993) Sports causing most injuries in Hong Kong. *Br J Sports Med* 27: 263-267.
4. Elias SR (2001) [10-year trend in USA Cup soccer injuries: 1988-1997. Med Sci Sports Exerc 33: 359-367.](#)
5. Schmidt-Olsen S, Bünemann LK, Lade V, Brassøe JO (1985) [Soccer injuries of youth. Br J Sports Med 19: 161-164.](#)
6. Yde J, Nielsen AB (1990) Sports injuries in adolescents' ball games: Soccer, handball and basketball. *Br J Sports Med* 24: 51-54.
7. Keller C, Noyes F, Buncher C (1987) [The medical aspects of soccer injury epidemiology. Am J Sports Med 15: 203-207.](#)
8. Ekstrand J, Gillquist J (1982) [The frequency of muscle tightness and injuries in soccer players. Am J Sports Med 10: 75-78.](#)
9. Radelet MA, Lephart SM, Rubinstein EN, Myers JB (2002) Survey of injury rate for children in community sports. *Pediatr* 110: 1-11.
10. Fuller C W, Ekstrand J, Junge A, Andersen T E, Bahr R, et al. (2006) British Consensus statement on injury definitions and data collection procedures in studies of football (soccer) injuries. *J Sports Med* 40: 193-201.
11. Junge A, Cheung K, Edwards T, Dvorak J (2004) [Injuries in youth amateur soccer and rugby players-comparison of incidence and characteristics. Br J Sports Med 38: 168-172.](#)

12. Wong P, Hong Y (2005) [Soccer injury in the lower extremities. Br J Sports Med 39: 473-482.](#)
13. Cluett J (2014) [Information about anatomy: Lower extremity.](#)
14. Dick R, Agel J, Marshall SW (2007) [National Collegiate Athletic Association Injury Surveillance System Commentaries: Introduction and methods. J Athl Train 42: 173-182.](#)
15. Chan KM, Fu F, Leung L (1984) [Sports injuries survey on university students in Hong Kong. Br J Sports Med 18: 195-202.](#)
16. Federation Internationale de Football Association (1956) [Amateur definition of the F.I.F.A.](#)
17. Carter S, Finkel A, Fortna P (2010) The west and beyond: New perspectives on an imagined region. Au Press, Athabasca University.
18. Popper KR, Weiss G (1959) [The logic of scientific discovery. Physics Today 12: 53-54.](#)
19. Harrop C (1981) Data collection in context: S. Ackroyd and J.A. Hughes. Longman London.
20. Cerin E, Lee KY, Barnett A, Sit CH, Cheung MC, et al. (2013) [Objectively-measured neighborhood environments and leisure-time physical activity in Chinese urban elders. Prev Med 56: 86-89.](#)
21. Lam MHS, Leung AYM (2016) [The effectiveness of health literacy oriented programs on physical activity behaviour in middle aged and older adults with type 2 diabetes: A systematic review. Health Psychol Res 4: 7-12](#)
22. Lee KY, Lee PH, Macfarlane D (2014) [Associations between moderate-to-vigorous physical activity and neighbourhood recreational facilities: The features of the facilities matter. Int J Environ Res Public Health 11: 12594-12610.](#)

23. Ho G, Yiu EYM, Lam MHS (2016) [The Hong Kong games in the eyes of local sports and recreation students. Int J Hist Sport 33: 1209-1225.](#)
24. Lam MHS (2016) [Exercise game exhilarates the elderly: A challenge to traditional training. J Athl Enhanc 5: 1-2.](#)
25. Lau PW, Lam MHS, Leung BW (2010) [National Identity and the Beijing Olympics: School Children's Responses in Mainland China, Taiwan and Hong Kong. Procedia Soc Behav Sci 2: 6729-6738.](#)
26. Lau PW, Lam MHS, Leung BW, Choi CR, Ransdell LB (2012) [The longitudinal changes of national identity in mainland China, Hong Kong and Taiwan before, during and after the 2008 Beijing Olympics Games. The International Journal of the History of Sport 29: 1281-1294.](#)
27. Lau PW, Lam MHS, Leung BW (2011) [The Beijing Olympics and Expressions of National Identity in China, Taiwan and Hong Kong. The Olympics in East Asia: Nationalism, Regionalism, and Globalism on the Center Stage of World Sports 147.](#)
28. Deng Y, Lee KY, Lam MHS, Lee PH (2016) Understanding socio-behavioral mitigators of depressive symptoms among US young adults. Behav Med 42: 217-226.
29. Lam MHS, Leung AYM, Chan SSC (2011) Psychological and cognitive determinants of the Health literacy on soon to be aged and older adults: A systematic review. Imanagers J Nurs 1: 46.
30. Fung L, Lam MHS (2012) [Effectiveness of a progressive stepping program on lower limb function in community dwelling older adults. J Exerc Sci Fit 10: 8-11.](#)
31. Lam MHS, Cheung SY, Chow BC (2011) [Effects of Tai Chi soft ball training on health-related quality of life of older adults with functional limitations. Asian J Gerontol Geriatr 6: 65-71.](#)

32. Lam MHS, Kok EYL, Louie HTL, Lee KY (2014). [External Chinese martial arts and health. Martial arts for health: Translating research into practice. OMICS Group eBook, California.](#)
33. Arnason A, Andersen TE, Holme I, Engebretsen L, Bahr R (2008) [Prevention of hamstring strains in elite soccer: An intervention study. Scand J Med Sci Sports 18: 40-48.](#)
34. Alentorn-Geli E, Myer G, Silvers H, Samitier G, Romero D, et al. (2009) [Prevention of non-contact anterior cruciate ligament injuries in soccer players. Part 1: Mechanisms of injury and underlying risk factors. Knee Surg Sports Traumatol Arthrosc 17: 705-729.](#)
35. Cerin E, Chan KW, Macfarlane DJ, Lee KY, Lai PC (2011) [Objective assessment of walking environments in ultra-dense cities: Development and reliability of the environment in Asia Scan Tool—Hong Kong version \(EAST-HK\). Health Place 17: 937-945.](#)
36. Cerin E, Lee KY, Barnett A, Sit CH, Cheung MC, et al. (2013) [Walking for transportation in Hong Kong Chinese urban elders: A cross-sectional study on what destinations matter and when. Int J Behav Nutr Phys Act 10: 78.](#)
37. Lee KY, Macfarlane DJ, Cerin E (2013) [Comparison of three models of actigraph accelerometers during free living and controlled laboratory conditions. Eur J Sport Sci 13: 332-339.](#)
38. Lee KY, Lam MHS, Deng Y. (2017) [Measuring post-concussive activity levels of patients: Step count or activity intensity? JAMA Pediatr.](#)
39. Lee KY, Lam MHS, Lee PH (2017) [Distance from home to the nearest tobacco outlet may not reflect the true accessibility. JAMA Intern Med 177: 287-287.](#)
40. Lee KY, Macfarlane D, Cerin E (2013) [Objective evaluation of recreational facilities: Development and reliability of the recreational facility audit tool. J Park Recreat Admi 31.](#)

41. Lam MHS, Leung AYM (2011) [The effects of Tai-Chi-Soft-Ball training on physical functional health of Chinese older adult. J Hum Sport Exerc 6: 540-553.](#)
42. Murphy DF, Connolly DAJ, Beynnon BD (2003) [Risk factors for lower extremity injury: A review of the literature. Br J Sports Med 37: 13-29.](#)