

## **HIIT: Current evidence and future application in cardiovascular rehabilitation**

NICHOLS, Simon <<http://orcid.org/0000-0003-0377-6982>>

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/21041/>

---

This document is the Presentation

### **Citation:**

NICHOLS, Simon (2017). HIIT: Current evidence and future application in cardiovascular rehabilitation. In: BACPR Annual Conference 2017, London, 5-6 October 2017. (Unpublished) [Conference or Workshop Item]

---

### **Copyright and re-use policy**

See <http://shura.shu.ac.uk/information.html>



# **HIIT: Current evidence and future application in cardiovascular rehabilitation**

Dr Simon Nichols

✉ [s.j.nichols@shu.ac.uk](mailto:s.j.nichols@shu.ac.uk)  
🐦 [@nichols87simon](https://twitter.com/nichols87simon)

BACPR Annual Conference 2017

**Conflicts of Interest: None**

# Outline


- Brief background of HIIT
- The aerobic fitness benefits(?) of HIIT
- HIIT in cardiac populations
- Emerging UK evidence

# What is HIIT?

Repeat bouts of short duration high intensity exercise interspersed with short duration low intensity active recovery periods

Some disagreement in intensity zones - typically submaximal efforts >80% of maximal HR/>80 peak work rate

## Variations in exercise volume/programme length



# Where did HIIT come from?

Reported as early as the 1920s in athletic populations

1954 - Sir Rodger Bannister used HIIT during as a medical student during his lunch break - achieved the 4 minute mile

1960's peer-reviewed literature begins to emerge in healthy/athletic populations

1970/80's— Evidence for interval and high intensity interval training in clinical populations begins to emerge

1996 – Katerina Meyer found that interval training in CHF was safe – Assessed catecholamine, cardiac/metabolic stress, & dyspnoea - CHF patients can tolerate HIIT.

Today – A vast volume of literature on the efficacy of HIIT in health and disease

# Is it safe?

**Table 1.** The number of patients, exercise-hours and the corresponding number of cardiovascular events associated with moderate- and high-intensity exercise, respectively.

Center	Patients (n)	Total training (hours)	Moderate-intensity (hours)	High-intensity (hours)
Ålesund	775	25 720 <sup>1</sup>	15 232	10 488 <sup>1</sup>
Feiring	2629	85 208 <sup>2</sup>	63 032 <sup>1</sup>	22 176 <sup>1</sup>
Røros	1442	64 892	51 192	13 700
<b>Total</b>	<b>4846</b>	<b>175 820</b>	<b>129 456</b>	<b>46 364</b>
<b>Event rates:</b>				
Cardiac arrest, fatal			1	0
Cardiac arrest, non-fatal			0	2
Myocardial infarction			0	0
<b>Risk of events</b>		<b>1/58 607</b>	<b>1/129 456</b>	<b>1/23 182</b>

The likelihood of a cardiac event in high risk individuals appears to be low when conducting either moderate, or high intensity exercise

# Is it Effective?

$VO_{2max}$ , insulin sensitivity and endothelial function all appear to improve to a greater extent during HIIT, compared to MISS

Findings can be variable


Improved endothelial  
function/increased  
NO

Improvements in muscle  
oxidative  
capacity/mitochondrial  
volume/quality

Greater depletion of  
muscle glycogen stores  
leading to enhanced  
muscle glycogen uptake  
→ improve insulin  
sensitivity

# Is it Really Effective?

## Higher Vol. — Bacon '13



Favours HIIT 0.51 L/min (43 to 0.60 L/min) up to 0.9 L for longer studies

## Lower Vol. — Weston '14

	Effect on VO <sub>2max</sub> (%)		Inference
	Mean	±90 % CL	
Effect on treatment groups <sup>a</sup>			
Sedentary males	10.0	±5.1	Possibly moderate ↑
Sedentary females	7.3	±4.8	Likely small ↑
Active non-athletic males	6.2	±3.1	Likely moderate ↑
Active non-athletic females	3.6	±4.3	Possibly moderate ↑
Athletic males	2.7	±4.6	Unclear
Controls	1.2	±2.0	Unclear

Weston et al (2014)

The effects of low vol. HIIT appear to favour less fit populations


There is no definitive consensus on whether HIIT is superior to *Well Prescribed* endurance exercise training apparently populations



# HIIT and Cardiometabolic Health

The effects of short-term HIIT also favour less fit populations

Batacan '17



Similar to VO<sub>2peak</sub>, HIIT appears more effective at improving cardiometabolic health when individuals are unfit or sedentary

# Adaptations in sedentary individuals

Type	Frequency	Time	Intensity	Result
MISS	Exercise 3 x p/week for 8 weeks	Exercise progressed from 20 to 35 minutes	~60% PPO	$VO_{2peak}$ - MISS 9% : HIIT 15% $a-VO_2$ diff – MISS ↑: HIIT ↑ $Q_{max}$ : MISS ~ : HIIT ↑ $VO_2$ Kinetics - MISS~ : HIIT: ↑ Exhaustion Time - MISS↑: MISS↑ ↑ Capillary/Fibre R - MISS↑: MISS↑
HIIT			4 min low / 1 minute 90% PPO	


Effect of interval versus continuous training on cardiorespiratory and mitochondrial functions: relationship to aerobic performance improvements in sedentary subjects

**Frédéric N. Daussin,<sup>1</sup> Joffrey Zoll,<sup>1</sup> Stéphane P. Dufour,<sup>1</sup> Elodie Ponsot,<sup>1</sup> Evelyne Lonsdorfer-Wolf,<sup>1</sup> Stéphane Doutreleau,<sup>1</sup> Bertrand Mettauer,<sup>1,2</sup> François Piquard,<sup>1</sup> Bernard Geny,<sup>1</sup> and Ruddy Richard<sup>1</sup>**

<sup>1</sup>CHRU of Strasbourg, Physiology and Functional Explorations Department, Civil Hospital, Strasbourg, France and University Louis Pasteur, Faculty of Medicine, Physiology Department, Strasbourg, France; and <sup>2</sup>Cardiology Department, Civil Hospital, Colmar, France

So what might this mean for cardiovascular rehabilitation programmes?

# Is HIIT Effective in Cardiac Populations?



Mean difference favours HIIT by 3.03 mL/kg/min (95% CI 2.00 to 4.07;  $p < 0.001$ )


Six out of 10 studies conducted by the same research group

Only 273 patients included

Other systematic reviews/meta-analyses show similar results

## Heart Failure Only

Mean difference favours HIIT by 2.14 mL/kg/min (95% CI 0.66 to 3.63)



# The Clinical Significance?

$\text{VO}_{2\text{peak}}$  is one of the strongest clinical prognosticators. Improvements in  $\text{VO}_{2\text{peak}}$  are consistently associated with improved survival:


- Kodama *et al.* (2009): ~103,000 patients; demonstrate 1 MET improvement in aerobic fitness confers 13% survival advantage
- Myers *et al.* (2002): ~6200 patients: 1 MET improvement in aerobic fitness confers 12% survival advantage
- Vanhees *et al.* (1995): 1% improvement in exercise training induced  $\text{VO}_{2\text{peak}}$  confers a 2% survival advantage in patients with CHD

# Low aerobic fitness indicates poor cardiometabolic health - Unpublished

- Lean body mass
- Android/gynoid ratio
- Ventilatory anaerobic threshold
- Heart rate recovery (1, 2, 3 and 6 minutes)
- Haemoglobin/Haematocrit

- BMI
- Waist to hip ratio
- Left ventricular ejection fraction
- Previous MI
- Lipid profiles

- NT-proBNP
- hs-CRP
- Glucose (non-fasting)
- VE/VCO<sub>2</sub> slope
- Oxygen uptake efficiency slope
- Resting heart rate



# The Role of HIIT in Cardiac Populations

## Exercise-based cardiac rehabilitation for coronary heart disease (Review)

Heran BS, Chen JMH, Ebrahim S, Moxham T, Oldridge N, Rees K, Thompson DR, Taylor RS



THE COCHRANE  
COLLABORATION®

## Exercise-Based Cardiac Rehabilitation for Coronary Heart Disease

### Cochrane Systematic Review and Meta-Analysis

Lindsey Anderson, PhD,\* Neil Oldridge, PhD,† David R. Thompson, PhD,‡ Ann-Dorthe Zwisler, MD,§ Karen Rees, PhD,|| Nicole Martin, MA,¶ Rod S. Taylor, PhD\*

## Changes in cardiorespiratory fitness in cardiac rehabilitation patients: A meta-analysis

Gavin Sandercock\*, Valentina Hurtado, Fernando Cardoso

## Rehabilitation after myocardial infarction trial (RAMIT): multi-centre randomised controlled trial of comprehensive cardiac rehabilitation in patients following acute myocardial infarction

Robert R West,<sup>1</sup> Dee A Jones,<sup>2</sup> Andrew H Henderson<sup>3</sup>

## Cardiorespiratory fitness changes in patients receiving comprehensive outpatient cardiac rehabilitation in the UK: a multicentre study

Gavin R H Sandercock,<sup>1</sup> Fernando Cardoso,<sup>1</sup> Meshal Almodhy,<sup>1</sup> Garyfallia Pepera<sup>2</sup>

## The minimum clinically important improvement in the incremental shuttle walk test following cardiac rehabilitation

Linzy Houchen-Wolloff, Sally Boyce and Sally Singh

# The Role of HIIT in Cardiac Populations

- Evidence for the application of HIIT in clinical practice is evolving
- There remains a lack of high quality 'pragmatic' real-world evidence
- A major research/logistical challenge in the UK? Prescribing HIIT when maximal exercise testing is not widely available.





# @HIITorMISSUK - Study update

Pragmatic multi-centre  
RCT – 510 patients

Eight weeks – 2 x per week

10 x high intensity bouts at 85 -  
90% PPO

10 x low intensity bouts at 20 -  
25% PPO

Control group – standard care  
at 40-70% HRR

Assessed following intervention  
~8 weeks and at 12 months

Primary outcome measure -  
 $VO_{2peak}$

Also assessing other  
physiological, psychosocial and  
economic outcomes


**BMJ Open High-intensity interval training versus moderate-intensity steady-state training in UK cardiac rehabilitation programmes (HIIT or MISS UK): study protocol for a multicentre randomised controlled trial and economic evaluation**

Gordon McGregor,<sup>1,2</sup> Simon Nichols,<sup>3</sup> Thomas Hamborg,<sup>4</sup> Lucy Bryning,<sup>5</sup> Rhiannon Tudor-Edwards,<sup>5</sup> David Markland,<sup>6</sup> Jenny Mercer,<sup>2</sup> Stefan Birkett,<sup>3</sup> Stuart Ennis,<sup>1,2</sup> Richard Powell,<sup>1</sup> Brian Begg,<sup>2,7</sup> Mark J Haykowsky,<sup>8</sup> Prithwish Banerjee,<sup>1,9</sup> Lee Ingle,<sup>3</sup> Rob Shave,<sup>2</sup> Karianne Backx<sup>2</sup>



@HIITorMISSUK 

# @HIITorMISSUK



# Have Your Say!

- We would like to understand how UK CR exercise programmes are currently operating.
- At the end of our survey, there is a section relating to the future role that HIIT may have in UK CR programmes.
- We just need one response from each team – Someone directly involved in exercise provision
- <https://northumbria.onlinesurveys.ac.uk/an-evaluation-of-exercise-provision-within-uk-cardiac-reha>

**Sheffield  
Hallam  
University**



# Summary

- Compared to MISS, HIIT appears to provide superior health benefits
- Variation in HIIT protocols and magnitude of benefit
- HIIT appears to be safe and effective in cardiac populations – HIITorMISSUK may provide much needed ‘real-world’ evidence.



# Thank You

✉ [s.j.nichols@shu.ac.uk](mailto:s.j.nichols@shu.ac.uk)  
🐦 [@nichols87simon](https://twitter.com/nichols87simon)

# References

- Meyer K, Samek L, Schwaibold M, Westbrook S, Hajric R, Lehmann M, Essfeld D and Roskamm H. Physical responses to different modes of interval exercise in patients with chronic heart failure--application to exercise training. *Eur Heart J* (1996); **17**: 1040-1047.
- Rognmo Ø, Moholdt T, Bakken H, Hole T, Mølsted P, Myhr NE, Grimsø J and Wisløff U. Cardiovascular Risk of High- Versus Moderate-Intensity Aerobic Exercise in Coronary Heart Disease Patients. *Circulation* (2012); **126**: 1436-1440.
- Daussin FN, Zoll J, Dufour SP, Ponsot E, Lonsdorfer-Wolf E, Doutreleau S, Mettauer B, Piquard F, Geny B and Richard R. Effect of interval versus continuous training on cardiorespiratory and mitochondrial functions: relationship to aerobic performance improvements in sedentary subjects. *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology* (2008); **295**: R264-R272
- Bacon AP, Carter RE, Ogle EA and Joyner MJ.  $\text{VO}_{2\text{max}}$  trainability and high intensity interval training in humans: a meta-analysis. *PloS one* (2013); **8**: e73182.
- Weston M, Taylor KL, Batterham AM and Hopkins WG. Effects of Low-Volume High-Intensity Interval Training (HIT) on Fitness in Adults: A Meta-Analysis of Controlled and Non-Controlled Trials. *Sports Med* (2014); **44**: 1005-1017.
- Weston KS, Wisløff U and Coombes JS. High-intensity interval training in patients with lifestyle-induced cardiometabolic disease: a systematic review and meta-analysis. *British Journal of Sports Medicine* (2014); **48**: 1227-1234.
- Haykowsky MJ, Timmons MP, Kruger C, McNeely M, Taylor DA and Clark AM. Meta-analysis of aerobic interval training on exercise capacity and systolic function in patients with heart failure and reduced ejection fractions. *The American journal of cardiology* (2013); **111**: 1466-1469.
- Kodama S, Saito K, Tanaka S and et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: A meta-analysis. *JAMA* (2009); **301**: 2024-2035.
- Myers J, Prakash M, Froelicher V, Do D, Partington S and Atwood JE. Exercise Capacity and Mortality among Men Referred for Exercise Testing. *New England Journal of Medicine* (2002); **346**: 793-801.
- Vanhees L, Fagard R, Thijs L and Amery A. Prognostic value of training-induced change in peak exercise capacity in patients with myocardial infarcts and patients with coronary bypass surgery. *The American journal of cardiology* (1995); **76**: 1014-1019.
- Heran BS, Chen J, Ebrahim S, Moxham T, Oldridge N, Rees K, Thompson DR and Taylor RS. Exercise-based cardiac rehabilitation for coronary heart disease. *Cochrane Database Syst Rev* (2011); **7**.
- Anderson L, Oldridge N, Thompson DR, Zwisler A-D, Rees K, Martin N and Taylor RS. Exercise-Based Cardiac Rehabilitation for Coronary Heart DiseaseCochrane Systematic Review and Meta-Analysis. *Journal of the American College of Cardiology* (2016); **67**: 1-12.
- Sandercock G, Hurtado V and Cardoso F. Changes in cardiorespiratory fitness in cardiac rehabilitation patients: A meta-analysis. *International Journal of Cardiology* (2011).
- West RR, Jones DA and Henderson AH. Rehabilitation after myocardial infarction trial (RAMIT): multi-centre randomised controlled trial of comprehensive cardiac rehabilitation in patients following acute myocardial infarction. *Heart* (2012); **98**: 637-644.
- Sandercock G, Cardoso F, Almodhy M and Pepera G. Cardiorespiratory fitness changes in patients receiving comprehensive outpatient cardiac rehabilitation in the UK: a multicentre study. *Heart* (2013); **99**: 785-790.
- Houchen-Wolloff L, Boyce S and Singh S. The minimum clinically important improvement in the incremental shuttle walk test following cardiac rehabilitation. *European Journal of Preventive Cardiology* (2014).
- McGregor G, Nichols S, Hamborg T, Bryning L, Tudor-Edwards R, Markland D, Mercer J, Birkett S, Ennis S and Powell R. High-intensity interval training versus moderate-intensity steady-state training in UK cardiac rehabilitation programmes (HIIT or MISS UK): study protocol for a multicentre randomised controlled trial and economic evaluation. *BMJ open* (2016); **6**: e012843.