

An observational study of temperature and thermal images of surgical wounds for detecting delayed wound healing within four days after surgery

SIAH, Chiew Jiat, CHILDS, Charmaine http://orcid.org/0000-0002-1558-5633, CHIA, Chung King and CHENG, Kin-Fong Karis

Available from Sheffield Hallam University Research Archive (SHURA) at:

https://shura.shu.ac.uk/23787/

This document is the Accepted Version [AM]

Citation:

SIAH, Chiew Jiat, CHILDS, Charmaine, CHIA, Chung King and CHENG, Kin-Fong Karis (2019). An observational study of temperature and thermal images of surgical wounds for detecting delayed wound healing within four days after surgery. Journal of clinical nursing, 28 (11-12), 2285-2295. [Article]

Copyright and re-use policy

See http://shura.shu.ac.uk/information.html

Table 1
Participants' Demographics Data Between the Non-infected and Infected Surgical Wound
Groups (N=60)

Groups (N=00)							
	Infected abdominal surgical			fected abdominal			
		wounds surgical wounds					
		n = 15		n = 45	_	p-value	
Characteristics	F	%	f	%	χ^2	p-value	
Gender:							
Female	5	33.3	22	48.9	0.7.	0 4 7 NS	
Male	10	66.7	23	51.1	0.56	$0.45^{\rm NS}$	
Ethnicity:							
Chinese	13	86.7	40	88.9			
Malay	1	6.7	4	8.9	0.74	$0.69^{\rm \ NS}$	
Others	1	6.7	1	2.2			
Diagnosis:							
Carcinoma	13	86.6	42	93.4			
Colon							
Diverticular	1	6.7	2	4.4	0.83	0.66^{NS}	
Disease							
Others	1	6.7	1	2.2			
Characteristics	Mean	Median (Range)	Mean	Median (Range)	Z	p-value	
	(SD)		(SD)			_	
Age (years)	66	65 (48–81)	64	65 (47-82)	-0.46	0.64 ^{NS}	
	(9)		(9)	03 (47-82)	-0.46	0.04	
Body core	36.3	26 4 (25 7 26 9)	36.4	26 4 (25 0 27 0)	0.26	0.79^{NS}	
temperature (°C)	(0.3)	36.4 (35.7–36.8)	(0.2)	36.4 (35.9- 37.0)	-0.26	0.79	
Peristomal skin	34.7	245 (22 2 26 9)	35.1	25 4 (22 7 26 2)	1.62	0.10^{NS}	
temperature (°C)	(1.04)	34.5 (33.3–36.8)	(0.7)	35.4 (33.7-36.3)	-1.63	0.10	

Clinical Data between the Non-infected and the Infected Surgical Wound Groups (N=60) Infected abdominal Non- infected abdominal surgical wounds surgical wounds N = 15N = 45 χ^2 F Characteristics % f % pvalue BMI category Underweight 2 13.3 3 6.7 Normal/Pre-Obese 12 80.0 40 88.9 $0.67\ ^{\text{NS}}$ 0.81 Overweight 1 2 4.4 6.7 Type of surgical procedure Closure of Ileostomy 11 73.3 34 75.6 $1.00\ ^{\rm NS}$ 0.03^{a} Closure of Colostomy 4 26.7 11 24.4 Site of Wound Right Lumbar 12 80.0 31 68.9 Umbilicus 0 0.0 2 4.4 $0.78\ ^{NS}$ Left Lumbar 2 13.3 8 17.9 1.77 Right Hypochondriac 0 0.0 2 4.4 2 Left Hypochondriac 1 6.7 4.4 Characteristics Mean (SD) Median Mean Median \overline{Z} p-

(Range)

(SD)

(Range)

value

Abdominal Girth (cm)	85.27 (11.35)	86.00 (60–106)	86.04 (9.87)	86.00 (63–111)	-0.06	0.95 ^{NS}
Adipose tissue thickness (mm)	15.60 (6.63)	18.00 (3–24)	18.11 (7.73)	18.00 (5–34)	-0.90	0.37^{NS}
Length of abdominal surgical wounds (cm)	5.93 (0.65)	6.00 (5–7)	6.10 (0.94)	6.00 (5–10)	-0.43	0.67 ^{NS}

Table 2 Temperature Readings of Surgical Site Surface between the Non-infected and Infected Surgical Wounds (N = 60)

		Temperature (°C)							
		Non-infected		ed	Infected				
		Overall				Overall			
		Median	IQR	Median	Median	IQR	Median	Z	p-value
				(IQR)			(IQR)		
Immediately	Body core temperature	36.3	35.8–37.3		36.3	35.4–37.3		44	.66 NS
after surgery	Highest skin surface temperature	35.1	31.6–37.1	31.6–37.1		33-36.9	34.3	-2.44	.80 ^{NS}
– Day 0	Lowest skin surface temperature	33.0	29–35.8	(33.1–34.8)	32.4	31.2–36	(33.3–35.3)	01	.99 ^{NS}
(N = 57)	Median skin surface temperature	33.8	30.8-36.4	(55.1–54.6)	33.8	32.1-36.3	(33.3–33.3)	19	.84 ^{NS}
After	Body core temperature	36.7	36-37.8		36.4	35.9-37.0		-1.21	.22 ^{NS}
surgery –	Highest skin surface temperature	36.0	32.6–38	35.2	35.5	33.8-36.6	34.6	-2.15	.03*
Day 1	Lowest skin surface temperature	34.4	31.5-36.8		33.4	31.3-35.8	(33.7–35.2)	-1.84	.06 ^{NS}
(N = 60)	Median skin surface temperature	35.3	32.4-37.7	(34.6–35.7)	34.6	32.1-36.2	(33.7–33.2)	-2.07	.03*
After	Body core temperature	36.5	35–38		36.6	36–37.2		53	.59 ^{NS}
surgery –	Highest skin surface temperature	36.1	34.3–37.5	35.5	35.8	33.9–36.4	34.8	-2.01	.04*
Day 2	Lowest skin surface temperature	34.8	32.3–36.9	(34.8–36.1)	33.9	31.9–34.9	(34.1–35.4)	-2.60	.01*
(N = 60)	Median skin surface temperature	35.5	32.6–37.2	(34.0-30.1)	34.9	33.2–35.6	(34.1–33.4)	-2.84	.01*
After	Body core temperature	36.3	36–37.4		36.2	36–37.4		-1.40	.16 NS
surgery –	Highest skin surface temperature	36.0	33.2–38	35.7	35.8	34.3–37.7	35.5	79	.43 ^{NS}
Day 3	Lowest skin surface temperature	35.2	31–37.4	(34.9–36.5)	34.7	32.5–36.3	(35.1–35.9)	-1.16	.24 ^{NS}
(N = 58)	Median skin surface temperature	35.7	32.2–37.6	(34.7–30.3)	35.3	34–36.5	(33.1–33.7)	-1.03	$.30^{\mathrm{NS}}$
After	Body core temperature	36.5	36–38.9		36.4	36.1–37.8		11	.91 NS
surgery –	Highest skin surface temperature	36.1	33.9–38.6	35.6	35.9	33.8–37	35.6	58	.56 NS
Day 4	Lowest skin surface temperature	35.3	31.7–37.8	(34.8–36.6)	34.9	31.7–35.8	(34.8–36.1)	85	.39 ^{NS}
(n = 50)	Median skin surface temperature	35.6	32.9–38.3	(34.0-30.0)	35.6	33.5–36.4	(34.0-30.1)	60	.54 ^{NS}

p-value determined by Mann-Whitney U test

NS = not significant

^{*} denotes significance on Days 2 to 4 in surgical site surface temperature between non-infected and infected surgical wounds