

Movement variability emerges in gait as adaptation to task constraints in dynamic environments

CABALLERO, Carla, DAVIDS, Keith <<http://orcid.org/0000-0003-1398-6123>>, HELLER, Ben <<http://orcid.org/0000-0003-0805-8170>>, WHEAT, Jonathan <<http://orcid.org/0000-0002-1107-6452>> and MORENO, Francisco J

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/24104/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

CABALLERO, Carla, DAVIDS, Keith, HELLER, Ben, WHEAT, Jonathan and MORENO, Francisco J (2019). Movement variability emerges in gait as adaptation to task constraints in dynamic environments. *Gait and posture*, 70, 1-5.

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

Movement Variability emerges in Gait as Adaptation to Task Constraints in Dynamic Environments

Abstract

Background. Motor variability has been related to motor control playing a functional role in human adaptive behaviours. However, the direction of the relationship between variability and motor control can be unclear. The specific relations that exist between task constraints and movement (re)organization could explain some of this controversy.

Research question. This study sought to understand whether manipulation of task constraints result in changes in the magnitude or structure of motor system variability observed in a basic walking task. We also investigated the relationship between performance in achieving task goals and the structure of motor variability.

Methods. Twenty volunteers walked around a circular track with binary combinations of 3 task constraints, providing 8 conditions. The manipulated task constraints were: 1) track width; 2) surface stiffness; and 3), walking direction. Performance was analysed using standard deviation (SD) of sacral displacement and its mean velocity (MV). Fuzzy Entropy (FE) and Detrended Fluctuation Analysis (DFA) were used to assess the kinematic variability structure.

Results. Individuals showed lower SD and MV walking on the narrower track. These changes were also followed by higher DFA values, indicating a more auto-correlated structure of variability. The foam surface was also associated with an increase in amplitude, velocity and irregularity (FE) of movement.

Significance. Results of this study describe how specific task constraints, such as the width of the walking track and the surface stiffness, shape emergent movement coordination patterns as participants search for functional information from the environment to regulate performance behaviors. Changes in variability structure could reveal the search for adaptive strategies during walking. Smaller movement fluctuations and higher velocity in gait patterns are related to greater irregularity and lower autocorrelation in the kinematic variability structure, demonstrating that a specific relationship emerges between system variability and movement performance, which is driven by task constraints.

Keywords: Dynamic balance; Task constraints; Adaptation; Nonlinear analyses; Motor variability

1. Introduction

Motor variability is inherent within neurobiological systems, playing a functional role in adaptive behaviours of humans [1, 2], characterized by refinements in movement performance during interactions with environmental contexts [3]. Humans need to use motor variability to drive adaptive behaviours in changing environments [1]. This process can be observed during performance of numerous everyday motor tasks, such as locomotion, in which one constantly needs to adapt each step to a preceding one and every step is different.

For this reason, motor variability has been studied in different performance contexts to establish the movement (re)organization being used to achieve task goals [4, 5] or to adapt to the effects of sensorimotor impairments

[6, 7]. However, the direction of the relationship between motor variability and motor control can be unclear. Some research on variability has associated it with performance impairment, such as when gait variability is related to balance deficits during walking (e.g. [8, 9]). However, other studies have proposed alternative explanations for such observations. For instance, Rosenblatt, Hurt [10] studied the relationship between variability of foot placement and stability of gait patterns. Using uncontrolled manifold analysis of the joint configuration variance, they distinguished two types of variability, defined as “good” (which did not affect the mediolateral trajectory of the foot in the frontal plane) and “bad” (which affected this trajectory). Their results suggested that larger amounts of good variability could improve stability. It suggests that motor variability could offer a window into the control structures that underlie behaviour regulation [11]. Humans appear to use different regulatory strategies to compensate for inherent motor limitations by exploiting motor system variability for successful task performance [12]. In other words, motor variability has a functional role to drive adaptive behaviours in movement systems, allowing the central nervous system to exploit the high dimensionality offered by the abundance of motor system degrees of freedom (DoF) [1].

During movement regulation, the role of perception and action is to support the (re)organization of intentional behaviours during interactions with environmental and task constraints [13]. Such constraints may shape adaptations to the (re)organization of motor system DoFs by structuring the state space of possible system configurations available. Some investigators have suggested that these continuous adaptations are reflected in the structure of motor variability [14], with the relationship between variability and motor

control being influenced by task constraints. This hypothesis has been tested during performance in manual force control tasks [14] or standing balance tasks [2]. Evidence implies that specificity of task constraints may shape emergent motor system behaviours. Indeed, the specific relations that exist between task constraints and movement (re)organization could explain some of the controversy explained above regarding the relationship between variability and motor control [2].

To examine this issue, we sought to understand whether manipulation of task constraints would result in changes in the magnitude or structure of motor system variability observed in a walking task. We also investigated the relationship between performance in achieving task goals and the structure of motor variability. Gait variability provides insights on neuromotor performance regulation [6], and has been studied to assess effects of aging [8, 15] and different disabilities [6, 9]. Here, we analysed emergent movement adaptations under varying task constraints. Our prediction was that the structure of motor variability would depend on the specific task constraints to be satisfied. Specifically, we expected that changes in properties of a locomotion task would lead to a reduction in the number of available configurations in participant motor systems. These changes were expected to result in greater regularity in gait, and increased auto-correlations within the structure of movement variability.

2. Methods

2.1. Participants.

Twenty healthy volunteers (8 females, 12 males) participated in this study (age = 26.6 ± 5.5 years; stature = 1.7 ± 0.1 m; mass = 69.4 ± 13.9 Kg). Exclusion criteria included current musculoskeletal injuries or balance deficits that impaired participants from walking safely along a track designed in a laboratory.

Written informed consent was obtained from participants prior to testing. The experimental procedures used were in accordance with the Declaration of Helsinki and were approved by Sheffield Hallam University Research Ethics Committee.

2.2. Experimental Procedure and Data Collection

To assess postural dynamic balance, kinematic data from the pelvis were recorded at 120 Hz using an electromagnetic tracking system (Polhemus G4, Polhemus, USA). Pelvis displacement, in three axes (antero-posterior (AP), medio-lateral (ML) and vertical (V)), was recorded through an electromagnetic sensor firmly taped to the sacrum of each participant.

Participants walked around a circular track on the floor (Figure 1) with binary combinations of 3 different task constraints, providing a total of 8 different conditions (Table 1). They walked barefoot to avoid any possible differences related to footwear. Manipulated task constraints were: 1) width of the walking track (0.15 or 0.30 m); 2) surface stiffness (foam or vinyl); and 3), walking direction (clockwise or anticlockwise). These task constraints were manipulated because they significantly impact on gait regulation in different ways: 1) the

width of the walking track can constrain participant behaviour by modifying the need for precision of foot placement. For example, Young and Dingwell [16] showed how stability increased when participants had to walk with wider steps. In this study, we manipulated the width of the track to assess how the width of the steps, limited by space available, affects the involvement of motor system DoFs; 2) surface stiffness can play a role regulating the perception of sensorimotor information from the soles of the feet when walking [17]. **The thickness of the foam used in this study was 5 cm of 'firm' foam**; and 3), direction of walking could be related to the intrinsic dynamics (movement organisation tendencies) of each individual in adapting gait regulation to clockwise and anti-clockwise directions, for example. Some studies have shown that people tend to show a turning preference during locomotion [18, 19]. We expected to find different strategies according to walking direction, since one is likely to be more stable than the other. Participants were instructed to walk in their typical way on the path and adjust their steps to a metronomic rhythm of 130 bpm, provided acoustically, in order to standardise step velocity. The duration of each test trial was 70 s and the rest period between trials was 1 min. Every condition was experienced twice by participants in a randomized order.

Table 1 around here

Figure 1 around here

2.3. Data analysis and reduction

An application written in Labview 2009 (National Instruments, Texas), developed in our laboratory, was used for data analysis. Data were already filtered by the G4 Polhemus tracking system with a single-pole, low-pass filter with an adaptive pole location. The pre-set filtering parameters were: sensitivity = 0.02; boundary (FLow) = 0.02; boundary (FHigh) = 0.8; Max transition rate = 0.95. Kinematic time series data were then down-sampled, by interpolation, from 120 Hz to 20 Hz. The first and last 5 s of each trial were discarded to avoid non-stationarity related to trial initiation and termination [20]. Time series length was 1200 data points.

Postural sway, used to assess task performance, was determined using standard deviation (SD) values of sacral displacement and its mean velocity (MV). Functional performance behaviour was defined in this study by lower dispersion and velocity of each participant's displacement trajectory. According to this definition, the best trial performed in each condition was selected for statistical analysis.

Variables used to assess the variability structure in kinematic data were Fuzzy Entropy (FE) and Detrended Fluctuation Analysis (DFA). Fuzzy Entropy values indicate the degree of irregularity in a signal: higher FE values indicate greater irregularity in the signal time domain, whereas lower FE values indicate greater regularity. To calculate this measure, the following parameter values were used: vector length, $m = 2$; tolerance window, $r = 0.2 \cdot SD$; and gradient, $n=2$. These parameter values have been shown to have high consistency, which underlie their frequent use [21, 22]. Detrended Fluctuation Analysis evaluates the presence of long-term correlations within a time series using a

parameter known as the scaling index, α [23]. The α value identifies the extent to which proceeding data are dependent on previous outcomes [24]. Different values of α indicate the following: $\alpha > 0.5$ implies persistence; $\alpha < 0.5$ implies anti-persistence; and $\alpha = 0.5$ implies an uncorrelated signal [25]. In the current study, this measure was computed according to procedures recommended by Peng, Havlin [23]. The slope α was obtained from the window range $4 \leq n \leq N/10$ to maximize the long-range correlations and reduce errors incurred by estimating α [26].

Dependent variables were calculated over the resultant distance (RD) kinematic time series, instead of the AP, ML and V axes, in order to obtain a global variable. RD is the vector distance from the centre of the circular track negotiated by participants (where the Polhemus G4 source was placed) to the sensor placed on the sacrum.

$$RD \text{ time series} = \sqrt{(X[n] - \bar{X})^2 + (Y[n] - \bar{Y})^2 + (Z[n] - \bar{Z})^2} \quad n = 1, 2, \dots, N.$$

where n is the number of data points in the kinematic time series. X , Y and Z correspond to values of the kinematic time series for AP, ML and V axes, successively and, \bar{X} , \bar{Y} and \bar{Z} correspond to the means of the kinematic time series for the AP, ML and V axes, respectively.

2.4. Statistical Analysis

Normality of variable distribution was evaluated using the Kolmogorov-Smirnov test with the Lilliefors correction. A repeated measures ANOVA with three intra-individual variables, width of a walking track, surface path and walking direction, was used to assess effects of constraints manipulations on performance outcome measures and nonlinear variables. Alpha levels were set

at $p < 0.05$. Partial eta squared (η_p^2) was calculated as a measure of effect size and to record the proportion of the overall variance attributable to each factor. Values of effect size above 0.64 were considered strong, between 0.25 and 0.64 for moderate and below 0.25 small [27]. Finally, Pearson Product Moment Correlation coefficients were calculated to assess relationships between performance variables (Displacement SD and MV) and nonlinear measures of variability (FE and DFA).

3. Results

Average values obtained in every walking condition are displayed in Table 2.

Table 2 around here

None of the dependent variables showed significant differences between clock and anti-clockwise walking directions (Table 3). The most sensitive variables to this constraint were the SD ($F_{1,16} = 4.194$, $p = .057$, $\eta_p^2 = .208$) and FE ($F_{1,16} = 4.236$, $p = .056$, $\eta_p^2 = .209$). When walking clockwise, SD values tended to be higher, being significantly different when negotiating the wide track with a solid surface (Figure 2). Under this same condition, FE values tended to be lower than in the anti-clockwise condition.

On the wide walking track, both SD and MV values were significantly higher than on the narrow track. Pairwise comparisons showed that this task constraint affected SD values most when walking clockwise on the solid surface. In contrast, MV values were affected regardless of walking direction or surface properties (Figure 2). Fuzzy Entropy values did not show any significant

differences, whilst DFA values were significantly lower when participants walked on the wide track (Table 3). Specifically, DFA differences between widths were most evident when participants walked on the solid surface (Figure 2).

Finally, walking track surface seemed to influence both performance variables. Surface properties did not have a significant effect on SD values (Table 3), in the clockwise direction when walking on the wide track. However, SD values decreased when the walking surface was solid (Figure 2). There were significant differences in MV values, regardless of track direction and width (Figure 2). Higher values of MV emerged when walking on the foam surface (Table 3). Concerning the nonlinear performance measures, FE values were significantly higher in the foam conditions, although these differences were only displayed when participants walked in the anti-clock wise direction, on the wide track (Figure 2). Detrended Fluctuation Analysis measures did not show any statistically significant differences in any conditions (Table 3).

Table 3 around here

Figure 2 around here

When effects of all interacting constraints were analysed, Pearson Product Moment Correlation coefficients were computed in order to assess the relationship between gait performance (SD and MV variables) and the structure of variability (FE and DFA) (Table 4). Both performance variables were not significantly correlated with each other, meaning that they contributed different information about walking performance. Regarding the structure of movement variability, SD values were negatively correlated with FE and positively related to DFA. The MV measure showed the opposite relationship (positively related to

FE and negatively related to DFA). These relationships were not statistically significantly correlated under all interacting conditions.

Table 4 around here

4. Discussion

Previous studies have argued that re-organisation of motor system DoFs depends on interactions between system intrinsic dynamics (coordination tendencies of individuals) and performance task constraints [14]. These system adaptations seem to reflect increases and decreases in motor pattern variability. Previously, we have highlighted how participants modify their postural control dynamics according to different constraints, particularly task difficulty and availability of biofeedback [2]. Results showed that FE reduced and DFA increased as task difficulty level increased in the presence of biofeedback. However, when biofeedback was unavailable, the opposite trend in FE and DFA values was observed. Additionally, higher FE and lower DFA values were observed when biofeedback was available, rather than not. Regardless, performance was related to the structure of the motor variability. Few studies have addressed this relationship in dynamic balance tasks, such as during locomotion. For example, variability has been associated to balance deficits during walking [8] providing insights on neuromotor performance regulation [6]. Nevertheless, most studies of gait variability have focused only on magnitude, and not time-dependent structure [24].

Here, we sought to understand whether manipulation of task constraints would result in changes in the magnitude or structure of motor system variability observed in a basic walking task. We also investigated the relationship between

performance in achieving task goals and the structure of motor variability. Three different constraints were manipulated during the experiment: walking direction, track width and surface properties.

First of all, it must be pointed out that the metronomic rhythm used is a constraint is not discuss by the authors as it was constant along all the trials, standardizing the possible effect it could cause. Knowing that performance in preferred conditions can lead to more irregular and less auto-correlated behaviours [24], the authors chose 130 bpm as metronomic rhythm in order to keep a cadence fast enough to be far from the preferred cadence of any participant.

Regarding the manipulated constraints, although the majority of participants were right-limb dominant, they did not display significant changes in their variability during task performance regarding the walking direction. However, the trend seems to indicate that the fluctuations were bigger and more regular in clock-wise conditions. Recent studies have provided clear evidence for a leftward (anti-clockwise) turning preference in right-handers, while non-right-handers show a bias towards the opposite turning direction (clockwise) [19]. Since most of the participants were right-limb dominant, the trend we found would support the findings in the literature, performance in preferred conditions can lead to more irregular and less auto-correlated behaviours [24].

Participants displayed lower SD and MV values when walking on the narrower track. This observation indicates that, because of the increased stability challenge this width implies, less variable and slower movements emerged in the narrower track condition. The need to maintain the centre of

mass (COM) lateral displacement inside a narrower track may have constrained their movements, and the structure of movement variability, with more auto-correlated movement (increase in the DFA values) being observed. However, similar movement regularity (FE) was recorded on the narrow walking track. The greater amount of auto-correlation displayed on the narrow walkway could indicate a reduction in the number of adjustments performed by participants [28], made without stepping outside the track. These observations imply that this task constraint caused a reduction in the number of available solutions to achieve the task goal. This restriction on the available solutions is reflected in the decrease of MV and DFA.

The foam surface was also associated with an increase in the amplitude and velocity of movement adaptations made. It is apparent that SD values changed significantly only during performance in the clockwise direction and on the wide walking track, being higher on the foam surface than on the solid surface. However, MV values were significantly lower when participants walked on the solid surface, regardless of the track direction and width. A possible reason for the increase in the magnitude of variability (SD) and movement velocity is that participants may have used it as part of a strategy for acquiring useful information from the environment to achieve their task goal [1, 29].

This idea would be supported by the fact that all the observed statistically significant differences were greater when walking on the solid surface. It is possible that, when walking on the foam surface, participants increased their movement variability (evidenced by higher SD values) and their movement velocity (higher MV), to create more information during performance to achieve the task goal, regardless of the width of the track. This can be related to the

phenomenon referred as “sensory reweighting”. Depending on environmental conditions, the relative contribution of each sensory system changes to achieve an appropriate adaptation [30]. During gait, the plantar sole of the foot detects changing pressure patterns which provide important information about the disposition and movement of the body’s COM [17]. Walking on foam surfaces may mask this capacity, decreasing the information provided by the environment (i.e. the floor). In order to enhance the information to control ankle movements and perform the task, participants may have had to increase their variability (SD) when they were walking on the foam surface. According to Davids, Glazier [1], this increase in movement variability can enhance perception of information to support motor performance. This exploratory process varies substantially from individual to individual and from task to task. This implies that the foam surface forced participants to increase their movement variability (SD) and movement velocity in order to perceive rich information to regulate their movements during task performance.

With regard to the structure of movement variability, on the foam surface FE mean values were higher. These differences were more clearly displayed when walking in the anti-clockwise condition and on the wide walking track.

The correlational analysis found that larger movement fluctuations are related to the structure of kinematic variability, with less irregularity and greater auto-correlations. On the other hand, the higher velocity in gait movements was related to greater irregularity and lower autocorrelation in the kinematic variability structure. These findings are aligned with results of previous studies [14], and are harmonious with findings from our study on static balance [2]. Taken together, these studies reveal that a specific relationship emerges

between system variability structure and movement performance, which is dependent on task constraints and can be observed in different types of balance tasks: static and dynamic.

In conclusion, this study suggests that specific task constraints lead to changes in the magnitude and structure of motor variability observed in locomotion. Specifically, the direction of walking has no effect on variability. However, the width and the compliance of the track changed both characteristics of variability. A narrower track reduced SD and MV and increased auto-correlation, suggesting a reduction in the number of available solutions to achieve the task goal. Conversely, a foam surface caused an increase in SD and MV, with higher values of FE. These results could be related to an exploratory strategy, of increased variability in order to enhance information perception.

References

- [1] Davids K, Glazier P, Araujo D, Bartlett R. Movement systems as dynamical systems: the functional role of variability and its implications for sports medicine. *Sports Med.* 2003;33:245-60.
- [2] Caballero C, Barbado D, Davids K, Moreno FJ. Variations in task constraints shape emergent performance outcomes and complexity levels in balancing. *Exp Brain Res.* 2016;234:1611-22.
- [3] Krakauer JW, Mazzoni P. Human sensorimotor learning: adaptation, skill, and beyond. *Curr Opin Neurobiol.* 2011;21:636-44.
- [4] Bauer HU, Schöllhorn W. Self-Organizing Maps for the Analysis of Complex Movement Patterns. *Neural Processing Letters.* 1997;5:193-9.
- [5] Mullineaux DR, Uhl TL. Coordination-variability and kinematics of misses versus swishes of basketball free throws. *Journal of Sports Sciences.* 2010;28:1017-24.
- [6] Balasubramanian CK, Neptune RR, Kautz SA. Variability in spatiotemporal step characteristics and its relationship to walking performance post-stroke. *Gait Posture.* 2009;29:408-14.
- [7] Torres EB. Atypical signatures of motor variability found in an individual with ASD. *Neurocase.* 2012;19:150-65.
- [8] Hausdorff JM, Rios DA, Edelberg HK. Gait variability and fall risk in community-living older adults: a 1-year prospective study. *Arch Phys Med Rehabil.* 2001;82:1050-6.
- [9] Schniepp R, Wuehr M, Schlick C, Huth S, Pradhan C, Dieterich M, et al. Increased gait variability is associated with the history of falls in patients with cerebellar ataxia. *J Neurol.* 2014;261:213-23.

- [10] Rosenblatt NJ, Hurt CP, Latash ML, Grabiner MD. An apparent contradiction: increasing variability to achieve greater precision? *Exp Brain Res.* 2014;232:403-13.
- [11] Riley MA, Turvey MT. Variability of determinism in motor behavior. *J Mot Behav.* 2002;34:99-125.
- [12] Black DP, Smith BA, Wu J, Ulrich BD. Uncontrolled manifold analysis of segmental angle variability during walking: preadolescents with and without Down syndrome. *Exp Brain Res.* 2007;183:511-21.
- [13] Shafizadeh M, Wheat J, Davids K, Ansari NN, Ali A, Garmabi S. Constraints on perception of information from obstacles during foot clearance in people with chronic stroke. *Exp Brain Res.* 2017;235:1665-76.
- [14] Vaillancourt DE, Newell KM. Aging and the time and frequency structure of force output variability. *J Appl Physiol (1985).* 2003;94:903-12.
- [15] Callisaya ML, Blizzard L, Schmidt MD, McGinley JL, Srikanth VK. Ageing and gait variability--a population-based study of older people. *Age Ageing.* 2010;39:191-7.
- [16] Young PMM, Dingwell JB. Voluntary changes in step width and step length during human walking affect dynamic margins of stability. *Gait & posture.* 2012;36:219-24.
- [17] Waddington G, Adams R. Football boot insoles and sensitivity to extent of ankle inversion movement. *Br J Sports Med.* 2003;37:170-4; discussion 5.
- [18] Lenoir M, Van Overschelde S, De Rycke M, Musch E. Intrinsic and extrinsic factors of turning preferences in humans. *Neurosci Lett.* 2006;393:179-83.
- [19] Mohr C, Landis T, Bracha HS, Brugger P. Opposite turning behavior in right-handers and non-right-handers suggests a link between handedness and cerebral dopamine asymmetries. *Behav Neurosci.* 2003;117:1448-52.
- [20] van Dieen JH, Koppes LL, Twisk JW. Postural sway parameters in seated balancing; their reliability and relationship with balancing performance. *Gait Posture.* 2010;31:42-6.
- [21] Chen W, Wang Z, Xie H, Yu W. Characterization of surface EMG signal based on fuzzy entropy. *IEEE Trans Neural Syst Rehabil Eng.* 2007;15:266-72.
- [22] Lake DE, Richman JS, Griffin MP, Moorman JR. Sample entropy analysis of neonatal heart rate variability. *Am J Physiol Regul Integr Comp Physiol.* 2002;283:R789-97.
- [23] Peng CK, Havlin S, Stanley HE, Goldberger AL. Quantification of scaling exponents and crossover phenomena in nonstationary heartbeat time series. *Chaos.* 1995;5:82-7.
- [24] Jordan K, Newell KM. The structure of variability in human walking and running is speed-dependent. *Exerc Sport Sci Rev.* 2008;36:200-4.
- [25] Roerdink M, De Haart M, Daffertshofer A, Donker SF, Geurts AC, Beek PJ. Dynamical structure of center-of-pressure trajectories in patients recovering from stroke. *Exp Brain Res.* 2006;174:256-69.
- [26] Chen Z, Ivanov P, Hu K, Stanley HE. Effect of nonstationarities on detrended fluctuation analysis. *Phys Rev E Stat Nonlin Soft Matter Phys.* 2002;65:041107.
- [27] Ferguson CJ. An effect size primer: A guide for clinicians and researchers. *Professional Psychology: Research and Practice.* 2009;40:532-8.
- [28] Barbado D, Caballero C, Moreside JM, Vera-García FJ, Moreno FJ. Can be the structure of motor variability predict learning rate? *Journal of Experimental Psychology: Human Perception and Performance.* 2017;43:596-607.
- [29] Herzfeld DJ, Shadmehr R. Motor variability is not noise, but grist for the learning mill. *Nat Neurosci.* 2014;17:149-50.
- [30] Assländer L, Peterka RJ. Sensory reweighting dynamics in human postural control. *Journal of neurophysiology.* 2014;111:1852-64.