

Barriers and motivators of physical activity participation in middle-aged and older-adults – a systematic review

SPITERI, Karl, BROOM, David <<http://orcid.org/0000-0002-0305-937X>>, BEKHET, Amira Hassan, DE CARO, John Xerri, LAVENTURE, Bob and GRAFTON, Kate

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/24699/>

This document is the Accepted Version [AM]

Citation:

SPITERI, Karl, BROOM, David, BEKHET, Amira Hassan, DE CARO, John Xerri, LAVENTURE, Bob and GRAFTON, Kate (2019). Barriers and motivators of physical activity participation in middle-aged and older-adults – a systematic review. *Journal of Aging and Physical Activity*. [Article]

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>



Barriers and motivators of physical activity participation in middle-aged and older adults – a systematic review

Journal:	<i>Journal of Aging and Physical Activity</i>
Manuscript ID	JAPA.2018-0343.R3
Manuscript Type:	Scholarly Review
Focus Area:	barriers < psychosocial perspectives, motivators < psychosocial perspectives
Statistical Methods:	qualitative research analysis
Free-Form Keywords:	physical activity, exercise



1
2
3
4
5
6

Barriers and Motivators of Physical Activity Participation in Middle-aged and Older-
adults – A Systematic Review

For Peer Review

1 Physical inactivity is responsible for morbidity, premature mortality and substantial
2 economic burden in which the cost was estimated to be 0.64% of global health expenditure in
3 2013 (Ding et al., 2016). An appropriate level of physical activity (PA) can prevent chronic
4 diseases (Booth, Roberts, & Laye, 2012) but the proportion of older adults is increasing with
5 older adults tending to be less physically active than younger age groups (Sun, Norman, &
6 While, 2013). PA happens differently throughout the day during work, travel and leisure
7 time. Its nature may be structured such as during sport or strength training; or unstructured
8 such as whilst doing household chores. Determinants of PA differ depending on the type of
9 activity (Koeneman, Verheijden, Chinapaw, & Hopman-Rock, 2011).

10 Previous systematic reviews have identified the barriers and motivators towards
11 physical activity or exercise participation among adults (30-64 years), older adults (60+ years)
12 or the older old (80 years and above) (Allender, Cowburn, & Foster, 2006; Baert, Gorus,
13 Mets, Geerts, & Bautmans, 2011; Burton et al., 2017; Franco et al., 2015). The most recent
14 review by Burton et al. (2017) highlighted barriers and motivators specific to resistance
15 training. Most of the barriers and motivators identified were similar to PA in general, with
16 some being specific to resistance training. Another systematic review which examined
17 barriers and motivators and included all age groups was limited to studies carried out in the
18 United Kingdom (UK) (Allender et al., 2006). The other two reviews did not explore
19 differences between middle-aged and young-old, which the current review aims to achieve to
20 address the gap.

21 PA plays a different role in middle-age and young-old (Spirduso, Francis, & MacRae,
22 2005). Reviews were identified which examined health promotion interventions around
23 middle-age and young-old (Baxter et al., 2016; King, Rejeski, & Buchner, 1998). In order to
24 conduct optimal life-style campaigns it is important to identify the motivators and barriers
25 specific to the population being targeted (Baert, et al., 2011). There are broad influences upon

1 PA (interpersonal, environmental, individual) and these might vary across the life course
2 (Bauman et al., 2012).

3 This review will explore differences and similarities in barriers and motivators
4 between the ages of 50-64 years and 65-70 years, adopting an aged approach similar to the
5 one used by Baxter et al. (2016). Synthesizing both qualitative and quantitative research on
6 barriers and motivators can be used to better inform policy and research direction (Dixon-
7 Woods, Argawal, Jones, Young, & Sutton, 2005). The Theoretical Domain Framework
8 version-2 (TDF) was used as the guiding theoretical framework for this study. This
9 framework was developed in 2005 and validated in 2012 (Atkins et al., 2017). TDF was
10 developed from theories of behavior and behavior change for easier use by health
11 professionals. The TDF has fourteen domains namely 1) knowledge, 2) skills, 3)
12 social/professional role and identity, 4) beliefs about capabilities, 5) optimism, 6) beliefs
13 about consequences 7) reinforcement, 8) intentions, 9) goals, 10) memory attention and
14 decision process, 11) environmental context and resources, 12) social influences, 13) emotion
15 14) emotional and behavioral regulation. Each of these domains has different constructs for
16 example the emotional domain includes stress, fear and anxiety. This framework has been
17 used previously in systematic reviews to synthesize the influences of behaviors across
18 different theoretical domains (Atkins et al., 2017). For example Rushforth et al. (2016) have
19 used the TDF to identify barriers in the management of diabetes in primary care.

20 The aim of this review was to synthesize the literature on barriers and motivators of
21 physical activity in middle aged (50-64 years) and older adults (65-70 years). The review
22 objectives were to identify a) what are the barriers, encountered by middle aged and older
23 adults, to being physical active and b) what motivates middle aged and older adults to be
24 physically active? In this review the terms physical activity and exercise are used
25 interchangeably.

1 Middleton, Henderson, & Girling, 2016). Facilitators and enablers are often used
2 synonymously as they assist or make it easier for behavior to occur, and are a form of
3 extrinsic motivation. In this review the search was limited to the term 'motivation' as it would
4 identify both intrinsic and extrinsic motivation.

5 **Information Source and Search Strategy**

6 Five databases (Pubmed, Medline, Embase, Psycinfo, CINAHL Complete, and
7 SPORTDiscus) were searched for articles. No date limit was set in order to identify all
8 literature and possible differences across time. The literature search was carried out in July
9 2018. A search for grey literature was undertaken using the internet search engines Google
10 and Google Scholar. This was done to identify any other evidence which was not indexed in
11 bibliographic databases which would meet the aforementioned eligibility criteria. References
12 lists from eligible studies and previous systematic reviews were reviewed to identify any
13 additional studies which might have been relevant. The search strategy used in Pubmed is
14 outlined in table 1. The search was modified according to the language used in the database.

15 **Study Selection**

16 The PRISMA state for reporting systematic reviews was used to ensure proper
17 reporting of the entire process (Moher et al, 2009). Figure 1 presents the study selection
18 process. Two authors (KS and AH) scanned all titles and abstracts and excluded studies that
19 did not meet the criteria. After title screening the authors did not agree on forty-eight articles
20 included in the abstracts to be reviewed. During abstract review there was disagreement on
21 fourteen articles which were to be included in the full text review. When disagreement
22 occurred in the full text review, KS and AH discussed until a consensus was reached mediated
23 by a third member of the team. Included articles were then assessed for quality.

1 **Study Quality**

2 The Critical Appraisal Skill Program (CASP) (CASP UK, 2018) and modified CASP
3 version were identified as potential quality assessment tools, however these do not cater for
4 cross-sectional and mixed-methods studies. The Mixed Method Appraisal Tool (MMAT)
5 does effectively cater for different study types so it was used in this review. This tool has
6 criteria to assess qualitative, quantitative (including observational studies) and mixed-methods
7 whilst positioning all types of studies on the same level (Pluye, Gagnon, Griffiths, & Johnson-
8 Lafleur, 2009). Using the same tool to assess all the studies made it easier to assess outcomes.
9 MMAT has reportedly good inter-rater reliability and is quick to use (Pace et al., 2012). It is
10 has already been used in more than fifty published systematic reviews (Souto et al., 2015).
11 The methodological quality of studies was assessed using the approaches described by
12 Dixon-Woods et al. (2004; 2006). Dixon-Woods et al. (2004) quality assessment addresses
13 study design, study reporting and relevancy. Studies classified as ‘fatally flawed’, i.e. those
14 studies considered to be of poor design and reporting, and of low relevance, were excluded
15 from the review. Articles achieving two out of the four criteria in the MMAT were included
16 in the review. If the article had a low score but was determined to be of relevance because it
17 contributed to the synthesis this was included. The MMAT score of all the articles is available
18 in supplementary material 1.

19 **Data Collection Process**

20 A data extraction sheet was developed and piloted on ten articles (five qualitative and
21 five quantitative) to ensure it met the needs of the analysis. The sheet was amended
22 accordingly and the following data was collected from all studies: Code, title, reference,
23 country, population, aim, theoretical perspective (example phenomenology), mode of data
24 collection, sample size, age range, age mean/standard deviation, study type, behavioral theory
25 used, PA measured, data sources/tools, response rate, follow-up, confounding variables

1 considered, barriers (50-64 years), motivators (50-64 years), barriers (65-70 years),
2 motivators (65-70 years). Data extraction was carried out via paper and pencil and then
3 transferred to Microsoft Word © to decrease legibility errors and to enable discussion
4 between researchers (Elamin et al., 2009).

5 **Data Analysis**

6 Theory driven thematic framework analysis was used for the data analysis (Dixon-
7 Woods et al., 2005). The TDF was used as a guiding theory as it includes domains from a
8 variety of behavior change theories (BCT). The analysis presents the frequency in which
9 domains within the TDF were identified in the different literature. Results were presented in a
10 narrative format using quotes to validate the relevance of the identified domains.

11 **Results**

12 **Study Selection**

13 Figure 1 highlights the study selection process. There were a total of 9400 articles after
14 the removal of duplicates. Following a review of abstracts a total of 298 articles met the
15 inclusion criteria, and following a full text review this was reduced to 162. Sixty-nine
16 quantitative articles were excluded as fifty percent of the population did not include in the age
17 range. Eighteen qualitative articles were excluded as they did not differentiate between the
18 two age groups being investigated. KS reviewed all the articles using MMAT whilst AM and
19 JXDC reviewed a twenty-five percent sample of the total articles after a full text review to
20 ensure the quality of the critical appraisal process. There was eighty-six percent agreement
21 with AH and ninety-three percent agreement with JXDC. Discrepancies were dealt with by
22 discussion between the reviewers until agreement was reached. The critical appraisal score are
23 in supplementary material 1. From a total of fifty-five articles, six contained quantitative data

1 (see supplementary material 2), and forty-nine contained qualitative data (see supplementary
2 material 3).

3 **Study Participants**

4 The majority of studies reviewed were from the World Health Organisation (WHO)
5 Americas region: USA (thirty-three studies), and Canada (nine), Brazil (three), and Colombia
6 (one). Thirteen studies were from different WHO European region: U.K. (six), Italy (two),
7 Belgium (one), Ireland (two), and Netherland (two). Ten studies from WHO Western Pacific
8 region: Australia (eight), New Zealand (one), and China (one). One study from WHO Eastern
9 Mediterranean Region: Iran. The qualitative studies had a total sample size of 1970
10 participants, with focus groups and interviews being the most commonly used methods of
11 data collection. Twelve studies used multi-method data collection. Twenty-three studies
12 framed their theoretical perspective. The quantitative studies included data from 3524
13 participants. One study was longitudinal, the remaining were cross-sectional. The average
14 sample size was thirty participants in qualitative and 587 for quantitative studies (range 472-
15 840). Questionnaires were the most common tool used in quantitative studies. The
16 questionnaire response rate ranged from thirty-nine to ninety-seven percent. The average
17 sample size for qualitative studies was thirty participants (range 7-78). Interviews and focus
18 groups were the most common tools used.

19 **Behavioral Theories**

20 Six quantitative studies and twenty-five qualitative studies adopted a BCT as the basis
21 for the investigation. Fourteen different theories were used in all studies. BCT presented in
22 table 2.

23 **Qualitative Data**

24 The results for the 50-64 year old group are presented in table 3; the results of the 65-
25 70 year old group are presented in table 4. The percentage studies identifying barriers and

1 motivators within each TDF domain are presented in Figures 2 and 3 respectively.
2 ‘Environmental context and resources’ were the most coded barriers in both groups, followed
3 by social influences. ‘Beliefs about capabilities’ had a higher percentage of studies coded
4 within the domain for the 65-70 year age group which identified as barrier then social
5 influences. ‘Social influences’ was the most coded domain as motivator for both groups. This
6 was followed by ‘goals’ for the 50-64 year old group, and reinforcement for 65-70 year old.
7 No studies were coded under the ‘Memory, attention and decision processes’ domain. The
8 extracted data is available in supplementary material 3.

9 **Quantitative Data**

10 Six studies met the inclusion criteria and achieved a 50% MMAT score. There were
11 four studies that assessed barriers to PA quantitatively. Gobbi et al., (2012) was the only study
12 with a higher proportion of participants in the 64-70 year age group. The study assessed
13 barriers using the ‘Barrier to PA Practice Questionnaire’ (Hirayama, 2006 as cited in Gobbi et
14 al., 2012). Analysis was carried out using age groups 60-69, 70-79 and more than 80 years of
15 age. Within the study a statistical difference was found in the mean number of barriers
16 reported between the 60-69 and 70-79 group.

17 Arras et al. (2006) examined health promoting behaviors among a population between
18 45-90 years. Barriers were measured using the ‘Barriers to Health Promoting Activities for
19 Disable People’ (Becker, Stuijferge, & Sands, 1991). Correlation between barriers and age
20 was carried out but no relationship was identified. Ayotte et al. (2010) carried out
21 meditational analysis to explore how PA is influenced using social cognitive theory. One of
22 the factors analyzed was perceived barriers using the ‘Perceived Barrier to Exercise Scale’
23 (Salmon, Owen, Crawford, Bauman, & Sallis, 2003). Within the model no correlation
24 between barriers and age was identified. In the final model barriers had an indirect
25 relationship with PA. Kowal & Fortier (2007) used a longitudinal study to investigate how

1 barriers and environmental characteristics influence PA behavior change. The authors used a
2 sample with a mean age of fifty-one years (SD 6.7). Fourteen barriers, identified from
3 previous literature, were assessed. Using a five year interval they examined differences in PA
4 barriers with age. The authors identified that woman at the age of sixty years reported low
5 level of physical barriers then other ages. Being too busy was reported more in the 50-54 year
6 olds compared to sixty plus years. Health problems were most common in women older than
7 sixty years.

8 Two studies which met the inclusion criteria investigated motivators (Dacey, Baltzell,
9 & Zaichkowsky, 2008; Kolt, Driver, & Giles, 2004). The studies included both populations
10 under study, but the percentage population could not be calculated. Kolt et al. (2004) had an
11 average age of 63.6 years (SD 7.8). The study used the 'Participation Motivation
12 Questionnaire for Older Adults' (Kirkby, Kolt, Habel, & Adams, 1999). It looked at reasons
13 for participation in PA in older Australian exercisers. The study examined different types of
14 motivators: social, fitness, recognition, challenges/benefits, medical, and involvement.
15 Statistical analysis was undertaken to check for differences between 55-64 year olds and 65-
16 74 year olds. Social reasons were rated higher in the middle-aged. There was a statistical
17 difference in involvement and medical factors. No other statistical differences were reported
18 with age group.

19 Dacey et al (2008) used the 'Exercise Motivation Inventory -2' (Markland &
20 Ingledew, 1997), aimed to examine differences in intrinsic and extrinsic motivation between
21 people with three different levels of exercise participation (Dacey et al., 2008). This study
22 looked at six types of motivators: health and fitness, social/emotional benefits, weight
23 management, stress management, enjoyment, and appearance. Statistical analysis was carried
24 out for age groups 50-59 years old, 60-69 year old and 70-79 year old. Using MANOVA a

1 significant score was found between 50-59 year old and 60-69 year old within the appearance
2 and stress management scores being higher in 50-59.

3 **Discussion**

4 The aim of this review was to synthesize the literature on barriers and motivators of
5 PA in middle aged (50-64 years) and older adults (65-70 years). With the barriers identified,
6 there were similarities with age groups but clear differences in the motivators were found.
7 The analysis of qualitative literature identified factors that acted as both a barrier and a
8 motivator. For example, previous experience of being physically active was identified as a
9 motivator, whilst the lack of it was identified as a barrier. Similar barriers and motivators
10 were extracted for both age groups from the qualitative studies.

11 However, in the quantitative literature, differences between the age groups were not
12 consistent which is partly explained by the use of different tools. This lack of statistical
13 difference might also be explained by the similar barriers identified between the two groups in
14 the qualitative literature. In analyzing the result of the qualitative studies, using the TDF, data
15 did not fit within certain domains such as 'Memory, attention and decision processes'. From
16 the analysis it is not possible to conclude whether these are not important or because of a lack
17 of research.

18 When compiling the results from the qualitative studies, differences in barriers
19 between the two age groups which were partly supported in the quantitative studies were
20 found. The 65-70 year old age groups identified these barriers which were not present in the
21 other group. In one study (Melillo et al., 2001) this group identified fears about safety to carry
22 out PA. Health problems were identified as a barrier more frequently in this group. This was
23 also identified in one quantitative study (Kowal & Fortier 2007). Lack of guidance from
24 healthcare professionals to start PA was another factor present only in this group. Being

1 labeled too old to carry out PA was a factor only identified within the young-old group. A
2 person's perception of self is influence by their social relations with others (Evans & Crust,
3 2015). The young-old mentioned fear of falling as barrier which was only present in this
4 group. In the middle-aged group work conditions were identified as barrier which was not
5 present in the other group. This is expected as people usually retire by the age of sixty-five
6 years (Ekerdt, 2010). Barriers with time management were identified in studies looking at
7 middle-aged.

8 Issues of being busy were reported more often in middle-aged also which was
9 highlighted in the qualitative study. Barriers within behavior regulation such as issues of
10 laziness were more prevalent in the younger group. Laziness was identified as idleness to
11 engage in PA and remaining sedentary. In a study (Belon, Nieuwendyk, Vallianatos, &
12 Nykiforuk, 2016) participants in this age group identified the difficulty in breaking the cycle
13 of using the car as a barrier to engaging in PA.

14 Differences were identified with respect to motivators between the two groups. A
15 motivator to be active identified only in the middle aged was the issue of being a role model
16 for their children. Another motivator mentioned by this age group only was fear of becoming
17 ill. Both groups identified health reasons as a motivator for engaging in PA, however, middle-
18 aged identified set targets of weight management or stress management. The young-old
19 mentioned staying health or staying independent as health motivator. Lifelong activity was a
20 motivator for the older group. Social motivators were identified by both groups, but being part
21 of a group and PA as an opportunity to socialize was mentioned more often in the young-old.
22 The young-old identified retirement as a motivator to become active. PA provided young-old
23 with a purpose in life, and another study identified PA as an opportunity to be busy. These
24 motivators where not identified in middle-aged.

1 **Comparison with previous reviews**

2 The findings of the current review are comparable to Schutzer & Graves (2004) in the
3 oldest old age group in that barriers and motivators were similar. Few behavior regulation
4 techniques were identified in the current review since it did not include interventional studies.
5 Schutzer & Graves (2004) had identified prompt such as the telephone interventions and the
6 use of music. Compared to reviews by Baert et al. (2011) and Burton et al. (2017) this review
7 used the TDF to analyze the findings whilst the other reviews used the socio-ecological
8 approach. This framework makes it possible to use results from the review in interventional
9 studies using different behavior change theory, as it incorporates concepts from different
10 theories. If an interventional study is set to target knowledge or goals these can be identified
11 through the results of this review. The list of barriers and motivators identified were similar to
12 those of the two previous reviews, however it was able to differentiate between the middle-
13 aged and young-old. The findings support previous findings from Allender et al. (2006).
14 Whilst their review was limited to UK studies this included findings from four WHO regions
15 and thirteen countries. This shows that factors might be transferrable across different
16 populations.

17 **The use of behavior change theory**

18 Nearly half of the reviewed studies, (forty-six percent), had a theoretical underpinning.
19 The socio-ecological theory was the most commonly used, probably because of the simplicity
20 of using this theory. Using a theoretical framework added context to what is being
21 investigated and to what type of influences are being identified (Atkins et al., 2017). Within
22 the analysis of this review certain TDF domains were not identified. It was not possible to
23 identify whether this was because the domain has not been researched or lack of importance
24 in the context.

1 **Strengths and Limitations**

2 The strengths of this review were the inclusion of both qualitative and quantitative
3 studies to get a broader perspective of barriers and motivators. There is a paucity of data on
4 the age groups included, so this review has extended the knowledge of what is an under
5 studied topic and population with, previous studies focusing on either middle-age adults (>60)
6 or the oldest old (80+). However, this review is not without its limitations.

7 The major limitation of this review is publication bias and the use of English language
8 studies only. The use of only one reviewer in the critical appraisal phase might increase error
9 (Buscemi, Hartling, Vandermeer, Tjosvold, & Klassen, 2006). This risk was reduced by
10 having two reviewers assessing an appropriate enough number of articles to ensure rigorous
11 standards were met. The systematic review used two reviewers, in the screening phase to
12 ensure that no studies were excluded in the review (Edwards et al., 2002). The use of one
13 quality appraisal tool (MMAT) might lack specificity since it is a generic tool which can be
14 used for all study. By using this tool the articles were assessed using the same measure,
15 including mixed-methods studies (Pace et al., 2012).

16 In the majority of studies both PA and exercise were used interchangeably. This might
17 have limited specificity of certain barriers or motivators. As Burton et al., (2017) suggested,
18 some barriers and motivators might be common across all forms of PA types but some might
19 be specific as in the case of strength training. As identified in previous literature the term
20 motivator was used in different context by different authors (Plonczynski, 2000). At times, the
21 term facilitator and enablers were used interchangeably in the studies. Given that these
22 definitions refer to a continuum making it clear at the beginning of the study to define what is
23 meant by motivation and what aspects are being examined would make comparison of studies
24 easier and specific.

1 **Implications of the findings**

2 This review identifies that barriers and motivators encountered by middle-aged and
3 young-old are similar. The review has achieved its aim of identifying barriers that prevent
4 people from being physically active. It has also identified motivators which health
5 professionals and PA promoters should have knowledge of to encourage behavior change in
6 these age groups. The identified barriers were similar, but there were differences in the
7 motivators between the two age groups which those developing and implementing
8 interventions need to be aware of.

9 The effects of the aging body were identified by studies as early as middle-aged
10 (George, Kolt, Rosenkranz, & Guagliano, 2014). This review identified environmental
11 barriers such as cost, weather, and facilities as barriers to undertaking PA for both age groups.
12 The social context is important, especially the family role, which can act either as a motivator
13 or barrier. Given the variety of barriers and motivators identified within the studies, health
14 professionals and physical activity promoters need to be aware of the broad characteristics
15 which might influence the uptake of PA or motivate its uptake.

16 This review highlights the complexity of PA influences with multiple factors being
17 identified from different domains of the TDF. Even though the review was limited to
18 community dwelling individuals and not specific patient groups, health problems were
19 identified as barriers for PA by both groups, especially young-old. Professionals need to be
20 aware of these limitations and teach individuals to be active within the constraints of their
21 health limitation. Any form of community intervention within these population age groups
22 needs to consider health problems and past injuries. The findings support the life course
23 approach to PA, as past master and previous experience of PA was identified as a motivator to
24 being physically active.

1 **Future Research Recommendations**

2 The issue of laziness mentioned in studies of middle-aged requires further exploration
3 in order to understand its meaning. Studying the link between different identified factors
4 would be of interest and would highlight the most influential factor. Meditational analysis
5 would help in identifying these factors. There is a need for further quantitative studies which
6 compare the importance of barriers and motivators at different age groups. Longitudinal
7 studies are also needed to assess how and whether there are any changes across an
8 individual's life span. Awareness of how motivators change with age can assist in keeping the
9 older adults active.

10 **Conclusion**

11 Kowal & Fortier (2007) highlighted it is important to identify age-specific PA barriers
12 as this can assist targeted interventions. This study has synthesized and extends the body of
13 knowledge on PA barriers and motivators in older adults by discussing the differences
14 between the age groups of 50-64 years and 65-70 years. For the 50-64 year group,
15 environmental and resources, social influence and difficulty in regulating behavior were the
16 key barriers. The key motivator was set goals, such as fitness, stress relief or taking dog out.
17 For the 65-70 years group the key motivators were different forms of reinforcement such as
18 peer encouragement and having fun, the social aspect of PA and support from health
19 professionals. The key barriers were lack of belief in capabilities, environmental and social
20 such as family roles or lack of support from family. It is therefore apparent that differences in
21 barriers and motivators between the two groups were minimal, but differences are present.

22 The majority of the key barriers and motivators were identified through qualitative
23 literature. It was not possible to identify the importance of these factors for each group
24 because of the lack of quantitative data over an extended period of time to ascertain if barriers

- 1 Ball, K., Salmon, J., Giles-Corti, B., & Crawford, D. (2006). How can socio-economic
2 differences in physical activity among women be explained? A qualitative study. *Women
3 & Health, 43*(1), 93–113. Retrieved from
4 <http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=106246058&site=ehost-live>
5
- 6 Bauman, A. E., Reis, R. S., Sallis, J. F., Wells, J. C., Loos, R. J. F., Martin, B. W., ... Group,
7 L. P. A. S. W. (2012). Correlates of physical activity: why are some people physically
8 active and others not?. *Lancet (London, England), 380*(9838), 258–271.
9 [https://doi.org/10.1016/S0140-6736\(12\)60735-1](https://doi.org/10.1016/S0140-6736(12)60735-1)
- 10 Baxter, S., Blank, L., Johnson, M., Everson-Hock, E., Woods, H. B., Goyder, E., ...
11 Mountain, G. (2016). Interventions to promote or maintain physical activity during and
12 after the transition to retirement: an evidence synthesis. *Public Health Research, 4*(4), 1–
13 138. <https://doi.org/10.3310/phr04040>
- 14 Becker, H., Stuijferge, A. K., & Sands, D. (1991). Development of a Scale to Measure
15 Barriers to Health Promotion Activities among Persons with Disabilities. *American
16 Journal of Health Promotion, 5*(6), 449–454.
- 17 Belon, A. P., Nieuwendyk, L. M., Vallianatos, H., & Nykiforuk, C. I. J. (2016). Community
18 Lenses Revealing the Role of Sociocultural Environment on Physical Activity. *American
19 Journal Of Health Promotion: AJHP, 30*(3), e92–e100.
20 <https://doi.org/http://dx.doi.org/10.4278/ajhp.140428-QUAL-169>
- 21 Booth, F., Roberts, C. K., & Laye, M. J. L. (2012). Lack of exercise is a major cause of
22 chronic diseases. *Comprehensive Physiology, 2*(2), 1143–1211.
23 <https://doi.org/10.1002/cphy.c110025.Lack>
- 24 Bopp, M., Lattimore, D., Wilcox, S., Laken, M., McClorin, L., Swinton, R., ... Bryant, D.
25 (2007). Understanding physical activity participation in members of an African

- 1 American church: A qualitative study. *Health Education Research*, 22(6), 815–826.
2 <https://doi.org/http://dx.doi.org/10.1093/her/cyl149>
- 3 Bruner, B., & Chad, K. (2013). Physical activity attitudes, beliefs, and practices among
4 women in a woodland cree community. *Journal of Physical Activity and Health*, 10(8),
5 1119–1127. Retrieved from
6 [http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed15&NEWS=N&](http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed15&NEWS=N&AN=1370395253)
7 [AN=1370395253](http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed15&NEWS=N&AN=1370395253)
- 8 Buman, M. P., Daphna Yasova, L., & Giacobbi, P. R. (2010). Descriptive and narrative
9 reports of barriers and motivators to physical activity in sedentary older adults.
10 *Psychology of Sport & Exercise*, 11(3), 223–230. Retrieved from
11 <http://search.ebscohost.com/login.aspx?direct=true&db=sph&AN=48604672&site=ehost>
12 -live
- 13 Burton, E., Farrier, K., Lewin, G., Pettigrew, S., Hill, A., Airey, P., ... Hill, K. D. (2017).
14 Motivators and Barriers for Older People Participating in Resistance Training: A
15 Systematic Review. *Journal of Aging and Physical Activity*, 25, 311–324.
16 <https://doi.org/10.1123/japa.2015-0289>
- 17 Buscemi, N., Hartling, L., Vandermeer, B., Tjosvold, L., & Klassen, T. P. (2006). Single data
18 extraction generated more errors than double data extraction in systematic reviews.
19 *Journal of Clinical Epidemiology*, 59(7), 697–703.
20 <https://doi.org/10.1016/j.jclinepi.2005.11.010>
- 21 Caperchione, C. M. C. M., Vandelanotte, C., Kolt, G. S. G. S., Duncan, M., Ellison, M.,
22 George, E., & Mummery, W. K. (2012). What a Man Wants: Understanding the
23 Challenges and Motivations to Physical Activity Participation and Healthy Eating in
24 Middle-Aged Australian Men. *American Journal of Men's Health*, 6(6), 453–461.
25 <https://doi.org/10.1177/1557988312444718>

- 1 Casey, M., Eime, R., Ball, K., & Payne, W. (2011). Characteristics of physically active and
2 inactive men from low socioeconomic communities and housing estates: a qualitative
3 study using the socioecological model. *Annals of Leisure Research, 14*(1), 1–21.
4 Retrieved from
5 <http://search.ebscohost.com/login.aspx?direct=true&db=sph&AN=98529850&site=ehost>
6 -live
- 7 CASP UK. (2018). CASP UK. Retrieved July 21, 2018, from <https://casp-uk.net/>
- 8 Cassou, A. C. N., Fermino, R., Añez, C. R. R., Santos, M. S., Domingues, M. R., & Reis, R.
9 S. (2011). Barriers to Physical Activity Among Brazilian Elderly Women From Different
10 Socioeconomic Status: A Focus-Group Study. *Journal of Physical Activity & Health,*
11 *8*(1), 126–132. Retrieved from
12 <http://search.ebscohost.com/login.aspx?direct=true&db=sph&AN=58103426&site=ehost>
13 -live
- 14 Ceria-Ulep, C. D., Serafica, R. C., & Tse, A. (2011). Filipino Older Adults' Beliefs About
15 Exercise Activity. *Nursing Forum, 46*(4), 240–250. <https://doi.org/10.1111/j.1744->
16 [6198.2011.00238.x](https://doi.org/10.1111/j.1744-6198.2011.00238.x)
- 17 Chatfield, S. L. (2015). Characteristics of lifelong physically active older adults. *Qualitative*
18 *Health Research, 25*(7), 966–973.
19 <https://doi.org/http://dx.doi.org/10.1177/1049732314554095>
- 20 Chaudhury, H., Mahmood, A., Michael, Y. L., Campo, M., & Hay, K. (2012). The influence
21 of neighborhood residential density, physical and social environments on older adults'
22 physical activity: An exploratory study in two metropolitan areas. *Journal of Aging*
23 *Studies, 26*(1), 35. Retrieved from
24 [http://lcproxy.shu.ac.uk/login?url=http://search.proquest.com/docview/910970995?accou](http://lcproxy.shu.ac.uk/login?url=http://search.proquest.com/docview/910970995?accountid=13827)
25 [ntid=13827](http://lcproxy.shu.ac.uk/login?url=http://search.proquest.com/docview/910970995?accountid=13827)

- 1 Cheng, H. G., & Phillips, M. R. (2014). Secondary analysis of existing data: opportunities and
2 implementation. *Shanghai Archives of Psychiatry*, 26(6), 371–375.
3 <https://doi.org/10.11919/j.issn.1002-0829.214171>
- 4 Chrisman, M., Nothwehr, F., Yang, G., & Oleson, J. (2015). Environmental influences on
5 physical activity in rural Midwestern adults: a qualitative approach. *Health Promotion
6 Practice*, 16(1), 142–148. <https://doi.org/http://dx.doi.org/10.1177/1524839914524958>
- 7 Cousins, S. O. (2003). A self-referent thinking model: how older adults may talk themselves
8 out of being physically active. *Health Promotion Practice*, 4(4), 439–448. Retrieved
9 from
10 <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed9&NEWS=N&A>
11 [N=137569678](http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed9&NEWS=N&AN=137569678)
- 12 Dacey, M., Baltzell, A., & Zaichkowsky, L. (2008). Older adults' intrinsic and extrinsic
13 motivation toward physical activity. *American Journal of Health Behavior*, 32(6), 570–
14 582. Retrieved from
15 <http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=105565930&site=eh>
16 [ost-live](http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=105565930&site=ehost-live)
- 17 Das, B. M., Sartore-Baldwin, M., & Mahar, M. . (2016). The Invisible Employee: University
18 Housekeeping Employees' Perceptions of Physical Activity. *Journal of Physical Activity
19 & Health*, 13(9), 952–956. <https://doi.org/http://dx.doi.org/10.1123/jpah.2015-0509>
- 20 Ding, D., Lawson, K. D., Kolbe-Alexander, T. L., Finkelstein, E. A., Katzmarzyk, P. T., van
21 Mechelen, W., & Pratt, M. (2016). The economic burden of physical inactivity: a global
22 analysis of major non-communicable diseases. *Lancet (London, England)*, 388(10051),
23 1311–1324. [https://doi.org/10.1016/S0140-6736\(16\)30383-X](https://doi.org/10.1016/S0140-6736(16)30383-X)
- 24 Dixon-Woods, M., Argawal, S., Jones, D., Young, B., & Sutton, A. (2005). Synthesising
25 qualitative and quantitative evidence: A review of possible methods. *Journal of Health*

- 1 *Services Research & Policy*, 10(1), 45–53. <https://doi.org/10.1258/1355819052801804>
- 2 Dixon-Woods, M., Cavers, D., Agarwal, S., Annandale, E. ., Antony, A., Harvey, J., ...
- 3 Sutton, A. J. (2006). Conducting a critical interpretive synthesis of the literature on
- 4 access to healthcare by vulnerable groups. *BMC Medical Research Methodology*, 6, 1–
- 5 13. <https://doi.org/10.1186/1471-2288-6-35>
- 6 Dixon-Woods, M., Shaw, R. L., Agarwal, S., & Smith, J. A. (2004). The problem of
- 7 appraising qualitative research. *Quality and Safety in Health Care*, 13(3), 223–225.
- 8 <https://doi.org/10.1136/qshc.2003.008714>
- 9 Edwards, P., Clarke, M., DiGuseppi, C., Pratap, S., Roberts, I., & Wentz, R. (2002).
- 10 Identification of randomized controlled trials in systematic reviews: Accuracy and
- 11 reliability of screening records. *Statistics in Medicine*, 21(11), 1635–1640.
- 12 <https://doi.org/10.1002/sim.1190>
- 13 Ekerdt, D. J. (2010). Frontiers of research on work and retirement. *The Journals of*
- 14 *Gerontology. Series B, Psychological Sciences and Social Sciences*, 65B(1), 69–80.
- 15 <https://doi.org/10.1093/geronb/gbp109>.
- 16 Elamin, M. B., Flynn, D. N., Bassler, D., Briel, M., Alonso-Coello, P., Karanicolas, P. J., ...
- 17 Montori, V. M. (2009). Choice of data extraction tools for systematic reviews depends
- 18 on resources and review complexity. *Journal of Clinical Epidemiology*, 62(5), 506–510.
- 19 <https://doi.org/10.1016/j.jclinepi.2008.10.016>
- 20 Evans, A. B., & Crust, L. (2015). ‘Some of these people aren’t as fit as us ...’: experiencing
- 21 the ageing, physically active body in cardiac rehabilitation. *Qualitative Research in*
- 22 *Sport, Exercise and Health*, 7(1), 13–36. <https://doi.org/10.1080/2159676X.2014.908945>
- 23 Evans, L. K. (2011). Rural Black women’s thoughts about exercise. *Applied Nursing*
- 24 *Research*, 24(4), 200–206. <https://doi.org/10.1016/j.apnr.2009.09.005>
- 25 Franco, M. R., Tong, A., Howard, K., Sherrington, C., Ferreira, P. H., Pinto, R. Z., & Ferreira,

- 1 M. L. (2015). Older people's perspectives on participation in physical activity: a
2 systematic review and thematic synthesis of qualitative literature. *British Journal of*
3 *Sports Medicine*, 49(19), 1268–1276. [https://doi.org/http://dx.doi.org/10.1136/bjsports-](https://doi.org/http://dx.doi.org/10.1136/bjsports-2014-094015)
4 2014-094015
- 5 Gallagher, N. A., Gretebeck, K. A., Robinson, J. C., Torres, E. R., Murphy, S. L., & Martyn,
6 K. K. (2010). Neighborhood factors relevant for walking in older, Urban, African
7 American adults. *Journal of Aging and Physical Activity*, 18(1), 99–115.
8 <https://doi.org/10.1123/japa.18.1.99>
- 9 George, E. S., Kolt, G. S., Rosenkranz, R. R., & Guagliano, J. M. (2014). Physical activity
10 and sedentary time: male perceptions in a university work environment. *American*
11 *Journal of Men's Health*, 8(2), 148–158. <https://doi.org/10.1177/1557988313497217>
- 12 Gobbi, S., Sebastiao, E., Papini, C. B., Nakamura, P. M., Valdanha Netto, A., & Gobbi, L. T.
13 B. (2012). Physical inactivity and related barriers: A study in a community dwelling of
14 older brazilians. *Journal of Aging Research*, 2012, 685190.
15 <https://doi.org/http://dx.doi.org/10.1155/2012/685190>
- 16 Gonzales, A., & Keller, C. (2004). Mi familia viene primero (My family comes first):
17 physical activity issues in older Mexican American women. *Southern Online Journal of*
18 *Nursing Research*, 5(4), 21p–21p. Retrieved from
19 [http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=106611319&site=eh](http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=106611319&site=ehost-live)
20 ost-live
- 21 Gray, P. M., Murphy, M. H., Gallagher, A. M., Simpson, E. E., P.M., G., M.H., M., ...
22 Simpson, E. E. (2016). Motives and Barriers to Physical Activity Among Older Adults
23 of Different Socioeconomic Status. *Journal Of Aging And Physical Activity*, 24(3), 419–
24 429. <https://doi.org/10.1123/japa.2015-0045>
- 25 Guell, C., Panter, J., Griffin, S., & Ogilvie, D. (2018). Towards co-designing active ageing

- 1 strategies: A qualitative study to develop a meaningful physical activity typology for
2 later life. *Health Expectations: An International Journal of Public Participation in*
3 *Health Care & Health Policy*. <https://doi.org/http://dx.doi.org/10.1111/hex.12686>
- 4 Guell, C., Shefer, G., Griffin, S., & Ogilvie, D. (2016). "Keeping your body and mind active":
5 An ethnographic study of aspirations for healthy ageing. *BMJ Open*, *6*(1), e009973.
6 <https://doi.org/http://dx.doi.org/10.1136/bmjopen-2015-009973>
- 7 Harley, A. E., Rice, J., Walker, R., Strath, S. J., & Quintiliani, L. M. (2014). Physically
8 Active, Low-Income African American Women: An Exploration of Activity
9 Maintenance in the Context of Sociodemographic Factors Associated with Inactivity.
10 *Women and Health*, *54*(4), 354–372.
11 <https://doi.org/http://dx.doi.org/10.1080/03630242.2014.896440>
- 12 Henwood, T., Tuckett, A., Edelstein, O., & Barlet, H. (2011). Exercise in later life: the older
13 adults' perspective about resistance training. *Ageing & Society*, *31*(8), 1330–1349.
14 Retrieved from
15 <http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=104473093&site=ehost-live>
16
- 17 Jancey, J. M., Clarke, A., Howat, P., Maycock, B., & Lee, A. H. (2009). Perceptions of
18 physical activity by older adults: A qualitative study. *The Health Education Journal*,
19 *68*(3), 196. Retrieved from
20 <http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=105327656&site=ehost-live>
21
- 22 Kalavar, J. M., Kolt, G. S., Giles, L. C., & Driver, R. P. (2004). Physical activity in older
23 Asian Indians living in the United States: barriers and motives. *Activities, Adaptation &*
24 *Aging*, *29*(1), 47–62. https://doi.org/http://dx.doi.org/10.1300/J016v29n01_04
- 25 Kamphuis, C. B. M., van Lenthe, F. J., Giskes, K., Brug, J., & Mackenbach, J. P. (2007).

- 1 Perceived environmental determinants of physical activity and fruit and vegetable
2 consumption among high and low socioeconomic groups in the Netherlands. *Health &*
3 *Place*, 13(2), 493–503.
4 <https://doi.org/https://dx.doi.org/10.1016/j.healthplace.2006.05.008>
- 5 Keegan, R., Middleton, G., Henderson, H., & Girling, M. (2016). Auditing the socio-
6 environmental determinants of motivation towards physical activity or sedentariness in
7 work-aged adults: a qualitative study. *BMC Public Health*, 16, 438.
8 <https://doi.org/10.1186/s12889-016-3098-6>
- 9 Kegler, M. C., Alcantara, I., Dubruiel, N., Veluswamy, J. K., Appelbaum, H., Handwerk, S.,
10 ... H., A. (2013). “Positive deviants”: a qualitative study of physically active adults in
11 rural environments. *The Journal Of Primary Prevention*, 34(1–2), 5–15.
12 <https://doi.org/10.1007/s10935-013-0291-6>
- 13 King, A. C., Rejeski, W. J., & Buchner, D. M. (1998). Physical Activity Interventions
14 Targeting Older Adults: A Critical Review and Recommendations. *American Journal of*
15 *Preventive Medicine*, 15(4), 316–333.
- 16 Kirkby, R. J., Kolt, G. S., Habel, K., & Adams, J. (1999). Exercise in older women: Motives
17 for participation. *Australian Psychologist*, 34(2), 122–127.
18 <https://doi.org/http://dx.doi.org/10.1080/00050069908257440>
- 19 Koeneman, M. A., Verheijden, M. W., Chinapaw, M. J. M., & Hopman-Rock, M. (2011).
20 Determinants of physical activity and exercise in healthy older adults: A systematic
21 review. *International Journal of Behavioral Nutrition and Physical Activity*, 8, 142.
22 <https://doi.org/http://dx.doi.org/10.1186/1479-5868-8-142>
- 23 Kolt, G. S., Driver, R. P., & Giles, L. C. (2004). Why older Australians participate in exercise
24 and sport. *Journal Of Aging And Physical Activity*, 12(2), 185–198. Retrieved from
25 <http://search.ebscohost.com/login.aspx?direct=true&db=cmedm&AN=15223886&site=e>

- 1 host-live
- 2 Kolt, G. S., Paterson, J. E., & Cheung, V. Y. M. (2006). Barriers to physical activity
3 participation in older Tongan adults living in New Zealand. *Australasian Journal on*
4 *Ageing*, 25(3), 119–125. [https://doi.org/http://dx.doi.org/10.1111/j.1741-](https://doi.org/http://dx.doi.org/10.1111/j.1741-6612.2006.00157.x)
5 6612.2006.00157.x
- 6 Kosteli, M. C., & Williams, S. E. (2016). Investigating the psychosocial determinants of
7 physical activity in older adults: A qualitative approach. *Psychology & Health*, 31(6),
8 730–749. <https://doi.org/http://dx.doi.org/10.1080/08870446.2016.1143943>
- 9 Kosteli, M. C., Williams, S. E., & Cumming, J. (2016). Investigating the psychosocial
10 determinants of physical activity in older adults: A qualitative approach. *Psychology &*
11 *Health*, 31(6), 730–749.
12 <https://doi.org/http://dx.doi.org/10.1080/08870446.2016.1143943>
- 13 Kowal, J., & Fortier, M. S. (2007). Physical activity behavior change in middle-aged and
14 older women: the role of barriers and of environmental characteristics. *Journal of*
15 *Behavioral Medicine*, 30(3), 233–242. Retrieved from
16 [http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=105828469&site=eh](http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=105828469&site=ehost-live)
17 ost-live
- 18 Leavy, B., & Aberg, A. (2010). “Not ready to throw in the towel”: perceptions of physical
19 activity held by older adults in Stockholm and Dublin. *Journal of Aging & Physical*
20 *Activity*, 18(2), 219–236. Retrieved from
21 [http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=105152356&site=eh](http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=105152356&site=ehost-live)
22 ost-live
- 23 Markland, D., & Ingledew, D. K. (1997). The measurement of exercise motives: Factorial
24 validity and invariance across gender of a revised Exercise Motivations Inventory.
25 *British Journal of Health Psychology*, 2(4), 361–376.

- 1 <https://doi.org/http://dx.doi.org/10.1111/j.2044-8287.1997.tb00549.x>
- 2 Mathews, E., Lakshmi, J. K., Ravindran, T. K. S., Pratt, M., & Thankappan, K. R. (2016).
3 Perceptions of barriers and facilitators in physical activity participation among women in
4 Thiruvananthapuram City, India. *Global Health Promotion, 23*(4), 27–36.
5 <https://doi.org/10.1177/1757975915573878>
- 6 McArthur, D., Dumas, A., Woodend, K., Beach, S., & Stacey, D. (2014). Factors influencing
7 adherence to regular exercise in middle-aged women: a qualitative study to inform
8 clinical practice. *BMC Women's Health, 14*(1), 49. [https://doi.org/10.1186/1472-6874-](https://doi.org/10.1186/1472-6874-14-49)
9 [14-49](https://doi.org/10.1186/1472-6874-14-49)
- 10 Melillo, K. D., Williamson, E., Houde, S. C., Futrell, M., Read, C. Y., & Campasano, M.
11 (2001). Perceptions of older Latino adults regarding physical fitness, physical activity,
12 and exercise. *Journal of Gerontological Nursing, 27*(9), 38–46. Retrieved from
13 [http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=107076816&site=eh](http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=107076816&site=ehost-live)
14 [ost-live](http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=107076816&site=ehost-live)
- 15 Miller, W., & Brown, P. R. (2017). Motivators, Facilitators, and Barriers to Physical Activity
16 in Older Adults: A Qualitative Study. *Holistic Nursing Practice, 31*(4), 216–224.
17 <https://doi.org/10.1097/HNP.0000000000000218>
- 18 Mitra, R., Siva, H., & Kehler, M. (2015). Walk-friendly suburbs for older adults? Exploring
19 the enablers and barriers to walking in a large suburban municipality in Canada. *Journal*
20 *of Aging Studies, 35*, 10–19.
21 <https://doi.org/http://dx.doi.org/10.1016/j.jaging.2015.07.002>
- 22 Mohamed, A. A., Hassan, A. M., Weis, J. A., Sia, I. G., Wieland, M. L., A.A., M., ...
23 Wieland, M. L. (2014). Physical Activity Among Somali Men in Minnesota: Barriers,
24 Facilitators, and Recommendations. *American Journal of Men's Health, 8*(1), 35–44.
25 <https://doi.org/http://dx.doi.org/10.1177/1557988313489132>

- 1 Mosquera, J., Parra, D. C., Gomez, L. F., Sarmiento, O., Schmid, T., & Jacoby, E. (2012). An
2 Inside Look at Active Transportation in Bogotá: A Qualitative Study. *Journal of*
3 *Physical Activity & Health*, 9(6), 776–785. Retrieved from
4 <http://search.ebscohost.com/login.aspx?direct=true&db=sph&AN=79553209&site=ehost>
5 -live
- 6 Nightingale, A. (2009). A guide to systematic literature reviews. *Surgery*, 27(9), 381–384.
7 <https://doi.org/10.1016/j.mpsur.2009.07.005>
- 8 Pace, R., Pluye, P., Bartlett, G., Macaulay, A. C., Salsberg, J., Jagosh, J., & Seller, R. (2012).
9 Testing the reliability and efficiency of the pilot Mixed Methods Appraisal Tool
10 (MMAT) for systematic mixed studies review. *International Journal of Nursing Studies*,
11 49(1), 47–53. <https://doi.org/10.1016/j.ijnurstu.2011.07.002>
- 12 Paluck, E. C., Allerdings, M., Kealy, K., & Dorgan, H. (2006). Health promotion needs of
13 women living in rural areas: an exploratory study. *Canadian Journal Of Rural Medicine:*
14 *The Official Journal Of The Society Of Rural Physicians Of Canada*, 11(2), 111–116.
15 Retrieved from
16 <http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=106341660&site=ehost-live>
17 ost-live
- 18 Plonczynski, D. J. (2000). Measurement of motivation for exercise. *Health Education*
19 *Research*, 15(6), 695–705. Retrieved from
20 <http://search.ebscohost.com/login.aspx?direct=true&db=cmedm&AN=11142077&site=ehost-live>
21 host-live
- 22 Pluye, P., Gagnon, M. P., Griffiths, F., & Johnson-Lafleur, J. (2009). A scoring system for
23 appraising mixed methods research, and concomitantly appraising qualitative,
24 quantitative and mixed methods primary studies in Mixed Studies Reviews.
25 *International Journal of Nursing Studies*, 46(4), 529–546.

- 1 <https://doi.org/10.1016/j.ijnurstu.2009.01.009>
- 2 Rathanaswami, K., Bengoechea, E. G., & Bush, P. L. (2016). Physical Activity in First
3 Generation South Asian Women Living in Canada: Barriers and Facilitators to
4 Participation. *Women in Sport & Physical Activity Journal*, 24(2), 110–119. Retrieved
5 from
6 <http://search.ebscohost.com/login.aspx?direct=true&db=sph&AN=120421030&site=eho>
7 [st-live](http://search.ebscohost.com/login.aspx?direct=true&db=sph&AN=120421030&site=ehost-live)
- 8 Romeike, K., Abidi, L., Lechner, L., de Vries, H., Oenema, A., K., R., ... H., de V. (2016).
9 Similarities and differences in underlying beliefs of socio-cognitive factors related to diet
10 and physical activity in lower-educated Dutch, Turkish, and Moroccan adults in the
11 Netherlands: a focus group study. *BMC Public Health*, 16(1), 813.
12 <https://doi.org/10.1186/s12889-016-3480-4>
- 13 Rosenkranz, R. R., Kolt, G. S., & Berentson-shaw, J. (2013). A review of enablers and
14 barriers to physical activity participation among older people of New Zealand and
15 international populations. *International SportMed Journal*, 14(4), 294–312.
- 16 Royce, S. W., Sharpe, P. A., Ainsworth, B. E., Greaney, M. L., Neff, L. J., & Henderson, K.
17 A. (2003). Conceptualising barriers and supports for physical activity: a qualitative
18 assessment. *International Journal of Health Promotion & Education*, 41(2), 49–56.
19 Retrieved from
20 <http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=106703912&site=eh>
21 [ost-live](http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=106703912&site=ehost-live)
- 22 Rushforth, B., McCrorie, C., Glidewell, L., Midgley, E., & Foy, R. (2016). Barriers to
23 effective management of type 2 diabetes in primary care: Qualitative systematic review.
24 *British Journal of General Practice*, 66(643), e114–e127.
25 <https://doi.org/10.3399/bjgp16X683509>

- 1 Salmon, J., Owen, N., Crawford, D., Bauman, A., & Sallis, J. F. (2003). Physical activity and
2 sedentary behavior: a population-based study of barriers, enjoyment, and preference.
3 *Health Psychology, 22*(2), 178–188. Retrieved from
4 <http://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=106838627&site=ehost-live>
5
- 6 Schutzer, K. A., & Graves, B. S. (2004). Barriers and motivations to exercise in older adults.
7 *Preventive Medicine, 39*(5), 1056–1061.
8 <https://doi.org/http://dx.doi.org/10.1016/j.ypmed.2004.04.003>
- 9 Sebastião, E., Chodzko-Zajko, W., Schwingel, A., Sebastiao, E., Chodzko-Zajko, W., &
10 Schwingel, A. (2015). An in-depth examination of perceptions of physical activity in
11 regularly active and insufficiently active older African American women: A participatory
12 approach. *PLoS ONE, 10*(11), 1–15. <https://doi.org/10.1371/journal.pone.0142703>
- 13 Sebastião, E., Ibe-Lamberts, K., Bobitt, J., Schwingel, A., & Chodzko-Zajko, W. (2014).
14 Employing a Participatory Research Approach to Explore Physical Activity among Older
15 African American Women. *Journal of Aging Research, 1*–8. <https://doi.org/2014/941019>
- 16 Souto, R. Q., Khanassov, V., Hong, Q. N., Bush, P. L., Vedel, I., & Pluye, P. (2015).
17 Systematic mixed studies reviews: Updating results on the reliability and efficiency of
18 the mixed methods appraisal tool. *International Journal of Nursing Studies, 52*(1), 500–
19 501. <https://doi.org/10.1016/j.ijnurstu.2014.08.010>
- 20 Spirduso, W. W., Francis, K., & MacRae, P. G. (2005). *Physical Dimensions of Aging*.
21 (HumanKinetics, Ed.) (2nd ed.). Leeds.
- 22 Steltenpohl, C. N., Shuster, M., Peist, E., Pham, A., & Mikels, J. A. (2018). Me Time, or We
23 Time? Age Differences in Motivation for Exercise. *The Gerontologist*.
24 <https://doi.org/10.1093/geront/gny038>
- 25 Sun, F., Norman, I. J., & While, A. E. (2013). Physical activity in older people: A systematic

- 1 review. *BMC Public Health*, 13(1). <https://doi.org/10.1186/1471-2458-13-449>
- 2 Van Dyck, D., Mertens, L., Cardon, G., De Cocker, K., & De Bourdeaudhuij, I. (2017).
3 Opinions Toward Physical Activity, Sedentary Behavior, and Interventions to Stimulate
4 Active Living During Early Retirement: A Qualitative Study in Recently Retired Adults.
5 *J Aging Phys Act*, 25(2), 277–286. <https://doi.org/10.1123/japa.2015-0295>
- 6 Wertman, A., Wister, A. V, Mitchell, B. A., A., W., A.V., W., Wertman, A., ... Mitchell, B.
7 A. (2016). On and off the mat: Yoga experiences of middle-aged and older adults.
8 *Canadian Journal on Aging = La Revue Canadienne Du Vieillissement*, 35(2), 190–205.
9 <https://doi.org/http://dx.doi.org/10.1017/S0714980816000155>

10




Figure 1. Study selection flowchart (PRISMA, 2015)

215x279mm (300 x 300 DPI)




Figure 2. Percentage of studies identifying barriers by TDF domain

297x210mm (300 x 300 DPI)




Figure 3. Percentage of studies identifying motivators by TDF domain

297x210mm (300 x 300 DPI)

Table 1

Search Strategy (using Pubmed terminology)

1	barrier* ti,ab.
2	motivat* ti,ab.
3	1 or 2
4	adult (MESH)
5	adult* ti,ab.
6	aged (MESH)
7	Older adult ti,ab.
8	4 or 5 or 6 or 7
9	exercise (MESH)
10	sport (MESH)
11	motor activity (MESH)
12	physical activity ti,ab.
13	exercise ti,ab.
14	sport* ti,ab.
15	9 or 10 or 11 or 12 or 13 or 14
16	3 and 8 and 15

Note: ti is title, ab is abstract

Table 2
Behavior change theories used in studies

Behavior Change Theory	Number of Studies	References
ANGELO framework	One	Belon, Nieuwendyk, Vallianatos, & Nykiforuk, 2016,
Health belief model	Two	Das, Sartore-Baldwin, & Mahar, 2016; Wertman et al., 2016
Integrated behaviour change	One	Arnautovska, O'Callaghan, & Hamilton, 2017
Kleinman's explanatory models	One	Evans, 2011
Pender's health promotion theory	One	Arras, Ogletree, & Welshimer, 2006
Positive deviance theory	Two	Harley, Rice, Walker, Strath, & Quintiliani, 2014; Kegler et al., 2013
Self-determination theory	Two	Dacey, Baltzell, & Zaichkowsky, 2008; Gray et al., 2016
Self-efficacy theory	Two	Kalavar, Kolt, Giles, & Driver, 2004; Kolt, Paterson, & Cheung, 2006
Self-regulating theories	One	Romeike et al., 2016
Socio-cognitive theory	Five	Ayotte, Margrett, & Hicks-Patrick, 2010; Chrisman, Nothwehr, Yang, & Oleson, 2015; Cousins, 2003; Gothe & Kendall, 2016; Kosteli & Williams, 2016
Socio-ecological theory	Thirteen	Ball, Salmon, Giles-Corti, & Crawford, 2006; Brittain, Baillargeon, McElroy, Aaron, & Gyurcsik, 2006; Casey, Eime, Ball, & Payne, 2011; Dave et al., 2015; Gobbi et al., 2012; Gonzales & Keller, 2004; Kaiser & Baumann, 2010; Kegler et al., 2013; Kowal & Fortier, 2007; Mathews, Lakshmi, Ravindran, Pratt, & Thankappan, 2016; Royce et al., 2003; Sebastião et al., 2015; Sebastião, Ibe-Lamberts, Bobitt, Schwingel, & Chodzko-Zajko, 2014
Socio-emotional selectivity theory	One	Steltenpohl, Shuster, Peist, Pham, & Mikels, 2018
Success model	One	Harley et al., 2014
Theory of planned behaviour	Two	Kamphuis, van Lenthe, Giskes, Brug, & Mackenbach, 2007; Romeike et al., 2016

Table 3
Barriers and motivators for 50-64 years old

TDF domain	Number of Studies	Barriers 50-64 years	Number of Studies	Motivators 50-64 years
1. Knowledge	Two	<ul style="list-style-type: none"> • Participants were not aware of the benefits of engage in regular PA and the need to be active (Das, Sartore-Baldwin, & Mahar, 2016). • PA at work was identified as enough to get the necessary benefits (Das et al., 2016). • Participants in the study expressed the perception that PA is not congruent with a busy lifestyle (Royce et al., 2003). 	One	<ul style="list-style-type: none"> • Knowledge of the need to participate in PA “You know it’s like what they say if you don’t use your muscles, you lose it” (Gray et al., 2016, p. 422).
2. Skills	Two	<ul style="list-style-type: none"> • Participants identified themselves as lacking the necessary skill to carry out exercise (Das et al., 2016; Mosquera et al., 2012). “I need direction based as far as this particular exercise you need to do x times a day to get these results. I need a plan” (Das et al., 2016, p. 954). 	Two	<ul style="list-style-type: none"> • Previous experience gave the necessary confidence for participants to continue practicing PA. A military background (Kegler et al., 2013) or because of participation since younger (Rathanaswami, Bengoechea, & Bush, 2016) “I guess it’s always been in, been embedded within me to, fitness...early on I just grew up with the desire to be physically fit and to stay active...” (Kegler et al., 2013, p. 11).
3. Social/Professional role and identity	Nil	<ul style="list-style-type: none"> • Nil 	Nil	<ul style="list-style-type: none"> • Nil
4. Beliefs about	Five	<ul style="list-style-type: none"> • Past injuries and medical conditions 	Nil	<ul style="list-style-type: none"> • Nil

capabilities		<p>were barriers which influenced the participants perceptions about their capability to carry out PA (Casey, Eime, Ball, & Payne, 2011; Evans, 2011; Van Dyck, Mertens, Cardon, De Cocker, & De Bourdeaudhuij, 2017). “While swimming is good for my back, the last few years I can’t do that. The pain has gone down my legs. I can’t even do swimming . . . It hurts so much because my life is not the same anymore. I can’t do physical activity” (Casey et al., 2011, p. 8).</p> <ul style="list-style-type: none"> • The aging body hindered the participants from participating in PA (George, Kolt, Rosenkranz, & Guagliano, 2014). • A history of sedentary lifestyle was seen to influence their capabilities to carry out PA (Harley, Rice, Walker, Strath, & Quintiliani, 2014). 		
5. Optimism	Nil	<ul style="list-style-type: none"> • Nil 	Nil	<ul style="list-style-type: none"> • Nil
6. Beliefs about Consequences	Six	<ul style="list-style-type: none"> • Pain during exercise or fear of pain after exercise (Das et al., 2016; Gonzales & Keller, 2004; Mathews, Lakshmi, Ravindran, Pratt, & Thankappan, 2016). • Fear of injury (Buman, Daphna Yasova, & Giacobbi, 2010). • Poor past exercise experience, either as lack of performance or labelling acted as a barrier toward PA (McArthur, Dumas, Woodend, Beach, 	Nine	<ul style="list-style-type: none"> • For health benefits (physical and mental) and disease prevention (Bopp et al., 2007; Caperchione et al., 2012; Chatfield, 2015; George et al., 2014; Gray et al., 2016; Harley et al., 2014; McArthur et al., 2014; Wertman et al., 2016). • Feeling good, energetic “Now that I have started walking and getting exercise, it makes me have more ambition so I just can’t sit. So actually,

		<ul style="list-style-type: none"> • & Stacey, 2014). • Exercise seen as a physical strain and therefore it is avoided (Harley et al., 2014). 		<p>what exercise is doing is making me be more energetic” (Kegler et al., 2013, p. 11).</p>
7. Reinforcement	Seven	<ul style="list-style-type: none"> • Due to a bad experience with the health system participants did not participate in PA, as health professionals did not support them (Casey et al., 2011). • Conditions at work was a barrier in one study (Ball, Salmon, Giles-Corti, & Crawford, 2006). • Lacking a partner or a group with whom encourage participation (Evans, 2011; Paluck, Allerdings, Kealy, & Dorgan, 2006; Royce et al., 2003; Van Dyck et al., 2017). • Attending classes which lacked structure was not stimulating (McArthur et al., 2014). 	Four	<ul style="list-style-type: none"> • Fun and enjoyment of participating in PA (McArthur et al., 2014). • Positive feeling when being active (McArthur et al., 2014) • Gratification “My doctor was very happy with me . . . She was like you’ve lost 8 pounds since the last time you were here . . . When I hear my doctor say that it just makes me want to keep going” (Harley et al., 2014, p. 362). • Encouragement from others (Keegan, Middleton, Henderson, & Girling, 2016; Steltenpohl, Shuster, Peist, Pham, & Mikels, 2018).
8. Intentions	Two	<ul style="list-style-type: none"> • PA identified not a priority but an inconvenience (Mohamed et al., 2014; Royce et al., 2003). “... people cannot keep it up. They won’t have time, are looking for food, or a job and pay check” (Mohamed et al., 2014, p. 39). 	Three	<ul style="list-style-type: none"> • PA was intentionally “me time” (Keegan et al., 2016, p. 8). • Faith was the motivation to participate in PA “It’s a known fact in order for you to have faith you have to have a healthy mind and to exercise your mind is exercising your faith” (Bopp et al., 2007, p. 818). • Being a role model for their children (Chrisman, Nothwehr, Yang, & Oleson, 2015).
9. Goals	Six	<ul style="list-style-type: none"> • No motivation to carry out PA (Ball et 	Ten	<ul style="list-style-type: none"> • Taking their dog out as their goal for

		al., 2006; Evans, 2011; George et al., 2014; Mathews et al., 2016; Romeike et al., 2016; Van Dyck et al., 2017).		undertaking PA (Belon, Nieuwendyk, Vallianatos, & Nykiforuk, 2016; Kamphuis, van Lenthe, Giskes, Brug, & Mackenbach, 2007; Kegler et al., 2013).
				<ul style="list-style-type: none"> • Health goals, either stress management, keep health, weight loss or fitness as their goal for undertaking PA (Caperchione et al., 2012; Chatfield, 2015; George et al., 2014; Harley et al., 2014; Kegler et al., 2013). "I think it's pretty ridiculous to get old and just turn into a little old man..." (Chatfield, 2015, p. 969). • Will power (Bopp et al., 2007). • Doing something different (McArthur et al., 2014). • A sense of competition (Mohamed et al., 2014).
10. Memory, attention and decision processes	Nil	<ul style="list-style-type: none"> • Nil 	Nil	<ul style="list-style-type: none"> • Nil
11. Environmental context and resources	Twenty-one	<ul style="list-style-type: none"> • Cost and affordability of carrying out PA was mentioned as a barrier (Buman et al., 2010; Casey et al., 2011; Harley et al., 2014; Mohamed et al., 2014; Van Dyck et al., 2017; Wertman et al., 2016). • Environmental barriers such as lack of safety, poor footpaths, aesthetics (Belon et al., 2016; Bruner & Chad, 2013; Kamphuis et al., 2007; Mosquera et al., 2012; Royce et al., 2003). 	Four	<ul style="list-style-type: none"> • Safe environment (Gray et al., 2016) • Appealing environment example walking paths, possibility to change scenery (George et al., 2014; Keegan et al., 2016; Mathews et al., 2016) • Favourable Weather (Gray et al., 2016).

- Weather extremes (Bruner & Chad, 2013; Chatfield, 2015; Evans, 2011; Kamphuis et al., 2007).
- Lack of facilities for particular groups (Casey et al., 2011; McArthur et al., 2014; Paluck et al., 2006; Royce et al., 2003; Van Dyck et al., 2017).
- Problems with accessibility due to distance (Mohamed et al., 2014; Paluck et al., 2006; Royce et al., 2003).
- Not enough time due to other obligations mostly work and family (Chatfield, 2015; Das et al., 2016; Evans, 2011; George et al., 2014; Mathews et al., 2016; Romeike et al., 2016; Van Dyck et al., 2017; Wertman et al., 2016).
- Temporal illness (Chatfield, 2015).

12. Social influences

Ten

- Culture was identified as barrier as it was not part of the culture (Casey et al., 2011; Mohamed et al., 2014; Mosquera et al., 2012; Royce et al., 2003). “If you give an old (Somali man) a pair of shorts and Somali women see him running, Somali women would say he is crazy.” (Mohamed et al., 2014, p. 39).
- Family and friends influence was a barrier (Evans, 2011; George et al., 2014; Gonzales & Keller, 2004).
- The family role was in competition with PA (Caperchione et al., 2012;

Eleven

- Role models (Keegan et al., 2016; Mohamed et al., 2014).
- Social aspect of PA (Belon et al., 2016; George et al., 2014; Harley et al., 2014; Mathews et al., 2016; Steltenpohl et al., 2018; Wertman et al., 2016). “The classes to me, motivated me . . . it was a group activity and I could be with people, and I could talk. . . that way I get a little social connection” (Harley et al., 2014, p. 362).
- Family and friends (George et al., 2014; Gonzales & Keller, 2004; Kegler

		Gonzales & Keller, 2004; Rathanaswami et al., 2016). “Some women take maybe a year to come out and try to do something for [themselves] because they always put their families, their husbands or whoever at home first before they attend to their needs to improve themselves . . .” (Rathanaswami et al., 2016, p. 113).		et al., 2013; Steltenpohl et al., 2018).
13. Emotion	Seven	<ul style="list-style-type: none"> • A negative past experience with exercise was a barrier towards PA (Buman et al., 2010; Casey et al., 2011; Mosquera et al., 2012; Royce et al., 2003). • Feelings of fatigue (Gonzales & Keller, 2004; McArthur et al., 2014; Royce et al., 2003). • Not in the mood (McArthur et al., 2014). • Embarrassment to exercise as a barrier (Mohamed et al., 2014). 	Three	<ul style="list-style-type: none"> • PA provided them with a sense of vitality good feeling (Bopp et al., 2007; Kegler et al., 2013). • Fear of becoming ill therefore they are active (Caperchione et al., 2012).
14. Behavioural regulation	Nine	<ul style="list-style-type: none"> • Break the habitual cycle of using the car (Belon et al., 2016). • Lacking time management skills either because of a lot of thing to do (Ball et al., 2006) or too much free time after retirement (Van Dyck et al., 2017). • Laziness prevented participants from getting into the routine of engaging in PA (Alizadeh & Salehi, 2015; Caperchione et al., 2012; Evans, 2011; George et al., 2014; Gonzales & 	Two	<ul style="list-style-type: none"> • Health professionals and a military background were the motivators for participating in PA “My oncologist tells me to exercise and stays on my butt about my weight” (Kegler et al., 2013, p. 10). • Guilty of not being active was the other behavioural regulation in this age group (George et al., 2014).

Keller, 2004). “I’ve actually seen a big difference in me from when I was 45 to where I am now, that I’ve actually become more lazy . . .in terms of doing physical activity” (George et al., 2014, p. 153). “Everyone knows that exercise is good for health but laziness don’t allow me to exercise..” (Alizadeh & Salehi, 2015, p. 293).

- Self-sacrifice to accommodate others (McArthur et al., 2014).

Note: TDF is Theoretical Domain Framework

Table 4

Barriers and motivators for 65-70 years old

TDF domain	Number of Studies	Barriers 65-70 years	Number of Studies	Motivators 65-70 years
1. Knowledge	Three	<ul style="list-style-type: none"> • Queries or fear about safety to carry out PA (Melillo et al., 2001). • Lacked knowledge on exercise (Gray et al., 2016; Kolt, Paterson, & Cheung, 2006). “We have to be educated by professional people before go ahead with our exercise” (Kolt et al., 2006, p. 121). • Not aware of the benefits of PA (Kolt et al., 2006). 	Two	<ul style="list-style-type: none"> • Knowledge of the benefits of PA (Mitra, Siva, & Kehler, 2015). • Media was considered a source of knowledge which motivated participate to engage in PA (Arnautovska, O’Callaghan, & Hamilton, 2017).
2. Skills	Three	<ul style="list-style-type: none"> • Lack of skill (Gray et al., 2016; Kolt et al., 2006). “I think it depends on the person you know, depends what they can do” (Gray et al., 2016, p. 424). • No past experience of exercise (Kalavar, Kolt, Giles, & Driver, 2004). 	Two	<ul style="list-style-type: none"> • Past mastery of activities (Cousins, 2003). • A life history of PA participation (Guell, Panter, Griffin, & Ogilvie, 2018).
3. Social/Professional role and identity	Nil	<ul style="list-style-type: none"> • Nil 	One	<ul style="list-style-type: none"> • Participants undertook PA for pride (Arnautovska et al., 2017).
4. Beliefs about capabilities	Twelve	<ul style="list-style-type: none"> • Health problems were barriers to PA. These limited their capability to participate PA example breathlessness and pain (Cassou et al., 2011; Guell et al., 2018; Guell, Shefer, Griffin, & Ogilvie, 2016; Jancey, Clarke, Howat, Maycock, & Lee, 2009; Kalavar et al., 2004; Kolt et al., 2006; Leavy & Aberg, 	Two	<ul style="list-style-type: none"> • Past experience of PA was a motivator which helped in adapting to new forms of PA as one got limited by aging changes (Kosteli & Williams, 2016). • Lifelong activities made them believe in their capabilities (Guell et al., 2016).

		2010; Miller & Brown, 2017; Mitra et al., 2015; Sebastião, Ibe-Lamberts, Bobitt, Schwingel, & Chodzko-Zajko, 2014; Sebastião et al., 2015). “. . .Oh need a new knee. I need a new knee replacement” (Sebastião et al., 2014, p. 8)		
		<ul style="list-style-type: none"> • Aging was an identified barrier (Arnautovska et al., 2017; Ceria-Ulep, Serafica, & Tse, 2011; Kalavar et al., 2004). • Poor body image of capabilities (Jancey et al., 2009) 		
5. Optimism	One	<ul style="list-style-type: none"> • Lack of a positive outlook toward participant in PA. “There’s no point in starting something if you don’t know if you can continue it. I have never been a candidate for anything that is too active” (Cousins, 2003a, p. 442). 	Nil	<ul style="list-style-type: none"> • Nil
6. Beliefs about Consequences	Three	<ul style="list-style-type: none"> • Participants’ believed that exercise could cause them pain or risk them injury (Arnautovska et al., 2017; Kalavar et al., 2004). “You have to do everything in moderation. I cannot begin exercise by lifting weights now suddenly. I will cause more problems than I already have” (Kalavar et al., 2004, p. 62). • Fear of falling (Gallagher et al., 2010). 	Five	<ul style="list-style-type: none"> • Participation in PA was for health benefits and well-being (Arnautovska et al., 2017; Gray et al., 2016; Guell et al., 2018; Henwood, Tuckett, Edelstein, & Barlet, 2011; Miller & Brown, 2017). “It (PA) has definitely changed my life. It has lifted the depression big time” (Gray et al., 2016, p. 422).
7. Reinforcement	Three	<ul style="list-style-type: none"> • Lack of support from friends or family (Cassou et al., 2011; Guell et al., 2016). • The presence of people was identified as 	Nine	<ul style="list-style-type: none"> • Fun and enjoyment in doing PA (Arnautovska et al., 2017; Gray et al., 2016; Kosteli & Williams, 2016; Miller

		a barrier when carrying out PA (Gallagher et al., 2010).		& Brown, 2017). “Aye you have to enjoy it encourages you to do it, when you are feeling good it spurs you on, gives you that confidence to know you are doing it right” (Gray et al., 2016, p. 422).
				<ul style="list-style-type: none"> • Feel good factor (Leavy & Aberg, 2010; Miller & Brown, 2017). • Encouragement (Cousins, 2003; Sebastião et al., 2014) • Peer encouragement (Kalavar et al., 2004). • Positive experience (Mitra et al., 2015).
8. Intentions	Nil	<ul style="list-style-type: none"> • Nil 	Three	<ul style="list-style-type: none"> • To stay health or maintain independence (Gray et al., 2016; Kosteli, Williams, & Cumming, 2016). • A conscious decision to participate in PA, “So when I retired I decided that I wasn’t going to go hankering after my old workmates but that I would create new relationships around. As a result I became a lot more active and a lot more involved. . . .” (Leavy & Aberg, 2010, p. 226).
9. Goals	Four	<ul style="list-style-type: none"> • No motivation (Ceria-Ulep et al., 2011; Guell et al., 2018; Kalavar et al., 2004; Kolt et al., 2006). 	Five	<ul style="list-style-type: none"> • To improve health (Henwood et al., 2011). • PA provided a purpose in life (Kosteli & Williams, 2016). • Sense of competition (Chatfield, 2015; Gray et al., 2016) • A sense of self-motivation (Kalavar et al., 2004).

10. Memory, attention and decision processes	Nil	<ul style="list-style-type: none"> • Nil 	Nil	<ul style="list-style-type: none"> • Nil
11. Environmental context and resources	Fifteen	<ul style="list-style-type: none"> • Cost (Cassou et al., 2011; Gray et al., 2016; Miller & Brown, 2017; Sebastião et al., 2015). • Retirement as a busy period (Guell et al., 2018) • Weather (Alizadeh & Salehi, 2015; Arnautovska et al., 2017; Cassou et al., 2011; Jancey et al., 2009; Kalavar et al., 2004; Mitra et al., 2015; Sebastião et al., 2014) • Lack of time (Arnautovska et al., 2017; Cassou et al., 2011; Gray et al., 2016; Kalavar et al., 2004) • Environmental such as traffic, lack of green areas (Alizadeh & Salehi, 2015; Chaudhury, Mahmood, Michael, Campo, & Hay, 2012; Gallagher et al., 2010; Kalavar et al., 2004; Leavy & Aberg, 2010; Mitra et al., 2015; Rathanaswami, Bengoechea, & Bush, 2016) • Lack of facilities (Gray et al., 2016) • Neighbourhood safety (Arnautovska et al., 2017; Cassou et al., 2011; Chaudhury et al., 2012; Gallagher et al., 2010; Kolt et al., 2006; Mitra et al., 2015; Sebastião et al., 2015, 2014) • Transport (Gray et al., 2016; Kolt et al., 2006). 	Two	<ul style="list-style-type: none"> • Environmental aesthetics (Arnautovska et al., 2017). • A sense of safety and nice environment “I love to hear the birds singing in the morning” (Gallagher et al., 2010, p. 104), “We walk around [the outside of] the hospital, because they have all kinds of cameras, and police there, and it’s safe, very safe” (Gallagher et al., 2010, p. 104).
12. Social influences	Ten	<ul style="list-style-type: none"> • Family as a barrier, because of their age they were over protective and did not 	Eleven	<ul style="list-style-type: none"> • The social component was important, being part of a group was an opportunity

- expect them to participate in PA not to hurt themselves (Kolt et al., 2006).
- Labelled as too old to carry out PA (Jancey et al., 2009).
 - PA not being part of culture “If we start now suddenly, then others will tease us that we are trying very hard to be youthful” (Kalavar et al., 2004, p. 60).
 - The family role (Cassou et al., 2011; Ceria-Ulep et al., 2011; Gray et al., 2016; Rathanaswami et al., 2016; Sebastião et al., 2015).
 - Social isolation (Cassou et al., 2011).
 - Not willing to mix with other cultures (Kolt et al., 2006).
 - Uncontrollable life circumstances (Miller & Brown, 2017).
- to meet people (Gallagher et al., 2010; Gray et al., 2016; Guell et al., 2018; Henwood et al., 2011; Kosteli & Williams, 2016; Leavy & Aberg, 2010; Miller & Brown, 2017; Sebastião et al., 2015). “I walk every morning. There are people out there at this time of year, at 5:30 or 6:00 [a.m.], as soon as it’s light. They will say, ‘Hey, you’re late!’ Everybody knows each other” (Gallagher et al., 2010, p. 103). “That would be the top reason for me, certainly, meeting friends and enjoying yourself, meeting new people” (Gray et al., 2016, p. 422).
- Family and friends involvement (Cousins, 2003; Miller & Brown, 2017; Sebastião et al., 2015, 2014). “Every fortnight, I have a friend I used to work with, and we go for quite a long walk. Maybe anything from sort of seven to fourteen miles we do on that” (Guell et al., 2016, p. 4).
 - Part of a group “to remove solitude” (Miller & Brown, 2017).
 - Comparison with others (Chatfield, 2015).
 - Part of the social norm (Guell et al., 2016).
 - Retirement as an opportunity to be active (Leavy & Aberg, 2010).
13. Emotion
- Four
- Participants mentioned guilt feelings of dedicating time to self rather than family as a barrier (Rathanaswami et al., 2016).
- One
- PA as a way to regulate their mood, “I think I would be quite depressed if I didn’t do physical exercise of some

Supplementary material 1

MMAT scores

Critical Appraisal Score

Reference	MMAT question number							
	Qual 1.1	Qual 1.2	Qual 1.3	Qual 1.4	Quan 1.1	Quan 4.2	Quan 4.3	Quan 4.4
Alizadeh & Salehi, (2015)	x		x					
Arnautovska, O'Callaghan, & Hamilton, (2017)	x	x	x	x				
Arras, Ogletree, & Welshimer, (2006)					x	x	x	
Ayotte, Margrett, & Hicks-Patrick, (2010)					x	x	x	
Ball, Salmon, Giles-Corti, & Crawford, (2006)	x	x	x					
Belon, Nieuwendyk, Vallianatos, & Nykiforuk, (2016)	x	x	x					
Bopp et al., (2007)	x	x	x					
Bruner, (2009)	x	x	x					
Buman, Daphna Yasova, & Giacobbi, (2010)	x	x	x					
Caperchione et al., (2012)	x	x	x					
Casey, Eime, Ball, & Payne, (2011)	x	x	x					
Cassou et al., (2011)	x	x	x					
Ceria-Ulep, Serafica, & Tse, (2011)	x	x	x					
Chatfield, (2015)	x	x		x				
Chaudhury, Mahmood, Michael, Campo, & Hay, (2012)	x	x						
Chrisman, Nothwehr, Yang, & Oleson, (2015)	x	x						
Cousins, (2003)	x	x	x					
Dacey, Baltzell, & Zaichkowsky, (2008)						x	x	
Das, Sartore-Baldwin, & Mahar, (2016)	x		x					
Evans, (2011)	x	x	x					
Gallagher et al., (2010)	x	x	x					
George, Kolt, Rosenkranz, & Guagliano, (2014)	x	x	x					
Gobbi et al., (2012)					x	x	x	
Gonzales & Keller, (2004)	x	x	x					

Gray et al., (2016)	x	x	x			
Guell, Panter, Griffin, & Ogilvie, (2018)	x	x	x			
Guell, Shefer, Griffin, & Ogilvie, (2016)	x	x	x			
Harley, Rice, Walker, Strath, & Quintiliani, (2014)	x	x	x			
Henwood, Tuckett, Edelstein, & Barlet, (2011)	x	x	x			
Jancey, Clarke, Howat, Maycock, & Lee, (2009)	x		x			
Kalavar, Kolt, Giles, & Driver, (2004)					x	x
Kamphuis, van Lenthe, Giskes, Brug, & Mackenbach, (2007)	x	x	x			
Keegan, Middleton, Henderson, & Girling, (2016)	x	x	x	x		
Kegler et al., (2013)	x	x	x			
Kolt, Driver, & Giles, (2004)					x	x
Kolt, Paterson, & Cheung, (2006)	x	x	x			
Kosteli, Williams, & Cumming, (2016)	x	x	x	x		
Kowal & Fortier, (2007)					x	x
Leavy & Aberg, (2010)	x	x				
Mathews, Lakshmi, Ravindran, Pratt, & Thankappan, (2016)	x	x	x			
McArthur et al., (2014)	x	x	x			
Melillo et al., (2001)	x	x	x			
Miller & Brown, (2017)	x	x				
Mitra, Siva, & Kehler, (2015)	x	x	x			
Mohamed et al., (2014)	x	x	x			
Mosquera et al., (2012)	x	x	x			
Paluck, Allerdings, & Kealy, (2006)		x	x			
Rathanaswami, Bengoechea, & Bush, (2016)	x	x	x			
Romeike et al., (2016)	x	x	x			
Royce et al., (2003)	x	x	x			
Sebastião et al., (2015)	x	x	x			
Sebastião, Ibe-Lamberts, Bobitt, Schwingel, & Chodzko-Zajko, (2014)	x	x	x			

Steltenpohl, Shuster, Peist, Pham, & Mikels, (2018)	x	x	x		
Van Dyck, Mertens, Cardon, De Cocker, & De Bourdeaudhuij, (2017)	x	x			
Wertman et al., (2016)	x	x	x	x	x

Note: 'x' denotes articles reached criteria.

MMAT is Mixed Methods Appraisal Tool

For Peer Review

Supplementary Material 2

Characteristics of studies containing quantitative data

Reference	Country	Sample	Age Range (years)	% Population within studied population	Behaviour theory	Barrier tool	Motivator tool
Arras et al., (2006)	USA	641	45-90	53.1	Health promotion model	Barriers to health promoting activities for disable people	Not assessed
Ayotte et al., (2010)	USA	472	50-75	100	Social cognitive theory	Perceived barriers to exercise scale	Not assessed
Dacey et al., (2008)	USA	703	50-79	53	Self-determination theory and transtheoretical model	Not assessed	Exercise motivation inventory-2
Gobbi et al., (2012)	Brazil	359	60+	49.3	Socio-ecological	Barrier to PA practice questionnaire	Not assessed
Kolt et al., (2004)	Australia	840	55-93	~60	No behavioural theory	Not assessed	Participation motivation questionnaire
Kowal & Fortier, (2007)	Canada	509	39-68	60	Ecological approach	Questionnaire based on previous literature	Not assessed

Supplementary Material 3

Qualitative data extracted and characteristics of studies containing qualitative data

Author	Country	Behavioural Theory	Age Range (years)	Sample Size	Barriers 50-64 years	Motivators 50-64 years	Barriers 65-70 years	Motivators 65-70 years
Alizadeh & Salehi, (2015)	Iran	No behaviour theory	60-97	60	Lack of motivation, physical health limitation, and customs.	No motivators reported in this age group.	Environmental safety, weather too hot.	No motivators reported in this age group.
Arnautovska et al., (2017)	Australia	Integrate Behaviour Change Model	67-87	20	No barriers reported in this age group.	No motivators reported in this age group.	Negative states (pain, fear and safety), past experience, perceptions about old age, time constraints, safety of footpaths, weather.	For well-being, obtain a health benefit, for fun, for pride, doctor, media, aesthetics of place.
Ball et al., (2006)	Australia	Ecological	18-65	56	Lack of time management, conditions at work, lack of motivation.	No motivators reported in this age group.	No barriers reported in this age group.	No motivators reported in this age group.
Belon et al., (2016)	Canada	ANGELO framework	25-64	35	Dirtiness, acts of vandalism, ugliness, safety, car	Social relationships, community organisation,	No barriers reported in this age group.	No motivators reported in this age group.

					dependence.	walking the dog.		
					Lack of time due to work, family and church. Lack of willpower or desire. Lack of knowledge on how to exercise, fatigue due to work schedule, health condition.			
					Exercise groups not constant, recreation places too crowded, neighbourhood safety concerns, no age appropriate programs.	Need to take care of body, health benefits of exercise, improve mental health, sense of vitality, being determined and have willpower.	No barriers reported in this age group.	No motivators reported in this age group.
Bopp et al., (2007)	USA	No behaviour theory		44				
					Cold weather, personal safety (primarily from animals),	No motivators reported in this age group.	No barriers reported in this age group.	No motivators reported in this age group.
Bruner & Chad, (2013)	Canada	No behaviour theory	15-55+	19				

Buman et al., (2010)	USA	No behaviour theory	50-75	26	laziness. History of traumatic experience, fear of injury, cost, time management.	Health concerns, prevention of disease, literature, witness other people getting sick, enjoyment.	No barriers reported in this age group.	No motivators reported in this age group.
Caperchione et al., (2012)	Australia	No behaviour theory	43.8 (±10.84)	30	Time, work, child care, family needs, laziness, lack of motivation.	Better health, weight management, disease prevention, set example to children, fear becoming ill.	No barriers reported in this age group.	No motivators reported in this age group.
Casey et al., (2011)	Australia	Socio-ecological model	25-64	25	Injuries and disabilities, pain, previous experience when younger, bad experience with health system, culture in social area, program with lack of	No motivators reported in this age group.	No barriers reported in this age group.	No motivators reported in this age group.

exclusivity for specific age groups, affordability of facilities, quality of facilities.

High socio-economic class: lack of social support and everyday obstacles, or barriers faced during their daily life, weather, social isolation and health conditions

Low socio-economic class women: cost, everyday obstacles, and household chores, lack of time, lack of safety.

No motivators reported in this age group.

No motivators reported in this

For Peer Review

Cassou et al., (2011) Brazil No behaviour theory 69.9 (±6.9) 25

Ceria-Ulep et al., (2011) Hawaii No behaviour 65-87 47

No barriers reported in this age group. No motivators reported in this age group.

No barriers reported in this No motivators reported in this

		theory		age group.	age group.	(family obligations, job constraints, community responsibilities), church religious, family obligations. Feeling tired and weak, lack of motivation and laziness, ageing.	age group.
Chatfield, (2015)	USA	No behaviour theory	53-70	16	Injuries, temporal illness, lack of time, weather extremes.	Mental health, manage daily stress, improve quality of life, Women: manage weight.	Change in motivation over time, competition motivation, comparison with others.
Chaudhury et al., (2012)	Canada	No behaviour theory	65-92	34	No barriers reported in this age group.	No motivators reported in this age group.	Uneven sidewalks or tripping hazard, busy street with high traffic volume, traffic congestion, feeling unsafe, crime, accessibility.
Chrisman et al., (2015)	USA	Social-cognitive	27-75	19	Narrow sidewalks, lack of resources	Children.	No barriers reported in this

		Theory			and facilities, busy streets, difficulty cycling, having to pay to use facilities, children, and crime.		age group.	age group.
Cousins, (2003)	USA	Social-cognitive Theory	55-92	41	No barriers reported in this age group.	No motivators reported in this age group.	Negative beliefs.	Social encouragement from family, friends, physicians, past mastery, habitual activity.
Das et al., (2016)	USA	Health Belief Model	51.3 (±11.6)	12	Perception of physical activity, work demands, believing they were physical active enough at work, family obligation, pain from PA ailments.	No motivators reported in this age group.	No barriers reported in this age group.	No motivators reported in this age group.
Evans, (2011)	USA	Kleinman's EMS	40-60	20	Physical condition, too tired, laziness	No motivators reported in this	No barriers reported in this	No motivators reported in this

					and lack of motivation, no partner, other obligations (work, family, social), lack of time, rain or excessive heat.	age group.	age group.	age group.
Gallagher et al., (2010)	USA	No behaviour theory	61-85	21	No barriers reported in this age group.	No motivators reported in this age group.	Rundown neighbourhood, presence of people, safety of area, traffic and sidewalks, bad weather, fear of falling.	Presence of people.
					Time and work commitments, family schedule, lack of available facilities, ageing and decline in physical condition, lack of motivation and laziness, not having something to	Maintaining and improve health, increase longevity, participate with physical activity with their children, be good role model to children, stress reliever, change of scenery, guilt of not being		
George et al., (2014)	Australia	No behaviour theory	34-64	15			No barriers reported in this age group.	No motivators reported in this age group.

					strive.	physical activity, belong to a social group, appearance and fitness, succeed in their sport of activity.		
Gonzales & Keller, (2004)	USA	Ecological	48-65	10	Family comes first, friends, laziness, fatigue, pain.	Family friends.	No barriers reported in this age group.	No motivators reported in this age group.
								Health benefits, maintain physical functioning, psychological well-being, enjoyment, satisfaction, opportunity to socialise, friendly competition.
Gray et al., (2016)	Northern Ireland	Self-determination Theory and Self-efficacy Theory	50+	28	Lack of recreational facilities, admission funds, lack of transport.	Low socio-economic class: health conditions, safety at night, knowledge on PA guidelines, weather.	High socio-economic class: time constraints, family commitments.	High socio-economic class: rehabilitation from health conditions, physical activity presented on the

For Peer Review

Guell et al., (2016)	UK	No behaviour theory	65-80	32	No barriers reported in this age group	No motivators reported in this age group	Physical limitation, ill health, lack of companions or friends.	media. Lifelong activities, social support, family, friends, neighbours, social norms.
Guell et al., (2018)	UK	No behaviour theory	65-80	40	No barriers reported in this age group.	No motivators reported in this age group.	Health problems, no motivation, retirement as a busy period in life.	Life history of participating if physical activity, health benefits, being part of a team, "busy ethic".
Harley et al., (2014)	USA	Success Model Approach and Positive Deviance	26-65	14	Financial, physical strain, history of sedentary relapse.	Getting or staying healthy, disease, weight loss, mental health, social connections and gratification.	No barriers reported in this age group.	No motivators reported in this age group.
Henwood et al., (2011)	Australia	No behaviour theory	65-81	18	No barriers reported in this age group.	No motivators reported in this age group.	No barriers reported in this age group.	Health benefits (physical and mental), group environment, social challenge to improve.

Jancey et al., (2009)	Australia	No behaviour theory	64-74	16	No barriers reported in this age group.	No motivators reported in this age group.	General aches and pains, loss of flexibility, lack of balance, confidence, and shortness of breath. Poor body image and incumbent weather. Social obstruction to exercise as they are labelled as old.	No motivators reported in this age group.
Kalavar et al., (2004)	USA	Self-efficacy Theory	66-79	10	No barriers reported in this age group.	No motivators reported in this age group.	Health problems, fear of injury, bad weather, inadequate environment (sidewalk, potholes), not something you do at old age, no past experience of exercise, not wanting to start something new, laziness, no motivation, not part of culture in	Peers, doctor, self-motivation.

For Peer Review

								US / daily routine, lack of time due to family.
					Bad weather and winter.			
					Low socio- economic class: poor neighbourhood aesthetics, feeling unsafe, cost for taking part in exercise.	High socio- economic class: walking the dog.	No barriers reported in this age group.	No motivators reported in this age group.
Kamphuis et al.,(2007)	Netherland	Theory of Planned Behaviour	29-81	38				
						Role models, awareness by others, activity together, moral support, 'me time', being part of group, physical environment.	No barriers reported in this age group.	No motivators reported in this age group.
Keegan et al., (2016)	UK	No behaviour theory	31-57	14	No barriers reported in this age group.			
		Positive Deviance and Socio- ecological	40-70	29	No barriers reported in this age group.	Managing health problems, advice from health care	No barriers reported in this age group.	No motivators reported in this age group.
Kegler et al., (2013)	USA							

For Peer Review

providers, manage or lose weight, health benefits of exercise, good feeling, and thinking of family, care for family members, and care for oneself. Pets, military background.

Lack of education about benefits and how to carry physical activity, lack of motivation, perceptions about safe walkways, weather, facilities, transport, extended family over protection of older generation, health and medical conditions, health professional not

No motivators reported in this age group.

No barriers reported in this age group.

No motivators reported in this age group.

Kolt et al., (2006) New Zealand Self-efficacy Theory 60-79 24

Kosteli et al., (2016)	UK	Social-cognitive Theory	54-79	37	Lack of time either perceived or real due to family or personal reasons, pain, lack of motivation to start exercising, less structure post-retirement, social comparison with younger generation, inclement weather, lack of exercise partner,	Prescription by physician, necessity to stay health, enjoyment, mood regulation, relaxation, social life, previous physical active at work, gives purpose in life.	No barriers reported in this age group.	No motivators reported in this age group.	instructed to exercise, embarrassed to exercise in front of other people, reluctance to participate with other cultural groups.
------------------------	----	-------------------------	-------	----	--	--	---	---	---

					financial.			
Leavy & Aberg, (2010)	Dublin and Stockholm	No behaviour theory	65-89	30	No barriers reported in this age group.	No motivators reported in this age group.	Environment being a barrier (heavy traffic, lack of green areas), joint pain and breathlessness.	Transition from working to retired life, self motivation feel good factor, being part of a club.
Mathews et al., (2016)	India	Socio-ecological model	25-60	28	Lack of time, motivation and interest, physical discomfort caused by walking.	Pleasant walking routes and sight of other walkers.	No barriers reported in this age group.	No motivators reported in this age group.
McArthur et al., (2014)	Canada	No behaviour theory	40-62	53	Poor mood, poor experience, self-sabotage, lack of structure, other demands, have necessary equipment ready, fatigue, self-sacrifice.	Positive feeling, enjoyment, doing something different, meaningful, perceived health benefits.	No barriers reported in this age group.	No motivators reported in this age group.
Melillo et al., (2001)	USA	No behaviour theory	59-76	17	No barriers reported in this	No motivators reported in this	Fear and a feeling of	No motivators reported in this

Author(s)	Country	Theory	Age Group	N	Barriers	Motivators	Other Findings
Miller & Brown, (2017)	USA	No behaviour theory	65-72	10	Health conditions, expensive, uncontrollable life events.	Health benefits, feel good factor, enjoyment, to remove solitude, part of routine, support from friends and family.	inappropriateness in their culture, negative effect on their health, shame at doing exercise,
Mitra et al., (2015)	Canada	No behaviour theory	65-74	14	No barriers reported in this age group	No motivators reported in this age group.	Personal safety, health condition, climate condition, sidewalks, wide roads, traffic, health benefits, previous life experience, health related problems.
Mohamed et al., (2014)	USA	No behaviour theory	28-65	17	Exercise not a top priority employment is.	Role models within their community,	No barriers reported in this

					Embarrassment to exercise, cultural barriers as in Somalia they as not use to exercising, cost of facilities and transport.	sense of competition.	age group.	age group.
					Lack of bike paths, physical infrastructure poor, past negative experience, hostile environment to cycle, cultural barrier to bicycle use, women perceived less safe than men, cultural barrier are they as not though at a young age.	No motivators reported in this age group.	No barriers reported in this age group.	No motivators reported in this age group.
Mosquera et al., (2012)	Colombia	No behaviour theory	20-64	44				
Paluck et al., (2006)	Canada	No behaviour theory	18-65+	44	Lack of facilities and resources, cold	No motivators reported in this	No barriers reported in this	No motivators reported in this

For Peer Review

					weather, distance to facilities, no formed groups.	age group.	age group.	age group.
Rathanaswami et al., (2016)	Canada	No behaviour theory	37-55	8	Family responsibility, upbringing, clothing, cost, language, societal issues.	No motivators reported in this age group.	Feeling guilty that they are taking time for themselves that does not benefit the whole family, location, immediate living surrounding and husband.	No motivators reported in this age group.
					Lack of time, busy schedule.			
					Male: irregular hours.			
					Female: work commitments, family commitments.			
Romeike et al., (2016)	Netherland	Theory of Planned Behaviour and Self- Regulating Theories	20-65	36	Lack of motivation, mood, tiredness.	No motivators reported in this age group.	No barriers reported in this age group.	No motivators reported in this age group.
Royce et al., (2003)	USA	Socio-	22-75	53	Perception that	No motivators	No barriers	No motivators

ecological

physical activity is inconvenient and not compatible with busy lifestyle. Tired at end of day, past experience with walking when younger, narrow view of what physical activity is, community values, gender difference in activity selection, lack of social support, no community value on competition and sport not participation, reliance on car transport, urban area makes

reported in this age group.

reported in this age group.

reported in this age group.

For Peer Review

					travelling long and distant, lack of available resources, poor walk ability, facilities opening hours, safety.			
Sebastião et al., (2015)	USA	Socio-ecological Perspective	60-80	20	No barriers reported in this age group.	No motivators reported in this age group.	Health issues, fixed income limited access to exercise, role as caregivers, neighbourhood unsafe.	Friends and family, social centre and church.
Sebastião et al., (2014)	USA	Socio-ecological	65-75	7	No barriers reported in this age group.	No motivators reported in this age group.	Chronic health problems, bad weather, lack of safety, lack of infra-structure.	Friends, encouragement.
Steltenpohl et al., (2018)	USA	Socio-emotional Selectivity Theory	18-26 and 59+	78	No barriers reported in this age group.	Encouragement from others, family and friends support, meet people.	No barriers reported in this age group.	No motivators reported in this age group.
Van Dyck et al., (2017)	Belgium	No behaviour theory	62.9 (± 1.9)	37	Physical barriers were	No motivators reported in this	No barriers reported in this	No motivators reported in this

					paving stones, tram rails (mainly dangerous for cyclists), wrongly parked cars, poorly maintained sidewalks, and dangerous crossings. Accessibility to sports centres, bad weather, no partner to be active with, lack of time, financial issues, no motivation, having too much time – difficult to organise free time, health barrier, insufficient opportunities.	age group.	age group.	age group.
Wertman et al., (2016)	Canada	Health Belief Model and Life Course	40-82	20	Not enough time, expensive.	Health benefits, new experiences,	No barriers reported in this age group.	No motivators reported in this age group.

For Peer Review

Theory

reduce
isolation,
individual
practice.

For Peer Review