

Reducing pain during wound dressings in burn care using virtual reality: a study of perceived impact and usability with patients and nurses.

FURNESS, Penelope <<http://orcid.org/0000-0003-4916-8800>>, PHELAN, Ivan, BABIKER, Nathan, FEHILY, Orla, THOMPSON, Andrew and LINDLEY, Shirley

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/24774/>

This document is the Accepted Version [AM]

Citation:

FURNESS, Penelope, PHELAN, Ivan, BABIKER, Nathan, FEHILY, Orla, THOMPSON, Andrew and LINDLEY, Shirley (2019). Reducing pain during wound dressings in burn care using virtual reality: a study of perceived impact and usability with patients and nurses. *Journal of Burn Care & Research*, 40 (6), 878-885. [Article]

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

1 **Title**

2 Reducing pain during wound dressing in burn care using VR: A study of perceived impact
3 and usability with patients and nurses.

4 **Abstract**

5 Burns patients often suffer severe pain during interventions such as dressing changes, even
6 with analgesia. Virtual Reality (VR) can be used to distract patients and reduce pain.
7 However, more evidence is needed from the patients and staff using the technology about its
8 use in clinical practice and the impact of different VR strategies. This small-scale qualitative
9 study explored patient and staff perceptions of the impact and usability of active and passive
10 VR during painful dressing changes. Five patients took part in three observed dressing
11 changes - one with an active VR scenario developed for the study, one with passive VR and
12 one with no VR - following which they were interviewed about their experiences. Three
13 nurses who performed the dressing changes participated in a focus group. Thematic analysis
14 of the resulting data generated four themes: 'Caution replaced by contentment', 'Distraction
15 and implications for pain and wound care', 'Anxiety, control and enjoyment' and 'Preparation
16 and communication concerns'. Results suggested that user-informed active VR was
17 acceptable to burn patients, helped manage their perceived pain, and was both usable and
18 desirable within the clinical environment. Further testing with larger samples is now required.

19

20 **Key words:** Burn Pain, Wound Care, Virtual Reality, Distraction, Usability, Acceptability,
21 Patient Perspectives, Staff Perspectives, Qualitative Methods.

22 **Introduction**

Burns patients often experience severe pain during interventions, such as when wound dressings are changed, combining the pain of treatment with the background pain of tissue damage^{1,2}. Opiates are routinely administered for burn pain³. However, opiates come with side effects⁴ and their effectiveness in managing the pain of procedures, such as dressing changes, has been questioned^{5,6}. Inadequate pain control has detrimental effects on psychological and physical wellbeing^{7,8,9}, patient confidence⁵ and compliance¹⁰. Therefore, evidence suggests other forms of analgesia should be considered. Pain theories, such as Gate Control Theory and neuromatrix theory^{11,12}, highlight the importance of psychological determinants of the pain experience, including perception, attention and anxiety. Interventions, such as hypnosis, which address these determinants, have proved effective in distracting patients⁶.

Virtual Reality (VR) as a clinical intervention can also act upon pain perception¹³. VR's 'artificial three-dimensional environment'¹⁴ works to increase demands upon attention¹⁵ and reduce cues to pain and anxiety before and during procedures¹⁶. When compared with analgesia alone, VR plus analgesia has been shown to achieve a significant reduction in procedural pain scores^{17,18}, and qualitative reports identify increased relaxation and cooperation, reduced pain and anxiety, and effective communication despite immersion in the VR technology¹⁸. Costs of VR technology are falling, and recent developments have both addressed shortcomings of earlier technology (such as nausea) and improved VR's applicability to the clinical area^{5, 19, 20}.

Based on dissatisfaction with current methods of pain control and a growing evidence base for the effectiveness of VR, reviewers have recommended its introduction to burn care and rehabilitation²¹. However, further detailed work is required to explore specific influential variables by considering the impact on different patient groups of different VR environments²². VR environments may need tailoring to specific groups for maximum

effect²³, for example, using ‘cold’ scenarios for burn patients, and developing different VR scenarios to suit children of different ages¹⁸. One variable of interest is the degree of immersion offered by the intervention.^{1,22, 19, 24} VR can offer active involvement for the user, or a passive experience of simply watching and listening. Tashjian et al. reported significantly greater reductions in pain when patients were involved in an active VR scenario via headset, compared with the passive experience of watching a video by the bed²⁵. However, given the differences between the two interventions, it was unclear to what extent whether the result was achieved through the active vs. passive element alone²⁶

A recent study conducted by the Authors (2018) developed user-informed scenarios based on active and passive VR and compared their effects on the experimental pain of a cold pressor test. Experimental pain studies offer greater variable control: participants can be administered the same pain stimulus and intervention, which makes it easier to distinguish the effects of the target variables on outcomes. Previous results have shown that experimental pain is lower with VR^{24,27,-28}. Our study supported these findings, demonstrating significant differences between VR conditions overall and the no-VR baseline in both pain threshold (the point at which pain was first experienced) and pain tolerance (the point at which the cold pressor pain became intolerable and participants removed their hand). In addition, findings showed that pain threshold was significantly higher in active, immersive VR conditions than passive ones. When results for active and passive scenarios were considered separately, significant differences from baseline were only demonstrated for the active condition. The small sample size is acknowledged; however these results indicated that the most effective form of VR in managing pain for this sample was an active, immersive experience (Authors, 2018).

Findings regarding VR - and especially immersive VR - in experimental pain relief are encouraging; however, experimental pain is relatively mild, of limited duration, escapable, and implies no health threat. It is not clear whether the effects on pain can be said to transfer

easily into the clinical environment²². Patients' types and levels of clinical pain are likely to differ, and their medical needs often influence how an intervention can be delivered²². It is therefore important that VR be trialled in the clinical arena to confirm its real world usability and effectiveness. The current study applied the VR interventions developed and trialled in our experimental pain trial to a small sample of burn inpatients undergoing regular dressing changes at a single UK Burns Unit. Approaching people who will actually use the intervention - patients and staff - has been described as a 'person-centred' approach which enhances the evidence base for intervention development and feasibility²⁹. The work was supported by a Medical Research Council Confidence in Concept grant [number will be supplied after blind review].

Aims

This study aimed to explore:

- patient and staff perceptions of the effect of active and passive VR on perceived pain and anxiety during painful dressings changes;
- patient perceptions of the usability, acceptability, engagement with active and passive VR scenarios;
- staff perceptions about the usability and implications of the VR technology within a Burns Unit inpatient setting.

Methods

Design

This was a small-scale qualitative usability study, employing qualitative methods in keeping with the person-centred approach to intervention development and feasibility work²⁹.

Review and Approval

The original study protocol was reviewed by the Patient and Public Involvement (PPI) Panel for the Directorate of Therapeutics and Palliative Care, [City] Teaching Hospitals NHS Foundation Trust, and their suggestions were followed. Ethical approvals for the trial as described were granted by The University Research Ethics Committee and NHS Research Ethics Committee (IRAS 221071).

Participants

Patients: Participants were adult inpatients at the local Burns Unit who were undergoing regular dressing changes during the study period. Exclusion criteria included head and neck burns, wound infection, current diagnosis of PTSD, active psychotic symptoms or high levels of distress. Suitable patients were briefly introduced to the study and supplied with a full information sheet, with details about aims, procedures and rights. Before taking written consent, participants were encouraged to try out a short VR experience. We aimed to recruit up to 10 participants, in keeping with similar intervention development and usability studies³⁰ Five patient participants were recruited during the time available. Hospital stays which were too brief for the trial, mental health problems, injury location and infection control problems were key factors in those who were not eligible or declined participation. Participant details are provided in Table 1.

TABLE 1 HERE

Staff: Three qualified (female) nurses who had been directly involved in the care of participating patients were invited to and participated in a short post-study focus group, to share their impressions of the VR technology, its impact, usability and acceptability.

Materials

Equipment: An Oculus Rift CV1 headset, PC and digital recorder.

VR Scenarios: From the four tested under experimental conditions (Authors, 2018), we offered participants a choice two active VR scenarios, both of which had proved effective. These were named 'Basket' and 'Flocker'. In Flocker the user-controlled character was engaged in herding sheep through various obstacles. Basket was an energetic scenario based on in which the user was involved making basketball shots and building up their score. As described in Authors (2018) these scenarios were developed by a games designer, following a consultative workshop which included burn survivors, games designers, clinical and academic psychologists. As described above, they were trialled under experimental conditions and proved acceptable and enjoyable to users, and effective in reducing perceived pain. As a passive VR experience, participants were offered a choice of videos from the Oculus video application, which included scenes such as seeing the world from the viewpoint of an eagle, swimming with dolphins, or exploring a space station.

Procedure

Patients took part in three observed dressing changes during the study - one without VR, one with an active VR scenario and one with the passive VR scenario. The order of dressing changes was altered between participants, as shown in Table 1. Decisions about the suitable timing of each were made between the patient, the clinical team and the researcher, and the order was varied between the five participants. IP spent time with the participant before, during and after the dressing. He prepared the equipment, provided instruction and facilitated short familiarisation sessions for the patients before they used each scenario. Dressings ranged from 12 minutes (P5, active VR) to 70 minutes (P3, active VR) in length, with most lasting between 25 and 40 minutes.

Data Collection

Patient Interviews: IP conducted interviews at the bedside following completion of the two observed VR dressing changes once participants were comfortable. Questions included such as 'How was your pain during the dressing change while you were in the VR environment?' 'How did you feel generally during the experience?' and 'How helpful did you find the VR during the dressing change?' IP conducted a second interview with each participant at the end of the study, to gather overview data, with questions such as, 'Which VR experience did you prefer and why?' and 'From your experience how does a dressing change under VR compare with one with no VR experience?'

Staff Focus group: PF conducted the staff focus group. It took place in a private room near the ward and was audio-recorded. Questions focused on staff members' experience, their sense of the patient experience, and their general impressions of the VR technology. Items included: 'How did the VR dressing changes differ, if at all, from the dressing change without VR?'; 'What do you think the patients' experience was of the VR dressing change?'; 'What have the difficulties or complications been when using this technology?' and 'On balance, do you feel this sort of intervention is beneficial; if so / if not, why?'

Analysis

Data from staff and patients were transcribed and anonymised. For example, nurses were identified by ns1, ns3, etc., and patient participants by pt2, pt4, etc.

Transcripts were analysed for themes using an in-depth inductive coding, thematic mapping and theme development process³¹. This was a semantic analysis, in which the focus was data content (rather than underlying assumptions) and interpretation involved identifying the significance and implications of themes and constituent data in the context of existing knowledge³¹. Themes were refined through constant comparative analysis within and between transcripts and then across the whole dataset. Key themes reflected what seemed to

be important aspects of the experience of VR among participants. PF acted as primary analyst, and themes were shared, discussed and refined through discussion with all authors.

Results

Four themes were generated from the combined dataset from patients and nurses: *Caution replaced by contentment*, *Distraction and implications for pain and wound care*’, *Anxiety, control and enjoyment*’ and *Preparation and communication concerns*’.

Caution replaced by contentment

This theme reflected how participants’ initial reluctance regarding VR had given way to positive perceptions. Two of the five participating patients initially decided against participating, but later changed their minds, based on the pain they had experienced without VR: *‘I didn’t want to, but it did good, and I’m glad I did’* (pt2). The novelty of and her unfamiliarity with VR technology initially caused pt5 anxiety and uncertainty; however, in retrospect, she commented, *‘I don’t think people should be afraid of doing it.’* It is not surprising that people experiencing the combined trauma of burn-injury, hospitalisation and severe pain were anxious and reluctant to take on something new. Nonetheless, these five participants had been willing to try VR and were unanimous that this had been a good idea. After the first VR trial, any initial anxiety had disappeared: as they approached the next VR trial, they were *‘excited to try it’* a second time (pt4).

Nurses were similarly impressed with how well VR had worked: *‘Generally my experience has been that the VR’s very helpful, very good at distracting’* (ns2). Both groups felt that nurses could *‘sell it more’* to patients, and one person suggested that hearing others’ positive experiences would help. Comments about VR and their experience of it from staff included *‘it was all positive’* (ns2), and from patients, *‘great’* (pt5), *‘brilliant’* (pt3, pt4), *‘it’s worth its*

189 *weight in gold'* (pt1) *'now I know what I want for Christmas'* (pt4), and *'If I get any money, I'll*
190 *get one of these'* (pt5). Based on their experience, patients wanted to use VR again for
191 dressing changes, even if this meant paying:

192 *'I will have it, and I would even say, as an option, you know. If people said, this is*
193 *early days, and you had to pay for it, I'd say, right then, I'd pay for it, I'd pay extra for*
194 *that. I would pay, rather than not have it.* (pt3).

195 Staff expressed their wish to be involved with any future funded research, were positive
196 about its future potential and impatient for it to be routinely available in the clinical arena.
197 Both groups suggested additional applications for VR in physiotherapy, rehabilitation,
198 childbirth, chronic pain and disabling conditions.

199 *'Distraction and its implications for pain and wound care*

200 This theme reflected the positive distracting effects of VR, and especially active scenarios,
201 which impacted on pain tolerance and gave nurses scope to do more and spend longer on
202 dressing changes. Additional nuanced data reflected the fluctuations in, and, sometimes,
203 increased pain resulting from more intensive wound care.

204 A key factor in reducing pain and increasing tolerance of wound care seemed to be the degree
205 of distraction created by VR:

206 *'It drags you off. It drags you off, definitely. They are picking off stuff where, say they*
207 *pick one or two off ... you'd be on it, wouldn't you, you're concentrating on the pain*
208 *all the time, where that does help me, it's distracting, the whole thing'* (pt3).

209 Active scenarios were more effective in distracting patients: *'[it was] better with VR; [but]*
210 *scenarios [were] better for taking mind off'* (pt1). In contrast, the relative slowness and
211 passivity of passive version facilitated only a limited degree of distraction for most

212 participants. Four spoke of feeling frustrated by the slowness and passivity of the experience
213 and needing better distraction from the pain. Immersion was further compromised during the
214 passive VR by swooping movements in videos, which induced dizziness and motion sickness
215 in some.

216 Patients were unanimous that they had achieved good levels of distraction (and no nausea) in
217 the active VR. Some spoke of awareness of pain and of what the nurses were doing - *'felt it*
218 *but not concentrating on it'* (pt2) - but their focus remained on the engaging scenario. Nurses
219 spoke of patients being *'amazed'* (ns2) by what they had done afterwards, and several patients
220 reported losing track of time, so immersed had they been in the virtual world: *'It seemed to go*
221 *much quicker than I thought'* (pt5).

222 In addition, wearing the headset and watching the scenario meant patients could not see the
223 wound and nursing activities: *'I didn't see what they were doing ... if I could see what they*
224 *were doing, I wouldn't let them'* (pt1). Without this distraction, normal behaviour involved
225 being drawn to and focusing on the wound and wound care, which increased pain. Not
226 watching meant reduced pain: *'Before you were thinking, it hurts, because watching them do*
227 *it makes it worse'* (pt2).

228 However, data suggested that the distraction of VR actually contributed towards pain in
229 unexpected ways. Participants' greater distraction from and tolerance of pain compared with
230 normal circumstances meant that nurses could spend longer on dressings and carry out more
231 intensive wound care, such as removal of numerous surgical staples and more extensive
232 debridement:

233 *'he was a lot better with the VR on and I did pick quite a lot ... he'd not allowed staff*
234 *to do what we would normally want to do because of the pain, whereas with the VR*
235 *he allowed me to do that'* (ns1).

236 This nurse commented that this patient's pain tolerance allowed her to remove more dead
237 tissue from the wound bed, with a potentially positive impact on healing and infection.
238 Without VR, the dressing change would therefore far more painful, yet with VR he had been
239 able to tolerate it and both he and the nurses were positive about the impact of VR on both
240 pain and wound care. However, pain relief and distraction for all patients came to an abrupt
241 end when the VR was removed after the dressing. A few patients - particularly where wound
242 care was more intensive - complained of lasting pain afterwards in both VR and non-VR
243 trials, as painkillers wore off. Participants suggested offering VR *after* a dressing, to extend
244 the positive distracting and analgesic effects.

245 Although there were reports of pain *after* dressings, perceived pain was clearly reduced
246 during the procedure with active VR. Nurses also believed patients had required less
247 analgesia with VR, but acknowledged the considerable variations brought about by
248 differences in the dressing change intervention and stage of healing, making it hard to
249 attribute this solely to VR:

250 Ns3: *'My patient didn't need any extra analgesia during, before or after the*
251 *dressing change. I think she probably would've liked some otherwise. I think*
252 *she felt she needed some, pre-dressing, and then she didn't.'*

253 Ns1: *'I get the feeling, on the whole, it did reduce it a little bit but then again ...*
254 *different dressing changes are different on the same person as things get*
255 *better.'*

256 This theme reflected the overall positive effects on pain and distraction of VR, and in
257 particular the active scenarios. That it might facilitate intensive wound care and potentially
258 affect post-procedural pain was not fully anticipated. These aspects are worthy of
259 consideration and will be discussed below.

260 *Anxiety, control and enjoyment*

261 This theme included data suggesting that VR had not only reduced negative psychological
262 effects of burns procedures, and had also created positive experiences, which were
263 unexpected. Participants believed that VR had reduced their pre-dressing anxiety before and
264 during their second trial of VR, because of their experience of distraction and its impact on
265 pain, especially in the active condition. Nurses' data were in agreement: their perception had
266 been '*lessened anxiety*' (ns1) and distress from patients during VR dressings. Some suggested
267 offering VR before (as well as during) a dressing change, to reduce anxiety, and on days
268 between dressings to reduce stress.

269 Most spoke of positive emotions in response to the VR. The active VR in particular was '*fun*',
270 '*challenging*', and '*enjoyable*' (various pts). Ns1 expressed surprise at participants' apparently
271 pleasurable engagement with the technology. She spoke about the '*laughter*', an outcome
272 rarely associated with painful dressing changes. Ns2 commented on occasional '*hilarity*' and
273 '*comical*' moments, noting that VR had '*lightened*' the experience for everyone.

274 One concern among eligible patients when deciding to take part was a fear of losing the
275 ability to talk easily with staff, for example, to ask them to stop, when engaged with the VR
276 scenarios. However, among those who actually participated, the technology had the opposite
277 effect: two described feeling they could control part of the otherwise passive and traumatic
278 dressing change experience when using VR. Having control meant retaining one's '*humanity*.'
279 The sense of having some control over the situation, along with the distraction and reduced
280 pain, helped some patients control their own emotional responses to the experience. For
281 example, pt5 spoke of '*trying to be a grown up*' despite the dreadful pain of her burns. The
282 VR, described as a '*crutch*,' meant that, rather than '*howling*' in response to dressing pain, she
283 had found '*something as trivial as a video was actually quite empowering for me because I*

284 *could take myself away'* (pt5). There was a sense of pride in her achievement of self-control
285 in circumstances which could otherwise be experienced as shameful, humiliating and
286 disempowering.

288 *Preparation and communication concerns*

289 Preparation and communication emerged as potentially problematic issues which impacted
290 primarily upon the nurses involved, but also by consequence upon the patients themselves. In
291 order to avoid burdening clinical staff, research team members took on the roles of preparing
292 participants for VR, managing the technology during dressing changes, and collecting data.
293 Therefore, although nurses were fully aware of the study, they did not receive training and
294 preparation in the technology. This limited their ability to discuss VR with patients before,
295 during and after its use between researcher visits. Both patients and staff commented that
296 greater staff knowledge would have helped: *'I thought the VR was really good but I didn't*
297 *know a lot about it before the dressing change. I hadn't got a clue how it worked'* (ns2). Both
298 patients and nurses suggested more preparation time (perhaps assisted by trained nurses)
299 would help, for example with *'the physicality of wearing it'* (pt5), or *'a practice with the VR*
300 *pre-dressing, so that they'd know what they'd like to do, what activity, and how to do it'*
301 (ns1). Greater direct involvement in the study could have allowed nurses to play a more
302 active role in preparing, supporting and informing VR users. Learning about the technology
303 together might also contribute towards development of closer staff-patient relationships.
304 Experienced burns staff may lose touch with the novelty of the experience of dressing
305 changes for patients. Shared unfamiliarity with and co-learning about VR in this context may
306 foster a greater empathy and understanding between staff and patients. Staff hopes in future

research for greater involvement with and '*training*' in VR use were mentioned in discussion, and will be considered below.

Practitioner-patient communication during procedures also emerged as a concern for the nursing staff. For optimal distraction, pain and anxiety relief effects, the user ideally requires deep immersion and minimal interruption from the outside world. Good nursing practice involves keeping the patient informed and involved:

'Normally when I'm doing a dressing, I'd explain what I'm doing, you know, explain things on their legs or whatever, how their wound is, what it looks like' (ns2).

Conflicting requirements placed nurses in a difficult position, caught between communication as interruption and communication as involvement: *'I couldn't kind of work out what my role was and what I should be doing... do you interrupt them when they're in that zone?'* (Ns2). Despite a sense of '*inadequacy*' in uncertain circumstances, these experienced practitioners navigated the situation well, opting to minimise their verbal interruptions to the most vital information, such as imminent body position changes etc. Nurses discussed how they might in future negotiate short breaks in the VR, when activities would temporarily cease to facilitate communication.

Discussion

This study explored the acceptability, perceived effectiveness and usability of active and passive VR scenarios in the clinical setting during inpatient dressing changes. Previous evidence has demonstrated reduced pain in burn patients when using VR, but detailed patient and staff perspectives have rarely been gathered. A recent mixed methods study set in a US burns outpatient clinic collected quantitative data from staff and quantitative and qualitative data from patients, which demonstrated satisfaction with and feasibility of the technology³³. Our findings add to what is already known, by providing in-depth qualitative evidence from

both staff and patients which demonstrated that VR was acceptable, feasible and welcomed by all participants when used during in-patient dressing changes. VR promoted distraction, reduced perceived pain during dressings, enhanced wound care, and improved wellbeing. Findings further suggested that immersive, active VR might be more useful in supporting pain and anxiety relief than more passive versions of the technology. O

Previous authors have recommended research focusing on the extent to which fun and presence contribute to effectiveness in VR interventions²². Our findings provide some insight into these aspects, indicating that user-informed immersive scenarios (e.g. those with increased presence and engagement) were particularly effective in distracting patients. They also suggest that, as well as reducing the negative impacts of dressing change on pain, anxiety and distress, immersive VR can create positive experiences of fun, challenge, hilarity and laughter, 'lightening' the experience for all parties. This study compared VR to normal care, which is minimal distraction, at best using a TV / video, but most often no pain relief beyond pharmacological methods. It has been noted that, while other distraction techniques, such as hypnosis, are effective, non-pharmacological interventions are rarely used in practice³⁴. A majority of European Burn Centres have expressed dissatisfaction with their current pain-management strategies for burns patients³⁵. This study contributes to a body of evidence demonstrating the potential for VR in addressing procedural pain.

Several unanticipated effects of the VR are worthy of discussion.

First, increased patient tolerance offered the nurses greater scope to provide intensive wound care, as reported elsewhere³², with positive potential for wound healing and recovery. This was tolerated well during the procedure but may have contributed to some reports of lasting pain afterwards. In addition, no matter how intensive the wound care, removing the VR also removes the distraction and analgesic effects. There will probably never be a way of

eradicating pain completely; however these unanticipated (negative) effects on the pain experience should be considered. It may mean the patient should be offered continued access to the VR afterwards, with the immersive experience gradually reduced rather than suddenly removed. It also suggests that VR and other forms of pain relief (such as analgesic medication) may be used in a complementary way, with one introduced before the other is withdrawn.

Second, communication during dressing changes is part of normal care, as a nurse informs the patient about what he/she is doing, answers questions, including about wound progress, and provides instruction to the patient, for example, about movements they need to assist with. Nurses were unsure how to manage this part of their role and activities in the present study, an issue which could be addressed more explicitly in future work. However, we noticed that, despite their uncertainty, nurses navigated this challenge very successfully. As a small team, the staff came to know their patients well and quickly developed an understanding of how to tailor communication to meet patient need. Individual preferences about communication could also be discussed with the patient, giving them an active role in decisions about their wound care, which should also support effective pain management³⁶.

Third, outcomes suggested that the decision to avoid burdening staff inadvertently limited their ability to support patients with its use. A recent mixed-methods study reported similar findings from its qualitative interviews³³. Short-term research projects led by funded research teams, in which researchers deliver the intervention, help demonstrate efficacy of an intervention^{33,37}, and indeed, our work suggested benefits to both staff and patients. However, more research needs to be done in which staff members are involved and empowered to engage, understand, and independently operate the equipment and explain the technology to patients. This helps ensure new treatments are properly costed and effectively integrated into the clinical setting after the research is finished. Markus et al.³⁸ trialled VR as an adjunct to

physiotherapy and found that the costs to staff in terms of time, setting up, managing and cleaning the equipment were so great, that they arguably outweighed the benefits to patients. Morris et al.³⁷ explored VR for burns physiotherapy in South Africa, and found, in contrast, that time spent managing the technology was not seen as problematic. Instead physiotherapists felt freed to focus more on movement than pain using VR, potentially benefitting patient recovery. This has resonance with our finding that nurses believed VR allowed them to focus more intensively on wound care (rather than pain management). The back-up systems, such as staff training, technical support, maintenance and cleaning of equipment, which would allow an intervention such as VR to support existing care without unduly burdening busy staff, simply aren't there³⁸. However, although systems are rarely in place yet, once set up and established, VR systems could be applied without great time and effort in routine clinical care of burn patients and others requiring dressing changes, such as those undergoing reconstructive surgery²². Indeed, if hospitals make the investment in the systems, there seems no reason why broader patient groups should not benefit, as suggested by the patients and staff in the current study.

Our study had methodological strengths and limitations. Strengths included user involvement in the development of the trialled active VR scenarios (for more detail, see Authors, 2018), which proved very acceptable and apparently effective in reducing perceived pain and anxiety. User involvement was recently recommended as a priority for burn rehabilitation research²¹. The qualitative approach was a strength: interview data from both staff and patients were very valuable in revealing unanticipated outcomes of this still relatively novel intervention, including unexpected experiential aspects, and detailed insights into implications of the technology for various stakeholders. This approach has been recommended in intervention feasibility and development work²⁹; however it is relatively unique in the field of VR research, which is dominated by quantitative approaches. Ford et

al.³³ gained some useful qualitative insights from patients but collected only quantitative data from staff, which limited its depth.

Limitations include the very small sample, which was constrained by the single-centre design, time limitations on funding use and clinical exclusion criteria. Future work should adopt multi-centre designs, allow longer for recruitment, and consider ways to reduce exclusions. For example, infection control concerns could be addressed by utilising replaceable foam inserts for use with the VR kit. Patients with head or neck burns were also excluded; however, one previous study found a way around this issue using arm-mounted VR equipment. While less immersive than a headset, authors found that those using the VR reported significantly lower pain than both passive distraction (watching a movie) and standard care³⁹. This was similar to our findings indicating the superiority of active VR. Having both head- and arm-mounted versions available would prevent excluding large sections of the burn population from accessing effective VR-based pain relief.

Finally, previous authors³⁹ have recommended physiological measures of pain, and, in keeping with its 'person-centred' approach²⁹, our study collected subjective perceptual data. Our sense is that, if patients themselves believe their pain is reduced and more tolerable, this should be sufficient recommendation. Indeed, pulse and BP ratings can increase under conditions of excitement (such as when playing an immersive scenario) as well as pain, so are open to misinterpretation. The patients' subjective experience and interpretation of their pain may be the most useful measure in improving their experience and reducing short and long-term impacts. Alternatively, if a more objective mode of pain assessment were required, one promising approach could be treating pharmacological analgesia use as a proxy for pain. A recent study found a 39% reduction in opioid requests under their immersive VR condition, despite no significant differences in pain and anxiety ratings⁴⁰. Like ours, their intervention was very positively evaluated, and 75% were willing to use it again. The finding of reduced

430 opiate analgesia during (and before and after) dressings due to lower pain perception⁴⁰ has
431 some support in our qualitative results. Reducing analgesia also reduces costs of care and
432 unwanted side effects. Side effects of opiates include respiratory depression, constipation,
433 sedation, nausea⁴¹⁻⁴³, and possibly even immunosuppression and infection⁴². Decreased use of
434 sedating, nauseating opiates may promote earlier mobilisation in recovery from burns²¹. VR
435 could have a role to play here, as suggested in physiotherapy studies^{37,38}, since it could enable
436 patients to focus on recovering movement, rather than on their pain.

437 This small study demonstrated the usability and acceptability of VR technology in a single
438 clinical setting, and the perceived effectiveness of active VR scenarios in managing the pain
439 and anxiety associated with dressing changes for five inpatients. Next steps would be to trial
440 on a multi-centre basis, using controlled approaches, as recommended by reviewers in the
441 area³⁴. Measures should also be taken to reduce exclusions, extend application of the
442 technology and recruit larger samples. Our experience suggests that future trials should
443 consider mixed methods because qualitative data help capture nuanced and unanticipated
444 outcomes. Staff preparation and involvement are important concerns, and teams should
445 consider the broader impact and analgesic potential of VR to address pain relief before,
446 during and after the procedure.

447 **References**

- 448 1. Hoffman HG, Patterson DR, Magula J, et al. Water-friendly virtual reality pain control
449 during wound care. *Int J Clin Psychol* 2004;60(2):189-95.
- 450 2. Pardesi O, Fuzaylov G. Pain Management in Pediatric Burn Patients: Review of Recent
451 Literature and Future Directions. *J Burn Care Res* 2017;38:335-47.
- 452 3. Patterson DR, Sharar SR. Burn Pain. In JD Loeser, SH Butler, CR Chapman et al. (eds.)
453 *Bonica's management of pain*. Philadelphia: Lippincott Williams and Wilkins; 2001, pp. 780-
454 787.
- 455 4. Cherny N, Ripamonti C, Pereira J. et al. Strategies to manage the adverse effects of oral
456 morphine: an evidence-based report. *J Clin Oncol*, 2001;19:2542-54.
- 457 5. Edwards J. Managing wound pain in patients with burns using soft silicone dressings.
458 *Wounds* 2011;7(4):1-4
- 459 6. Hoffman HG, Patterson DR, Carougher GJ et al. Effectiveness of Virtual Reality-Based
460 Pain Control With Multiple Treatments. *Clin J Pain* 2001;17:229-235.
- 461 7. Stoddard FJ, Saxe G, Ronfeldt H et al. Acute stress symptoms in young children with
462 burns. *J Am Acad Child Adolesc Psychiatry* 2006;45:87-93.
- 463 8. Macleod R, Shepherd L, Thompson AR. Posttraumatic Stress Symptomatology and
464 Appearance Distress Following Burn Injury: An Interpretative Phenomenological Analysis.
465 *Health Psychol* 2016;35(11):1197-1204.
- 466 9. Berger MM, Davadent M, Marin C, et al. Impact of a pain protocol including hypnosis in
467 major burns. *Burns* 2010;36:639-46.

468 10. Richardson P, Mustard L. The management of pain in the burns unit. *Burns* 2009; 35:921-
469 36.

470 11. Melzack R. Pain and the neuromatrix in the brain. *J Dent Educ* 2001;65:1378-1382.

471 12. Melzack R. Evolution of the neuromatrix of pain. The Prithvi Raj Lecture: presented at
472 the third World Congress of World Institute of Pain, Barcelona 2004. *Pain Pract* 2005;5:85-
473 94.

474 13. Gold JJ, Belmont KA, Thomas DA. The neurobiology of virtual reality pain attenuation.
475 *Cyberpsychol Behav* 2007;10:536-544.

476 14. Garrett B, Taverner T, Masinde W et al. Rapid Evidence Assessment of immersive
477 Virtual Reality as an Adjunct Therapy in Acute Pain Management in Clinical Practice. *Clin J*
478 *Pain* 2014;30(12):1089-1098.

479 15. Hoffman HG, Prothero J, Wells M et al. Virtual chess: the role of meaning in the
480 sensation of presence. *J Hum-Comput Interaction* 1998;10:251-263.

481 16. Hoffman HG, Patterson DR, Carouger GJ. Use of virtual reality for adjunctive treatment
482 of adult burn pain during physical therapy: a controlled study. *Clin J Pain* 2000;16(3):244-
483 250.

484 17. Farrar JT, Portenoy RK, Berlin JA et al. Defining the clinically important difference in
485 pain outcome measures. *Pain* 2000;88:287-294.

486 18. Das DA, Grimmer KA, Sparnon AL et al. (2005) The efficacy of playing a virtual reality
487 game in modulating pain for children with acute burn injuries: A randomised controlled trial.
488 *BMC Pediatrics* 2005;5:1.

- 489 19. Hoffman HG, Patterson DR, Seibel E et al. Virtual Reality Pain Control During Burn
490 Wound Debridement in the Hydrotank. *Clin J Pain* 2008;24(4):299-304.
- 491 20. Jones, T., Moore, T., Choo, J. The impact of virtual reality on chronic pain. *PLoS ONE*
492 2016;11(12):e0167523.
- 493 21. Schneider JC. Rehabilitation Research Priorities: The Next 10 Years. *J Burn Care Res*
494 2017;38:e635-e637.
- 495 22. Malloy KM, Milling LS. The effectiveness of virtual reality distraction for pain reduction:
496 A systematic review. *Clin Psychol Rev* 2010;30:1011-1018.
- 497 23. Hua Y, Qiu R, Yao W et al.. The Effect of Virtual Reality Distraction on Pain Relief
498 During Dressing Changes in Children with Chronic Wounds on Lower Limbs. *Pain Manage*
499 *Nurs* 2015;16(5):685-691.
- 500 24. Dahlquist LM, McKenna KD, Jones KK et al. Active and passive distraction using a
501 head-mounted display helmet on cold pressor pain in children. *Health Psychol* 2007;26:794-
502 801.
- 503 25. Tashjian VC, Mosadeghi S, Howard AR et al. Virtual Reality of Management of Pain in
504 Hospitalized Patients: Results of a Controlled Trial. *JIMR Health* 2017;4(1):e9
- 505 26. Taylor TF. The influence of shame on posttrauma disorders: have we failed to see the
506 obvious? *Eur J Psychotraumatol*. 2015; 6: 10.3402/ejpt.v6.28847.
- 507 27. Hoffman HG, Seibel EJ, Richards TL et al. Virtual reality helmet display influences the
508 magnitude of virtual reality analgesia. *J Pain* 2006;7:843-850.

- 509 28. Patterson DR, Hoffman HG, Palacious AG et al. Analgesic effects of posthypnotic
510 suggestions and virtual reality distraction on thermal pain. *J Abnorm Psychol* 2006;115:834-
511 841.
- 512 29. Yardley L1, Morrison L, Bradbury K et al. The person-based approach to intervention
513 development: application to digital health-related behavior change interventions.
514 *J Med Internet Res*. 2015; 30;17(1):e30. doi: 10.2196/jmir.4055.
- 515 30. Riiser, K., Løndal, K., Ommundsen, Yet al. - Development and Usability Testing of an
516 Internet Intervention to Increase Physical Activity in Overweight Adolescents. *JMIR Res*
517 *Protoc*. 2013; 2(1): e7, doi: 10.2196/resprot.2410
- 518 31. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2009;
519 3(2):77-101.
- 520 32. Faber AW, Patterson DR, Bremer M. Repeated Use of Immersive Virtual Reality therapy
521 to Control Pain During Wound Dressing Changes in Pediatric and Adult Burn Patients. *J*
522 *Burn Care Res* 2013;34:563-568.
- 523 33. Ford CG, Manegold EM, Randall CL et al. Assessing the feasibility of implementing low-
524 cost virtual reality therapy during routine burn care. *Burns* 2018;
525 <https://doi.org/10.1016/j.burns.2017.11.020>.
- 526 34. Sheffler M, Koranyi S, Meissner W et al. Efficacy of non-pharmacological interventions
527 for procedural pain relief in adults undergoing burn wound care: A systematic review and
528 meta-analysis of randomized controlled trials. *Burns* 2017.
529 <https://doi.org/10.1016/j.burns.2017.11.019>.
- 530 35. Retrouvey H, Shahrokhi S. Pain and the Thermally Injured patient - A Review of Current
531 Therapies. *J Burn Care Res* 2015;36:315-323.

532 36. Solowiej K, Mason V, Upton D. Psychological stress and pain in wound care, part 3:
533 management. *Journal of Wound Care*, 2010; 19(4); 153-155.

534 37. Morris LD, Louw QA, Crous LC. Feasibility and potential effect of a low-cost virtual
535 reality system on reducing pain and anxiety in adult burn injury patients during physiotherapy
536 in a developing country. *Burns* 2010;36:859-64.

537 38. Markus LA, Willems KE, Maruna CC et al. Virtual reality: Feasibility of implementation
538 in a regional burn center. *Burns* 2009;35:967-69.

539 39. Jeffs D, Dorman D, Brown S, et al. Effect of Virtual Reality of Adolescent Pain During
540 Burn Wound Care. *J Burn Care Res* 2014;35:395-408.

541 40. McSherry T, Atterbury M, Gartner S et al. Randomized, Crossover Study of Immersive
542 Virtual Reality to Decrease Opioid Use During Painful Wound Care Procedures in Adults. *J*
543 *Burn Care Res* 2017; doi:10.1097/BCR.0000000000000589.

544 41. Christie MJ, Connor M, Vaughan CW et al. Cellular actions of opioids and other
545 analgesics: implications for synergism in pain relief. *Clin Exp Pharmacol Physiol*
546 2000;27:520-3.

547 42. Rittner HL, Roewer N, Brack A. The clinical (ir)relevance of opioid-induced immune
548 suppression. *Curr Opin Anaesthesiol* 2010;23:588-92.

549 43. Maani CV, Hoffman HG, Fowler M et al. Combining ketamine and virtual reality pain
550 control during severe burn wound care: one military and one civilian patient. *Pain Med*
551 2011;12:673-8.

552