

## **Exercise Training as a Mediator for Enhancing Coronary Collateral Circulation: A Review of the Evidence**

NICKOLAY, Thomas <<http://orcid.org/0000-0002-1481-7949>>, NICHOLS, Simon <<http://orcid.org/0000-0003-0377-6982>>, INGLE, Lee and HOYE, Angela

Available from Sheffield Hallam University Research Archive (SHURA) at:  
<https://shura.shu.ac.uk/25036/>

---

This document is the Accepted Version [AM]

### **Citation:**

NICKOLAY, Thomas, NICHOLS, Simon, INGLE, Lee and HOYE, Angela (2019). Exercise Training as a Mediator for Enhancing Coronary Collateral Circulation: A Review of the Evidence. *Current Cardiology Reviews*, 15. [Article]

---

### **Copyright and re-use policy**

See <http://shura.shu.ac.uk/information.html>

# REFERENCES

- [1]  
Zimarino M, D'Andreanmatteo M, Waksman R, Epstein SE, De Caterina R. The dynamics of the coronary collateral circulation. *Nat Rev Cardiol* 2014; 11(4): 191-7.  
[<http://dx.doi.org/10.1038/nrcardio.2013.207>] [PMID: [24395049](#)]
- [2]  
Werner GS, Surber R, Ferrari M, Fritzenwanger M, Figulla HR. The functional reserve of collaterals supplying long-term chronic total coronary occlusions in patients without prior myocardial infarction. *Eur Heart J* 2006; 27(20): 2406-12.  
[<http://dx.doi.org/10.1093/eurheartj/ehl270>] [PMID: [17003048](#)]
- [3]  
Cui K, Lyu S, Song X, *et al.* Effect of Coronary Collaterals on Prognosis in Patients Undergoing Primary Percutaneous Coronary Intervention for Acute ST-Segment Elevation Myocardial Infarction: A Meta-Analysis. *Angiology* 2018; 69(9): 803-11.  
[<http://dx.doi.org/10.1177/0003319718768399>] [PMID: [29656656](#)]
- [4]  
Baroldi G, Mantero O, Scomazzoni G. The collaterals of the coronary arteries in normal and pathologic hearts. *Circ Res* 1956; 4(2): 223-9.  
[<http://dx.doi.org/10.1161/01.RES.4.2.223>] [PMID: [13293825](#)]
- [5]  
Fulton WFM. The Time Factor in the Enlargement of Anastomoses in Coronary Artery Disease. *Scott Med J* 1964; 9(1): 18-23.  
[<http://dx.doi.org/10.1177/003693306400900104>] [PMID: [14103801](#)]
- [6]  
Zoll PM, Wessler S, Schlesinger MJ. Interarterial coronary anastomoses in the human heart, with particular reference to anemia and relative cardiac anoxia. *Circulation* 1951; 4(6): 797-815.  
[<http://dx.doi.org/10.1161/01.CIR.4.6.797>] [PMID: [14879489](#)]
- [7]  
Meier P, Hemingway H, Lansky AJ, Knapp G, Pitt B, Seiler C. The impact of the coronary collateral circulation on mortality: a meta-analysis. *Eur Heart J* 2012; 33(5): 614-21.  
[<http://dx.doi.org/10.1093/eurheartj/ehr308>] [PMID: [21969521](#)]
- [8]  
Koerselman, van der Graaf, de Jaegere, Grobbee. Coronary Collaterals. *Circulation* 2003; 107(19): 2507-11.  
[PMID: [12756191](#)]
- [9]  
Pipp F, Boehm S, Cai WJ, *et al.* Elevated fluid shear stress enhances postocclusive collateral artery growth and gene expression in the pig hind limb. *Arterioscler Thromb Vasc Biol* 2004; 24(9): 1664-8.  
[<http://dx.doi.org/10.1161/01.ATV.0000138028.14390.e4>] [PMID: [15242864](#)]
- [10]  
Van Royen N, Piek JJ, Schaper W, Bode C, Buschmann I. Arteriogenesis: mechanisms and modulation of collateral artery development. *J Nucl Cardiol* 2001; 8(6): 687-93.  
[<http://dx.doi.org/10.1067/mnc.2001.118924>] [PMID: [11725265](#)]
- [11]  
Baeyens N, Bandyopadhyay C, Coon BG, Yun S, Schwartz MA. Endothelial fluid shear stress sensing in vascular health and disease. *J Clin Invest* 2016; 126(3): 821-8.  
[<http://dx.doi.org/10.1172/JCI83083>] [PMID: [26928035](#)]
- [12]  
Schaper. Collateral circulation. *Basic Res Cardiol* 2009; 104(1): 5-21.  
[<http://dx.doi.org/10.1007/s00395-008-0760-x>] [PMID: [19101749](#)]
- [13]  
Scholz D, Cai WJ, Schaper W. Arteriogenesis, a new concept of vascular adaptation in occlusive disease. *Angiogenesis* 2001; 4(4): 247-57.  
[<http://dx.doi.org/10.1023/A:1016094004084>] [PMID: [12197469](#)]
- [14]  
Seiler C, Stoller M, Pitt B, Meier P. The human coronary collateral circulation: development and clinical importance. *Eur Heart J* 2013; 34(34): 2674-82.  
[<http://dx.doi.org/10.1093/eurheartj/ehz195>] [PMID: [23739241](#)]
- [15]  
Carmeliet P. Mechanisms of angiogenesis and arteriogenesis. *Nat Med* 2000; 6(4): 389-95.  
[<http://dx.doi.org/10.1038/74651>] [PMID: [10742145](#)]
- [16]  
Degen A, Millenaar D, Schirmer SH. Therapeutic approaches in the stimulation of the coronary collateral circulation. *Curr Cardiol Rev* 2014; 10(1): 65-72.  
[<http://dx.doi.org/10.2174/1573403X113099990027>] [PMID: [23721076](#)]
- [17]  
Eitenmüller I, Volger O, Kluge A, *et al.* The range of adaptation by collateral vessels after femoral artery occlusion. *Circ Res* 2006; 99(6): 656-62.  
[<http://dx.doi.org/10.1161/01.RES.0000242560.77512.dd>] [PMID: [16931799](#)]
- [18]  
Qin X, Deng Y, Wu D, Yu L, Huang R. Does Enhanced External Counterpulsation (EECP) Significantly Affect Myocardial Perfusion?: A Systematic Review & Meta-Analysis. *PLoS One* 2016; 11(4): e0151822  
[<http://dx.doi.org/10.1371/journal.pone.0151822>] [PMID: [27045935](#)]
- [19]  
Gloeckler S, Meier P, de Marchi SF, *et al.* Coronary collateral growth by external counterpulsation: a randomised controlled trial. *Heart* 2010; 96(3): 202-7.  
[<http://dx.doi.org/10.1136/hrt.2009.184507>] [PMID: [19897461](#)]
- [20]

- Duncker DJ, Bache RJ. Regulation of coronary blood flow during exercise. *Physiol Rev* 2008; 88(3): 1009-86.  
[<http://dx.doi.org/10.1152/physrev.00045.2006>] [PMID: [18626066](#)]
- [21]  
Telkova IL, Teplakov AT. [Relationships between changes of coronary blood flow, energy metabolism of the myocardium, and hyperinsulinemia in patients with ischemic heart disease]. *Kardiologia* 2005; 45(8): 61-8.  
[PMID: [16091647](#)]
- [22]  
Magnani S, Roberto S, Sainas G, *et al.* Metaboreflex-mediated hemodynamic abnormalities in individuals with coronary artery disease without overt signs or symptoms of heart failure. *Am J Physiol Heart Circ Physiol* 2018; 314(3): H452-63.  
[<http://dx.doi.org/10.1152/ajpheart.00436.2017>] [PMID: [29127237](#)]
- [23]  
Schirmer SH, Degen A, Baumhäkel M, *et al.* Heart-rate reduction by If-channel inhibition with ivabradine restores collateral artery growth in hypercholesterolemic atherosclerosis. *Eur Heart J* 2012; 33(10): 1223-31.  
[<http://dx.doi.org/10.1093/eurheartj/ehr255>] [PMID: [21821843](#)]
- [24]  
Gloekler S, Traupe T, Stoller M, *et al.* The effect of heart rate reduction by ivabradine on collateral function in patients with chronic stable coronary artery disease. *Heart* 2014; 100(2): 160-6.  
[<http://dx.doi.org/10.1136/heartjnl-2013-304880>] [PMID: [24186565](#)]
- [25]  
Eckstein RW. Effect of exercise and coronary artery narrowing on coronary collateral circulation. *Circ Res* 1957; 5(3): 230-5.  
[<http://dx.doi.org/10.1161/01.RES.5.3.230>] [PMID: [13427133](#)]
- [26]  
Scheel KW, Ingram LA, Wilson JL. Effects of exercise on the coronary and collateral vasculature of beagles with and without coronary occlusion. *Circ Res* 1981; 48(4): 523-30.  
[<http://dx.doi.org/10.1161/01.RES.48.4.523>] [PMID: [7460222](#)]
- [27]  
Heaps CL, Parker JL. Effects of exercise training on coronary collateralization and control of collateral resistance. *J Appl Physiol* 2011; 111(2): 587-98.  
[<http://dx.doi.org/10.1152/japplphysiol.00338.2011>] [PMID: [21565987](#)]
- [28]  
Nolewajka AJ, Kostuk WJ, Rechnitzer PA, Cunningham DA. Exercise and human collateralization: an angiographic and scintigraphic assessment. *Circulation* 1979; 60(1): 114-21.  
[<http://dx.doi.org/10.1161/01.CIR.60.1.114>] [PMID: [445714](#)]
- [29]  
Niebauer J, Hambrecht R, Marburger C, *et al.* Impact of intensive physical exercise and low-fat diet on collateral vessel formation in stable angina pectoris and angiographically confirmed coronary artery disease. *Am J Cardiol* 1995; 76(11): 771-5.  
[[http://dx.doi.org/10.1016/S0002-9149\(99\)80224-0](http://dx.doi.org/10.1016/S0002-9149(99)80224-0)] [PMID: [7572652](#)]
- [30]  
Fujita M, Sasayama S, Asanoi H, Nakajima H, Sakai O, Ohno A. Improvement of treadmill capacity and collateral circulation as a result of exercise with heparin pretreatment in patients with effort angina. *Circulation* 1988; 77(5): 1022-9.  
[<http://dx.doi.org/10.1161/01.CIR.77.5.1022>] [PMID: [2834115](#)]
- [31]  
Belardinelli R, Georgiou D, Ginzton L, Cianci G, Purcaro A. Effects of moderate exercise training on thallium uptake and contractile response to low-dose dobutamine of dysfunctional myocardium in patients with ischemic cardiomyopathy. *Circulation* 1998; 97(6): 553-61.  
[<http://dx.doi.org/10.1161/01.CIR.97.6.553>] [PMID: [9494025](#)]
- [32]  
Werner GS. The role of coronary collaterals in chronic total occlusions. *Curr Cardiol Rev* 2014; 10(1): 57-64.  
[<http://dx.doi.org/10.2174/1573403X1066614031123814>] [PMID: [24611646](#)]
- [33]  
Zbinden R, Zbinden S, Windecker S, Meier B, Seiler C. Direct demonstration of coronary collateral growth by physical endurance exercise in a healthy marathon runner. *Heart* 2004; 90(11): 1350-1.  
[<http://dx.doi.org/10.1136/hrt.2003.023267>] [PMID: [15486146](#)]
- [34]  
Zbinden R, Zbinden S, Meier P, *et al.* Coronary collateral flow in response to endurance exercise training. *Eur J Cardiovasc Prev Rehabil* 2007; 14(2): 250-7.  
[<http://dx.doi.org/10.1097/HJR.0b013e3280565dee>] [PMID: [17446804](#)]
- [35]  
Möbius-Winkler S, Uhlemann M, Adams V, *et al.* Coronary Collateral Growth Induced by Physical Exercise: Results of the Impact of Intensive Exercise Training on Coronary Collateral Circulation in Patients With Stable Coronary Artery Disease (EXCITE) Trial. *Circulation* 2016; 133(15): 1438-48.  
[<http://dx.doi.org/10.1161/CIRCULATIONAHA.115.016442>] [PMID: [26979085](#)]
- [36]  
Pijls NHJ, De Bruyne B, Peels K, *et al.* Measurement of fractional flow reserve to assess the functional severity of coronary-artery stenoses. *N Engl J Med* 1996; 334(26): 1703-8.  
[<http://dx.doi.org/10.1056/NEJM199606273342604>] [PMID: [8637515](#)]
- [37]  
Laskey WK. Brief repetitive balloon occlusions enhance reperfusion during percutaneous coronary intervention for acute myocardial infarction: a pilot study. *Catheter Cardiovasc Interv* 2005; 65(3): 361-7.  
[<http://dx.doi.org/10.1002/ccd.20397>] [PMID: [15945105](#)]
- [38]  
Crisafulli A, Melis F, Tocco F, *et al.* Exercise-induced and nitroglycerin-induced myocardial preconditioning improves hemodynamics in patients with angina. *Am J Physiol Heart Circ Physiol* 2004; 287(1): H235-42.  
[<http://dx.doi.org/10.1152/ajpheart.00989.2003>] [PMID: [15016635](#)]
- [39]

Togni M, Gloekler S, Meier P, *et al.* Instantaneous coronary collateral function during supine bicycle exercise. *Eur Heart J* 2010; 31(17): 2148-55.

[<http://dx.doi.org/10.1093/eurheartj/ehq202>] [PMID: 20584776]

[40]

Lin S, Lu X, Chen S, *et al.* Human coronary collateral recruitment is facilitated by isometric exercise during acute coronary occlusion. *J Rehabil Med* 2012; 44(8): 691-5.

[<http://dx.doi.org/10.2340/16501977-0989>] [PMID: 22729798]

[41]

Shu W, jing J, Fu LC, *et al.* The relationship between diastolic pressure and coronary collateral circulation in patients with stable angina pectoris and chronic total occlusion. *Am J Hypertens* 2013; 26(5): 630-5.

[<http://dx.doi.org/10.1093/ajh/hps096>] [PMID: 23391622]

[42]

MacDougall JD, Tuxen D, Sale DG, Moroz JR, Sutton JR. Arterial blood pressure response to heavy resistance exercise. *J Appl Physiol* 1985; 58(3): 785-90.

[<http://dx.doi.org/10.1152/jap.1985.58.3.785>] [PMID: 3980383]

[43]

McCartney N. Acute responses to resistance training and safety. *Med Sci Sports Exerc* 1999; 31(1): 31-7.

[<http://dx.doi.org/10.1097/00005768-199901000-00007>] [PMID: 9927007]