

Reproducibility of the Online Food4Me Food-Frequency Questionnaire for Estimating Dietary Intakes across Europe

MARSHALL, Steven J., LIVINGSTONE, Katherine M., CELIS-MORALES, Carlos, FORSTER, Hannah, FALLAIZE, Rosalind, O'DONOVAN, Clare B., WOOLHEAD, Clara, MARSAUX, Cyril F.M., MACREADY, Anna L., NAVAS-CARRETERO, Santiago, SAN-CRISTOBAL, Rodrigo, KOLOSSA, Silvia, TSIRIGOTI, Lydia, LAMBRINOU, Christina P., MOSCHONIS, George, GODLEWSKA, Magdalena, SURWIŁŁO, Agnieszka, DREVON, Christian A., MANIOS, Yannis, TRACZYK, Iwona, MARTÍNEZ, J. Alfredo, SARIS, Wim H., DANIEL, Hannelore, GIBNEY, Eileen R., BRENNAN, Lorraine, WALSH, Marianne C., LOVEGROVE, Julie A., GIBNEY, Mike and MATHERS, John C.

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/25685/>

This document is the Accepted Version [AM]

Citation:

MARSHALL, Steven J., LIVINGSTONE, Katherine M., CELIS-MORALES, Carlos, FORSTER, Hannah, FALLAIZE, Rosalind, O'DONOVAN, Clare B., WOOLHEAD, Clara, MARSAUX, Cyril F.M., MACREADY, Anna L., NAVAS-CARRETERO, Santiago, SAN-CRISTOBAL, Rodrigo, KOLOSSA, Silvia, TSIRIGOTI, Lydia, LAMBRINOU, Christina P., MOSCHONIS, George, GODLEWSKA, Magdalena, SURWIŁŁO, Agnieszka, DREVON, Christian A., MANIOS, Yannis, TRACZYK, Iwona, MARTÍNEZ, J. Alfredo, SARIS, Wim H., DANIEL, Hannelore, GIBNEY, Eileen R., BRENNAN, Lorraine, WALSH, Marianne C., LOVEGROVE, Julie A., GIBNEY, Mike and MATHERS, John C. (2016). Reproducibility of the Online Food4Me Food-Frequency Questionnaire for Estimating Dietary Intakes across Europe. *The Journal of Nutrition*, 146 (5), 1068-1075. [Article]

Copyright and re-use policy

Sheffield Hallam University Research Archive

<http://shura.shu.ac.uk>

See <http://shura.shu.ac.uk/information.html>

Title

Reproducibility of the online Food4Me food frequency questionnaire for estimating dietary intakes across Europe^{1,2,3}

Footnotes

1, This work was supported by the European Commission under the Food, Agriculture, Fisheries and Biotechnology Theme of the 7th Framework Programme for Research and Technological Development [265494].

2, Author disclosures: None of the authors had a personal or financial conflict of interest.

3, Supplemental Tables and Figures are available from the “Online Supporting Material” link in the online posting of the article and from the same link in the online table of contents at <http://jn.nutrition.org>.

Author names

Steven J. Marshall^{*1}, Katherine M. Livingstone^{*1}, Carlos Celis-Morales¹, Hannah Forster², Rosalind Fallaize³, Clare B. O'Donovan², Clara Woolhead², Cyril F.M. Marsaux⁴, Anna L. Macready³, Santiago Navas-Carretero⁵, Rodrigo San-Cristobal⁵, Silvia Kolossa⁶, Lydia Tsirigoti⁷, Christina P. Lambrinou⁷, George Moschonis⁷, Magdalena Godlewska⁸, Agnieszka Surwillo⁸, Christian A. Drevon⁹, Yannis Manios⁷, Iwona Traczyk⁸, J. Alfredo Martinez⁵, Wim H. Saris⁴, Hannelore Daniel⁶, Eileen R. Gibney², Lorraine Brennan², Marianne C. Walsh², Julie A. Lovegrove³, Mike Gibney², John C. Mathers¹, on behalf of the Food4Me Study.

Author affiliations

1, Human Nutrition Research Centre, Institute of Cellular Medicine, Newcastle University, Newcastle Upon Tyne, UK (KML, katherine.livingstone@newcastle.ac.uk; CCM, carlos.celis@newcastle.ac.uk; JCM, john.mathers@newcastle.ac.uk; SJM steven.j.marshall@northumbria.ac.uk)

2, UCD Institute of Food and Health, University College Dublin, Belfield, Dublin 4, Republic of Ireland (CBD, clare.odonovan@ucdconnect.ie; HF, hannah.forster@ucdconnect.ie; CW, clara.woolhead@ucdconnect.ie; ERG, eileen.gibney@ucd.ie; LB, lorraine.brennan@ucd.ie; MCW, marianne.walsh@ucd.ie; MG, mike.gibney@ucd.ie)

3, Hugh Sinclair Unit of Human Nutrition and Institute for Cardiovascular and Metabolic Research, University of Reading, Reading, UK (ALM, a.l.macready@reading.ac.uk; RF, r.fallaize@reading.ac.uk; JAL, j.a.lovegrove@reading.ac.uk)

4, Department of Human Biology, NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University Medical Centre, Maastricht, The Netherlands (CFMM, c.marsaux@maastrichtuniversity.nl; WHS, w.saris@maastrichtuniversity.nl)

5, Center for Nutrition Research, University of Navarra, Pamplona, Spain; CIBER Fisiopatología Obesidad y Nutrición (CIBERObn), Instituto de Salud Carlos III, Madrid, Spain (SNC, snavas@unav.es; RSC, rsan.1@alumni.unav.es; JAM, jalfmtz@unav.es)

6, ZIEL Research Center of Nutrition and Food Sciences, Biochemistry Unit, Technical University of Munich, Germany (SK, silvia.kolossa@tum.de; HD, hannelore.daniel@tum.de)

7, Department of Nutrition and Dietetics, Harokopio University, Athens, Greece (LT, tsirigoti.lydia@gmail.com; CPL, cplambrinos@gmail.com; GM, gmoschi@hua.gr; YM, manios@hua.gr)

8, National Food & Nutrition Institute (IZZ), Poland (MG, mgodlewska@izz.waw.pl; AS, asurwillo@izz.waw.pl; IT, itraczyk@izz.waw.pl)

9, Department of Nutrition, Institute of Basic Medical Sciences, Faculty of Medicine,
University of Oslo, Oslo, Norway (CAD, c.a.drevon@medisin.uio.no)

*SJM and KML are joint first authors

Pubmed indexing: Marshall; Livingstone; Celis-Morales; Foster; Fallaize; O'Donovan;
Woolhead; Marsaux; Macready; Navas-Carretero; San-Cristobal; Kolossa; Tsirigoti;
Lambrinou; Moschonis; Godlewska; Surwiłło; Drevon; Manios; Traczyk; Martinez; Saris;
Daniel; Gibney; Brennan; Walsh; Lovegrove; Gibney; Mathers

Corresponding author; request for reprints

Professor John C. Mathers

Human Nutrition Research Centre

Institute of Cellular Medicine

Newcastle University

Biomedical Research Building

Campus for Ageing and Vitality

Newcastle upon Tyne

NE4 5PL

john.mathers@newcastle.ac.uk

Tel: +44 (0) 1912081133 Fax: +44 (0) 1912481101

Word Count - 5303

Number of figures - 1

Number of tables - 3

Running title: Online food frequency questionnaire reproducibility

OSM available**Abbreviations**

Food frequency questionnaire (FFQ); Limits of agreement (LOA); Monounsaturated fatty acid (MUFA); Omega-3 fatty acid (n-3 FA); Pearson's correlation coefficient (PCC); Physical activity level (PAL); Polyunsaturated fatty acid (PUFA); Randomized controlled trial (RCT); Saturated fatty acid (SFA); Sedentary behavior (SB); Spearman's correlation coefficient (SCC, rho)

Abstract

Background: Accurate dietary assessment is key to understanding nutrition-related outcomes and is essential for estimating dietary change in nutrition-based interventions.

Objective: The objective of this study was to assess the pan-European reproducibility of the Food4Me FFQ in assessing the habitual diet of adults.

Methods: Participants were included from the Food4Me study, a 6-mo, internet-based, randomized controlled trial of personalized nutrition conducted in the UK, Ireland, Spain, the Netherlands, Germany, Greece and Poland. Screening and baseline data (both prior to commencement of the intervention) were used in the present analyses and participants were only included if they completed FFQs at screening and at baseline within a one-month timeframe prior to the commencement of the intervention. Socio-demographic (e.g. sex and country) and lifestyle (e.g. BMI and physical activity) characteristics were collected. Linear regression, correlation coefficients, concordance (%) in quartile classification and Bland-Altman plots for daily intakes were used to assess reproducibility.

Results: 567 participants (age 38.7 ± 13.4 y; 59% female; BMI 25.4 ± 4.8 kg/m²) completed both FFQs within one-month (mean 19.2 ± 6.2 d). Exact plus adjacent classification of total energy intake in participants was highest in Ireland (94%) and lowest in Poland (81%). Spearman Correlation Coefficients (ρ) in total energy intake between FFQs ranged from 0.50 for obese participants to 0.68 and 0.60 in normal and overweight participants respectively. Bland-Altman plots showed a mean difference between FFQs of 210 kcal/d, with the agreement deteriorating as energy intakes increased. There was little variation in reproducibility of total energy intakes between sex and age groups.

23 Conclusions: The online Food4Me FFQ was shown to be reproducible across 7 European
24 countries when administered within a one month period to a large number of participants.
25 The results support the utility of the online Food4Me FFQ as a reproducible tool across
26 multiple European populations.

27 Trial registration – Clinicaltrials.gov NCT01530139

28 **Key words:** Food frequency questionnaire; reproducibility; online; dietary intakes; European

29 Introduction

30 Given that poor diet is a predominant cause of the growing burden of non-communicable
31 diseases, more effective strategies for improving diet are of increasing importance (1) . In
32 tandem, accurate dietary assessment tools are essential for evaluating the efficacy of lifestyle
33 interventions (2) but all current methods of assessing habitual dietary intakes (including
34 weighed-dietary intakes, 24-hour dietary recall and food frequency questionnaires (FFQ)) are
35 subjective (3). Although weighed dietary recalls are considered the most accurate of the three
36 (4), retrospective recalls (24-hour recalls and FFQs) offer the advantages of lower costs and
37 lower-responder burden (5) and are therefore widely used in large scale epidemiological and
38 intervention studies.

39 With more than 70% of Europeans now Internet users (6), Internet-based diet and lifestyle
40 interventions, including Internet-based FFQs, are an attractive, cost-effective and scalable
41 alternative to face-to-face interventions (7). However, self-reported dietary assessment is
42 prone to respondent bias (8), which may limit reproducibility of the FFQ, resulting in poor
43 measures of dietary change and in chance associations with disease outcomes (9, 10). It is
44 therefore essential to evaluate the measurement error and reproducibility of FFQs to ensure
45 confidence in the precision of any diet-related outcomes.

46 The online Food4Me FFQ used in this study was validated previously against a weighed food
47 record over a 4-wk period (n=49) and showed moderate agreement (correlation coefficient
48 0.47) for assessing energy and nutrient intake (11), and a good agreement (0.60) against the
49 EPIC-Norfolk printed FFQ (n=113) (12). Furthermore the reproducibility of the online
50 Food4Me FFQ was assessed in the UK (n=100) and showed good agreement, with mean
51 cross-classification into "exact agreement plus adjacent" at 92% for both nutrient and food
52 group intakes (11). The aim of our present investigation was to verify that the online

Food4Me FFQ was reproducible across 7 European countries by comparing estimated intakes of foods, energy and nutrients between screening and baseline in the Food4Me study.

Methods

Study design

The Food4Me study was a 6-mo, internet-based, randomized controlled trial (RCT) of personalized nutrition designed to improve diet and PA behaviors, which was conducted across 7 European countries (n=1607). Recruitment was via the Food4Me website (13) from the following sites: University College Dublin (Ireland), Maastricht University (The Netherlands), University of Navarra (Spain), Harokopio University (Greece), University of Reading (United Kingdom, UK) and National Food and Nutrition Institute (Poland), Technical University of Munich (Germany). Individuals with ill-health, food intolerances, or special nutritional requirements (e.g. pregnancy) were ineligible to participate. Body mass index (BMI) was estimated from self-reported body weight and height (14). Participants self-reported smoking habits and occupation. Physical activity level (ratio between total energy expenditure and basal metabolic rate; PAL) and sedentary behavior (SB; min/d) were estimated from tri-axial accelerometers (TracmorD, Philips Consumer Lifestyle, The Netherlands). The Research Ethics Committees at each University or Research Centre granted ethical approval for the study. All participants signed online consent forms. The Food4Me trial was registered as a RCT (NCT01530139) at Clinicaltrials.gov. Full details on the study design are available elsewhere (14).

Food4Me FFQ

The Food4Me FFQ is an online, semi-quantitative FFQ, which was administered to individuals at screening, baseline and at follow-up timepoints following randomization. For the purposes of this reproducibility study, screening and baseline were used, as no change in diet was expected. FFQs were available in the language of the country, with respondents asked to report mean consumption over the previous month for 157 items in the UK and Ireland (based on the 130-item printed EPIC-Norfolk FFQ (version CAMB/PQ/6/1205) (12, 15)), with additional country-specific foods added to capture intakes in the other 5 recruitment countries (e.g. “stroopwafels” was added to the Dutch FFQ). A total of 11 food categories were included: 1) cereal, 2) bread and savory biscuits, 3) potatoes, rice and pasta, 4) meat and fish, 5) dairy products, 6) fats and spreads, 7) sweets and snacks, 8) soups, sauces and spreads, 9) drinks, 10) fruit and 11) vegetables (Table S1). Frequency of consumption of each food item was estimated by selecting one of the following options: never or less than once/mo, 1-3 times/mo, once/wk, 2-4 times/wk, 5-6 times/wk, once/d, 2-3 times/d, 5-6 times/d or >6 times/d. The online Food4Me FFQ included photographs of the foods and participant selected the appropriate portion size from the following options: very small, small, small/medium, medium, medium/large, large or very large. Food intake (g/d) was then calculated by multiplying portion size by frequency of consumption. For the purpose of comparing food group intakes, the 11 food categories were subdivided into 35 food groups based on previous validation by Forster *et al.* (12). Further details on the Food4Me FFQ are provided elsewhere (14).

Statistical Analysis

Statistical analysis was performed using STATA (version 12; StataCorp, College Station, TX, USA) and MedCalc Statistical Software (version 12.2.1.0; Medcalc, Mariakerke, Belgium).

ANOVA (continuous data) and logistic regression (categorical) tested for overall differences in anthropometric and socio-demographic characteristics (dependent variable) between countries (independent variable) and were adjusted for age and sex. Post hoc Tukey's tests and logistic regression (adjusted for age and sex) investigated differences in characteristics (dependent variable) between a given country and the overall mean for all countries (independent variable) (**Table 1**). FFQ reproducibility was determined by comparing dietary intakes at screening and baseline (mean 2.7 ± 0.9 wk apart). As the FFQ was designed to assess dietary intakes over a 1-month period, participants were excluded from the current analysis if the time period between completion of FFQs was > 1 month (16). Participants with implausible energy intakes were excluded based on the upper limit of sustained energy expenditure defined by the Scientific Advisory Committee for Nutrition: energy intake > 2.5 x Basal Metabolic Rate (17). Multiple linear regression was used to determine differences in total energy, nutrient and food group intakes (dependent variable) between FFQs (independent variable) and were adjusted for age, sex, country, time of FFQ completion and total energy intake at screening. Normality of data was assessed using the Shapiro-Wilk test and, depending on the outcome, comparison of energy, nutrients and food group intake was assessed using Pearson's product moment correlation coefficients (PCC) or Spearman's correlation coefficient (SCC, rho). Energy-adjusted correlation coefficients were estimated using the residual method (18). Briefly, residuals from the regression analysis (energy intake as independent variable and nutrient intake as dependent) were added to the expected nutrient value for the mean energy intake of the sample (**Table 2** and **Table 3**). The coefficient of reproducibility between methods was calculated (19). Concordance (%) in quartile classification estimated the relative agreements between FFQs. Quartiles of intakes of nutrients and food groups were used to determine changes in classification between timepoints. The percentages of participants classified into the correct quartile (exact

classification), adjacent quartile (exact classification plus adjacent), two quartiles apart (misclassification) or three quartiles apart (extreme misclassification) were estimated (**Supplemental Table 2 and Supplemental Table 3**). Bland-Altman plots determined clinical relevance of any difference in total energy and nutrients between methods based on the mean difference between methods (bias), trends, variability and widths of the limits of agreement (LOA; **Figure 1**). Reproducibility of total energy intakes was assessed according to age (<45 y and ≥ 45 y), sex, country, completion period between FFQs (short: 0-15.6 d; medium: 15.6-22.6 d; long: 22.6-31 d) and BMI at screening (underweight: BMI <18.5 kg/m²; normal weight: 18.5 to 24.9 kg/m²; overweight: 25 to 29.9 kg/m²; obese: ≥ 30 kg/m²) using regression analyses, SCC and concordance (%) in quartile classification (**Supplemental Table 4**).

Sensitivity analysis

Sensitivity analyses excluded participants who over- or under-reported energy intakes (**Supplemental Fig. 1**). Under-reporting was operationalized as an energy intake < 1.1 multiplied by predicted basal metabolic rate (using the Henry equation (20)) (21), and energy intakes > 4500 kcal/d were classified as over-reporting (22).

Results

Of the 1607 randomized participants, 1480 completed the FFQ at screening and at baseline and 665 completed the FFQs within one month of each other. Spain was excluded from all analyses due to insufficient numbers completing the FFQs within the 1-month timeframe (n=5). A further 93 participants were excluded based on implausible energy intakes.

Individuals from Greece had higher BMI, WC, more participants in routine and manual work, less students and more participants not currently working than the overall mean across all countries. Less Polish participants were in routine and manual employment and more Polish participants were females, while more Dutch were leaner, than the overall mean. Less participants from the United Kingdom were Caucasian, while there were less female participants from Ireland than the overall mean. No significant differences in PAL, BW or SB were identified (Table 1).

Reproducibility of nutrient intakes

Total energy intakes and intakes of protein, carbohydrate, total fat, saturated (SFA), mono- (MUFA) and polyunsaturated fatty acids (PUFA), omega-3 (n-3 FA), sugar, salt, calcium, folate, iron, carotene, riboflavin, fiber, sodium and vitamins B-6, C, A, D and E were lower at baseline than at screening ($P<0.05$; Table 2). There were no significant differences between timepoints for percentage energy intakes from total fat, MUFAs, PUFAs, protein, carbohydrate and sugars or for intakes of alcohol, vitamin B-12, thiamine and retinol. Shapiro-Wilk tests revealed that data were not normally distributed therefore SCC was used to examine correlations. Unadjusted SCCs ranged from 0.59 for total fat (g/d) to 0.89 for alcohol (mean 0.67; $P<0.001$), while energy adjusted SCCs ranged from 0.59 for total fat to 0.89 for alcohol (0.69; $P<0.001$; Table 2).

The percentage of participants whose dietary intakes were classified exactly at baseline, compared with screening, ranged from lowest for total fat to highest for alcohol (mean 50%; Supplemental Table 2). In total, 88% of participants were classified into the same or adjacent quartile, 10% were misclassified and 2% were extremely misclassified.

Bland-Altman plots comparing intakes of energy, total fat, protein and carbohydrate between timepoints are shown in Figure 1. The bias (mean difference) for total energy, carbohydrate, protein and fat intake was 210 kcal/d, 11.4%, 9.1% and 9.0% respectively. A positive trend indicated a lower agreement in intakes between timepoints for those who reported higher energy intakes (>4500 kcal/d) and who were classified as over-reporters in the sensitivity analyses. The amount consumed did not affect the agreement between intakes of carbohydrate, protein and fat.

Reproducibility of food group intakes

Reported intakes of wholemeal bread, biscuits, other fruits, meat products and soups, sauces and miscellaneous foods were lower at baseline compared with screening ($P<0.05$; Table 3). Unadjusted SCC ranged from 0.42 for tinned fruit or vegetables to 0.89 for alcoholic beverages (mean 0.71, $P<0.001$), while energy adjusted SCCs ranged from 0.45 for rice, pasta, grains and starches to 0.87 for alcoholic beverages (mean 0.69; $P<0.001$).

As shown in Supplemental Table 3, the percentage of participants correctly classified into the same quartile for food group intakes was lowest for rice, pasta, grains and starches and highest for alcoholic beverages. For all food groups, the mean percentages of participants who were misclassified and extremely misclassified were 8% and 2% respectively.

Sub group analysis: reproducibility of total energy intakes

As summarized in Supplemental Table 4, energy intake was lower at baseline than at screening for Greece, Poland and Germany. Correlations in energy intakes between timepoints were highest for the Netherlands and lowest for Greece, while the percentage

energy intakes correctly classified was lowest in Germany and the United Kingdom and highest in the Netherlands. Energy intake was lower at baseline compared with screening for those with short and medium time between assessments but not for the longest. For participants with the longest period of time between completing FFQs, SCC of energy intakes were poorest (Table S3). Energy intake was lower at baseline than at screening for normal and overweight participants but not for obese participants. SCCs were lower and the percentage of individuals misclassified was higher in overweight and obese participants than normal weight participants (Table S3). Energy intake was lower at baseline than at screening for participants both \geq and < 45 y. SCCs for energy intakes between timepoints were higher for participants ≥ 45 y, with similar proportions of individuals correctly classified and extremely misclassified. Energy intakes at baseline were lower than at screening for both male and females. Although more females than males were correctly classified into the same quartile, more females than males were misclassified (Table S3).

Sensitivity Analysis

Analyses were repeated in valid reports (n=437) after the removal of over- (n=8) and under-reporters (n=122). Supplemental Fig. 1 summarizes the delta between timepoints for percentage energy from fat, carbohydrates and protein in the total cohort and in valid reporters. This difference between timepoints is consistently smaller for the valid reporters in comparison with the whole cohort. After exclusion of mis-reporters, differences between timepoints in reported intakes of total fat, SFAs, MUFAs, PUFAs, n-3 FA, protein, calcium, carotene, riboflavin and vitamins C, A, biscuits, other fruits and soups, sauces and miscellaneous foods were not significant. For nutrients, SCC ranged from 0.60 for total fat and SFA g/day to 0.91 for alcohol and for food groups from 0.52 for rice, pasta, grains and

starches to 0.91 for alcoholic beverages ($P<0.001$). Bland-Altman analysis on valid reports produced a higher agreement between timepoints for total energy intake (bias reduced from 210 kcal/d to 88.5 kcal/d), carbohydrate (11.4% to 5.3%), protein (9.2% to 2.3%) and fat (9.5% to 2.4%). The coefficient of reproducibility in valid reports was reduced by 780kcal/d for energy intake, 14.4% for percentage energy from carbohydrate, 12.7% for protein intake and 13.3% for fat intake.

Discussion

Main findings

Our main findings indicate that the online Food4Me FFQ is reproducible for estimation of nutrient and food group intakes by adults across 7 European countries.

Comparison with other studies

An earlier study investigated the reproducibility of the online Food4Me FFQ by asking 100 participants within a single country (UK) to complete the FFQ on two occasions 4 wk apart. In that study, Fallaize *et al.* (11) reported higher mean correlation coefficients than in the present study for total energy intake (0.77 vs 0.61), nutrients (0.75 vs 0.67) and food group intakes (0.75 vs 0.71). Cross classification analysis for nutrients was also higher, with 92% of participants classified into the same or adjacent quartile, compared with 88% in the current paper. Bland-Altman analysis indicated a lower mean difference for total energy intake in the study by Fallaize *et al.* (11) compared with ours (135 kcal/d vs 210 kcal/d), however, the removal of mis-reporters lowered the mean difference in the current study to 89kcal/d. In the current study, the online Food4Me FFQ was administered to a much larger and more diverse

group of participants across 7 European countries who, in addition to completing the FFQ, were responding to a wider range of questionnaires. Furthermore, FFQ reproducibility in the study by Fallaize *et al.* (11) was assessed in conjunction with validation against a 4-day weighed food diary, which may have increased the participants awareness of their habitual intake and, thus, they may have been more likely to report similar intakes. The observed lower agreement between repeated administrations of the FFQ in the current study may be because the participants were less focused on the FFQ per se. Previous studies of the reproducibility of FFQs have reported correlation coefficients for total energy intake of 0.66 and 0.65 (8, 23, 24), which are very similar to our observations. The much higher correlation of 0.82 reported by Beasley *et al.* (25) was for an internet-based FFQ repeated within a short time interval (one wk) and thus subject to less variation (26). The shortest interval between FFQ administrations in the current study (0-15.65 d) produced a correlation of 0.64, lower than the 0.82 reported by Beasley *et al.* (25). However reproducibility in Beasley *et al.* (25) was also accompanied by a validation study against a 4-d weighed food diary, which may have improved correlations by increasing the participants awareness of their diet. Cross-classification analyses in the current study showed agreements that were comparable with previous studies for energy, nutrients and food groups (27-29). We observed that reported energy intakes were lower in the second FFQ, which confirms findings from other reproducibility studies (11, 25, 28, 30) and may be attributed to the learning effect of repeated measure. Alternatively, this observation may be due to fatigue caused by overburdening participants who had recently completed the initial FFQ (31). However, when mis-reporters were excluded, most differences between screening and baseline were no longer significant.

Previous FFQ reproducibility studies using repeated assessments within one month have reported coefficient ranges of 0.58-0.86 for energy intake between several countries (11, 23, 25, 28, 29, 32). Inter-country variations in SCCs in the Food4Me FFQ were similar,

suggesting that this dietary assessment tool has wide applicability across several European countries. The disparity between the cross-classifications and SCC in the UK may have been due to the presence of dietary mis-reporters and following exclusion of mis-reporters, these measures of reproducibility were more closely aligned. Our gender-dependent findings are consistent with a previous study (33), reporting higher reproducibility for a 240-item FFQ in males than in females (PCC 0.70 and 0.65, respectively). The reproducibility of the online Food4Me FFQ was similar for both older and younger participants. The lower reported energy intake at baseline compared with screening was significant for both normal weight and overweight participants but not for obese participants. This is probably due to a smaller sample size of obese individuals (n=79) compared with normal weight (n=296) and overweight (n=192) individuals as when assessed by SCC, reproducibility was lowest in the obese group. These findings confirm previous results, where obese individuals are more likely to mis-report their dietary intakes (34, 35). Self-administered dietary assessment tools should thus be interpreted with caution when applied to a population of predominantly obese subjects.

Previous studies on the validation and reproducibility of the Food4Me FFQ excluded under- and over-reporters prior to the main analysis (11, 12). The current study included the whole cohort. The percentage of people under-reporting (21.5%) was higher than that of over-reporters (1.4%), a common occurrence that has been previously reported (36). A sensitivity analysis following removal of misreports improved the reproducibility of the Food4Me FFQ.

Strengths and limitations

The main strength of this study is large number of participants from 7 European countries, which enabled stratification according to country, age, sex, obesity status and time interval

between FFQs. However, by excluding participants who did not complete FFQs within a 1-month period, we had too few participants from Spain (n=5) to allow comparisons with this country. Nonetheless, another strength of this study is that it was possible to assess the FFQ reproducibility between valid and mis-reporters in a European population. As recommended by Cade *et al.* (16), we applied the cut off of < 1 month between repeated FFQs to avoid confounding by real temporal changes in food intake. With a short time between the FFQs, it is conceivable that participants might remember and, therefore replicate, their previous FFQ responses (16). However, the comprehensive nature of the online Food4Me FFQ would make this unlikely and a 1-month period is considered an optimal time-period to assess reproducibility (16), whilst minimizing any influence of dietary change over time (11).

Conclusion

The Food4Me FFQ is moderately reproducible when administered to a large cohort of European adults. Variations in reproducibility between countries were small, thus providing confidence in the utility of the method for reporting intakes of energy, nutrients and food groups across multiple European countries.

Authors' contributions

The authors' responsibilities were as follows: YM, IT, CAD, ERG, LB, JAL, JAM, WHS, HD, MG and JCM contributed to the research design. JCM was the Proof of Principle study leader. CCM, CFMM, HF, CBO, CW, ALM, RF, SNC, RSC, SK, LT, CPL, MG, AS, MCW, ERG, LB and JCM contributed to the developing the Standardised Operating Procedure for

313 the study. CCM, SNC, RSC, CW, CBO, HF, CFMM, ALM, RF, SK, LT, CPL, MG, AS,
314 MCW and JCM conducted the intervention. CCM, CFMM and WHS contributed to physical
315 activity measurements. SJM and KML drafted the paper and performed the statistical
316 analysis for the manuscript and are joint first authors. All authors contributed to a critical
317 review of the manuscript during the writing process. All authors approved the final version to
318 be published.

319 **Figure Legends**

320 **Figure 1.** Bland-Altman plots for reproducibility between screening and baseline intakes of
321 A. total energy, B. fat, C. protein and D. carbohydrate (n=567) in European adults. The solid
322 line represents the mean difference, the dashed line represents the limits of agreement and the
323 dotted line represents the trend in agreement.

References

1. Beaglehole R, Bonita R, Horton R, Adams C, Alleyne G, Asaria P, Baugh V, Bekedam H, Billo N, Casswell S, et al. Priority actions for the non-communicable disease crisis. *Lancet*. 2011;377:1438-47.
2. Adamson AJ, Mathers JC. Effecting dietary change. *Proc Nutr Soc*. 2004;63:537-47.
3. Dhurandhar NV, Schoeller D, Brown AW, Heymsfield SB, Thomas D, Sorensen TI, Speakman JR, Jeansson M, Allison DB. Energy balance measurement: When something is not better than nothing. *Int J Obes*. 2015;39:1109-13.
4. Boushey C, Coulston AM. Nutrition in the prevention and treatment of disease. Amsterdam: Academic Press; 2008.
5. Subar AF. Developing dietary assessment tools. *J Am Diet Assoc*. 2004;104:769-70.
6. Miniwatts Marketing Group. World internet usage and population statistics [cited 2015 April 11]. Available from: <http://www.internetworldstats.com/stats.htm>.
7. Steele RM, Mummery WK, Dwyer T. A comparison of face-to-face or internet-delivered physical activity intervention on targeted determinants. *Health Educ Behav*. 2009;36:1051-64.
8. Fernandez-Ballarath JD, Lluís Pinol J, Zazpe I, Corella D, Carrasco P, Toledo E, Perez-Bauer M, Angel Martinez-Gonzalez M, Salas-Salvado J, Martin-Moreno JM. Relative validity of a semi-quantitative food-frequency questionnaire in an elderly mediterranean population of Spain. *Br J Nutr*. 2010;103:1808-16.
9. Marks GC, Hughes MC, van der Pols JC. Relative validity of food intake estimates using a food frequency questionnaire is associated with sex, age, and other personal characteristics. *J Nutr*. 2006;136:459-65.
10. Schatzkin A, Kipnis V, Carroll RJ, Midthune D, Subar AF, Bingham S, Schoeller DA, Troiano RP, Freedman LS. A comparison of a food frequency questionnaire with

- a 24-hour recall for use in an epidemiological cohort study: Results from the biomarker-based observing protein and energy nutrition (open) study. *Int J Epidemiol.* 2003;32:1054-62.
11. Fallaize R, Forster H, Macready AL, Walsh MC, Mathers JC, Brennan L, Gibney ER, Gibney MJ, Lovegrove JA. Online dietary intake estimation: Reproducibility and validity of the food4me food frequency questionnaire against a 4-day weighed food record. *J Med Internet Res.* 2014;16:e190.
 12. Forster H, Fallaize R, Gallagher C, O'Donovan CB, Woolhead C, Walsh MC, Macready AL, Lovegrove JA, Mathers JC, Gibney MJ, et al. Online dietary intake estimation: The food4me food frequency questionnaire. *J Med Internet Res.* 2014;16:e150.
 13. Food4Me. An integrated analysis of opportunities and challenges for personalised nutrition [cited 2015 December 12]. Available from: <http://www.food4me.org/>.
 14. Celis-Morales C, Livingstone KM, Marsaux CFM, Forster H, O'Donovan CB, Woolhead C, Macready AL, Fallaize R, Navas-Carretero S, San-Cristobal R, et al. Design and baseline characteristics of the food4me study: A web-based randomised controlled trial of personalised nutrition in seven european countries. *Genes Nutr.* 2015;10:450.
 15. Bingham SA, Gill C, Welch A, Cassidy A, Runswick SA, Oakes S, Lubin R, Thurnham DI, Key TJA, Roe L, et al. Validation of dietary assessment methods in the uk arm of epic using weighed records, and 24-hour urinary nitrogen and potassium and serum vitamin c and carotenoids as biomarkers. *Int J Epidemiol.* 1997;26:S137-S51.
 16. Cade J, Thompson R, Burley V, Warm D. Development, validation and utilisation of food-frequency questionnaires - a review. *Public Health Nutr.* 2002;5:567-87.

17. Scientific Advisory Committee on Nutrition. Dietary reference values for energy 2011 [cited 2016 19 February]. Available from:
https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/339317/SACN_Dietary_Reference_Values_for_Energy.pdf.
18. Willett WC, Howe GR, Kushi LH. Adjustment for total energy intake in epidemiologic studies. *Am J Clin Nutr*. 1997;65:1220-8.
19. Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet*. 1986;1:307-10.
20. Henry CJK. Basal metabolic rate studies in humans: Measurement and development of new equations. *Public Health Nutr*. 2005;8:1133-52.
21. Goldberg GR, Black AE, Jebb SA, Cole TJ, Murgatroyd PR, Coward WA, Prentice AM. Critical-evaluation of energy-intake data using fundamental principles of energy physiology: 1. Derivation of cutoff limits to identify under-recording. *Eur J Clin Nutr*. 1991;45:569-81.
22. Hebert JR, Peterson KE, Hurley TG, Stoddard AM, Cohen N, Field AE, Sorensen G. The effect of social desirability trait on self-reported dietary measures among multi-ethnic female health center employees. *Ann Epidemiol*. 2001;11:417-27.
23. Vereecken CA, De Bourdeaudhuij I, Maes L, Grp HS. The helena online food frequency questionnaire: Reproducibility and comparison with four 24-h recalls in belgian-flemish adolescents. *Eur J Clin Nutr*. 2010;64:541-8.
24. Johansson L, Solvoll K, Opdahl S, Bjorneboe GEA, Drevon CA. Response rates with different distribution methods and reward, and reproducibility of a quantitative food frequency questionnaire. *Eur J Clin Nutr*. 1997;51:346-53.
25. Beasley JM, Davis A, Riley WT. Evaluation of a web-based, pictorial diet history questionnaire. *Public Health Nutr*. 2009;12:651-9.

26. Tsubono Y, Nishino Y, Fukao A, Hisamichi S, Tsugane S. Temporal change in the reproducibility of a self-administered food frequency questionnaire. *Am J Epidemiol.* 1995;142:1231-5.
27. Filippi AR, Amodio E, Napoli G, Breda J, Bianco A, Jemni M, Censi L, Mammina C, Tabacchi G. The web-based ASSO-food frequency questionnaire for adolescents: Relative and absolute reproducibility assessment. *Nutr J.* 2014;13:
28. Labonte ME, Cyr A, Baril-Gravel L, Royer MM, Lamarche B. Validity and reproducibility of a web-based, self-administered food frequency questionnaire. *Eur J Clin Nutr.* 2012;66:166-73.
29. Nurul-Fadhilah A, Teo PS, Foo LH. Validity and reproducibility of a food frequency questionnaire (ffq) for dietary assessment in Malay adolescents in Malaysia. *Asia Pac J Clin Nutr.* 2012;21:97-103.
30. Zhao W-H, Huang Z-P, Zhang X, He L, Willett W, Wang J-L, Hasegawa K, Chen J-S. Reproducibility and validity of a Chinese food frequency questionnaire. *Biomed Environ Sci.* 2010;23:1-38.
31. Wong JE, Parnell WR, Black KE, Skidmore PML. Reliability and relative validity of a food frequency questionnaire to assess food group intakes in New Zealand adolescents. *Nutr J.* 2012;11:
32. Sarmento RA, Antonio JP, Riboldi BP, Montenegro KR, Friedman R, de Azevedo MJ, de Almeida JC. Reproducibility and validity of a quantitative ffq designed for patients with type 2 diabetes mellitus from Southern Brazil. *Public Health Nutr.* 2014;17:2237-45.
33. Kesse-Guyot E, Castetbon K, Touvier M, Hercberg S, Galan P. Relative validity and reproducibility of a food frequency questionnaire designed for French adults. *Ann Nutr Metab.* 2011;57:153-62.

34. Johansson L, Solvoll K, Bjorneboe GEA, Drevon CA. Under- and overreporting of energy intake related to weight status and lifestyle in a nationwide sample. *Am J Clin Nutr.* 1998;68:266-74.
35. Pfrimer K, Vilela M, Resende CM, Scagliusi FB, Marchini JS, Lima NKC, Moriguti JC, Ferriolli E. Under-reporting of food intake and body fatness in independent older people: A doubly labelled water study. *Age Ageing.* 2015;44:103-8.
36. Lutonski JE, van den Broeck J, Harrington J, Shiely F, Perry IJ. Sociodemographic, lifestyle, mental health and dietary factors associated with direction of misreporting of energy intake. *Public Health Nutr.* 2011;14:532-41.




Figure 1.

Table 1 Anthropometric and socio-demographic characteristics of European adults by country at the time of completing the screening Food4Me food frequency questionnaire¹

	Total (n=567)	Country						P ²
		Greece (n=160)	Ireland (n=70)	Netherlands (n=108)	Poland (n=153)	UK (n=49)	Germany (n=27)	
Age, y	38.7 ± 13.4	38.3 ± 11.2	39.6 ± 13.2	42.7 ± 16.6	35.0 ± 12.1	38.4 ± 12.6	43.4 ± 15.4	<0.001
Sex, Female, %	58.9	58.1	41.4*	50.9	70.1*	67.4	59.3	0.03
Ethnicity, Caucasian, %	97.5	99.4	97.1	95.4	100	87.8*	100	0.04
Occupation, %								
Professional and managerial	31.2	31.3	40.0	36.1	19.0	46.9	29.6	0.98
Intermediate occupations	29.1	28.1	21.4	17.6	46.4	12.2	33.3	0.47
Routine and manual	11.6	18.1*	14.3	8.3	5.9*	14.3	7.4	0.02
Student	17.1	7.5*	15.7	24.1	22.9	18.4	14.8	0.048
Not currently working	10.9	15.0*	8.6	13.9	5.9	8.2	14.8	0.04
Anthropometrics								
BMI, kg/m ²	25.4 ± 4.8	26.7 ± 5.5*	26.0 ± 4.6	24.4 ± 3.9*	24.7 ± 4.7	25.3 ± 4.3	24.5 ± 3.0	<0.001
Waist circumference, cm	85.5 ± 14.1	89.3 ± 14.8*	87.5 ± 14.1	84.6 ± 12.5	81.6 ± 14.2	84.2 ± 11.7	85.4 ± 13.0	<0.001
Body weight, kg	75.0 ± 15.4	76.9 ± 15.7	78.3 ± 16.3	74.7 ± 13.5	71.1 ± 16.2	72.7 ± 14.1	75.0 ± 12.1	0.13
Physical activity								
PAL	1.7 ± 0.2	1.7 ± 0.1	1.8 ± 0.2	1.7 ± 0.2	1.7 ± 0.2	1.7 ± 0.2	1.7 ± 0.2	0.07
SB, min/d	745 ± 78.0	744 ± 89.4	755 ± 72.1	753 ± 72.1	741 ± 79.2	725 ± 59.2	762 ± 66.7	0.36

1, Values represent means ± SD or percentages; PAL, physical activity level (ratio between total energy expenditure and basal metabolic rate); SB, sedentary behavior.

2, ANOVA and logistic regression were used to test for significant differences across countries in continuous and categorical variables, respectively. Analyses were adjusted for age and sex; * Post hoc Tukey tests (continuous data) and logistic regression (categorical) were used to test for significant differences between a given country and the overall study mean across all countries, P<0.05.

Table 2 Differences in total energy and nutrient intakes in European adults between screening and baseline as assessed using multiple linear regression and correlation coefficients¹

	Timepoint ²		P ³	Correlation coefficient ⁴	
	Screening	Baseline		Crude	Energy adjusted
Total energy, kcal/d	2455 ± 685	2246 ± 730	<0.001	0.61	-
Total fat, g/d	96.4 ± 32.2	89.2 ± 32.8	<0.001	0.59	0.59
Total fat, % energy	35.4 ± 6.2	35.7 ± 5.8	0.423	0.61	0.61
SFA, g/d	38.1 ± 14.6	35.1 ± 14.6	0.001	0.61	0.64
SFA, % energy	13.9 ± 3.2	14 ± 3.1	0.78	0.65	0.65
MUFA, g/d	36.7 ± 13.8	33.9 ± 13.2	<0.001	0.62	0.69
MUFA, % energy	13.5 ± 3.5	13.6 ± 3.2	0.54	0.72	0.71
PUFA, g/d	15.3 ± 5.4	14.3 ± 5.8	0.004	0.67	0.67
PUFA, % energy	5.6 ± 1.3	5.8 ± 1.4	0.13	0.68	0.68
Omega-3 FA, g/d	1.8 ± 0.7	1.7 ± 0.7	0.004	0.65	0.68
Protein, g/d	104 ± 34.3	95 ± 33.1	<0.001	0.63	0.68
Protein, % energy	17.1 ± 3.4	17.2 ± 3.4	0.49	0.71	0.70
Carbohydrate, g/d	288 ± 96.7	259 ± 96.1	<0.001	0.64	0.63
Carbohydrate, % energy	46.8 ± 7.6	46 ± 7.4	0.11	0.65	0.66
Total sugars, g/d	128 ± 47.8	117 ± 48.0	<0.001	0.66	0.72
Total sugars, % energy	21.1 ± 6.1	21 ± 5.9	0.83	0.73	0.73
Fiber, g/d	29.8 ± 12.1	26.8 ± 11.5	<0.001	0.71	0.73
Alcohol, g/d	10.4 ± 12.8	10.3 ± 13.7	0.83	0.89	0.89
Calcium, g/d	1225 ± 478	1111 ± 462	<0.001	0.63	0.69
Folate, µg/d	370 ± 131	338 ± 130	<0.001	0.65	0.70
Iron, mg/d	15.6 ± 5.1	14.2 ± 5	<0.001	0.62	0.63
Carotene, mg/d	6393 ± 5895	5546 ± 4103	0.005	0.7	0.71
Riboflavin, mg/d	2.3 ± 0.9	2.1 ± 0.9	0.001	0.71	0.76
Thiamin, mg/d	2.5 ± 2.3	2.4 ± 2.3	0.34	0.62	0.59
Vitamin B-6, mg/d	2.7 ± 0.9	2.5 ± 0.9	<0.001	0.67	0.69
Vitamin B-12, µg/d	7.7 ± 4.1	7.3 ± 4.1	0.06	0.73	0.75
Vitamin C, mg/d	167 ± 99.7	155 ± 94.3	0.04	0.73	0.76
Vitamin A, mg/d	1658 ± 1083	1506 ± 886	0.008	0.67	0.68
Retinol, µg/d	593 ± 451	582 ± 496	0.65	0.65	0.62
Vitamin D, µg/d	3.8 ± 2.3	3.5 ± 2	0.04	0.67	0.66
Vitamin E, mg/d	11.4 ± 4.3	10.4 ± 4.4	<0.001	0.67	0.70
Salt, g/d	7.2 ± 2.9	6.5 ± 2.7	<0.001	0.65	0.67
Sodium, mg/d	2896 ± 1144	2606 ± 1094	<0.001	0.65	0.67

1, Values represent means ± SD or percentages n=567; SFA, saturated fatty acids; MUFA, monounsaturated fatty acids; PUFA, polyunsaturated fatty acids; Omega-3 FA, Omega- 3 fatty acid; RE, retinol equivalents.

2, Mean difference between screening and baseline questionnaires was 2.7 ± 0.9 wk.

3, Multiple linear regression between screening and baseline FFQs adjusted for country, time of FFQ completion, age, sex and total energy at screening.

4, Spearman correlation coefficient (rho) between screening and baseline FFQs. All results were significant to P<0.001.

Table 3 Differences in the Food4Me food frequency questionnaire food group intakes (g/d) in European adults between screening and baseline as assessed using multiple linear regression and correlation coefficients¹

	Timepoint ²		P ³	Correlation coefficient ⁴	
	Screening	Baseline		Crude	Energy adjusted
Rice, pasta, grains and starches	76.2 ± 57.8	70.2 ± 56.5	0.08	0.52	0.45
Savouries (lasagne, pizza)	36.6 ± 33.3	34.7 ± 35.4	0.34	0.65	0.65
White bread (rolls, tortillas, crackers)	53 ± 95.4	44.2 ± 73.9	0.07	0.76	0.76
Wholemeal, brown breads and rolls	103 ± 131	86.3 ± 102	0.01	0.75	0.69
Breakfast cereals and porridge	56.9 ± 73	52.8 ± 73.4	0.35	0.81	0.80
Biscuits	28.1 ± 46.1	22.4 ± 40.8	0.03	0.61	0.60
Cakes, pastries and buns	15.7 ± 17.4	14.6 ± 16.8	0.34	0.57	0.54
Milk	185 ± 215	170 ± 199	0.21	0.7	0.66
Cheeses	38.5 ± 36.7	35.7 ± 35.5	0.17	0.64	0.67
Yogurts	70.9 ± 89.4	76.6 ± 119	0.27	0.66	0.61
Ice cream, creams and desserts	21.9 ± 22	21.5 ± 25.4	0.74	0.61	0.59
Eggs and egg dishes	30.8 ± 49.4	29.2 ± 41.9	0.55	0.75	0.68
Fats and oils (e.g. butter, low-fat spreads)	19.7 ± 17.3	18.5 ± 15.1	0.16	0.7	0.69
Potatoes and potato dishes	55.4 ± 56.6	53.1 ± 51.5	0.46	0.74	0.71
Chipped, fried & roasted potatoes	14.8 ± 16.8	15.5 ± 17.5	0.49	0.77	0.75
Peas, beans, lentils, vegetable dishes	31.9 ± 33.8	33.1 ± 47.5	0.56	0.79	0.78
Green vegetables	43.6 ± 49.9	38.9 ± 39.5	0.07	0.68	0.70
Carrots	22.6 ± 36.2	19.4 ± 20.5	0.11	0.67	0.66
Salad vegetables (e.g. lettuce)	51.2 ± 57.4	47.5 ± 46.6	0.06	0.77	0.78
Other vegetables (e.g. onions)	55.2 ± 50.2	51.8 ± 47.3	0.24	0.75	0.74
Tinned fruit or vegetables	2.2 ± 8.8	1.9 ± 6.3	0.45	0.42	0.46
Bananas	41.1 ± 50.5	37.6 ± 43.8	0.26	0.81	0.82
Other fruits (e.g. apples pears oranges)	246 ± 214	218 ± 196	0.02	0.8	0.81
Nuts and seeds, herbs and spices	4.8 ± 7.6	4.9 ± 9.4	0.91	0.68	0.67
Fish and fish products/dishes	48.3 ± 40.2	47 ± 42.2	0.60	0.75	0.73
Bacon and ham	18.1 ± 24.9	17.8 ± 27.3	0.81	0.76	0.73
Red meat (e.g. beef, veal, lamb, pork)	38.4 ± 36	36.8 ± 33.4	0.40	0.74	0.73
Poultry (chicken and turkey)	26.2 ± 36.2	22.7 ± 24.5	0.05	0.59	0.58
Meat products (e.g. burgers and sausages)	46 ± 53.1	40 ± 37.8	0.03	0.65	0.64
Alcoholic beverages	134 ± 173	139 ± 207	0.69	0.89	0.87
Sugars, syrups, preserves and sweeteners	4.7 ± 10.3	4.5 ± 9.1	0.66	0.85	0.81
Confectionary and savory snacks	16.7 ± 20.8	15.9 ± 21.7	0.56	0.71	0.64
Soups, sauces and miscellaneous foods	103 ± 80.9	92.7 ± 75.4	0.03	0.71	0.68
Teas and coffees	593 ± 505	579 ± 484	0.67	0.81	0.68
Other beverages (e.g. fruit juices, squash)	238 ± 289	223 ± 277	0.35	0.75	0.74

1, Values represent mean ± SD, n=567.

2, Mean difference between screening and baseline questionnaires was 2.7 ± 0.9 wk.

3, Multiple linear regression between screening and baseline FFQs adjusted for country, time of FFQ completion, age, sex and energy intake at screening;

4, Spearman correlation coefficient (rho) between screening and baseline FFQs. All results were significant to P<0.001.

Online Supporting Material

Supplemental Table 1 Food items included within each food category in the Food4Me FFQ

Food category	Number of food items listed within the group	Examples
Cereal	4	Porridge, readybrek Breakfast cereals, wholegrain e.g. branflakes
Bread and savoury biscuits	8	White bread Brown bread and seeded bread
Potatoes, rice and pasta	12	Potatoes - mashed, instant, roast Potatoes - boiled, jacket
Meat and Fish	24	Beef, venison (roast, steak, mince) Pork (roast, chops)
Dairy Products	19	Full-fat/whole milk, buttermilk Low-fat or semi-skimmed milk
Fats and Spreads	7	Butter Block/hard margarine e.g. stork/krona
Sweets and snacks	18	Sweet biscuits, chocolate e.g. digestive, cookies Plain cakes e.g. fruit, sponge, scones, gingerbread
Soups, sauces and spreads	10	Creamy soups e.g. chowder, cream of mushroom Non-creamy soups e.g. minestrone, vegetable
Drinks	15	Tea (black, green, fruit, herbal) Coffee, milky, latte, cappuccino
Fruit	12	Apples Pears
Vegetables	28	Carrots Butternut squash, pumpkin

Online Supporting Material

Supplemental Table 2 Concordance (%) in quartile classification of total energy and nutrient intakes between administration of the Food4Me food frequency questionnaire at screening and baseline¹

	Exact classification ²		Exact classification plus adjacent ³		Misclassification ⁴		Extreme misclassification ⁵	
	Crude	Energy adjusted	Crude	Energy adjusted	Crude	Energy adjusted	Crude	Energy adjusted
Total energy, kcal/d	44.8	-	86.1	-	10.8	-	3.2	-
Total fat, g/d	43.6	44.6	83.4	85.0	14.1	11.6	2.5	3.4
Total fat, % energy	44.3	46.2	84.7	84.8	13.4	12.9	1.9	2.3
SFA, g/d	47.3	48.0	85.4	87.7	12.5	9.7	2.1	2.6
SFA, % energy	48.7	49.1	86.2	86.7	11.1	11.5	2.6	1.8
MUFA, g/d	45.5	53.3	86.1	89.6	11.8	8.6	2.1	1.8
MUFA, % energy	51.0	50.8	90.3	90.7	8.1	7.9	1.6	1.4
PUFA, g/d	45.9	45.9	86.9	86.9	11.8	11.8	1.2	1.2
PUFA, % energy	47.3	48.1	88.4	88.5	10.1	9.9	1.6	1.6
Omega-3 FA, g/d	50.4	50.1	85.7	89.4	12.2	9.0	2.1	1.6
Protein, g/d	44.3	51.7	86.4	89.1	12.0	8.8	1.6	2.1
Protein, % energy	53.1	51.1	91.0	90.5	7.4	8.3	1.6	1.2
Carbohydrate, g/d	50.8	46.6	85.4	86.6	12.2	11.1	2.5	2.3
Carbohydrate, % energy	49.7	48.7	88.0	88.0	9.3	9.7	2.6	2.3
Total sugars, g/d	48.0	53.3	86.9	89.6	10.8	9.3	2.3	1.1
Total sugars, % energy	52.2	51.9	91.5	91.0	7.2	7.6	1.2	1.4
Fiber, g/d	51.0	55.2	89.6	92.1	9.5	6.3	0.9	1.6
Alcohol, g/d	70.0	66.5	98.1	97.5	1.8	2.5	0.2	0.0
Calcium, mg/d	47.1	48.9	87.3	88.5	9.5	9.2	3.2	2.3
Folate, µg/d	48.7	50.6	86.9	90.7	11.1	7.8	1.9	1.6
Iron, mg/d	46.7	47.3	84.7	87.5	12.7	10.4	2.6	2.1
Carotene, µg/d	52.6	50.8	89.8	89.9	9.3	8.6	0.9	1.4
Riboflavin, mg/d	50.8	56.3	91.0	92.9	7.9	6.3	1.1	0.7
Thiamin, mg/d	49.7	51.7	86.1	85.7	11.1	9.9	2.8	4.4
Vitamin B6, mg/d	47.1	51.7	89.2	91.0	9.2	7.1	1.6	1.9
Vitamin B12, µg/d	55.6	54.1	91.7	91.9	7.4	7.1	0.9	1.1
Vitamin C, mg/d	54.1	54.5	90.5	92.2	8.5	7.1	1.1	0.7
Vitamin A RE, µg/d ¹	52.7	53.4	87.7	88.0	10.9	10.2	1.4	1.8
Retinol, mcg/d	51.7	52.2	87.7	85.7	8.8	10.4	3.5	3.9
Vitamin D, µg/d	50.4	50.6	87.5	89.8	10.8	8.1	1.8	2.1
Vitamin E, mg/d	49.0	52.0	87.7	89.9	10.4	7.8	1.9	2.3
Salt, g/d	47.8	52.7	85.9	88.5	12.7	8.8	1.4	2.6
Sodium, mg/d	47.8	52.0	85.9	87.7	12.7	9.0	1.4	3.3

1, SFA, saturated fatty acids; MUFA, monounsaturated fatty acids; PUFA, polyunsaturated fatty acids; Omega-3 FA, Omega- 3 fatty acid; RE, retinol equivalents. Values represent percentages n=567. Mean difference between screening and baseline questionnaires was 2.7 ± 0.9 wks. The expected value for a 4-category model is 25% exact agreement by chance.

2, Percentage of participants classified into the same quartile.

3, Percentage of participants classified into the same plus the adjacent quartile.

4, Percentage of participants classified two quartiles apart.

5, Percentage of participants classified three quartiles apart.

Online Supporting Material

Supplemental Table 3 Concordance (%) in quartile classification of the Food4Me Food frequency questionnaire food group intakes between screening and baseline¹

	Exact classification ²		Exact classification plus adjacent ³		Misclassification ⁴		Extreme misclassification ⁵	
	Crude	Energy adjusted	Crude	Energy adjusted	Crude	Energy adjusted	Crude	Energy adjusted
Rice, pasta, grains and starches	45.3	43.0	81.3	79.9	14.5	15.3	4.2	4.8
Savouries (lasagne, pizza)	48.7	46.7	88.7	88.5	9.5	9.9	1.8	1.6
White bread (rolls, tortillas, crackers)	55.0	57.1	91.7	91.5	7.6	7.9	0.7	0.5
Wholemeal, brown breads and rolls	57.0	52.0	91.7	90.1	7.8	8.5	0.5	1.4
Breakfast cereals and porridge	67.4	65.8	93.8	94.9	4.8	3.5	1.4	1.6
Biscuits	50.6	48.7	85.7	85.7	10.9	11.5	3.4	2.8
Cakes, pastries and buns	47.1	47.4	85.0	84.1	11.8	11.6	3.2	4.2
Milk	52.2	49.4	88.5	86.9	10.4	11.8	1.1	1.2
Cheeses	48.5	48.3	86.9	87.1	10.8	11.0	2.3	1.9
Yogurts	53.6	51.5	88.2	86.2	10.1	10.4	1.8	3.4
Ice cream, creams and desserts	47.8	47.3	86.1	85.9	11.5	10.4	2.5	3.7
Eggs and egg dishes	57.5	52.2	93.8	87.8	6.0	10.4	0.2	1.8
Fats and oils (e.g. butter, low-fat spreads)	53.8	52.9	87.3	87.3	11.6	11.1	1.1	1.6
Potatoes and potato dishes	55.6	53.4	91.0	89.9	8.3	8.8	0.7	1.2
Chipped, fried & roasted potatoes	59.6	54.9	93.1	90.8	6.3	9.2	0.5	0.0
Peas, beans, lentils, vegetable dishes	60.8	58.2	92.9	93.8	6.0	5.1	1.1	1.1
Green vegetables	52.4	49.8	88.0	100.0	10.2	0.0	1.8	0.0
Carrots	54.9	52.6	89.1	87.7	9.2	10.1	1.8	2.3
Salad vegetables (e.g. lettuce)	54.9	56.8	93.5	93.1	5.5	5.8	1.1	1.1
Other vegetables (e.g. onions)	53.3	55.0	92.4	90.8	6.9	7.9	0.7	1.2
Tinned fruit or vegetables	54.8	45.9	80.6	81.7	12.9	11.8	5.9	6.5
Bananas	63.7	61.6	95.1	93.5	4.6	5.6	0.4	0.9
Other fruits (e.g. apples pears oranges)	61.9	63.3	94.2	95.4	4.9	3.9	0.9	0.7
Nuts and seeds, herbs and spices	55.9	52.2	87.3	87.1	10.8	11.1	1.9	1.8
Fish and fish products/dishes	53.3	53.8	91.0	90.7	8.5	8.5	0.5	0.9
Bacon and ham	56.1	52.0	93.3	92.1	5.6	7.1	1.1	0.9
Red meat (e.g. beef, veal, lamb, pork)	54.1	54.0	92.1	89.9	7.2	9.3	0.7	0.7
Poultry (chicken and turkey)	53.3	48.7	87.5	83.4	10.4	12.5	2.1	4.1
Meat products (e.g. burgers and sausages)	49.2	50.6	85.5	86.1	12.3	12.7	2.1	1.2
Alcoholic beverages	70.5	64.6	97.9	97.0	1.9	3.0	0.2	0.0
Sugars, syrups, preserves and sweeteners	60.3	60.2	93.3	93.2	5.9	5.6	0.8	1.2
Confectionary and savoury snacks	52.9	49.0	90.8	87.6	7.1	10.4	2.1	2.0
Soups, sauces and miscellaneous foods	51.1	50.6	89.8	89.4	9.0	8.8	1.2	1.8
Teas and coffees	64.6	64.6	94.2	94.2	5.1	5.1	0.7	0.7
Other beverages (e.g. fruit juices, squash)	56.4	56.3	91.7	91.0	6.7	7.8	1.6	1.2

1, Values represent percentages n=567. Mean difference between screening and baseline questionnaires was 2.7 ± 0.9 weeks. The expected value for a 4-category model is 25% exact agreement by chance.

2, Percentage of participants classified into the same quartile.

3, Percentage of participants classified into the same plus the adjacent quartile.

4, Percentage of participants classified two quartiles apart.

5, Percentage of participants classified three quartiles apart.

Online Supporting Material

Supplemental Table 4 Differences in total energy intake between the Food4Me food frequency questionnaire at screening and baseline, Spearman correlation coefficients (SCC, rho) and cross-classifications of quartiles by subgroup¹

	n	Energy intake (kcal/day)		P ³	SCC ⁴	Quartiles %			
		Screening	Baseline			Exact classification ⁵	Exact classification plus adjacent ⁶	Misclassification ⁷	Extreme misclassification ⁸
Country ²									
Greece	160	2376 ± 676	2056 ± 708	<0.001	0.54	44.4	81.3	13.1	5.6
Ireland	70	2625 ± 642	2546 ± 652	0.476	0.61	42.9	92.9	7.1	0
Netherlands	108	2556 ± 695	2393 ± 736	0.057	0.73	49.1	94.4	4.6	0.9
Poland	153	2411 ± 740	2201 ± 796	0.006	0.60	43.8	81.0	14.4	4.6
United Kingdom	49	2353 ± 584	2285 ± 584	0.528	0.70	42.9	87.8	12.2	0
Germany	27	2518 ± 562	2178 ± 530	0.015	0.59	42.9	85.7	7.1	7.1
FFQ completion period ⁹									
Short	189	2342 ± 671	2233 ± 714	0.001	0.64	43.9	85.7	11.1	3.2
Medium	189	2468 ± 688	2207 ± 705	<0.001	0.66	45.0	87.3	9.5	3.2
Long	189	2465 ± 699	2296 ± 771	0.057	0.61	45.5	85.2	11.6	3.2
BMI category									
Underweight & normal	296	2331 ± 602	2164 ± 662	<0.001	0.68	42.2	85.8	11.5	2.7
Overweight	192	2528 ± 753	2259 ± 739	<0.001	0.60	45.3	87.5	9.4	3.1
Obese	79	2743 ± 794	2520 ± 878	0.065	0.50	53.2	83.5	11.4	5.1
Age group									
Under 45 years	359	2456 ± 690	2230 ± 718	<0.001	0.65	43.7	85.5	11.4	3.1
Over 45 years	208	2453 ± 679	2273 ± 752	0.007	0.62	46.6	87.0	9.6	3.4
Sex									
Male	233	2803 ± 696	2542 ± 817	<0.001	0.62	45.9	87.6	9.9	2.6
Female	334	2213 ± 563	2039 ± 581	<0.001	0.61	44.0	85.0	11.4	3.6


1, Values represent means ± SD or percentages. Mean difference between screening and baseline questionnaires was 2.7 ± 0.9 weeks. The expected value for a 4-category model is 25% exact agreement by chance.

2, Spain was excluded from the analysis due to a lack of participants (n=5) completing the two FFQs within the acceptable time frame (1 month).

3, Multiple linear regression tested for significant differences in energy intakes between screening and baseline FFQs (analyses were stratified by country, FFQ completion period, BMI category, age group and sex). Models were adjusted for country, time of FFQ completion, age and sex (except when used a stratifying variable).

Online Supporting Material

- 4, Unadjusted Spearman correlation coefficients (ρ) between screening and baseline FFQs. All results were significant to $P < 0.001$.
- 5, Percentage of participants classified into the same quartile.
- 6, Percentage of participants classified into the same plus the adjacent quartile.
- 7, Percentage of participants classified two quartiles apart.
- 8, Percentage of participants classified three quartiles apart.
- 9, Short: 0-15.65 days; Medium: 15.66 – 22.63 days; Long: 22.64 – 31 days.



Supplemental Figure 1 Differences in percentage of energy from fat, carbohydrates and protein between administration of the Food4Me food frequency questionnaire at screening and baseline. Data represent delta in the total cohort (n=567) and in valid reporters (n=437). TE, Total energy; CHO, carbohydrate. Mean difference between screening and baseline questionnaires was 2.7 ± 0.9 weeks.