

Exploring the feasibility of an exercise programme including aerobic and resistance training in people with limited cutaneous systemic sclerosis.

MITROPOULOS, Alexandros, GUMBER, Anil, CRANK, Helen, AKIL, Mohammed and KLONIZAKIS, Markos <<http://orcid.org/0000-0002-8864-4403>>

Available from Sheffield Hallam University Research Archive (SHURA) at:
<https://shura.shu.ac.uk/25706/>

This document is the Supplemental Material

Citation:

MITROPOULOS, Alexandros, GUMBER, Anil, CRANK, Helen, AKIL, Mohammed and KLONIZAKIS, Markos (2020). Exploring the feasibility of an exercise programme including aerobic and resistance training in people with limited cutaneous systemic sclerosis. *Clinical Rheumatology*, 39 (6), 1889-1898. [Article]

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

Appendix A

Feeling Scale (FS)

While participating in exercise, it is common to experience changes in mood. Some individuals find exercise pleasurable, whereas others find it to be unpleasant. Additionally, feeling may fluctuate across time. That is, one might feel good and bad a number of times during exercise. Scientists have developed this scale to measure such responses.

+5 Very good

+4

+3 Good

+2

+1 Fairly good

0 Neutral

-1 Fairly bad

-2

-3 Bad

-4

-5 Very bad