

The acute physiological effects of high- and low-velocity resistance exercise in older adults

RICHARDSON, DL, DUNCAN, MJ, JIMENEZ GUTIERREZ, Alfonso, JONES, VM, JURIS, PM and CLARKE, ND

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/26558/>

This document is the Accepted Version [AM]

Citation:

RICHARDSON, DL, DUNCAN, MJ, JIMENEZ GUTIERREZ, Alfonso, JONES, VM, JURIS, PM and CLARKE, ND (2018). The acute physiological effects of high- and low-velocity resistance exercise in older adults. *European Journal of Ageing*, 15, 311-319. [Article]

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

The Acute Physiological Effects of High and Low Velocity Resistance Exercise in Older Adults

Darren L. Richardson ^a, Michael J. Duncan ^a, Alfonso Jimenez, ^a, Victoria M. Jones, ^a Paul M. Juris, ^b Neil D. Clarke, ^a

^a Centre for Applied Biological & Exercise Sciences, School of Life Sciences, Coventry University, Coventry, UK.

^b Department of Kinesiology, University of Massachusetts Amherst, 30 Eastman Lane, Amherst, MA

Corresponding Author: Richardson, D.L. **Email:** Richa190@uni.coventry.ac.uk

Postal Address: Life Sciences, Faculty Health and Life Sciences, Coventry University, Priory Street, CV1 5FB, UK

Word Count: 3,894

Abstract

The aim of the present study was to determine if workload matched, high velocity (HVE) and low velocity (LVE) resistance exercise protocols, elicit differing acute physiological responses in older adults. 10 older adults completed three sets of eight exercises on six separate occasions (three HVE and three LVE sessions). Systolic blood pressure, diastolic blood pressure and blood lactate were measured pre- and post-exercise, heart rate was measured before exercise and following each set of each exercise. Finally, rating of perceived exertion was measured following each set of each exercise. There were no significant differences in blood lactate ($F_{(1,9)}=0.028$; $P=0.872$; $\eta_p^2 = 0.003$), heart rate ($F_{(1,9)}= 0.045$; $P=0.837$; $\eta_p^2 = 0.005$), systolic blood pressure ($F_{(1,9)}= 0.023$; $P=0.884$; $\eta_p^2 = 0.003$) or diastolic blood pressure ($F_{(1,9)}= 1.516$; $P=0.249$; $\eta_p^2 = 0.144$) between HVE and LVE. However, LVE elicited significantly greater ratings of perceived exertion compared to HVE ($F_{(1,9)}=13.059$; $P=0.006$; $\eta_p^2 = 0.592$). The present workload matched HVE and LVE protocols produced comparable physiological responses, although greater exertion was perceived during LVE.

Keywords: Ageing; Physical activity; Health education; Older adults

1 Introduction

2
3 Ageing is associated with the loss of skeletal muscle mass known as sarcopenia and the loss of
4 muscle strength known as dynapenia (Clark and Manini 2008), both of which contribute to
5 disability, frailty, comorbidities, hospital admissions and death in older adults (Yu 2015). In
6 addition to ageing, a lack of physical activity has been identified as playing a significant role
7 in the loss of muscle size and strength (Cruz-Jentoft and Landi 2014), contributing to functional
8 decline and loss of independence in older adults (Clark and Manini 2008). To effectively
9 address such issues requires a multidisciplinary approach, comprising aspects of both exercise
10 prescription and nutritional strategies (Cruz-Jentoft and Landi 2014). Within exercise
11 prescription, one approach that has been explored is resistance exercise. Resistance exercise
12 has been shown to be effective in attenuating age related declines in muscle strength (Liu and
13 Latham 2009), whilst having beneficial effects on functional status, health and quality of life
14 in older adults (Hunter et al. 2004).

15 The fact that resistance exercise has been shown to have these positive effects, has led to major
16 health organisations such as the American College of Sports Medicine (ACSM), developing
17 resistance exercise guidelines for older adults. These ACSM guidelines state that 10-15
18 repetitions of 8-10 exercises that target the major muscle groups should be performed on two
19 or more nonconsecutive days per week, partnered with other activities that improve flexibility
20 and balance (Nelson et al. 2007). These are similar to the physical activity guidelines in the
21 United kingdom (UK) (Bull et al. 2010). However, as these guidelines are so brief, it is
22 unsurprising that few older adults in the UK are meeting them (Jefferis et al. 2014). Therefore,
23 there is a need for these physical activity guidelines to be expanded upon to provide more
24 guidance to older adults.

25 An important step in providing more guidance, is to understand the most pertinent mode of
26 resistance exercise for producing positive effects on functional status, strength and muscle
27 mass in older adults. Early investigation into resistance exercise identified the importance of
28 muscle strength for functional performance in older adults (Aniansson et al. 1980). More
29 recently, it has been highlighted that muscle power may be more relevant to functional
30 performance, as being able to move a limb fast against a low external resistance (e.g. moving
31 a limb quickly to stabilise to avoid a fall) is more useful than being able to move a limb slowly
32 against a high external resistance (Sayers and Gibson 2014). This has led to investigation into
33 the influence of high velocity (HVE) and/or low velocity (LVE) resistance exercise (also

referred to as power and strength training respectively) on functional performance (Ramirez-Campillo et al. 2014), muscle mass (Van Roie et al. 2013) and strength gains (Marsh et al. 2009). Yet, despite numerous investigations, the most effective mode of resistance exercise remains unclear (Tschopp et al. 2011).

Surprisingly, it appears there has been little consideration of the acute physiological changes that resistance exercise may facilitate in older adults, with the few studies that have, focusing on hormonal changes (Hakkinen and Pakarinen 1995; Marcell et al. 1999). It is well reported that the physiological mechanisms that are stimulated during resistance exercise are dependent on the nature of that exercise (e.g. sets, repetitions, velocity, mode etc.) with repeated exposure to a certain exercise stimulus, facilitating specific adaptations of those physiological mechanisms (Kraemer et al. 1988). It has been shown that the assessment of acute physiological responses to resistance exercise protocols can aid in understanding how they differ (Kraemer et al. 1996) and may be useful in explaining the mechanisms of potential adaptations (Ramirez-Campillo et al. 2014). Such investigation is important to better understand the utility and safety of each type of resistance exercise for exercise prescription in older adults.

As ageing negatively influences the structure and function of the cardiovascular system, arteries, peripheral circulation and the autonomic nervous system (Queiroz et al. 2010), the effect resistance exercise can have on blood pressure is a significant concern for older adults. At the time of performing resistance exercise, there can be very large increases in blood pressure (MacDougall et al. 1985) but following cessation, blood pressure can decrease below that of baseline, also known as post-exercise hypotension (Hurley and Gillin 2015). However, it is unclear if factors such as: frequency, intensity, time, mode and volume have an effect on blood pressure following exercise (Hurley and Gillin 2015), meaning the differences between HVE and LVE in older adults are not well understood. Additionally, other useful measures can be derived from blood pressure data, such as mean arterial pressure which has been shown to be a predictor of cardiovascular disease (Sesso et al. 2000) and combined with heart rate, rate pressure product which can be used as a measure of myocardial oxygen demand and cardiac workload (Hermida et al. 2001).

Differing intensity, load and velocity of resistance exercise has been shown to have a varying influence on blood lactate responses in young men, with greater exercise intensity showing a

greater increase in blood lactate than low intensity (Arazi et al. 2014). However, between studies it is hard to compare physiological responses, as protocols have varied in combinations of intensity, number of sets, rest times and velocity of movement (Arazi et al. 2014). Mazzetti et al. (2007) observed that LVE elicited a greater lactate response than HVE whereas, Nitzsche et al. (2017) observed that both blood lactate and heart rate responses were similar following three different resistance exercise protocols that varied in load, repetitions, number of sets and rest times.

As prior research has not fully considered whether velocity of resistance exercise elicits different acute physiological responses in older adults, the optimal prescription of resistance exercise in this population remains to be fully elucidated. Therefore, an important first step is to examine acute physiological markers such as heart rate, blood pressure and blood lactate. Furthermore, perception of exercise intensity is related to physiological demand, and the subsequent feelings of exertion that occur as a consequence of exercise intensity, may influence exercise adherence (Ekkekakis et al. 2005). Hence, monitoring rating of perceived exertion (RPE) would provide guidance on the perceptual response to both HVE and LVE. Such data is key in better refining resistance exercise programming for older adults, and informing health care professionals on how physiological and perceptual responses vary with velocity of resistance exercise. Therefore, the aims of this study were to measure the physiological and perceptual responses of a group of older adults to workload matched HVE and LVE protocols. We hypothesised that both physiological responses and RPE would be greater during LVE compared to HVE.

Materials and Methods

Participants

The present study used a randomised, counterbalanced, crossover study design and following institutional ethics approval by the local ethics committee, 10 recreationally active older adults (five males and five females; Table 1) were recruited by word of mouth for participation. All participants were made aware of the exercise protocols and associated risks before providing informed consent, and completing a health screen questionnaire prior to each trial. After providing details of any current medications, each participant was required to meet strict inclusion criteria, namely: the absence of cognitive impairment (Mini-Mental State Examination score < 23) (Folstein et al. 1975), acute or terminal illness, myocardial infarction, upper or lower extremity fracture in the previous six months, symptomatic coronary artery

disease, congestive heart failure, uncontrolled hypertension (>150/90 mmHg), neuromuscular disease and not undergoing hormone replacement therapy (Reid et al. 2015). Finally, participants should not have had participated in any purposeful strength or power training in the previous six months (de Vos et al. 2005) to be eligible to take part.

Table 1. Participant characteristics

Participant Information	Males ($n=5$)	Females ($n=5$)
Age (years)	66 \pm 3	68 \pm 2
Age Range (years)	63-71	67-71
Height (cm)	174.5 \pm 5.4	162.6 \pm 5.8
Body Mass (kg)	89.4 \pm 13.6	70.9 \pm 10.7
Body Mass Index (kg/m ²)	29 \pm 4	27 \pm 3
Baseline Systolic Blood Pressure (mmHg)	141 \pm 9	140 \pm 7
Baseline Diastolic Blood Pressure (mmHg)	80 \pm 6	81 \pm 6
Baseline Mean Arterial Pressure (mmHg)	100 \pm 7	101 \pm 6
Medications Taken	1 \pm 1	1 \pm 1
Mini-Mental State Examination score (0-30)	29 \pm 1	29 \pm 1

Values are means \pm SD; n = number of participants

Familiarisation

Prior to familiarisation and all trials, participants were asked to refrain from caffeine use for a minimum of 12 hours (Syed et al. 2005) and any other fatiguing exercise or physical activity for 24 hours. Firstly, height (cm) and mass (kg) were recorded (Seca Instruments, Hamburg, Germany). Participants then completed a warm-up protocol which consisted of five minutes self-selected paced cycling (Marsh et al. 2009) followed by five dynamic stretches which targeted the main muscle groups and joints used in the programme (Miszko et al. 2003). This warm-up was repeated before all subsequent trials. Following the warm-up, the correct,

individual anthropometric setup for each exercise was noted on each piece of Cybex exercise equipment (Cybex, Medway, MA, USA). The correct technique for all exercises were then demonstrated to participants and practiced. Finally, participants were taken through a predictive 1-RM (one repetition maximum) testing protocol for each exercise, which provided a prediction of the maximum amount of weight, that could be lifted for just one repetition. Participants performed repetitions on a weight they felt was challenging but manageable. The resistance was progressively increased until no more than 10 repetitions could be performed with correct form. If a participant could complete more than 10 repetitions before failure, three minutes of rest was given, the weight increased by 10-15% and the process repeated. Weight lifted and number of repetitions completed were used to provide an estimation of 1-RM for each exercise (Table 2) using the prediction equation: $(\text{weight lifted} \div (1.0278 - (0.0278 \times \text{number of repetitions performed})))$ (Brzycki 1993).

Exercise protocol

For clarity, when discussing the exercise protocols, the word trial is used to describe each visit to the sports centre, and set is used to describe the collection of single repetitions (one complete movement of an exercise). Participants were randomised to complete one of the two workload matched protocols (identical total weight lifted) displayed in Table 2. Both protocols consisted of three sets of eight different exercises (chest press, leg press, leg extension, leg curl, calf raise, seated row, bicep curl and tricep extension). Participants had three days of rest between each of the three trials for each velocity of training and a week ‘washout period’ before crossing over to the other protocol, meaning the trial period lasted approximately five weeks. The exercise protocols used in the present study (described in Table 2) were based on others that have previously demonstrated a positive impact on functional performance in older adults (Beltran Valls et al. 2014; Brochu et al. 2002; Kalapotharakos et al. 2005; Reid et al. 2015) with the number of sets and repetitions being similar to others that have attempted to match workloads (Hortobagyi et al. 2001; Sayers and Gibson 2014).

The concentric phase (lifting of the weight) in the HVE group was performed “as fast as possible” without causing dangerous fly away (unloading) of the weight stack, and the eccentric phase (lowering of the weight) was performed over three seconds (Henwood et al. 2008). The LVE group performed the concentric phase over two seconds and the eccentric phase over three seconds (Van Roie et al. 2013). A metronome was used to provide the cadence for exercise, except during the concentric phase of the HVE protocol. Each participant

performed all their trials as near to the same time of day as possible to reduce fluctuations in strength due to circadian variation (Duncan and Oxford 2011).

Table 2. 1-RM data and details of the exercise protocols

Exercises	1-RM Males (kg)	1-RM Females (kg)	HVE Protocol	LVE Protocol
Leg Press	130.2 ± 29.5	78.9 ± 12.5	40% 1-RM	80% 1-RM
Seated Row	62.6 ± 7.5	33.8 ± 4.8	3 sets	3 sets
Chest Press	54.4 ± 5.3	21.4 ± 2.6	14 repetitions	7 repetitions
Leg Extension	58.8 ± 16.1	29.1 ± 7.2	Concentric phase “as fast as possible” with 3 second eccentric	2 second concentric phase and 3 second eccentric phase
Leg Curl	51.6 ± 9.1	25.6 ± 4.0		
Calf Raise	117.7 ± 27.2	89.1 ± 19.9	2 minutes rest between sets	2 minutes rest between sets
Tricep Extension	36.0 ± 6.9	15.5 ± 6.8	3 minutes between exercises	3 minutes between exercises
Bicep Curl	30.3 ± 7.6	12.5 ± 6.4		

Values are means ± SD; HVE = High Velocity Exercise; LVE = Low Velocity Exercise; 1-RM = One repetition maximum

Physiological measurements

Systolic and diastolic blood pressure were measured with an automatic blood pressure monitor (Omron M3 Intellisense HEM-7200-E, Omron Matsusaka Co Ltd, Kyoto, Japan) from the left arm, while seated in an upright position, prior to every trial and immediately following the last exercise of each session. Mean arterial pressure ($2 \times \text{diastolic blood pressure} + \text{systolic blood pressure}$)/3 and rate pressure product ($\text{systolic blood pressure} \times \text{heart rate}$) were calculated prior to and post-exercise using the blood pressure data. A fingertip blood sample was collected via a capillary tube, prior to each session and immediately at the end of the session. Samples were then analysed using a blood lactate analyser (Biosen C-line clinic, EKF Diagnostics, Magdeburg, Germany). Finally, heart rate was measured using heart rate telemetry (Polar Electro Oy, Kempele, Finland) before exercise and immediately following each set of each exercise.

Perceptual measure

RPE (Borg 1982) was used to assess the intensity that participants perceived during each of the exercise protocols. The 15-point RPE scale ranges from 6 (no exertion) to 20 (maximal exertion) and was presented to participants following each set of each exercise for both HVE and LVE, so that a value from the scale could be given to represent the exertion they perceived in that moment. Figure 1 displays the approximate timescale of the sessions and when various

measures were collected. As the LVE protocol had half the amount of repetitions as the HVE protocol, each LVE trial was approximately eight minutes shorter in duration than the HVE trials.




Figure 1. A schematic diagram of the experimental protocol

X3 = Collected following all three sets

Statistical Analysis

All data was analysed using IBM SPSS Statistics for Windows, Version 22.0 (Armonk, NY: IBM Corp) and descriptive statistics are presented as mean \pm SD and 95% confidence intervals (95% CI). Factorial analysis of variance (ANOVA) with repeated measures were used to compare the dependent variables heart rate, blood pressure, blood lactate and RPE with the independent variable, exercise velocity. Within group changes were further investigated using repeated measures ANOVA and T-tests with Bonferroni correction where necessary. When Mauchley's test of sphericity was significant and the Greenhouse-Geisser level of violation was >0.75 , degrees of freedom were corrected using Huynh-Feldt adjustment. When violation was <0.75 , Greenhouse-Geisser correction was used. Where any differences were found, pairwise comparisons with Bonferroni correction were used to show exactly where they lay. Significance was determined by a P value of <0.05 and reported as exact values unless below $P=0.001$. Effect size was used to quantify the meaningfulness of any differences found between conditions, it was calculated using η_p^2 and defined as: trivial (<0.1), small ($0.1-0.29$), moderate ($0.3-0.49$) or large ($0.5>$) (Hopkins et al. 2009). An *a priori* power calculation suggested that

a sample size of ten participants would be necessary to detect a statistical difference given an estimated effect size of 0.25, a $1-\beta$ error probability of 0.90 and a P value significance level less than 0.05.

Results

Blood lactate

There were trivial differences in blood lactate concentrations between HVE and LVE ($F_{(1,9)}=0.028$; $P=0.872$; 95% CI: -0.7, 0.6; $\eta_p^2=0.003$; Table 3) but large increases in blood lactate concentrations from pre- to post-exercise regardless of velocity ($F_{(1,9)}=13.828$; $P=0.005$; 95% CI: 0.9, 3.7; $\eta_p^2=0.61$).

Systolic Blood Pressure

There were trivial differences in systolic blood pressure between HVE and LVE ($F_{(1,9)}=0.023$; $P=0.884$; 95% CI: -5.6, 4.9; $\eta_p^2=0.003$; Table 3) and moderate increases in systolic blood pressure from pre- to post-exercise regardless of velocity ($F_{(1,9)}=4.068$; $P=0.074$; 95% CI: -0.6, 10.3; $\eta_p^2=0.31$).

Diastolic Blood Pressure

There were small differences in diastolic blood pressure during HVE and LVE ($F_{(1,9)}=1.516$; $P=0.249$; 95% CI: -1.1, 3.6; $\eta_p^2=0.14$; Table 3) and small differences between pre- and post-exercise regardless of velocity ($F_{(1,9)}=2.010$; $P=0.190$; 95% CI: -4.8, 1.1; $\eta_p^2=0.18$).

Mean Arterial Pressure

There were trivial differences in mean arterial pressure between HVE and LVE ($F_{(1,9)}=0.408$; $P=0.539$; 95% CI: -2.1, 3.5; $\eta_p^2=0.04$; Table 3) and trivial differences in mean arterial pressure between pre- and post-exercise regardless of velocity ($F_{(1,9)}=0.074$; $P=0.792$; 95% CI: -2.7, 3.4; $\eta_p^2=0.01$).

Rate Pressure Product

There were trivial differences in rate pressure product between HVE and LVE ($F_{(1,9)}=0.580$; $P=0.466$; 95% CI: -1329, 660; $\eta_p^2=0.06$; Table 3) and trivial differences between pre- and post-exercise regardless of velocity ($F_{(1,9)}=0.867$; $P=0.376$; 95% CI: -922, 2213; $\eta_p^2=0.09$).


Table 3. Physiological measures for both HVE and LVE for all trials

	HVE		LVE	
	Pre-Exercise	Post-Exercise	Pre-Exercise	Post-Exercise
Blood Lactate (mmol/l)	2.3 ± 1.2	4.3 ± 2.1	2.0 ± 0.8	4.6 ± 2.8
Systolic Blood Pressure (mmHg)	131.8 ± 14.5	138.7 ± 18.7	133.5 ± 17.4	136.3 ± 18.4
Diastolic Blood Pressure(mmHg)	75.0 ± 7.3	72.5 ± 7.3	75.6 ± 6.6	74.4 ± 8.9
Mean Arterial Pressure(mmHg)	93.9 ± 8.9	94.6 ± 9.6	94.9 ± 9.4	95.0 ± 10.2
Rate Pressure Product (mmHg.bpm)	12383 ± 1846	12720 ± 2853	11740 ± 2425	12694 ± 2392

Values are means ± SD; HVE = High Velocity Exercise; LVE = Low Velocity Exercise

Heart Rate

There was a significant interaction between velocity of exercise and different exercises ($F_{(7,63)}=8.841$; $P<0.001$; $\eta_p^2=0.50$; Figure 2). Repeated measures ANOVA revealed that there were significant differences in heart rate between exercises for both HVE ($F_{(7,63)}=10.202$; $P<0.001$; $\eta_p^2=0.53$) and LVE ($F_{(7,63)}=12.263$; $P<0.001$; $\eta_p^2=0.58$). Further investigation with T-tests revealed heart rate during the leg press ($P<0.001$; 95% CI: -8.4, -5.4) and seated row ($P<0.001$; 95% CI: -4.1, -1.2) were significantly higher during LVE compared to HVE. But for both the chest press ($P<0.001$; 95% CI: 3.4, 7.5) and leg extension ($P<0.001$; 95% CI: 2.6, 6.6), heart rate was significantly higher during HVE.



1 /

Figure 2. Heart rate (mean \pm SD) for all participants during HVE and LVE

*= LVE significantly greater than HVE

†= HVE significantly greater than LVE

HVE = High Velocity Exercise; LVE = Low Velocity Exercise

RPE

There was a significant interaction between velocity and perception of exercises ($F_{(7,63)}=6.184$; $P<0.001$; $\eta_p^2=0.41$; Figure 3). The interaction plot revealed that all exercises were perceived as harder during LVE compared with HVE except for the chest press. T-tests revealed that during LVE, participants rated RPE significantly greater for leg press ($P<0.001$; 95% CI: -3.4, -2.5), seated row ($P<0.001$; 95% CI: -2.0, -1.4), leg curl ($P<0.001$; 95% CI: -2.4, -1.5), and calf raise ($P<0.001$; 95% CI: -2.0, -1.1) than during HVE.




Figure 3. RPE (mean \pm SD) for all participants during HVE and LVE

*= LVE significantly greater than HVE

RPE = Rating of Perceived Exertion; HVE = High Velocity Exercise; LVE = Low Velocity Exercise

Discussion

The present study is novel as it reports the physiological and perceptual responses to workload matched HVE and LVE in a sample of older adults. These measures are important in understanding how the ageing biological system responds to these modes of resistance exercise. This information can then feed forward, recognising the effect of exercise is multifaceted and multidisciplinary in nature. We hypothesised that LVE would elicit both a greater physiological response and a greater RPE response than HVE. This hypothesis must be rejected as physiological responses were similar, but the RPE response was significantly greater during LVE. The findings of the present study suggest there are no significant differences between workload matched HVE and LVE in blood lactate, systolic blood pressure, diastolic blood pressure, mean arterial pressure or rate pressure product responses in older adults. As would be expected, heart rate varied between exercises significantly, due to body position (Achten and Jeukendrup 2003) and the varying blood demands of active muscle (Peçanha et al. 2013). The leg press and seated row elicited significantly greater heart rate responses during LVE, while the chest press and leg extension elicited significantly greater heart rate responses during HVE.

Although changes were not significant, HVE produced increases in systolic blood pressure of

1 approximately 10 mmHg in trials one and two from pre- to post-exercise, whereas LVE saw a
2 10 mmHg increase in trial one and trivial changes in trials two and three. A similar trend was
3 observed by da Silva et al. (2007) who examined acute systolic blood pressure changes
4 following three sets of maximum velocity bench press exercise in untrained older women, the
5 authors reported that blood pressure was significantly lower at baseline than after the first,
6 second, and third sets. This potential increase in systolic blood pressure is something that
7 practitioners should be aware of when designing resistance exercise programmes for older
8 adults, especially in populations at risk.

9 Previously, it has been reported that resistance exercise can have a post-exercise hypotensive
10 effect (Hardy and Tucker 1998). Although changes were not significant, it is important to note
11 that diastolic blood pressure decreased from pre- to post-exercise following both HVE and
12 LVE in the present study. As the participants were normotensive, and individuals with an
13 elevated blood pressure are those who experience the greatest post-exercise hypotensive effect
14 of resistance exercise (Cardoso et al. 2010), it is unsurprising that only insignificant decreases
15 were observed.

16 The main differences within the present study lay within the patterns observed for RPE. Despite
17 comparable physiological strain, RPE was significantly greater for four of the eight exercises
18 during LVE compared to HVE. These findings are consistent with Gearhart et al. (2002) who
19 also observed that rating of perceived exertion was significantly greater when workload
20 matched, heavier resistance exercise was performed for fewer repetitions compared with lighter
21 resistance exercise for more repetitions. Therefore, the findings of the present study may have
22 particular implications for exercise adherence, as the American College of Sports Medicine
23 state that when intensity of exercise is higher, exercise adherence is generally lower (Whaley
24 et al. 2006). Furthermore, other affective responses such as enjoyment of exercise, have been
25 shown to predict long-term adherence to exercise programmes (Ekkekakis et al. 2011;
26 Williams et al. 2008) meaning it would be beneficial for future studies to examine the affective
27 responses of HVE and LVE in older adults to establish the likelihood of long-term adherence.
28 Affective responses to exercise are particularly important to consider as it has been suggested
29 that individuals differ in the exercise intensities they can tolerate and prefer (Ekkekakis et al.
30 2005), meaning that the mode of resistance exercise that should be prescribed may also need
31 to consider individual preference.

Methodological considerations

It may have been useful to measure blood pressure during each exercise to observe if there were differences in blood pressure between LVE and HVE in addition to pre- and post-trial. Furthermore, monitoring blood pressure throughout recovery could have been useful to examine any potential post-exercise hypotensive effects. Lastly, measurement of the velocity of the HVE and LVE protocols would have assured an appreciable difference between protocols and provided some guidance on the range of velocity older adults are able to produce. This is especially important as it has recently been reported that there is a large variation in self-selected maximal limb velocity in such exercise protocols and improvements in functional performance might be optimised in individuals with the highest training velocities (Sayers et al. 2016).

Conclusion

Workload matched HVE and LVE produced comparable physiological responses in a group of older adults. While physiological responses were similar between velocities, LVE was perceived as harder, meaning it is possible that the affective responses to these velocities of exercise were different. Clear recommendations cannot be drawn from the findings of the present study, but HVE might be a more appealing mode of resistance exercise to propose to older adults, as it may produce the same physiological stimulus as LVE while being perceived as less exerting. Exercise practitioners and those working in community settings with older adults might therefore want to employ HVE preferentially given the link between RPE and continuation of exercise in the longer term. However, the investigation of the affective responses to both HVE and LVE would be useful in further clarifying general recommendations for older adults.

Conflict of Interest

None declared

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- Achten J, Jeukendrup AE (2003) Heart rate monitoring: applications and limitations Sports medicine (Auckland, NZ) 33:517-538
- Aniansson A, Rundgren A, Sperling L (1980) Evaluation of functional capacity in activities of daily living in 70-year-old men and women Scandinavian journal of rehabilitation medicine 12:145-154
- Arazi H, Mirzaei B, Heidari N (2014) Neuromuscular and metabolic responses to three different resistance exercise methods Asian journal of sports medicine 5:30
- Beltran Valls MR, Dimauro I, Brunelli A, Tranchita E, Ciminelli E, Caserotti P, Duranti G, Sabatini S, Parisi P, Parisi A, Caporossi D (2014) Explosive type of moderate-resistance training induces functional, cardiovascular, and molecular adaptations in the elderly Age (Dordrecht, Netherlands) 36:759-772 doi:10.1007/s11357-013-9584-1
- Borg GA (1982) Psychophysical bases of perceived exertion Medicine and science in sports and exercise 14:377-381
- Brochu M, Savage P, Lee M, Dee J, Cress ME, Poehlman ET, Tischler M, Ades PA (2002) Effects of resistance training on physical function in older disabled women with coronary heart disease Journal of applied physiology (Bethesda, Md : 1985) 92:672-678 doi:10.1152/jappphysiol.00804.2001
- Brzycki M (1993) Strength testing—predicting a one-rep max from reps-to-fatigue Journal of Physical Education, Recreation & Dance 64:88-90
- Bull F, Biddle S, Buchner D, Ferguson R, Foster C, Fox K, Haskell B, Mutrie N, Murphy M, Reilly J, Riddoch C, Skelton D, Stratton G, Tremblay M, Watts C (2010) Physical activity guidelines in the UK: review and recommendations. 2010 URL: http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_127931
- Cardoso CG, Jr., Gomides RS, Queiroz AC, Pinto LG, da Silveira Lobo F, Tinucci T, Mion D, Jr., de Moraes Forjaz CL (2010) Acute and chronic effects of aerobic and resistance exercise on ambulatory blood pressure Clinics (Sao Paulo, Brazil) 65:317-325 doi:10.1590/s1807-59322010000300013
- Clark BC, Manini TM (2008) Sarcopenia≠ dynapenia The Journals of Gerontology Series A: Biological Sciences and Medical Sciences 63:829-834
- Cruz-Jentoft AJ, Landi F (2014) Sarcopenia Clinical Medicine 14:183-186 doi:10.7861/clinmedicine.14-2-183
- da Silva RP, Novaes J, Oliveira RJ, Gentil P, Wagner D, Bottaro M (2007) High-velocity resistance exercise protocols in older women: effects on cardiovascular response Journal of sports science & medicine 6:560

- 1 de Vos NJ, Singh NA, Ross DA, Stavrinou TM, Orr R, Fiatarone Singh MA (2005) Optimal
2 load for increasing muscle power during explosive resistance training in older adults The
3 journals of gerontology Series A, Biological sciences and medical sciences 60:638-647
- 4 Duncan MJ, Oxford SW (2011) The effect of caffeine ingestion on mood state and bench
5 press performance to failure Journal of strength and conditioning research / National Strength
6 & Conditioning Association 25:178-185 doi:10.1519/JSC.0b013e318201bddd
- 7 Ekkekakis P, Hall EE, Petruzzello SJ (2005) Some like It Vigorous: Measuring Individual
8 Differences in the Preference for and Tolerance of Exercise Intensity Journal of Sport and
9 Exercise Psychology 27:350-374 doi:10.1123/jsep.27.3.350
- 10 Ekkekakis P, Parfitt G, Petruzzello SJ (2011) The pleasure and displeasure people feel when
11 they exercise at different intensities: decennial update and progress towards a tripartite
12 rationale for exercise intensity prescription Sports medicine (Auckland, NZ) 41:641-671
- 13 Folstein MF, Folstein SE, McHugh PR (1975) "Mini-mental state". A practical method for
14 grading the cognitive state of patients for the clinician Journal of psychiatric research 12:189-
15 198
- 16 Gearhart RF, Jr., Goss FL, Lagally KM, Jakicic JM, Gallagher J, Gallagher KI, Robertson RJ
17 (2002) Ratings of perceived exertion in active muscle during high-intensity and low-intensity
18 resistance exercise Journal of strength and conditioning research / National Strength &
19 Conditioning Association 16:87-91
- 20 Hakkinen K, Pakarinen A (1995) Acute hormonal responses to heavy resistance exercise in
21 men and women at different ages International journal of sports medicine 16:507-513
22 doi:10.1055/s-2007-973045
- 23 Hardy DO, Tucker LA (1998) The effects of a single bout of strength training on ambulatory
24 blood pressure levels in 24 mildly hypertensive men American journal of health promotion :
25 AJHP 13:69-72
- 26 Henwood TR, Riek S, Taaffe DR (2008) Strength versus muscle power-specific resistance
27 training in community-dwelling older adults The journals of gerontology Series A, Biological
28 sciences and medical sciences 63:83-91
- 29 Hermida RC, Fernandez JR, Ayala DE, Mojon A, Alonso I, Smolensky M (2001) Circadian
30 rhythm of double (rate-pressure) product in healthy normotensive young subjects
31 Chronobiology international 18:475-489
- 32 Hopkins WG, Marshall SW, Batterham AM, Hanin J (2009) Progressive statistics for studies
33 in sports medicine and exercise science Medicine and science in sports and exercise 41:3-13
34 doi:10.1249/MSS.0b013e31818cb278
- 35 Hortobagyi T, Tunnel D, Moody J, Beam S, DeVita P (2001) Low- or high-intensity strength
36 training partially restores impaired quadriceps force accuracy and steadiness in aged adults
37 The journals of gerontology Series A, Biological sciences and medical sciences 56:B38-47
- 38 Hunter GR, McCarthy JP, Bamman MM (2004) Effects of resistance training on older adults
39 Sports medicine (Auckland, NZ) 34:329-348

- 1 Hurley BF, Gillin AR (2015) Can Resistance Training Play a Role in the Prevention or
2 Treatment of Hypertension? In: Pescatello LS (ed) Effects of Exercise on Hypertension:
3 From Cells to Physiological Systems. Springer International Publishing, Cham, pp 25-46.
4 doi:10.1007/978-3-319-17076-3_2
- 5 Jefferis BJ, Sartini C, Lee I-M, Choi M, Amuzu A, Gutierrez C, Casas JP, Ash S, Lennnon
6 LT, Wannamethee SG (2014) Adherence to physical activity guidelines in older adults, using
7 objectively measured physical activity in a population-based study BMC public health 14:382
- 8 Kalapotharakos VI, Michalopoulos M, Tokmakidis SP, Godolias G, Gourgoulis V (2005)
9 Effects of a heavy and a moderate resistance training on functional performance in older
10 adults Journal of strength and conditioning research / National Strength & Conditioning
11 Association 19:652-657 doi:10.1519/15284.1
- 12 Kraemer WJ, Deschenes MR, Fleck SJ (1988) Physiological adaptations to resistance
13 exercise. Implications for athletic conditioning Sports medicine (Auckland, NZ) 6:246-256
- 14 Kraemer WJ, Fleck SJ, Evans WJ (1996) Strength and power training: physiological
15 mechanisms of adaptation Exercise and Sport Sciences Reviews 24:363-397
- 16 Liu CJ, Latham NK (2009) Progressive resistance strength training for improving physical
17 function in older adults The Cochrane database of systematic reviews:Cd002759
18 doi:10.1002/14651858.CD002759.pub2
- 19 MacDougall JD, Tuxen D, Sale DG, Moroz JR, Sutton JR (1985) Arterial blood pressure
20 response to heavy resistance exercise Journal of applied physiology (Bethesda, Md : 1985)
21 58:785-790
- 22 Marcell TJ, Wiswell RA, Hawkins SA, Tarpenning KM (1999) Age-related blunting of
23 growth hormone secretion during exercise may not be solely due to increased somatostatin
24 tone Metabolism: clinical and experimental 48:665-670
- 25 Marsh AP, Miller ME, Rejeski WJ, Hutton SL, Kritchevsky SB (2009) Lower extremity
26 muscle function after strength or power training in older adults Journal of aging and physical
27 activity 17:416-443
- 28 Mazzetti S, Douglass M, Yocum A, Harber M (2007) Effect of explosive versus slow
29 contractions and exercise intensity on energy expenditure Medicine and science in sports and
30 exercise 39:1291-1301 doi:10.1249/mss.0b013e318058a603
- 31 Miszko TA, Cress ME, Slade JM, Covey CJ, Agrawal SK, Doerr CE (2003) Effect of
32 strength and power training on physical function in community-dwelling older adults The
33 journals of gerontology Series A, Biological sciences and medical sciences 58:171-175
- 34 Nelson ME, Rejeski WJ, Blair SN, Duncan PW, Judge JO, King AC, Macera CA, Castaneda-
35 Sceppa C (2007) Physical activity and public health in older adults: recommendation from
36 the American College of Sports Medicine and the American Heart Association Medicine and
37 science in sports and exercise 39:1435-1445 doi:10.1249/mss.0b013e3180616aa2
- 38 Nitzsche N, Baumgärtel L, Weigert M, Neuendorf T, Fröhlich M, Schulz H (2017) Acute
39 Effects of Three Resistance Exercise Programs on Energy Metabolism International Journal
40 of Sports Science 7:22-35 doi:10.5923/j.sports.20170702.02.

- 1 Peçanha T, Vianna JM, Sousa ÉDd, Panza PS, Lima JRPd, Reis VM (2013) Influence of the
2 muscle group in heart rate recovery after resistance exercise *Revista Brasileira de Medicina*
3 *do Esporte* 19:275-279
- 4 Queiroz AC, Kanegusuku H, Forjaz CL (2010) Effects of resistance training on blood
5 pressure in the elderly *Arquivos brasileiros de cardiologia* 95:135-140
- 6 Ramirez-Campillo R, Castillo A, de la Fuente CI, Campos-Jara C, Andrade DC, Alvarez C,
7 Martinez C, Castro-Sepulveda M, Pereira A, Marques MC, Izquierdo M (2014) High-speed
8 resistance training is more effective than low-speed resistance training to increase functional
9 capacity and muscle performance in older women *Experimental gerontology* 58:51-57
10 doi:10.1016/j.exger.2014.07.001
- 11 Reid KF, Martin KI, Doros G, Clark DJ, Hau C, Patten C, Phillips EM, Frontera WR,
12 Fielding RA (2015) Comparative effects of light or heavy resistance power training for
13 improving lower extremity power and physical performance in mobility-limited older adults
14 *The journals of gerontology Series A, Biological sciences and medical sciences* 70:374-380
15 doi:10.1093/gerona/glu156
- 16 Sayers SP, Gibson K (2014) High-speed power training in older adults: a shift of the external
17 resistance at which peak power is produced *Journal of strength and conditioning research /*
18 *National Strength & Conditioning Association* 28:616-621
19 doi:10.1519/JSC.0b013e3182a361b8
- 20 Sayers SP, Gibson K, Bryan Mann J (2016) Improvement in functional performance with
21 high-speed power training in older adults is optimized in those with the highest training
22 velocity *European journal of applied physiology* doi:10.1007/s00421-016-3484-x
- 23 Sesso HD, Stampfer MJ, Rosner B, Hennekens CH, Gaziano JM, Manson JE, Glynn RJ
24 (2000) Systolic and diastolic blood pressure, pulse pressure, and mean arterial pressure as
25 predictors of cardiovascular disease risk in Men *Hypertension (Dallas, Tex : 1979)* 36:801-
26 807
- 27 Syed SA, Kamimori GH, Kelly W, Eddington ND (2005) Multiple dose pharmacokinetics of
28 caffeine administered in chewing gum to normal healthy volunteers *Biopharmaceutics &*
29 *drug disposition* 26:403-409 doi:10.1002/bdd.469
- 30 Tschopp M, Sattelmayer MK, Hilfiker R (2011) Is power training or conventional resistance
31 training better for function in elderly persons? A meta-analysis *Age and ageing* 40:549-556
32 doi:10.1093/ageing/afr005
- 33 Van Roie E, Delecluse C, Coudyzer W, Boonen S, Bautmans I (2013) Strength training at
34 high versus low external resistance in older adults: effects on muscle volume, muscle
35 strength, and force-velocity characteristics *Experimental gerontology* 48:1351-1361
36 doi:10.1016/j.exger.2013.08.010
- 37 Whaley MH, Brubaker PH, Otto RM, Armstrong LE (2006) *ACSM's Guidelines for Exercise*
38 *Testing and Prescription* Lippincott Williams & Wilkins, Philadelphia
- 39 Williams DM, Dunsiger S, Ciccolo JT, Lewis BA, Albrecht AE, Marcus BH (2008) Acute
40 Affective Response to a Moderate-intensity Exercise Stimulus Predicts Physical Activity

- 1 Participation 6 and 12 Months Later Psychology of sport and exercise 9:231-245
- 2 doi:10.1016/j.psychsport.2007.04.002
- 3 Yu J (2015) The etiology and exercise implications of sarcopenia in the elderly International
- 4 Journal of Nursing Sciences 2:199-203 doi:<http://dx.doi.org/10.1016/j.ijnss.2015.04.010>
- 5