

An injectable, self-healing and MMP-inhibiting hyaluronic acid gel via iron coordination

GAO, Ziyu, YANG, Xuebin, JONES, Elena, BINGHAM, Paul
<<http://orcid.org/0000-0001-6017-0798>>, SCRIMSHIRE, Alex
<<http://orcid.org/0000-0002-6828-3620>>, THORNTON, Paul and TRONCI, Giuseppe

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/27389/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

GAO, Ziyu, YANG, Xuebin, JONES, Elena, BINGHAM, Paul, SCRIMSHIRE, Alex, THORNTON, Paul and TRONCI, Giuseppe (2020). An injectable, self-healing and MMP-inhibiting hyaluronic acid gel via iron coordination. *International Journal of Biological Macromolecules*, 165 (B), 2022-2029.

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

Supporting information

An injectable, self-healing and MMP-inhibiting hyaluronic acid gel via iron coordination

Table S1 Formulation of different samples

Sample name	HA (2 wt.%, mL)	Fe ³⁺ -GSH (10 mg/ml, μ L)	HCl (120 mM, μ L)	H ₂ O (μ L)
Fe 100	1	100	0	400
Fe 200	1	200	0	300
Fe 300	1	300	0	200
Fe 400	1	400	0	100
Fe 500	1	500	0	0
Ctrl 0	1	0	0	500
Ctrl 100	1	0	100	400
Ctrl 200	1	0	200	300
Ctrl 300	1	0	300	200
Ctrl 400	1	0	400	100
Ctrl 500	1	0	500	0

*Solvent information: HA (H₂O), Fe³⁺-GSH (120 mM HCl), HCl (H₂O).




Fig.S1 Shear modulus of HA-HCl control sample measured via strain sweep.




Fig.S2 Optical images of ATDC 5 cells after 1-day culture with Fe^{3+} -GSH gel, (a) 0 μL (TCPs), (b) 5 μL , (c) 10 μL , (d) 20 μL , (e) 30 μL , (f) 40 μL and (g) 50 μL per well. Scale bar: 100 μm .




Fig.S3 Optical images of ATDC 5 cells after 5-day culture with Fe³⁺-GSH gel, (a) 0 μ L (TCPs), (b) 5 μ L, (c) 10 μ L, (d) 20 μ L, (e) 30 μ L, (f) 40 μ L and (g) 50 μ L per well. Scale bar: 100 μ m.