

When It HIITs, You Feel No Pain: Psychological and Psychophysiological Effects of Respite-Active Music in High-Intensity Interval Training.

KARAGEORGHIS, Costas I., JONES, Leighton http://orcid.org/0000-0002-7899-4119, HOWARD, Luke W., THOMAS, Rhys M., MOULASHIS, Panayiotis and SANTICH, Sam J.

Available from Sheffield Hallam University Research Archive (SHURA) at:

https://shura.shu.ac.uk/27951/

This document is the Accepted Version [AM]

Citation:

KARAGEORGHIS, Costas I., JONES, Leighton, HOWARD, Luke W., THOMAS, Rhys M., MOULASHIS, Panayiotis and SANTICH, Sam J. (2020). When It HIITs, You Feel No Pain: Psychological and Psychophysiological Effects of Respite-Active Music in High-Intensity Interval Training. Journal of Sport and Exercise Psychology, 1-12. [Article]

Copyright and re-use policy

See http://shura.shu.ac.uk/information.html

1	Running Head: RESPITE-ACTIVE MUSIC IN HIIT
2	
3	
4	When it HIITs, You Feel No Pain: Psychological and Psychophysiological Effects of
5	Respite–Active Music in High-Intensity Interval Training
6	
7	Costas I. Karageorghis ¹ , Leighton Jones ² , Luke W. Howard ¹ , Rhys M. Thomas ¹ ,
8	Panayiotis Moulashis ¹ , & Sam J. Santich ³
9	
10	¹ Department of Life Sciences, Brunel University London, UK
11	² Academy of Sport and Physical Activity, Sheffield Hallam University, UK
12	³ Department of Exercise Science and Sports Studies, Springfield College, USA
13	

1	
2	W

Abstract

2 e investigated the effects of respite-active music (i.e., music used for active recovery in 3 between high-intensity exercise bouts) on psychological and psychophysiological outcomes. 4 Participants (N = 24) made four laboratory visits for a habituation, medium- and fast-tempo 5 music conditions, and a no-music control. A HIIT protocol comprising 8×60 -s exercise 6 bouts at 100% Wmax with 90 s active recovery was administered. Measures were taken at the 7 end of exercise bouts and recovery periods (RPE, state attention, core affect), then upon 8 cessation of the protocol (enjoyment and remembered pleasure). Heart rate (HR) was 9 measured throughout. Medium-tempo music enhanced affective valence during exercise and 10 recovery, while both music conditions increased dissociation (only during recovery), 11 enjoyment, and remembered pleasure relative to control. Medium-tempo music lowered RPE 12 relative to control but the HR results were inconclusive. As predicted, medium-tempo music 13 in particular, had a meaningful effect on a range of psychological outcomes. 14

1

2

When it HIITs You, You Feel No Pain: Psychological and Psychophysiological Effects of Respite–Active Music in High-Intensity Interval Training

3 Recovery can be defined as the organism's return to baseline or resting state (Kellman 4 & Beckmann, 2017). The enhancement of recovery has implications for the degree to which 5 recreationally active individuals both enjoy and adhere to exercise (Martin & Woods, 2012). 6 Most of the literature addressing the psychological and psychophysiological effects of music 7 in the exercise domain has focused upon pretask and in-task applications (see e.g., Terry, 8 Karageorghis, Curran, Martin, & Parsons-Smith, 2020 for a meta-analysis); scant attention 9 has been given to investigation of the use of music for recovery. One important distinction, 10 not always made explicit by researchers, has entailed the use of music for movement-based 11 recovery, known as "active recovery", and static-based recovery, known as "passive 12 recovery" (Karageorghis, 2017).

13 Jones, Tiller, and Karageorghis (2017) coined the term "respite music" for music 14 applied during periods of recovery *within* an exercise session but there is a need for greater 15 conceptual clarity given the different ways in which respite music can be applied. In the Jones et al. study, music was applied in the recovery periods in between high-intensity bouts 16 17 when participants were taking passive (i.e., static) recovery. We propose that the term, 18 "respite-passive music" represents greater precision for such instances, whereas "respite-19 active music" more accurately reflects the present application (i.e., participants cycled at a 20 low intensity during a recovery period while listening to music).

High-intensity interval training (HIIT) is a short-duration form of exercise that entails
a series of short, high-intensity efforts, punctuated by recovery periods (Stork, Banfield,
Gibala, & Martin Ginis, 2017). Studies have shown that several weeks of HIIT can bring
benefits to physical health that are analogous to those derived from the long-duration,
aerobic-type exercise that is more commonly adopted by the general public (Batacan,
Duncan, Dalbo, Tucker, & Fenning, 2017). Such health benefits can also be realized by at-

risk and diseased populations (Gibala et al., 2014; Quindry, Franklin, Chapman, Humphrey,
& Mathis, 2019). Albeit there is a growing body of evidence showing the physiological
benefits of HIIT, from a public health/adherence perspective, there is a significant
shortcoming. It is widely acknowledged that people can find HIIT to be rather unpleasant; a
factor that can undermine long-term participation in such activity (see e.g., Decker &
Ekkekakis, 2017; Ekkekakis, 2020).

Given the aforementioned health benefits of interval-type exercise, coupled with the
likelihood of it being perceived as unpleasant, there has been growing interest in the
application of music as a means by which to ameliorate negative psychological responses,
such as affective decline, low enjoyment, and high levels of perceived exertion (e.g., Jones et
al., 2017; Stork, Kwan, Gibala, & Martin Ginis, 2015). Hypothetically, using music to
enhance the experience of interval-type exercise may have positive consequences for future
participation (i.e., exercise adherence; see Stork, Karageorghis, & Martin Ginis, 2019).

The last decade has seen the emergence of an extensive corpus of work addressing the effects of music during high-intensity exercise (e.g., Jones et al., 2017; Karageorghis et al., 2009; Karageorghis et al., 2018; Stork et al., 2015). In HIIT protocols, the use of environmental stimuli to enhance recovery or respite periods has yet to be investigated in a systematic manner and provides tantalizing opportunities for researchers in the realm of sport and exercise psychology.

Recent work has shown the degree to which affective memory and affective forecasting can influence physical activity behaviors (e.g., Ekkekakis, Zenko, Ladwig, & Hartman, 2018). Moreover, the collection of physiological measures (e.g., heart rate) alongside psychological measures (e.g., self-reported affective responses) is likely to proffer a better understanding of the effects that environmental stimuli can have during HIIT. Among the body of work that has explored how bodily pulses can resonate with music pulses, Jones et al. (2017) found that there were minimal effects associated with the application of slow1 and fast-tempo music on breathing and heart rates.

2 In their examination of recuperative music, Karageorghis et al. (2018) indicated that 3 fast-tempo music inhibited HR recovery following exhaustive exercise. If a targeted music 4 intervention can expedite physiological recovery during an interval-type session, intuitively, 5 this is likely to engender a more pleasant exercise experience. Nonetheless, a salient 6 consideration in any such session is that, an intervention that engenders a positive effect on 7 recovery, does not compromise performance in the next exercise bout. Therefore, a balance 8 needs to be struck between the amelioration of negative affect during a recovery period, and 9 maintenance of the degree of psychomotor arousal needed to optimize power output in the 10 ensuing exercise bout. That is why, in the present study, medium-tempo music was used for 11 the recovery periods and not slow-tempo music, as in past work of a similar nature (e.g., 12 Hutchinson & O'Neil, 2020; Jones et al., 2017).

Music tempo is central to the biomusicological phenomenon of *entrainment*, which concerns bodily pulses such as heart/respiration rate and brainwaves being drawn into a common oscillation with musical tempo (Terry et al., 2020). This phenomenon has yet to be examined in the context of high-intensity exercise with the application of respite–active music. Accordingly, the purpose of the present study was to investigate the effects of respite– active music on the psychological experiences and physiological responses of recreationally active individuals who were administered a HIIT protocol in a laboratory setting.

Based on findings from previous research (Eliakim, Bodner, Meckel, Nemet, &
Eliakim, 2012; Hutchinson & O'Neil, 2019; Karageorghis & Jones, 2014), we hypothesized
that recovery periods accompanied by medium-tempo, respite–active music (120–125 bpm)
would be superior in terms of psychological (i.e., higher affective
valence/enjoyment/pleasure scores and less association) and psychophysical (i.e., lower RPE)
outcome measures when compared to fast-tempo, respite–active music (135–140 bpm), and a

26 no-music control condition (H_1) . In accord with related literature pertaining to the application

of recuperative music following exhaustive exercise (Karageorghis et al., 2018), we expected lower average heart rate and the lowest absolute heart rate in recovery periods accompanied by medium-tempo music when compared to fast-tempo music and control conditions (H_2). We did not expect any differences among conditions in terms of average and peak heart rate during the high-intensity exercise bouts and so the null hypothesis was tested (H_3). We did not expect the experimental manipulations–applied only to recovery periods–to have any significant effect on measures taken at the end of the exercise bouts (H_4).

8

Method

9 Study design and Power Analysis

10 A fully counterbalanced, within-subjects design was employed with one control 11 condition (no music) and two experimental conditions (medium-tempo and fast-tempo 12 music). Using a more conservative effect size than that derived from Jones et al. (2017; $\eta_p^2 =$ (0.08) owing to a more subtle tempo manipulation between experimental conditions, an a13 *priori* power analysis using a small-to-medium effect size ($\eta_p^2 = 0.05$), an alpha level of .05, 14 15 power at .8, indicated that a sample of 21 participants would be required to detect differences in a twoway repeated-measures (RM) 3 (Condition) × 8 (Time [HIIT exercise bouts and 16 17 active recovery periods]) ANOVA of Feeling Scale scores (the primary outcome measure). 18 An additional three participants were recruited to guard against deletions due to outliers and 19 to enable full counterbalancing of conditions.

20 **Participants**

With institutional ethics approval, 24 participants (12 women and 12 men; M_{age} 22.5 years, SD = 1.7 years; $M_{height} = 175.3$ cm, SD = 9.5 cm; M_{mass} 80.7 kg, SD = 12.6 kg; M_{weekly} vigorous activity 60 min, SD = 25.8 min; $M_{weekly moderate activity} = 130$ min; SD = 70.5 min) were recruited through word of mouth at a university in ______ and all provided written informed consent. Participant inclusion criteria were that they: a) were aged 18–25 years; b) could fully comprehend spoken and written English; c) were healthy; and d) participated in at least 150 min/week of moderate physical activity (PA) and/or more than 75 min/week of
vigorous PA over the previous 3 months (World Health Organization, 2010). Participants
were of a similar age in order to limit the effects of age on music preference, thereby
addressing a potential confound (Karageorghis & Terry, 1997). Similarly, participants were
recruited in such a way that there was a degree of homogeneity in their weekly levels of
physical activity. This is important given that physical activity status is known to influence
affective responses to exercise (Magnan, Kwan, & Bryan, 2013).

8 **Procedures**

9 Potential volunteers were screened to ensure they met the study inclusion criteria and 10 determine any significant health problems that might prevent them from engaging in a HIIT 11 protocol. The Physical Activity Readiness Questionnaire (PAR-Q) and the International 12 Physical Activity Questionnaires (IPAQ) were administered prior to each participant's first 13 laboratory visit. Participants did not report any significant health concerns and weekly active 14 minutes confirmed that participants met the aforementioned inclusion criteria.

15 Baseline fitness testing and familiarization. We recorded the participant's height 16 and mass then explained the procedures associated with completion of a maximal exercise 17 test on an electronically-braked cycle ergometer (Lode Excalibur Sport; Lode B.V., 18 Groningen, the Netherlands). The test comprised a 4-min warm-up at 50 W followed by a 19 ramped protocol with 20 W increases each minute until volitional exhaustion. Peak wattage 20 was recorded and used in subsequent experimental trials. The protocol adhered to criteria for 21 maximal tests (see Porszasz, Casaburi, Somfay, Woodhouse, & Whipp, 2003); specifically 22 that the protocol: a) included a low initial metabolic demand; b) provided a constant linear 23 increase in work rate; and c) brought participants to the limit of tolerance in ~10 min. 24 Following a 10-min recovery period, the participant completed a HIIT habituation session 25 with 4×60 s bouts interspersed with 90 s recovery (no music). The workload for the exercise 26 bouts of the HIIT session was 100% Wmax.

1 The participant was asked to report her/his rating of perceived exertion by use 2 of the CR-10 RPE scale (Borg, 1998), state attention by use of the Attention Scale (Tammen, 3 1996), pleasure by use of the Feeling Scale (Hardy & Rejeski, 1989), and perceived 4 activation by use of the Felt Arousal Scale (Svebak & Murgatroyd, 1985) in the final 15 s of 5 each 60-s high-intensity exercise bout and before commencement of any music. The same 6 measures were administered in the final 15 s of each 90-s recovery period but in the reverse 7 order to minimize common method variance. Remembered pleasure was assessed using a 8 200-point scale (Zenko, Ekkekakis, & Ariely, 2016), which ranges from -100 (very 9 *unpleasant*) to 100 (*very pleasant*) and 0 represents a neutral response. Enjoyment was 10 measured using the Physical Activity Enjoyment Scale (PACES; Kendzierski & DeCarlo, 11 1991); scores can range from 18 to 126, with 72 representing the midpoint and thus a neutral 12 score. These scales were administered immediately following the exercise bout, once the participant had dismounted the cycle ergometer. At the end of the session, the participant was 13 14 led through a warm-down and encouraged to ask any questions s/he might have. 15 Experimental trials. After a minimum of 48 hours following her/his pretest 16 familiarization visit to the laboratory, the participant completed a HIIT session under three 17 conditions: a) medium-tempo music (120–125 bpm); b) fast-tempo music (135–140 bpm); 18 and c) a no-music control. The three HIIT test conditions were administered ~48–72 hours 19 apart. Each participant was instructed to maintain her/his typical habits in terms of both sleep 20 and diet. Moreover, the participant was asked to desist from any other form of physical

and diet. Moreover, the participant was asked to desist from any other form of physical
activity for the entire day when s/he was due to make a visit to the laboratory. Prior to
initiating any of the physical tasks, each participant had the procedures explained to her/him
in full, and was afforded an opportunity to ask questions. Each participant's heart rate was
recorded throughout by use of a chest-strap transmitter linked to a wristwatch (Polar H10;
Polar Electro Oy, Kempele, Finland), with exercise bout peak and average HR values as well
as recovery low and average values recorded for each HIIT stage. Eight HIIT stages were

included, each with an exercise bout and an ensuing recovery period. The stages were treated
 as distinct units for analytical purposes.

The HIIT session comprised a 4-min warm-up at 50 W followed immediately by the first 60-s exercise bout (100% Wmax) at 75–80 rpm. The exercise bout was followed by a 90-s recovery period at 50 W wherein the participant continued to cycle at 65–70 rpm. A further seven exercise bouts were completed, each separated by a 90-s recovery period. Following the final exercise bout, the participant completed a 90-s recovery period at 50 W and then dismounted the cycle ergometer.

9 It should be noted that the HIIT protocol that we adopted is nonstandard; such 10 protocols normally comprise of 10 exercise bouts, rather than eight, and the recovery periods 11 are either of a 60-s or 75-s duration (Stork et al., 2017). In terms of the number of bouts, a 12 nonstandard protocol was purposefully selected given that participants were required to visit 13 the laboratory on four occasions, and we did not wish to render each visit highly unpleasant 14 and thus run a risk of high participant attrition. In terms of the slightly extended recovery 15 period, this was incorporated to allow sufficient scope for bodily pulses to entrain with the 16 rhythmical qualities of the respite-active music selections (e.g., Khalfa, Roy, Rainville, Dalla 17 Bella, & Peretz, 2008). Each participant was scheduled at the same time of day for their 18 experimental and control trials in order to reduce diurnal variation in HIIT performance (see 19 e.g., Atkinson & Reilly, 1996).

The participant was prompted to report RPE, state attention, FS, and FAS in the last 15 s of each of the eight bouts of high-intensity exercise and in the last 15 s of each recovery period. Perceived enjoyment and remembered pleasure were reported immediately following cool-down and 5 min postexercise. At the end of laboratory visit #4, each participant was asked two open-ended questions as a form of manipulation check to evaluate their perception of differences in the audio content between the two experimental conditions. It is important within such protocols that differences in music tempi across conditions are discernible given that, in real-world conditions, exercisers are able to decide upon just how stimulating or
energizing a music program will be (Karageorghis & Jones, 2014). Each participant was also
administered a music liking item (see Karageorghis & Jones, 2014) to gauge whether there
were differences between the music programs used in each of the two experimental
conditions.

6 Music Choice and Delivery

7 Respite-active music was played by use of a cellphone and Bluetooth speaker 8 (Libratone Zipp Mini 2; Libratone, Nordhavn, Denmark; to enable verbal interaction between 9 the participant and experimenters) at a sound intensity of 70 dBA for the entire 90-s recovery 10 period following each bout of exercise. There was no music playing during the warm-up or 11 each high-intensity exercise bout. The music tracks were selected by the experimenters based 12 on the criteria outlined by Karageorghis et al. (2006) and with reference to the Karageorghis 13 (2016) theoretical model pertaining to music selection in the domain of exercise and sport. 14 Each participant was administered a contemporary pop-music playlist in the two 15 experimental conditions. The playlist for each experimental condition was developed by the 16 experimenters with reference to the aforementioned tempo criteria but entailed the use of a 17 selection panel comprised of participants who were not involved in the experimental phase of 18 the study. The panel (N = 6; three women and three men) had a similar sociocultural 19 background and were in the same age range as the experimental participants (see 20 Karageorghis & Terry, 1997). These playlists were 12 min in length to match the total 21 duration of the recovery sessions and were rated by use of the Brunel Music Rating 22 Inventory-3 (Karageorghis & Terry, 2011) to ensure they were invariant in terms of their 23 motivational qualities (the two playlists can be viewed in Supplementary File 1).

24 Data Analysis

Following data screening and checks for the relevant parametric assumptions,
differences in theoretically linked dependent variables (e.g., affect and arousal) measured

1 over time were analyzed by use of mixed-model 3 (Condition) \times 8 (Time) MANOVAs. 2 Differences in remembered pleasure and exercise enjoyment were analyzed by use of 3 oneway, RM ANOVAs. Maximal HR was calculated using the Gellish et al. (2007) formula 4 and percentage of maximal heart rate was calculated for the exercise bouts and associated 5 recovery periods. Analysis of HR data was conducted by means of 3 (Condition) \times 8 (Time) 6 ANOVAs. In all ANOVAs and MANOVAs, Greenhouse–Geisser-corrected F values were 7 used in the case of sphericity violations and pairwise comparisons were subject to Bonferroni 8 adjustment. Music liking differences were analyzed using a paired-samples t test.

9

Results

10 **Data Diagnostics**

11 Data screening revealed 51 univariate outliers of which 45 were accounted for by one 12 case that was deleted prior to the analyses. The remaining six outliers, all associated with HR measures, were adjusted until they came within the range $z \pm 3.29$ (see Tabachnick & Fidell, 13 14 2018). Normality tests indicated that HR, in particular, exhibited instances of non-normality 15 (negative skewness) and square root transformations were applied but did not entirely remedy 16 the non-normality; accordingly, the non-normalized data were retained for analysis. To aid 17 interpretation of the results presented herein, descriptive statistics for all dependent measures, 18 across conditions and throughout the HIIT protocol, are provided in Table 1.

19 In-Task Psychological Variables

The 3 (Condition) × 8 (Stage) MANOVA for RPE and state attention at the end of the exercise bouts indicated significant omnibus statistics, Pillai's Trace = 0.19, F(28, 616) = 2.26, p < .001, $\eta_p^2 = .93$. Stepdown *F* tests indicated a significant interaction effect for RPE (see Table 2). Examination of standard errors (± 2 *SE*) showed differences from Stage 5 onwards between medium-tempo music and control (see Figure 1), with medium-tempo music yielding lower RPE. Differences also emerged between medium- and fast-tempo music at Stage 5 and Stage 8 (medium < fast). The twoway interaction for state attention was 1 nonsignificant (see Table 2 and Figure 2).

A main effect of condition emerged for RPE (see Table 2) with pairwise comparisons indicating differences between control and medium-tempo music (control > medium), and between medium- and fast-tempo music (medium < fast; see Figure 1). Also, a main effect of stage emerged for RPE (see Table 2), with pairwise comparisons indicating differences between the early and late stages of HIIT (see Figure 1). Only a main effect of stage emerged for state attention (see Table 2), with pairwise comparisons indicating differences between the early and late stages of HIIT (see Figure 2).

9 The 3 (Condition) \times 8 (Stage) MANOVA for RPE and state attention at the end of the 10 recovery period indicated significant omnibus statistics (Pillai's Trace = 0.17, F(28, 616) = 2.09, p < .001; $\eta_p^2 = .09$). Stepdown F tests indicated a significant interaction effect for RPE 11 12 (see Table 2). Examination of standard errors showed differences from Stage 5 onwards 13 between control and medium-tempo music, with the latter yielding lower RPE. There were 14 also differences between medium- and fast-tempo music from Stages 6–8 (medium < fast; see 15 Figure 1). The twoway interaction for state attention was nonsignificant (see Table 2 and 16 Figure 2).

17 A main effect of condition emerged for RPE with pairwise comparisons indicating 18 differences between control and medium-tempo music (control > medium), as well as 19 between medium- and fast-tempo music (medium < fast; see Table 2). Also, a main effect of 20 stage emerged for RPE with pairwise comparisons indicating differences between the early 21 and late stages of HIIT (see Table 2 and Figure 1). A main effect of condition also emerged 22 for state attention (Table 2), with pairwise comparisons indicating differences between 23 control and both music conditions; the latter yielded higher dissociation scores (see Figure 2). 24 A main effect of stage emerged for state attention (see Table 2), with pairwise comparisons indicating differences between the early and late stages of HIIT (i.e., association increased 25 26 through the HIIT stages; see Figure 2).

1

In-Task Affective Measures

2 The 3 (Condition) \times 8 (Stage) MANOVA for affective measures (valence and 3 arousal) at the end of the exercise bouts indicated significant omnibus statistics, Pillai's Trace = 0.24, F(28, 616) = 3.00, p < .001, $\eta_p^2 = .12$, and stepdown F tests indicated a significant 4 5 interaction effect for arousal (see Table 2). Examination of standard errors showed 6 differences at Stage 2 between medium-tempo music and control (medium < control), and at 7 Stage 8 between fast-tempo music and control (fast > control; see Figure 3). The twoway 8 interaction for affective valence was nonsignificant (see Table 2 and Figure 3). 9 There was a main effect of condition for affective measures (valence and arousal) at the end of the exercise bouts, Pillai's Trace = 0.46, F(4.88) = 6.65, p < .001, $\eta_p^2 = .23$. 10

Stepdown F tests indicated a significant effect only for affective valence (see Table 2), with 11 pairwise comparisons showing differences between medium-tempo music and control 12 (medium > control), medium-tempo music and fast-tempo music (medium > fast), and fast-13 14 tempo music and control (fast > control; see Figure 3). There was also a main effect of stage, Pillai's Trace = 1.02, F(14, 308) = 22.70, p < .001, $\eta_p^2 = .51$. Stepdown F tests indicated a 15 significant effect for affective valence (see Table 2), with pairwise comparisons showing a 16 gradual decline through the stages (see Figure 3). Stepdown F tests indicated a significant 17 18 effect for arousal (see Table 2), with pairwise comparisons showing a gradual increase 19 through the stages (see Figure 3).

The 3 (Condition) × 8 (Stage) MANOVA for affective measures (valence and arousal) at the end of the recovery period yielded significant omnibus statistics, Pillai's Trace $= 0.14, F(28, 616) = 1.69, p = .015, \eta_p^2 = .07$. Stepdown *F* tests indicated a significant twoway interaction effect for arousal (see Table 2). Examination of standard errors showed differences at Stage 1 between fast-tempo music and control (fast < control), and at Stage 6 between fast-tempo music and control (fast > control; see Figure 3). The twoway interaction for affective valence was nonsignificant (see Table 2 and Figure 3).

1 There was a main effect of condition for affective measures (valence and arousal) at the end of the recovery period, Pillai's Trace = 0.51, F(4, 88) = 7.54, p < .001, $n_p^2 = .26$). Step 2 3 down F tests indicated a significant effect only for affective valence (see Table 2), with 4 pairwise comparisons showing differences between medium-tempo music and control 5 (medium > control), and medium-tempo and fast-tempo music (medium > fast; see Figure 3). 6 There was also a main effect of stage, Pillai's Trace = 0.88, F(14, 308) = 17.35, p < .001, η_p^2 7 = .44). Stepdown F tests indicated a significant effect for affective valence (see Table 2), 8 with pairwise comparisons showing a gradual decline through the stages (see Figure 3). 9 Stepdown F tests indicated a significant effect for arousal (see Table 2), with pairwise 10 comparisons showing an increase between the early and late stages of HIIT (see Figure 3).

11 Heart Rate

12 The 3 (Condition) \times 8 (Stage) ANOVA for average HR during the exercise bouts was significant (see Table 2), with an examination of standard errors showing a difference 13 14 between medium-tempo music and control (medium > control), only at Stage 1. There was a 15 main effect of condition; albeit no significant differences emerged in the ensuing pairwise 16 comparisons (see Table 2). There was a main effect of stage (see Table 2), with pairwise 17 comparisons showing a gradual increase in HR through the stages (see Table 1). The 3 18 (Condition) \times 8 (Stage) ANOVA for peak HR during the exercise bouts was nonsignificant, 19 as was the main effect of condition (see Table 2). A main effect of stage emerged, with 20 pairwise comparisons showing differences from Stages 1–8 (i.e., see Table 1 and Table 2). 21 The 3 (Condition) \times 8 (Stage) ANOVA for average HR during the recovery periods 22 was significant (see Table 2), with an examination of standard errors showing that at Stages 23 4–5, there was a difference between fast-tempo music and control (fast < control). There was 24 no main effect of condition; however, there was a main effect of stage, with pairwise 25 comparisons showing a gradual increase in HR through to Stage 7 (see Table 2). The 3 26 (Condition) \times 8 (Stage) ANOVA for lowest recorded HR during the recovery periods was

1	significant (see Table 2). Examination of standard errors showed differences between control
2	and fast-tempo music at Stage 4 and Stage 5 (control > fast tempo). There was a main effect
3	of condition; albeit pairwise comparisons showed no significant differences (see Table 2).
4	There was also a main effect of stage, with pairwise comparisons indicating differences
5	among all stages through to Stage 7 (i.e., lowest recorded HR increased throughout; see Table
6	2).
7	Post-Task Psychological Measures
8	The RM ANOVA for exercise enjoyment (PACES) was significant. Pairwise
9	comparisons showed differences across all conditions, with medium-tempo music eliciting
10	the highest scores followed by fast-tempo and control (see Table 2). The RM ANOVA for
11	remembered pleasure was significant, with pairwise comparisons showing that medium-
12	tempo music elicited higher scores than control, as did fast-tempo music (see Table 2).
13	Music Liking
14	The <i>t</i> test for the music liking scores between the medium-tempo ($M = 5.92$, $SD =$
15	0.63) and fast-tempo ($M = 5.58$, $SD = 0.75$) music conditions showed that there was no
16	significant difference, $t(22) = 2.05$, $p = .052$.
17	Manipulation Check
18	Responses from the open-ended questions indicated that 23 participants were
19	cognizant that one music condition was faster than the other.
20	Discussion
21	The main purpose of the study was to investigate the effects of respite-active music
22	on psychological and psychophysiological responses both during and immediately after an
23	eight-stage HIIT protocol. In regard to H_1 , it appears that medium-tempo music was generally
24	superior during recovery periods in terms of psychological outcomes (higher affective
25	valence/enjoyment/pleasure scores, and lower RPE), with the exception of state attention in
26	which dissociation scores were similar across the two music conditions (see Figure 2), and so

1 it is partially accepted. With reference to H_2 , which related to recovery HR, there was no 2 clear differentiation across conditions and so it is not accepted. It is notable, however, that a 3 main effect of condition was identified, albeit this was not associated with significant 4 pairwise comparisons (see Table 2). H_3 is accepted, as there was no Condition \times Time 5 interaction for either average or peak HR in the exercise bouts. H₄ concerned the expected 6 lack of effect of the music manipulation on exercise bout-related measures and was only 7 partially accepted, given that medium-tempo music elicited lower RPE scores as the HIIT 8 stages progressed (see Table 2 and Figure 1).

9 In-Task Psychological Measures

10 The core affect data show how the medium-tempo condition assuaged the inevitable 11 displeasure that is experienced in the latter stages of both the exercise bouts and recovery 12 periods of HIIT. This concurs with the findings of recent related studies (Eliakim et al., 2012; 13 Hutchinson & O'Neil, 2020) and it is clear that the effects of music were seemingly more 14 potent as fatigue levels increased. Recent mechanistic work suggests that the presence of 15 music inhibits communication across somatosensory regions of the brain during cycle 16 ergometer exercise (Bigliassi, Karageorghis, Wright, Orgs, & Nowicky, 2017). Such dulling 17 of afferent signals might account for the affective benefits that emerged in the present study 18 (see Figure 3).

19 An interesting and somewhat unexpected finding to emerge from the RPE data is that, 20 from Stage 4, there is a marked reduction in RPE at the end of the exercise bouts with 21 medium-tempo music, compared to the other two conditions (see Figure 1). This hints at a 22 cumulative carryover effect of the medium-tempo music that was delivered during the 23 recovery periods. The presence of a carryover effect counters what Crust (2004) reported 24 when examining carryover effects of music in an isometric strength task. The strength test to 25 failure required high effort and participants experienced localized muscle fatigue in a manner 26 akin to HIIT protocols. Albeit Crust (2004) reported ergogenic effects of music during the

1 task, music exposure *prior* to the task did not have any such effects. In the present study,

there seemed to be some carryover effects of respite–active music on perceptions of exertion
during exercise bouts (see Figure 1).

The aforementioned differences in RPE were not, however, matched in state attention for the exercise bouts; there was no differentiation between the two music conditions, both of which yielded higher scores for association than control in the recovery periods (see Table 1 and Figure 2). It is noteworthy that the present findings are indicative of the notion that RPE and state attention are not phenomenologically isomorphic (Razon, Hutchinson, &

9 Tenenbaum, 2012).

10 Heart Rate (HR) Data

The main effect of condition for recovery low HR was significant but the associated pairwise comparisons did not exhibit any differences (see Table 2). There was an anomaly evident in this subset of the data given that, overall, fast-tempo music elicited slightly lower HRs (see Table 1). As expected, HR data indicated that physiological workload increased as the HIIT session progressed. The interaction effects identified in the average HR during exercise bouts, and during recovery periods, do not point to any sort of meaningful trend, owing to the rather sporadic nature of such differences.

Given that the application of respite-active music had no bearing on the three HR 18 19 indices, we can deduce that, from a physiological arousal perspective, there is no potential 20 adverse effect on bouts of high-intensity exercise. This point holds both for medium-tempo 21 and fast-tempo music (see also Jones et al., 2017). Nonetheless, the medium-tempo music did 22 engender some psychological benefits and such benefits in recovery did not appear to prompt 23 participants to exert less effort—in physiological terms-during subsequent exercise bouts 24 (see Table 1 and Table 2). This finding suggests that respite–active music of a tempo > 120 25 bpm can facilitate active recovery without inhibiting subsequent effort exertion.

26 The lack of a main effect of condition contrasts with the recent findings of

Karageorghis et al. (2018). It could be that the expected differences in HR recovery might only be evident following longer exposure to music in the context of either high-intensity or exhaustive exercise. Longer exposure to a musical work renders the entrainment process more likely and so, the use of musical excerpts, rather than entire musical works, may have precluded HR entrainment (see Terry et al., 2020). Nonetheless, in the context of HIIT recovery, the use of entire musical works is not possible, unless such works are of very short duration.

8 Post-Task Psychological Measures

9 The results for exercise enjoyment, as measured by PACES (see Table 2), provided 10 the most defined differentiation across conditions. The medium-tempo music yielded the 11 highest scores, which suggests that in terms of a gestalt assessment of the HIIT experience, 12 such music is superior to both fast-tempo music and a no-music control. Similarly, 13 remembered pleasure showed medium-tempo music to be superior to control but, in this 14 instance, differences between music conditions did not emerge. Exercise practitioners might 15 consider either type of music (i.e., medium or fast tempo) with a view to enhancing people's 16 recall of high-intensity exercise (see Hutchinson & O'Neil, 2019; Jones et al., 2017). 17 Notably, fast-tempo music administered across the entire course of an interval-type session 18 (including exercise bouts and recovery periods) has also been shown to enhance enjoyment 19 compared to a no-music control condition (Stork et al., 2019).

20 Strengths and Limitations

There have been over 300 published studies that examined pretask and in-task music applications in exercise or sport, but only a handful that have examined post-task applications (see Terry et al., 2020). This study is the first to empirically examine the relevance of music tempo during the application of respite–active music and provides insight into how the benefits of this mode of music application can be maximized. A postexperiment manipulation check indicated that 23 out of 24 participants were able to identify that there were tempo differences between the two music conditions. This is a strength given that participants
correctly perceived the independent variable manipulation, which allows more accurate
conclusions to be drawn in regard to how the manipulation influenced the dependent
variables (Thomas, Nelson, & Silverman, 2015). Moreover, as the check took place at the end
of all experimental trials, it could not influence participants' responses or cognitions (e.g., via
an experimenter effect) during the trials.

7 The affective responses to exercise evident in healthy and recreationally active 8 participants, such as those tested in the present study, differ from obese and inactive 9 individuals (see e.g., Decker & Ekkekakis, 2017). The applicability of the present data is 10 limited to active/healthy populations. Accordingly, further examination of respite-active 11 music with at-risk populations (e.g., obese, type 2 diabetics) and people who are 12 insufficiently active, appears warranted. The inclusion criterion pertaining to the recruitment only of young adults precludes the generalization of the present findings to other subgroups 13 14 of the population for which the effects of medium-tempo music might be different. Previous 15 research examining music preferences in an exercise context has detailed how music-tempo 16 preferences vary in accord with age (e.g., Priest & Karageorghis, 2008).

17 Respite music is an atypical music application and would require considerable 18 planning and track editing for a playlist to contour the demands of an exercise or training 19 session (Karageorghis, 2017). The present data indicate some utility in the adoption of 20 playlists to match the peaks and troughs of a HIIT session with a view to positively 21 influencing psychological responses. Note that the atypical nature of the present application 22 of music led to the selection of a recovery period (90 s) that was longer than the typical 23 recovery period associated with HIIT protocols (60-75 s; see e.g., Little, Safdar, Wilkin, 24 Tarnopolsky, & Gibala, 2010). This slightly extended recovery was adopted to give the respite-active music sufficient time to take effect, but the use of brief musical excerpts (e.g., 25 26 60 s) may not be as efficacious as the 90-s excerpts used in the present study. Moreover, the

tempo-related recommendations presented herein should only be considered for HIIT
protocols that include active recovery periods. Passive recovery is integral to many HIIT
protocols and further research to examine possible differences in terms of optimal musictempo prescription appears warranted.

5 Attitudes toward HIIT (see e.g., Stork & Martin Ginis, 2017) were not assessed before 6 and after the experiment, and thus might be considered for inclusion by future investigators. 7 Similarly, HR was not measured as a manipulation check during the baseline test and future 8 researchers might combine physiological measures with Wmax. A further measurement-9 related limitation is that the affect measures taken at the end of the exercise bouts and 10 recovery periods (FS and FAS) were not also taken prior to the start of each test protocol. 11 Such measures might have been useful in gauging participants' pretask affective state and 12 whether this had any bearing on subsequent psychological and psychophysiological 13 outcomes. Nonetheless, the use of a within-subjects design to reduce the influence of 14 between-subjects error (see Tabachnick & Fidell, 2018), served to assuage any potential 15 threat to internal validity posed by pretask affective state. The order reversal of the four 16 single-item scales (RPE, state attention, FS, and FAS) for the recovery period may have 17 affected participant responses to a small degree (e.g., through temporal differences vis-à-vis 18 the exercise bout administration). It should be stressed, however, that the order reversal was 19 undertaken to mitigate response order effects.

As expected, a significant difference for liking scores did not emerge between the two music conditions; nonetheless, the medium-tempo music program attracted slightly higher scores than its fast-tempo counterpart ($M_{diff.} = 0.34$). Such aesthetic differences between medium-tempo and fast-tempo are well documented in the psychomusicological literature (e.g., Berlyne, 1971) and music-in-exercise literature (e.g., Karageorghis et al., 2011). Reasons advanced for the difference include people's general familiarity with medium tempi, as most popular pieces are recorded in the medium-tempo range (i.e., 100–125 bpm), and the 1 ease with which musical works at medium tempi can be processed by the human brain.

2 **Implications for Practice**

3 Over the last decade, a vibrant debate has raged in exercise psychology regarding the 4 displeasure associated with high-intensity exercise regimens such as HIIT, and whether 5 promotion of such regimens might be counterproductive from a public health perspective (see 6 e.g., Ekkekakis, Hartman, & Ladwig, 2020). The present findings do not, by any means, 7 suggest that music-related applications represent a "magic bullet" in countering sedentary 8 lifestyles. What is clear, however, is that judicious application of music to active recovery 9 enhances pleasure and reduces perceived exertion. Accordingly, practitioners should consider 10 coordinating interval-type exercise sessions with an emphasis on musical accompaniment for 11 recovery. Whether the benefits reported herein would accrue with shorter musical excerpts 12 (e.g., 60 s) is presently unknown and so future work could explore the minimum exposure 13 time required for respite-active music to be efficacious.

14 On balance, medium-tempo music appears to yield the most benefits when used in 15 between high-intensity exercise bouts and it should be noted that slow-tempo music can have 16 a detrimental effect on subsequent anaerobic performance (see e.g., Hutchinson & O'Neil, 2020). Individual exercisers might consider editing playlists, using freely available software 17 18 (e.g., Audacity), for HIIT sessions that include epochs of music and silence to match their 19 desired HIIT protocol. If the efficacy of this type of music application is established, an 20 algorithm could be developed that creates playlists that are punctuated by periods of silence 21 to coincide with exercise bouts.

22

Conclusions and Recommendations

Medium-tempo respite–active music showed the greatest capacity to influence psychological responses both during and after a HIIT session. Notably, perceptions of exertion during the exercise bouts (when no music was played) *and* recovery periods in the latter stages of a HIIT session appeared to be reduced through the use of medium-tempo

1 respite-active music when compared to a no-music control. Nonetheless, the present data 2 clearly indicate that a decline in pleasure (i.e., exercise-related affect) is inevitable during 3 HIIT-type protocols but music used during recovery periods can make the session seem less 4 unpleasant. There is a need for such interventions to be studied in longitudinal studies to 5 further understanding of how in-session manipulation of the exercise experience influences 6 adherence in the longue durée. Respite-active music offers a novel music application that is 7 inexpensive and might be considered by practitioners seeking to improve affective responses 8 to HIIT sessions.

1	References
2	Atkinson, G., & Reilly, T. (1996). Circadian variation in sports performance. Sports
3	Medicine, 21, 292-312. doi:10.2165/00007256-199621040-00005
4	Batacan, R. B., Duncan, M. J., Dalbo, V. J., Tucker, P. S., & Fenning, A. S. (2017). Effects
5	of high-intensity interval training on cardiometabolic health: A systematic review and
6	meta-analysis of intervention studies. British Journal of Sports Medicine, 51, 494-503.
7	doi:10.1136/bjsports-2015-095841
8	Berlyne, D. E. (1971). Aesthetics and psychobiology. New York, NY: Appleton Century
9	Crofts.
10	Bigliassi, M., Karageorghis, C. I., Wright, M. J., Orgs, G., & Nowicky, A. V. (2017). Effects
11	of auditory stimuli on electrical activity in the brain during cycle ergometry. Physiology
12	& Behavior, 177, 135-147. doi:10.1016/j.physbeh.2017.04.023
13	Borg, G. (1998). Borg's perceived exertion and pain scales. Champaign, IL: Human Kinetics.
14	Crust, L. (2004). Carry-over effects of music in an isometric muscular endurance task.
15	Perceptual & Motor Skills, 98, 985-991. doi:10.2466/pms.98.3.985-991
16	Decker, E. S., & Ekkekakis, P. (2017). More efficient, perhaps, but at what price? Pleasure
17	and enjoyment responses to high-intensity interval exercise in low-active women with
18	obesity. Psychology of Sport & Exercise, 28, 1-10.
19	doi:10.1016/j.psychsport.2016.09.005
20	Ekkekakis, P., Hartman, M. E., & Ladwig, M. A. (2020). Affective responses to exercise. In
21	G. Tenenbaum & R. C. Eklund (Eds.), Handbook of sport psychology (4th ed., pp. 233-
22	253). New York, NY: Wiley.
23	Ekkekakis, P., Zenko, Z., Ladwig, M. A., & Hartman, M. E. (2018). Affect as a potential
24	determinant of physical activity and exercise: Clinical appraisal of an emerging
25	research field. In D. Williams, R. Rhodes, & M. Conner (Eds.), Affective determinants
26	of health behavior (pp. 237–239), Oxford, UK: Oxford University Press.
27	Eliakim, M., Bodner, E., Eliakim, A., Nemet, D., & Meckel, Y. (2012). Effect of
28	motivational music on lactate levels during recovery from intense exercise. The Journal
29	of Strength & Conditioning Research, 26, 80-86. doi:10.1519/JSC.0b013e31821d5f31

1	Gellish, R. L., Goslin, B. R., Olson, R. E., McDonald, A., Russi, G. D., & Moudgil, V. K.
2	(2007). Longitudinal modeling of the relationship between age and maximal heart rate.
3	Medicine & Science in Sports & Exercise, 39, 822–829.
4	doi:10.1097/mss.0b013e31803349c6
5	Gibala, M. J., Gillen, J. B., & Percival, M. E. (2014). Physiological and health-related
6	adaptations to low-volume interval training: influences of nutrition and sex. Sports
7	Medicine, 44, 127-137. doi:10.1007/s40279-014-0259-6
8	Hardy, C. J., & Rejeski, W. J. (1989). Not what, but how one feels: The measurement of
9	affect during exercise. Journal of Sport & Exercise Psychology, 11, 304-317.
10	doi:10.1123/jsep.11.3.304
11	Hutchinson, J. C., Jones, L., Vitti, S. N., Moore, A., Dalton, P. C., & O'Neil, B. J. (2018).
12	The influence of self-selected music on affect-regulated exercise intensity and
13	remembered pleasure during treadmill running. Sport, Exercise, and Performance
14	Psychology, 7, 80-92. 7, 80-92. doi:10.1037/spy0000115
15	Hutchinson, J. C., Karageorghis, C. I., & Jones, L. (2014). See hear: Psychological effects of
16	music and music-video during treadmill running. Annals of Behavioral Medicine, 49,
17	199–211. doi:10.1007/s12160-014-9647-2
18	Hutchinson, J. C., & O'Neil, B. J. (2019). Effects of respite music during recovery between
19	bouts of intense exercise. Sport, Exercise, and Performance Psychology. Advance
20	online publication. doi:10.1037/spy0000161
21	Jones, L., Tiller, N. B., & Karageorghis, C. I. (2017). Psychophysiological effects of music
22	on acute recovery from high-intensity interval training. Physiology & Behavior, 170,
23	106–114. doi:10.1016/j.physbeh.2016.12.017
24	Karageorghis, C. I. (2016). The scientific application of music in exercise and sport: Towards
25	a new theoretical model. In A. M. Lane (Ed.), Sport and exercise psychology (2nd ed.,
26	pp. 274–320). Abingdon, UK: Taylor & Francis.
27	Karageorghis, C. I. (2017). Applying music in exercise and sport. Champaign, IL: Human
28	Kinetics.

1	Karageorghis, C. I., Bruce, A. C., Pottratz, S. T., Stevens, R. C., Bigliassi, M., & Hamer, M.
2	(2018). Psychological and psychophysiological effects of recuperative music
3	postexercise. Medicine & Science in Sports & Exercise, 50, 739–746.
4	doi:10.1249/MSS.0000000 000001497
5	Karageorghis, C. I., Ekkekakis, P., Bird, J. M., & Bigliassi, M. (2017). Music in the exercise
6	and sport domain: Conceptual approaches and underlying mechanisms. In M. Lesaffre,
7	PJ. Maes, & M. Leman (Eds.), The Routledge companion to embodied music
8	interaction (pp. 284–293). New York, NY: Routledge.
9	Karageorghis, C. I., Jones, L. Priest, D. L., Akers, R. I., Clarke, A., Perry, J. M., et al. (2011).
10	Revisiting the relationship between exercise heart rate and music tempo preference.
11	Research Quarterly for Exercise and Sport, 82, 274–284.
12	Karageorghis, C. I., Mouzourides, D. A., Priest, D. L., Sasso, T. A., Morrish, D. J., & Walley,
13	C. L. (2009). Psychophysical and ergogenic effects of synchronous music during
14	treadmill walking. Journal of Sport & Exercise Psychology, 31, 18-36.
15	doi.org/10.1123/jsep.31.1.18
16	Karageorghis, C. I., Priest, D. L., Terry, P. C., Chatzisarantis, N. L., & Lane, A. M. (2006).
17	Redesign and initial validation of an instrument to assess the motivational qualities of
18	music in exercise: The Brunel Music Rating Inventory-2. Journal of Sports Sciences,
19	24, 899–909. doi:10.1080/02640410500298107
20	Karageorghis, C. I., & Terry, P. C. (2011). Inside sport psychology. Champaign, IL: Human
21	Kinetics.
22	Karageorghis, C. I., & Terry, P. C. (1997). The psychophysical effects of music in sport and
23	exercise: A review. Journal of Sport Behavior, 20, 54-68.
24	Kellmann, M., & Beckmann, J. (Eds.). (2017). Sport, recovery, and performance:
25	Interdisciplinary insights. New York, NY: Routledge.
26	Kendzierski, D., & DeCarlo, K. J. (1991). Physical Activity Enjoyment Scale: Two
27	validation studies. Journal of Sport & Exercise Psychology, 13, 50-64.
28	doi:10.1123/jsep.13.1.50

1	Khalfa, S., Roy, M., Rainville, P., Dalla Bella, S., & Peretz, I. (2008). Role of tempo
2	entrainment in psychophysiological differentiation of happy and sad music?
3	International Journal of Psychophysiology, 68, 17–26.
4	doi:10.1016/j.ijpsycho.2007.12.001
5	Little, J. P., Safdar, A., Wilkin, G. P., Tarnopolsky, M. A., & Gibala, M. J. (2010). A
6	practical model of low-volume high-intensity interval training induces mitochondrial
7	biogenesis in human skeletal muscle: potential mechanisms. The Journal of Physiology,
8	588, 1011–1022. doi:10.1113/jphysiol.2009.181743
9	Magnan, R. E., Kwan, B. M., & Bryan, A. D. (2013). Effects of current physical activity on
10	affective response to exercise: Physical and social-cognitive mechanisms. Psychology
11	& Health, 28, 418-433. doi:10.1080/08870446.2012.733704
12	Martin, A. M., & Woods, C. B. (2012). What sustains long-term adherence to structured
13	physical activity after a cardiac event? Journal of Aging and Physical Activity, 20, 135-
14	147. doi:10.1123/japa.20.2.135
15	Porszasz, J., Casaburi, R., Somfay, A., Woodhouse, L. J., & Whipp, B. J. (2003). A treadmill
16	ramp protocol using simultaneous changes in speed and grade. Medicine & Science in
17	Sports & Exercise, 35, 1596–1603. doi:10.1249/01.MSS.0000084593.56786.DA
18	Priest, D. L., & Karageorghis, C. I. (2008). A qualitative investigation into the characteristics
19	and effects of music accompanying exercise. European Physical Education Review, 14,
20	347–366. doi:10.1177/1356336X08095670
21	Quindry, J. C., Franklin, B. A., Chapman, M., Humphrey, R., & Mathis, S. (2019). Benefits
22	and risks of high-intensity interval training in patients with coronary artery disease. The
23	American Journal of Cardiology, 123, 1370–1377. doi:10.1016/j.amjcard.2019.01.008
24	Razon, S., Hutchinson, J. C., & Tenenbaum, G. (2012). Effort perception. In G. Tenenbaum,
25	R. Eklund, & A. Kamata (Eds.), Measurement in sport and exercise psychology (pp.
26	265–275). Champaign, IL: Human Kinetics.
27	Stork, M. J., Banfield, L. E., Gibala, M. J., & Martin Ginis, K. A. (2017). A scoping review
28	of the psychological responses to interval exercise: Is interval exercise a viable
29	alternative to traditional exercise? Health Psychology Review, 11, 324-344.
30	doi:10.1080/17437199.2017.1326011

1	Stork, M. J., Karageorghis, C. I., & Martin Ginis, K. A. (2019). Let's Go! Psychological,
2	psychophysical, and physiological effects of music during sprint interval exercise.
3	Psychology of Sport and Exercise, 45, 101547. doi:10.1016/j.psychsport.2019.101547
4	Stork, M. J., Kwan, M. Y., Gibala, M. J., & Martin Ginis, K. A. (2015). Music enhances
5	performance and perceived enjoyment of sprint interval exercise. Medicine & Science
6	in Sports & Exercise, 47, 1052–1060. doi:10.1016/j.psychsport.2019.101547
7	Stork, M. J., & Martin Ginis, K. A. (2017). Listening to music during sprint interval exercise:
8	The impact on exercise attitudes and intentions. Journal of Sports Sciences, 35, 1940-
9	1946. doi:10.1080/02640414.2016.1242764
10	Svebak, S., & Murgatroyd, S. (1985). Metamotivational dominance: A multimethod
11	validation of reversal theory constructs. Journal of Personality and Social Psychology,
12	48, 107–116. doi:10.1037/0022-3514.48.1.107
13	Tabachnick, B.G., & Fidell, L.S. (2018). Using multivariate statistics (7th ed.). Boston, MA:
14	Pearson.
15	Tammen, V. V. (1996). Elite middle and long distance runner's associative/dissociative
16	coping. Journal of Applied Sport Psychology, 8, 1-8. doi:10.1080/10413209608406304
17	Terry, P. C., Karageorghis, C. I., Curran, M. L., Martin, O. V., & Parsons-Smith, R. L.
18	(2020). Effects of music in exercise and sport: A meta-analytic review. Psychological
19	Bulletin, 146, 91-117. doi:10.1037/bul0000216
20	Thomas, J. R., Nelson, J. K., & Silverman, S. J. (2015) Research methods in physical activity
21	(7th ed.). Champaign, IL: Human Kinetics.
22	World Health Organization. (2010). Global recommendations on physical activity for health.
23	World Health Organization. https://www.who.int/dietphysicalactivity/publications/9789241599979/en/
24	Zenko, Z., Ekkekakis, P., & Ariely, D. (2016). Can you have your vigorous exercise and
25	enjoy it too? Ramping intensity down increases postexercise, remembered, and
26	forecasted pleasure. Journal of Sport & Exercise Psychology, 38, 149–159.
27 28	doi:10.1123/jsep.2015-028

1 Table 1

2 Descriptive Statistics (M [SD]) for Dependent Measures Across Conditions for Exercise

			Condition	
	Stage	Control	Medium tempo	Fast tempo
RPE	~81			
	E1	3.74 (1.63)	3.96 (1.15)	3.52 (1.16)
	R1	3.61 (1.62)	2.39 (0.89)	3.13 (1.22)
	E2	5.13 (1.36)	4.87 (1.22)	4.52 (1.08)
	R2	3.78 (1.13)	3.04 (0.93)	3.17 (0.89)
	E3	5.65 (1.23)	4.91 (1.00)	5.00 (1.24)
	R3	4.52 (1.12)	3.52 (0.99)	3.87 (0.82)
	E4	6.43 (1.27)	5.65 (1.07)	6.22 (0.95)
	R4	4.57 (1.04)	3.74 (1.18)	4.65 (0.94)
	E5	6.74 (0.86)	5.17 (1.40)	6.43 (0.95)
	R5	5.35 (1.11)	4.26 (0.86)	5.04 (1.15)
	E6	7.61 (1.27)	6.48 (1.16)	7.17 (1.40)
	R6	5.52 (1.28)	4.22 (1.24)	5.65 (1.34)
	E7	8.09 (1.20)	6.78 (1.13)	7.70 (1.22)
	R7	7.13 (1.33)	5.04 (1.11)	6.52 (1.24)
	E8	9.04 (0.88)	7.43 (1.08)	8.87 (1.10)
	R8	7.26 (1.63)	4.74 (1.74)	7.13 (1.60)
State Attention			. ,	
	E1	58.70 (17.66)	53.48 (15.84)	51.74 (15.57)
	R 1	55.22 (16.20)	69.57 (15.52)	68.26 (14.66)
	E2	48.26 (17.23)	50.00 (11.28)	49.13 (15.35)
	R2	53.74 (15.83)	69.13 (10.84)	65.65 (19.03)
	E3	43.48 (11.52)	45.22 (9.94)	41.74 (14.35)
	R3	47.83 (17.57)	63.48 (11.52)	56.52 (15.55)
	E4	37.39 (14.21)	37.39 (12.51)	40.87 (11.64)
	R4	49.57 (16.37)	56.52 (12.29)	57.39 (16.02)
	E5	33.04 (13.63)	32.61 (11.76)	36.96 (13.63)
	R5	43.48 (13.69)	51.30 (13.92)	48.26 (16.42)
	E6	26.52 (12.29)	29.13 (10.41)	27.83 (10.43)
	R6	39.57 (15.52)	46.96 (17.17)	43.91 (14.06)
	E7	21.74 (9.84)	25.22 (12.01)	21.30 (10.58)
	R7	28.26 (11.54)	41.74 (15.86)	40.00 (15.95)
	E8	17.83 (8.51)	20.87 (9.96)	20.00 (12.79)
	R8	36.52 (18.24)	44.78 (15.04)	41.30 (20.74)
Feeling Scale				
-	E1	1.26 (1.71)	2.57 (1.56)	1.48 (1.31)
	R1	1.39 (1.12)	2.09 (1.08)	1.74 (1.21)
	E2	0.57 (1.08)	1.57 (1.24)	0.7 (1.46)
	R2	1.17 (1.11)	1.91 (0.79)	1.26 (1.39)
	E3	-0.43 (1.41)	0.78 (0.90)	0.13 (1.29)
	R3	0.39 (0.94)	1.39 (1.12)	1 (1.24)
	E4	-1.04 (1.22)	0 (1.21)	-0.22 (1.20)
	R4	-0.17 (1.15)	1 (1.21)	0.22 (1.38)
	E5	-1.74 (1.01)	-0.13 (1.66)	-0.61 (0.99)
	R5	-1.04 (1.26)	0.43 (1.12)	-0.43 (1.41)
				(continued)

3 Bouts (E) and Recovery Periods (R) Through Eight Stages of HIIT

Table 1 (continued)					
		Condition			
	Stage	Control	Medium tempo	Fast tempo	
	E6	-2.78 (0.90)	-0.74 (1.18)	-2.17 (1.44)	
	R6	-1.57 (1.24)	0.04 (1.11)	-0.96 (1.40)	
	E7	-2.78 (1.13)	-1.57 (1.04)	-2.04 (1.55)	
	R7	-2.48 (1.04)	-0.61 (1.23)	-1.91 (1.51)	
	E8	-3.43 (1.08)	-2.17 (1.50)	-2.78 (1.35)	
	R8	-2.13 (1.69)	-1.17 (1.72)	-1.65 (1.72)	
Felt Arousal Scale					
	E1	2.30 (1.11)	2.13 (0.97)	1.96 (0.88)	
	R1	3.30 (0.97)	2.61 (0.58)	2.48 (0.90)	
	E2	3.78 (1.41)	2.65 (0.83)	3.04 (1.11)	
	R2	2.96 (0.71)	2.83 (0.83)	2.87 (1.01)	
	E3	3.43 (0.95)	3.48 (1.16)	2.87 (1.06)	
	R3	3.00 (1.00)	2.96 (0.64)	2.83 (0.89)	
	E4	3.35 (1.15)	3.39 (0.84)	3.17 (0.83)	
	R4	2.87 (1.22)	3.26 (0.81)	3.22 (0.74)	
	E5	3.87 (1.18)	3.65 (0.89)	3.39 (1.16)	
	R5	3.26 (1.21)	3.35 (0.89)	3.48 (0.67)	
	E6	3.65 (1.53)	3.57 (1.08)	4.35 (0.78)	
	R6	3.39 (1.16)	3.83 (0.65)	4.35 (0.94)	
	E7	4.04 (1.43)	4.39 (1.08)	4.83 (0.98)	
	R7	3.65 (1.34)	3.96 (0.88)	4.13 (0.92)	
	E8	4.00 (1.17)	4.78 (0.95)	4.91 (0.67)	
	R8	3.57 (1.27)	3.52 (1.16)	3.78 (1.09)	
Heart Rate (%max)					
	E1	67.77 (4.44)	74.02 (4.20)	70.69 (6.14)	
	R1	69.90 (6.12)	71.44 (3.14)	73.04 (7.19)	
	E2	77.75 (4.82)	77.27 (3.17)	76.72 (6.36)	
	R2	77.84 (4.97)	76.92 (2.87)	74.87 (6.05)	
	E3	83.07 (3.59)	82.11 (3.30)	78.90 (5.69)	
	R3	81.89 (4.70)	80.10 (3.03)	78.35 (6.38)	
	E4	85.31 (4.19)	84.38 (3.41)	81.25 (5.81)	
	R4	85.70 (4.58)	83.45 (2.71)	80.80 (6.65)	
	E5	87.67 (4.12)	86.40 (3.52)	84.21 (6.78)	
	R5	87.85 (4.00)	85.45 (3.36)	82.67 (6.87)	
	E6	90.27 (3.86)	88.86 (3.73)	85.32 (7.29)	
	R6	89.62 (4.94)	88.37 (3.73)	85.39 (7.80)	
	E7	92.09 (4.36)	91.41 (4.70)	89.49 (6.56)	
	R7	91.98 (4.54)	91.75 (4.40)	89.47 (6.56)	
	E8	95.33 (4.71)	95.08 (5.00)	92.50 (7.31)	
	R8	91.68 (4.38)	91.11 (3.65)	89.92 (8.73)	
PACES		60.43 (3.87)	78.13 (6.98)	68.96 (5.09)	
Remembered Pleasure		-23.91 (31.15)	13.04 (28.03)	0.87 (29.22)	

Table 1 (continued)

1 Table 2

2 Inferential Statistics for Analyses on Psychological and Psychophysiological Measures
 3

	F	df	р	η_p^2	Source of Difference
RPE (E)					
Condition × Stage	3.49	6.68, 146.96	.002	.14	Medium < Control from Stage 5 onward; Medium < Fast at Stage 5 and 8
Condition	12.40	2, 88	< .001	.36	Control > Medium; Medium < Fast
Stage	151.82	7, 308	< .001	.87	Stage $1-4 < $ Stage $5-8$
RPE (R) Condition × Stage	3.61	6.92, 152.23	< .001	.14	Medium < Control from Stage 5 onward; Medium < Fast Stage 6–8
Condition	31.00	2, 88	< .001	.59	Control > Medium; Medium < Fast
Stage	80.20	4.47, 98.29	< .001	.79	Incr. from Stage 1–8
State Attention (E) Condition × Stage	1.09	6.53, 143.61	.374	.047	
Condition	.102	2, 154	.903	.005	
Stage	69.14	3.58, 143.61	< .001	.76	Decr. from Stage 1-8
State Attention (R) Condition × Stage	.70	14, 308	.770	.31	_
Condition	11.34	11.34, 35.11	< .001	.34	Control < Medium, Fast
Stage	39.24	3.97, 87.22	< .001	.64	Decr. from Stage 2–5; Decr. from Stage 6–8
Feeling Scale (E) Condition × Stage	1.37	6.02, 142.18	.231	.06	
Condition	18.60	2, 44	< .001	.46	Medium > Control, Fast;
Stage	135.17	4.03, 88.69	< .001	.86	Decr. from Stage 1–4; Decr. from Stage 5–8
Feeling Scale (R) Condition × Stage	1.22	6.78, 149.13	.294	.05	_
Condition	20.28	2, 44	< .001	.48	Medium > Control, Fast
Stage	86.91	7, 154	< .001	.80	Decr. from Stage 1–7
Felt Arousal Scale (E) Condition × Stage	4.75	14, 308	< .001	.18	Medium < Control at Stage
Condition	.152	2, 44	.859	.007	2, rast > Control at stage 8
Stage	49.78	3.98, 88.69	< .001	.69	Incr. from Stage 1–7 (continued)

Table 2 (continued)					
	F	df	р	η_p^2	Source of Difference
Felt Arousal Scale (R) Condition × Stage	2.21	14, 308	.007	.09	Fast < Control at Stage 1;
Condition	.550	2, 44	.581	.024	—
Stage	16.11	7, 154	< .001	.42	Incr. between Stages 1 and 3; Incr. between Stages 6 and 8
Heart Rate Average (%max; E)					
Condition × Stage	9.87	3.69, 81.28	< .001	.31	Medium > Control at Stage 1
Condition	4.62	1.53, 33.69	.25	.17	—
Stage	300.50	1.97, 43.43	< .001	.93	Incr. from Stage 1–8
Heart Rate Average $\binom{96}{2}$ max: P)					
Condition \times Stage	6.63	3.94, 86.70	< .001	.23	Fast < Control at Stages 4–5
Condition	3.63	1.46, 32.04	.051	.14	
Stage	187.13	1.42, 31.31	< .001	.90	Incr. from Stage 1–7
Heart Rate Peak					
(% max; E) Condition × Stage	1.92	5.10, 112.27	.095	.08	
Condition	1.61	1.43, 31.52	.217	.07	
Stage	303.21	2.33, 51.36	<.001	.93	Incr. from Stage 1–7
Heart Rate Low					
(%) Condition × Stage	244.16	4.10, 90.13	< .001	.08	Control > Fast at Stage 4–5
Condition	3.72	2, 44	.032	.15	—
Stage	256.98	1.44, 31.73	< .001	.92	Incr. from Stage 1–7
PACES	14.62	2, 44	< .001	.79	Medium > Control, Fast; Fast > Control
Remembered Pleasure	14.62	1.52, 33.35	< .001	.40	Medium > Control; Fast > Control

Note. E = Exercise bout, R = Recovery bout, Incr = Increase, Decr = Decrease.







Figure 2. State attention responses across conditions during HIIT exercise bouts (top) and
 recovery periods (bottom). T-bars denote ± 2 standard errors.



- 23 24
- 24 25
- *Figure 3*. Feeling Scale (*x*-axis) and Felt Arousal Scale (*y*-axis) scores plotted on the Circumplex Model of Affect for three conditions. The arrows denote the "affective journey" through the eight stages of the HIIT protocol. The first stage is indicated with an "s" and the final stage
- with an "e".

Supplementary Table 1

Artist	Track Name	HIIT Recovery Period
Medium-tempo music (120–125 bpm)		
Mike Mago, Dragonette	Secret Stash	1
Nyte	Fall For You	2
Loud Luxury	Body	3
Otto Knows (ft. Lindsey Stirling and		
Alex Aris	Dying For You	4
Robin Shulz (ft. Erika Sirola)	Speechless	5
Jack Wins (ft. Amy Grace)	Forever Young	6
Sigala, Ella Eyre	Came Here For Love	7
Calvin Harris, Sam Smith	Promises	8
Fast-tempo music (135–140 bpm)		
Davey Asprey	Fallout	1
Gareth Emery, Emma Hewitt	Take Everything	2
Markus Schulz	Facedown	3
Paul van Dyk, Ronald van Gelderen		
(ft. Eric Lumiere)	Everyone Needs Love	4
Saad Ayub (ft. Christina Novelli)	The Only One (Uplifting Mix)	5
Superfitness	Feel This Moment	6
Superfitness	Havana Remix	7
Above & Revond	Sun & Moon	8

Track Lists for Experimental Conditions

Above & BeyondSun & Moon8Note. The tempo of Sun & Moon was digitally altered to 136 bpm. HIIT = high-intensity
interval training.8