

The impact of COVID-19 on health behaviour, well-being, and long-term physical health

MCBRIDE, Emily <<http://orcid.org/0000-0001-9926-429X>>, ARDEN, Madelynne <<http://orcid.org/0000-0002-6199-717X>>, CHATER, Angel <<http://orcid.org/0000-0002-9043-2565>> and CHILCOT, Joseph <<http://orcid.org/0000-0001-6427-4690>>

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/28503/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

MCBRIDE, Emily, ARDEN, Madelynne, CHATER, Angel and CHILCOT, Joseph (2021). The impact of COVID-19 on health behaviour, well-being, and long-term physical health. *British Journal of Health Psychology*, 26 (2), 259-270.

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>



Editorial

The impact of COVID-19 on health behaviour, well-being, and long-term physical health

Statement of contribution Emily McBride and Joseph Chilcot conceived and drafted the article. All authors contributed to the final version and approved the submission.

Background

The COVID-19 pandemic has led to substantial changes in population behaviour and health-related outcomes. Approximately 10% of people who test positive for COVID-19 experience long-term health effects lasting for more than 12 weeks post-infection, including persistent physical, neuropsychological, and mental health-related symptoms (termed Long COVID; Office for National Statistics, 2020). Although outcomes are improving due to better understanding of COVID-19 and clinical advancements, a significant proportion of people with severe disease require hospitalization and experience substantially impaired quality of life (Dennis, McGovern, Vollmer, & Mateen, 2021). There are also indirect effects on health and well-being resulting from unemployment, trauma and stress related to the disease, bereavement, and the implementation of national and local restriction measures to control infection (Arden et al., 2020; Brooks et al., 2020; Cabarkapa, Nadjidai, Murgier, & Ng, 2020; LeRoy, Robles, Kilpela, & Garcini, 2020).

Prior to the pandemic, less than optimal engagement in preventative health behaviours (such as physical activity) and enacting risk behaviours (such as tobacco use) were estimated to account for up to 40% of the determinants of premature mortality (McGinnis, Williams-Russo, & Knickman, 2002; Mokdad, Marks, Stroup, & Gerberding, 2004; Puterman et al., 2020). Furthermore, the combined impact of multiple preventative healthy behaviours appears to further attenuate mortality risk (Ford, Bergmann, Boeing, Li, & Capewell, 2012). Recent data from the UK Biobank reports that engagement in healthy behaviours was associated with increased life expectancy by up to 7.6 years for women and 6.3 years for men. These findings were consistent in people with and without multi-morbidity (Chudasama et al., 2020).

Since the onset of COVID-19, negative changes to some health behaviours have been inevitable with large segments of the population under prolonged conditions of self-isolation, shielding, and physical distancing. Socio-economic factors such as reduced

income and job loss, as well as increased caring responsibilities (particularly for women), have posed significant barriers to engaging with health promoting behaviours, as has the impact of poorer physical and mental health. Based on UK risk factor prevalence estimates, the combined impact of physical inactivity, smoking, and obesity accounted for up to 51% of the population attributable fraction of severe COVID-19 (Hamer, Kivimäki, Gale, & Batty, 2020). Whilst the long-term impacts of COVID-19 on population health and well-being are unknown, most health experts anticipate that the effects will be profound with the most disadvantaged people in society disproportionately burdened. Here, we consider the impact of the pandemic on population health behaviours, well-being, and long-term physical health (including the prevention and detection of chronic disease). We also discuss the need for upstream behaviour change interventions and long-term strategies to mitigate downstream physical and mental health consequences.

COVID-19 vaccination uptake

A UK population survey of 12,035 adults, between 24 November and 1 December 2020, indicated that 18% reported being hesitant (unlikely or very unlikely) to receive a COVID-19 vaccine (Robertson et al., 2021). General mistrust in vaccines and concerns about future side effects have been listed as key barriers to optimizing population immunity (Paul, Steptoe, & Fancourt, 2020). Another UK survey between September and October 2020 reported that 11.7% of the 5,114 respondents were hesitant to receive a COVID-19 vaccine. A large proportion of the variance (86%) in COVID-19 vaccine hesitancy was explained by perceptions of lower collective importance, lower vaccine efficacy, potential side effects, and concerns regarding the speed of vaccine development. Vaccine hesitancy was associated with socio-demographic factors including female gender, younger age, lower income, and ethnicity, but overall, these only explained 9.8% of the variance in the hesitation to get the vaccine (Freeman et al., 2020). In contrast, data in clinically vulnerable populations confirm high willingness for vaccination. For example, in UK patients with chronic respiratory disease, 86% of respondents wanted to receive the vaccine (Williams et al., 2020). Willingness to receive the vaccine has been associated with beliefs regarding the value to personal health, health consequences for others, concerns regarding vaccine safety, and perceptions of COVID-19 severity (Paul et al., 2020; Williams et al., 2020). Since the vaccine was rolled out across England in priority populations, preliminary data have shown lower vaccination rates among ethnic minority groups (those with Black ethnicity displaying lowest rates), people living in areas of higher deprivation, and those with severe mental illness or learning disabilities (MacKenna et al., 2021).

Eating behaviour and nutrition

Data from a large study of 22,374 UK adults found that one third reported changes to quantities of food consumed through the first lockdown period. Subgroups of the population were found to persistently eat more, whilst others reported eating less (particularly those who were already underweight; Herle, Smith, Bu, Steptoe, & Fancourt, 2021). Another study found that 56% of participants reported snacking more frequently (Robinson et al., 2021). Food products consumed also appear to have shifted, with a UK study finding that one third of participants reported eating less healthily than usual (McAtamney, Mantzios, Egan, & Wallis, 2021). A recent study indicated that people have

consumed on average one portion of fruit and vegetables less per day than before the pandemic (Naughton et al., 2021). A potential mechanism for changes to eating behaviours may be emotion dysregulation, with those reporting changes to their eating behaviours during the pandemic also displaying greater levels of depression (Herle et al., 2021; McAtamney et al., 2021). Barriers to weight management have, for example, included reports of lower motivation to eat healthily and lower control over eating (Robinson et al., 2021).

Physical activity and sedentary behaviour

An online UK survey of 9,190 adults found that a quarter reported lower levels of physical activity since COVID-19 (Rogers et al., 2020). Vulnerable populations reported engaging in around half a day less of 30 min of moderate-to-vigorous physical activity each week through the first lockdown period (Naughton et al., 2021). Another UK survey found that 57% had either maintained or increased their levels of physical activity during lockdown. However, only a third met recommended physical activity guidelines of 150 min of moderate-to-vigorous physical activity per week (Spence et al., 2021). Changes in physical activity were found to be associated with components of the COM-B model (Michie, van Stralen, & West, 2011), notably reduced physical opportunity and reflective motivation (Spence et al., 2021). Sedentary behaviour, such as prolonged periods of sitting, has also risen due to government mandates to stay at home (Wedig, Duelge, & Elmer, 2020). Sedentary behaviour is now acknowledged as an independent risk factor for poor cardiometabolic health that should be considered in addition to physical inactivity (Dempsey, Owen, Yates, Kingwell, & Dunstan, 2016). Daily sitting time increases the risk of type 2 diabetes, cardiovascular disease, and mortality (Bailey, Hewson, Champion, & Sayegh, 2019; Katzmarzyk, Church, Craig, & Bouchard, 2009), independent of physical activity levels. There is good evidence that breaking up sitting can benefit cardiometabolic health (Bailey, Mugridge, Dong, Zhang, & Chater, 2020; Brierley, Chater, Smith, & Bailey, 2019) and also improve mood and psychological well-being (Bailey et al., 2020).

Tobacco smoking, alcohol consumption, and substance use

A national survey of 2,254 adults in the United Kingdom indicated that one third (31%) reported drinking more alcohol through COVID-19 than normal (King's College London, 2020). Around 24% of people with pre-existing alcohol disorders, at high risk of relapse, reported increasing their alcohol intake (Kim et al., 2020). Data regarding substance use are more limited, with preliminary reports suggesting increased rates of drug use for substances such as cannabis, prescription benzodiazepines, and prescription opioids. Also, higher relapse rates have been observed for those in recovery from addiction (Society for the Study of Addiction, 2020).

In contrast to other behaviours, emerging evidence has suggested that tobacco quit rates and attempted quit rates have improved since the pandemic. The charity Action on Smoking and Health reported that over 1 million people had stopped smoking by July 2020, and 440,000 had attempted to quit. Increased quit rates may be due to increased motivation as a result of evidence and awareness that there is a higher risk of severe COVID-19 and hospitalization in smokers (Action on Smoking & Health, 2020). This suggests that the pandemic may present an opportune moment for promoting tobacco control policies and smoking cessation services.

Preventative health services and help-seeking

Many preventative health services have seen declines in attendance and delays in help-seeking for symptoms (Jones et al., 2020; Rees et al., 2020). For example, bowel, breast, and cervical cancer screening programmes in the United Kingdom were temporarily paused or have been operating a reduced service at different stages of the pandemic. By September 2020, an estimated 3 million people in the United Kingdom were overdue for screening (Cancer Research UK, 2020). The numbers urgently referred to secondary care with suspected cancer symptoms had also starkly declined at the height of the pandemic; though figures are now improving, they are still lower than before the pandemic. This was largely due to fewer people visiting their GP with suspected cancer symptoms but also because many GPs were reluctant to risk sending people to hospitals (Cancer Research UK, 2020). Furthermore, there have been reductions in patient engagement across general health services such as visits to Accidents and Emergency (Thornton, 2020). Some reasons reported for non-attendance have included fear of COVID-19 infection, not wanting to burden the health system, and practical barriers (Jo's Cervical Cancer Trust, 2020; NHS England, 2020).

Sleep hygiene

Increased prevalence of sleep disorders through COVID-19 has been highlighted in the research literature across different countries. Studies have examined the effect on sleep of SARS-CoV-2 infection and explored confounders related to isolation, quarantine, anxiety, stress, or financial losses (Partinen, 2021). It is thought that symptoms of insomnia could be related to psychosocial factors and the impact of confinement (Altena et al., 2020). A large UK study found poor quality sleep was associated with the occurrence of adverse events through the pandemic. Adverse events included, for example, illness with COVID-19, financial difficulty, loss of paid work, problems with sourcing medicine, difficulties accessing food, and perceived threats to personal safety (Wright, Steptoe, & Fancourt, 2020).

Mental health and well-being

The impact of the pandemic has had a profound effect upon mental health and well-being (Horesh, Kapel Lev-Ari, & Hasson-Ohayon, 2020; Lades, Laffan, Daly, & Delaney, 2020; McElroy et al., 2020; Pierce et al., 2020). Data in both the United Kingdom and the United States suggest that symptoms of psychological distress have increased during the pandemic, with people aged between 18 and 24 years showing the greatest deterioration (McGinty, Presskreischer, Han, & Barry, 2020; Pierce et al., 2020). Whilst increased psychological distress is a normal and understandable response for many given these challenging circumstances, the consequences and impact of COVID-19 provide significant risk factors for clinically relevant anxiety and depression (Holmes et al., 2020).

Increased prevalence of social isolation and loneliness (MQ: Transforming Mental Health & the Academy of Medical Sciences, 2021) through COVID-19 remains considerable risk factors for poorer mental health (O'Connor & Kirtley, 2018). Furthermore, social isolation and loneliness are associated with physical health outcomes, including being risk factors for type 2 diabetes (Hackett, Hudson, & Chilcot, 2020), obesity (Whisman, 2010), and incident coronary heart disease (Valtorta, Kanaan, Gilbody, Ronzi, & Hanratty, 2016). Many individuals with existing long-term conditions (LTCs) are

extremely clinically vulnerable and have been shielding for a considerable period of time, with patient surveys reporting heightened feelings of psychological distress and loneliness. Longitudinal data studies from the United States report that perceptions regarding health risks related to COVID-19 partially explained the variability of psychological distress during the pandemic (Robinson & Daly, 2020). Furthermore, people living with LTCs have had disruptions to the accessibility and delivery of their usual care, compounded further in those needing to shield. Compared with the general population, people living with LTCs are more likely to have symptoms of depression and anxiety, which are associated with adverse clinical outcomes and events (The King's Fund, 2012). Whilst the longer-term mental health impact of COVID-19 on those with pre-existing LTCs is not yet known, research and clinical interventions are needed to better understand and mitigate these effects.

Impact of COVID-19 on inequalities

The pandemic and COVID-19 are disproportionately burdening low socio-economic and black and minority ethnic (BAME) groups, as well as women and others with caring responsibilities (Chadeau-Hyam et al., 2020; Power, 2020; Raisi-Estabragh et al., 2020). As has been highlighted in various Government reports and the recent COVID-19 Marmot Review, there is particular concern around rapidly widening health and social inequalities (Marmot, Allen, Goldbatt, Herd, & Morrison, 2020; UK Government, 2020). Several COVID-19 and wider research studies have already shown that socio-economic status, ethnicity, gender, age, and education are predictive of poorer health behaviours and health-related outcomes (Bann et al., 2020; Marteau, Rutter, & Marmot, 2021). In addition, people in these groups display the highest risk of SARS-CoV-2 infection, hospitalization, and death from COVID-19 (Marmot et al., 2020). The combined effects of worsening behavioural outcomes paired with higher risk of non-communicable disease and severe COVID-19 are likely to accentuate existing inequities with long-lasting systemic effects.

The need for tailored interventions and long-term prevention strategies

Overall, the pandemic has been associated with adverse population trends across most health behaviours, as well as poorer mental health and well-being outcomes. Although these trends are so far preliminary, the UK Government and international bodies anticipate long-term systemic health and economic effects, far beyond the direct effects of COVID-19 (UK Government, 2020). Poor health behaviours and disengagement with, or lack of access to, health services are likely to contribute to subsequent downstream physical and mental health consequences, carrying significant wider social and economic implications. Action must now be taken to mitigate the impact of COVID-19 on health behaviours, mental well-being, and the prevention of LTCs. Emphasis should be placed on developing, testing, and implementing cost-effective interventions that support health behaviour change and promote health and well-being in individuals, communities, populations, and systems. Particular consideration should be given to meeting the needs of high-risk groups and underserved populations where inequities are already known and further anticipated. Optimal modes of delivery for interventions also need to be carefully assessed to ensure that preventative measures and support services reach all the populations they are targeting, whilst factoring in potential restrictions or access barriers in the COVID-19 context (e.g., telephone, digital applications, online, in-person).

Evidence from the existing health psychology and behavioural science literature can be used as a foundation and springboard to inform interventions and translate them into the COVID-19 context. For example, already there have been systematic reviews on effective messaging which encourages vaccination uptake (Lawes-Wickwar et al., 2021) and on psychological and behavioural responses to testing positive for other viruses (Luo, Chua, Xiong, Ho, & Ho, 2020; McBride et al., 2020). There is a wealth of established behaviour change techniques and interventions targeting health behaviours such as physical activity and healthy eating (Howlett, Trivedi, Troop, & Chater, 2019; Michie, Ashford, et al., 2011; Rhodes, Boudreau, Josefsson, & Ivarsson, 2020); tobacco smoking (Michie, Hyder, Walia, & West, 2011; Roberts, Kerr, & Smith, 2013); and alcohol consumption (Michie et al., 2012). Strategies to increase engagement with preventative health services, such as cancer screening, have been considered extensively (Camilloni et al., 2013; Sabatino et al., 2012; Tsipa et al., 2020). Similar principles can be carried forward and integrated with the proliferating and emerging COVID-19 literature to design context-specific tailored behaviour change and psychological interventions. These can also be used to inform the development of potential services or treatments for Long COVID.

Importantly, in tandem with evidence-based practice, nationally recommended inclusivity and diversity frameworks such as 'The INCLUDE Ethnicity Framework' should be used to guide research design and intervention testing (Trial Forge; Witham et al., 2020). Guidelines produced by networks such as 'EQUATOR' can be used to promote transparent and accurate reporting of health research (EQUATOR Network). Patient and public involvement, which adopt representative sampling and ensure inclusion from high-risk and underserved groups, will also be key to improving intervention quality and acceptability (Greenhalgh et al., 2019; Holmes et al., 2019).

Conclusion

Long-term disease prevention and detection strategies should be prioritized now that most of the UK have been living under policy restrictions for over a year, and the COVID-19 vaccine is rolling out nationally. Cost-effective and evidence-based interventions that support the uptake of healthy behaviours and promote engagement with preventative health services are needed to offset indirect consequences of COVID-19 on disease prevention and mental well-being. Emphasis should be placed on ensuring future research and intervention development align to meet the needs of high-risk groups and underserved populations, to help reduce widening health and social inequalities.

Emily McBride*¹ , **Madelyne A. Arden**² , **Angel Chater**³  and **Joseph Chilcot***⁴ 

¹Institute of Epidemiology and Health Care, University College London, UK; ²Centre for Behavioural Science and Applied Psychology, Sheffield Hallam University, UK; ³Centre for Health, Wellbeing and Behaviour Change, University of Bedfordshire, Bedford, UK and

⁴Institute of Psychiatry, Psychology & Neuroscience, King's College London, UK

E-mail: e.mcbride@ucl.ac.uk

References

- Action on Smoking and Health. (2020). A million people have stopped smoking since the COVID pandemic hit Britain. Retrieved from <https://ash.org.uk/media-and-news/press-releases-media-and-news/pandemicmillion/>
- Altena, E., Baglioni, C., Espie, C. A., Ellis, J., Gavrilloff, D., Holzinger, B., . . . Riemann, D. (2020). Dealing with sleep problems during home confinement due to the COVID-19 outbreak: Practical recommendations from a task force of the European CBT-I Academy. *Journal of Sleep Research, 29*(4), e13052. <https://doi.org/10.1111/jsr.13052>
- Arden, M. A., Byrne-Davis, L., Chater, A., Hart, J., McBride, E., & Chilcot, J. (2020). The vital role of health psychology in the response to COVID-19. *The British Journal of Health Psychology, 25*, 831–838. <https://doi.org/10.1111/bjhp.12484>
- Bailey, D. P., Hewson, D. J., Champion, R. B., & Sayegh, S. M. (2019). Sitting time and risk of cardiovascular disease and diabetes: A systematic review and meta-analysis. *American Journal of Preventive Medicine, 57*, 408–416. <https://doi.org/10.1016/j.amepre.2019.04.015>
- Bailey, D. P., Mugridge, L. H., Dong, F., Zhang, X., & Chater, A. M. (2020). Randomised controlled feasibility study of the MyHealthAvatar-diabetes smartphone app for reducing prolonged sitting time in type 2 diabetes mellitus. *International Journal of Environmental Research and Public Health, 17*, 4414. <https://doi.org/10.3390/ijerph17124414>
- Bann, D., Villadsen, A., Maddock, J., Hughes, A., Ploubidis, G., Silverwood, R., & Patalay, P. (2020). Changes in the behavioural determinants of health during the coronavirus (COVID-19) pandemic: Gender, socioeconomic and ethnic inequalities in 5 British cohort studies. *medRxiv, 2020.2007.2029.20164244*. <https://doi.org/10.1101/2020.07.29.20164244>
- Brierley, M. L., Chater, A. M., Smith, L. R., & Bailey, D. P. (2019). The effectiveness of sedentary behaviour reduction workplace interventions on cardiometabolic risk markers: A systematic review. *Sports Medicine, 49*, 1739–1767. <https://doi.org/10.1007/s40279-019-01168-9>
- Brooks, S. K., Webster, R. K., Smith, L. E., Woodland, L., Wessely, S., Greenberg, N., & Rubin, G. J. (2020). The psychological impact of quarantine and how to reduce it: Rapid review of the evidence. *Lancet, 395*, 912–920. [https://doi.org/10.1016/S0140-6736\(20\)30460-8](https://doi.org/10.1016/S0140-6736(20)30460-8)
- Cabarkapa, S., Nadjidai, S. E., Murgier, J., & Ng, C. H. (2020). The psychological impact of COVID-19 and other viral epidemics on frontline healthcare workers and ways to address it: A rapid systematic review. *Brain, Behavior, & Immunity – Health, 8*, 100144. <https://doi.org/10.1016/j.bbih.2020.100144>
- Camilloni, L., Ferroni, E., Cendales, B. J., Pezzarossi, A., Furnari, G., Borgia, P., . . . Rossi, P. G. (2013). Methods to increase participation in organised screening programs: A systematic review. *BMC Public Health, 13*, 464. <https://doi.org/10.1186/1471-2458-13-464>
- Cancer Research UK. (2020). What's happened to cancer services during the COVID-19 pandemic? Retrieved from <https://scienceblog.cancerresearchuk.org/2020/09/11/whats-happened-to-cancer-services-during-the-covid-19-pandemic/>
- Chadeau-Hyam, M., Bodinier, B., Elliott, J., Whitaker, M. D., Tzoulaki, I., Vermeulen, R., . . . Elliott, P. (2020). Risk factors for positive and negative COVID-19 tests: A cautious and in-depth analysis of UK biobank data. *International Journal of Epidemiology, 49*, 1454–1467. <https://doi.org/10.1093/ije/dyaa134>
- Chudasama, Y. V., Khunti, K., Gillies, C. L., Dhalwani, N. N., Davies, M. J., Yates, T., & Zaccardi, F. (2020). Healthy lifestyle and life expectancy in people with multimorbidity in the UK Biobank: A longitudinal cohort study. *PLOS Medicine, 17*(9), e1003332. <https://doi.org/10.1371/journal.pmed.1003332>
- Dempsey, P. C., Owen, N., Yates, T. E., Kingwell, B. A., & Dunstan, D. W. (2016). Sitting less and moving more: Improved glycaemic control for type 2 diabetes prevention and management. *Current Diabetes Reports, 16*(11), 114. <https://doi.org/10.1007/s11892-016-0797-4>
- Dennis, J. M., McGovern, A. P., Vollmer, S. J., & Mateen, B. A. (2021). Improving survival of critical care patients with coronavirus disease 2019 in England: A National Cohort Study, March to June 2020*. *Critical Care Medicine, 49*(2), 209–214. <https://doi.org/10.1097/CCM.0000000000004747>

- EQUATOR Network. Enhancing the QUALity and Transparency Of health Research. Retrieved from <https://www.equator-network.org/>
- Ford, E. S., Bergmann, M. M., Boeing, H., Li, C., & Capewell, S. (2012). Healthy lifestyle behaviors and all-cause mortality among adults in the United States. *Preventive Medicine*, *55*(1), 23–27. <https://doi.org/10.1016/j.ypmed.2012.04.016>
- Freeman, D., Loe, B. S., Chadwick, A., Vaccari, C., Waite, F., Rosebrock, L., . . . Lambe, S. (2020). COVID-19 vaccine hesitancy in the UK: The Oxford coronavirus explanations, attitudes, and narratives survey (Oceans) II. *Psychological Medicine*, 1–15. <https://doi.org/10.1017/S0033291720005188>
- Greenhalgh, T., Hinton, L., Finlay, T., Macfarlane, A., Fahy, N., Clyde, B., & Chant, A. (2019). Frameworks for supporting patient and public involvement in research: Systematic review and co-design pilot. *Health Expectations*, *22*, 785–801. <https://doi.org/10.1111/hex.12888>
- Hackett, R. A., Hudson, J. L., & Chilcot, J. (2020). Loneliness and type 2 diabetes incidence: Findings from the English Longitudinal Study of Ageing. *Diabetologia*, *63*, 2329–2338. <https://doi.org/10.1007/s00125-020-05258-6>
- Hamer, M., Kivimäki, M., Gale, C. R., & Batty, G. D. (2020). Lifestyle risk factors, inflammatory mechanisms, and COVID-19 hospitalization: A community-based cohort study of 387,109 adults in UK. *Brain, Behavior, and Immunity*, *87*, 184–187. <https://doi.org/10.1016/j.bbi.2020.05.059>
- Herle, M., Smith, A. D., Bu, F., Steptoe, A., & Fancourt, D. (2021). Trajectories of eating behavior during COVID-19 lockdown: Longitudinal analyses of 22,374 adults. *Clinical Nutrition ESPEN*. <https://doi.org/10.1016/j.clnesp.2021.01.046>
- Holmes, E. A., O'Connor, R. C., Perry, V. H., Tracey, I., Wessely, S., Arseneault, L., . . . Bullmore, E. D. (2020). Multidisciplinary research priorities for the COVID-19 pandemic: A call for action for mental health science. *Lancet Psychiatry*, *7*, 547–560. [https://doi.org/10.1016/S2215-0366\(20\)30168-1](https://doi.org/10.1016/S2215-0366(20)30168-1)
- Holmes, L., Cresswell, K., Williams, S., Parsons, S., Keane, A., Wilson, C., . . . Starling, B. (2019). Innovating public engagement and patient involvement through strategic collaboration and practice. *Research Involvement and Engagement*, *5*(1), 30. <https://doi.org/10.1186/s40900-019-0160-4>
- Horesh, D., Kapel Lev-Ari, R., & Hasson-Ohayon, I. (2020). Risk factors for psychological distress during the COVID-19 pandemic in Israel: Loneliness, age, gender, and health status play an important role. *The British Journal of Health Psychology*, *25*, 925–933. <https://doi.org/10.1111/bjhp.12455>
- Howlett, N., Trivedi, D., Troop, N. A., & Chater, A. M. (2019). Are physical activity interventions for healthy inactive adults effective in promoting behavior change and maintenance, and which behavior change techniques are effective? A systematic review and meta-analysis. *Translational Behavioral Medicine*, *9*(1), 147–157. <https://doi.org/10.1093/tbm/iby010>
- Jones, D., Neal, R. D., Duffy, S. R. G., Scott, S. E., Whitaker, K. L., & Brain, K. (2020). Impact of the COVID-19 pandemic on the symptomatic diagnosis of cancer: The view from primary care. *Lancet Oncology*, *21*, 748–750. [https://doi.org/10.1016/S1470-2045\(20\)30242-4](https://doi.org/10.1016/S1470-2045(20)30242-4)
- Jo's Cervical Cancer Trust. (2020). Impact of COVID-19 on cervical screening. Retrieved from <https://www.jostrust.org.uk/about-us/news-and-blog/blog/impact-covid-19-cervical-screening>
- Katzmarzyk, P. T., Church, T. S., Craig, C. L., & Bouchard, C. (2009). Sitting time and mortality from all causes, cardiovascular disease, and cancer. *Medicine and Science in Sports and Exercise*, *41*, 998–1005. <https://doi.org/10.1249/MSS.0b013e3181930355>
- Kim, J. U., Majid, A., Judge, R., Crook, P., Nathwani, R., Selvapatt, N., . . . Lemoine, M. (2020). Effect of COVID-19 lockdown on alcohol consumption in patients with pre-existing alcohol use disorder. *The Lancet Gastroenterology & Hepatology*, *5*, 886–887. [https://doi.org/10.1016/S2468-1253\(20\)30251-X](https://doi.org/10.1016/S2468-1253(20)30251-X)
- King's College London. (2020). Nearly a third of UK public drinking more alcohol than usual during the pandemic [Press release]. Retrieved from <https://www.kcl.ac.uk/news/nearly-a-third-of-uk-public-drinking-more-alcohol-than-usual-during-the-pandemic>

- Lades, L. K., Laffan, K., Daly, M., & Delaney, L. (2020). Daily emotional well-being during the COVID-19 pandemic. *The British Journal of Health Psychology*, *25*, 902–911. <https://doi.org/10.1111/bjhp.12450>
- Lawes-Wickwar, S., Ghio, D., Tang, M. Y., Keyworth, C., Stanescu, S., Westbrook, J., . . . Epton, T. (2021). A rapid systematic review of public responses to health messages encouraging vaccination against infectious diseases in a pandemic or epidemic. *Vaccines (Basel)*, *9*(2), 72. <https://doi.org/10.3390/vaccines9020072>
- LeRoy, A. S., Robles, B., Kilpela, L. S., & Garcini, L. M. (2020). Dying in the face of the COVID-19 pandemic: Contextual considerations and clinical recommendations. *Psychological Trauma*, *12*(S1), S98–S99. <https://doi.org/10.1037/tra0000818>
- Luo, Y., Chua, C. R., Xiong, Z., Ho, R. C., & Ho, C. S. H. (2020). A systematic review of the impact of viral respiratory epidemics on mental health: An implication on the coronavirus disease 2019 pandemic. *Frontiers in Psychiatry*, *11*, 565098. <https://doi.org/10.3389/fpsy.2020.565098>
- MacKenna, B., Curtis, H. J., Morton, C. E., Inglesby, P., Walker, A. J., Morley, J., Goldacre, B. (2021). Trends, regional variation, and clinical characteristics of COVID-19 vaccine recipients: A retrospective cohort study in 23.4 million patients using OpenSAFELY. *medRxiv*, 2021.2001.2025.21250356. <https://doi.org/10.1101/2021.01.25.21250356>
- Marmot, M., Allen, J., Goldbatt, P., Herd, E., & Morrison, J. (2020). *Build Back Fairer: The COVID-19 Marmot Review*. Retrieved from <http://www.instituteofhealthequity.org/resources-reports/build-back-fairer-the-covid-19-marmot-review/build-back-fairer-the-covid-19-marmot-review-full-report.pdf>
- Marteau, T. M., Rutter, H., & Marmot, M. (2021). Changing behaviour: An essential component of tackling health inequalities. *BMJ*, *372*, n332. <https://doi.org/10.1136/bmj.n332>
- McAtamney, K., Mantzios, M., Egan, H., & Wallis, D. J. (2021). Emotional eating during COVID-19 in the United Kingdom: Exploring the roles of alexithymia and emotion dysregulation. *Appetite*, *161*, 105120. <https://doi.org/10.1016/j.appet.2021.105120>
- McBride, E., Tatar, O., Rosberger, Z., Rockliffe, L., Marlow, L. M., Moss-Morris, R., . . . Waller, J. O. (2020). Emotional response to testing positive for human papillomavirus at cervical cancer screening: A mixed method systematic review with meta-analysis. *Health Psychology Review*, 1–35. <https://doi.org/10.1080/17437199.2020.1762106>
- McElroy, E., Patalay, P., Moltrecht, B., Shevlin, M., Shum, A., Creswell, C., & Waite, P. (2020). Demographic and health factors associated with pandemic anxiety in the context of COVID-19. *The British Journal of Health Psychology*, *25*, 934–944. <https://doi.org/10.1111/bjhp.12470>
- McGinnis, J. M., Williams-Russo, P., & Knickman, J. R. (2002). The case for more active policy attention to health promotion. *Health Affairs*, *21*(2), 78–93. <https://doi.org/10.1377/hlthaff.21.2.78>
- McGinty, E. E., Presskreischer, R., Han, H., & Barry, C. L. (2020). Psychological distress and loneliness reported by US adults in 2018 and April 2020. *JAMA*, *324*(1), 93–94. <https://doi.org/10.1001/jama.2020.9740>
- Michie, S., Ashford, S., Sniehotta, F. F., Dombrowski, S. U., Bishop, A., & French, D. P. (2011). A refined taxonomy of behaviour change techniques to help people change their physical activity and healthy eating behaviours: The CALO-RE taxonomy. *Psychology & Health*, *26*, 1479–1498. <https://doi.org/10.1080/08870446.2010.540664>
- Michie, S., Hyder, N., Walia, A., & West, R. (2011). Development of a taxonomy of behaviour change techniques used in individual behavioural support for smoking cessation. *Addictive Behaviors*, *36*, 315–319. <https://doi.org/10.1016/j.addbeh.2010.11.016>
- Michie, S., van Stralen, M. M., & West, R. (2011). The behaviour change wheel: A new method for characterising and designing behaviour change interventions. *Implementation Science*, *6*, 42. <https://doi.org/10.1186/1748-5908-6-42>
- Michie, S., Whittington, C., Hamoudi, Z., Zarnani, F., Tober, G., & West, R. (2012). Identification of behaviour change techniques to reduce excessive alcohol consumption. *Addiction*, *107*, 1431–1440. <https://doi.org/10.1111/j.1360-0443.2012.03845.x>

- Mokdad, A. H., Marks, J. S., Stroup, D. F., & Gerberding, J. L. (2004). Actual causes of death in the United States, 2000. *JAMA*, *291*, 1238–1245. <https://doi.org/10.1001/jama.291.10.1238>
- MQ: Transforming Mental Health and the Academy of Medical Sciences. (2021). *Survey results: Understanding people's concerns about the mental health impacts of the COVID-19 pandemic*. Retrieved from <https://acmedsci.ac.uk/file-download/99436893>
- Naughton, F., Ward, E., Khondoker, M., Belderson, P., Marie Minihane, A., Dainty, J., . . . Notley, C. (2021). Health behaviour change during the UK COVID-19 lockdown: Findings from the first wave of the C-19 health behaviour and well-being daily tracker study. *The British Journal of Health Psychology*. <https://doi.org/10.1111/bjhp.12500>
- NHS England (2020). Help us help you: NHS urges public to get care when they need it. Retrieved from <https://www.england.nhs.uk/2020/04/help-us-help-you-nhs-urges-public-to-get-care-when-they-need-it/>
- O'Connor, R. C., & Kirtley, O. J. (2018). The integrated motivational-volitional model of suicidal behaviour. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *373*, 20170268. <https://doi.org/10.1098/rstb.2017.0268>
- Office for National Statistics. (2020). The prevalence of long COVID symptoms and COVID-19 complications [Press release]. Retrieved from <https://www.ons.gov.uk/news/statementsandletters/theprevalenceoflongcovidsymptomsandcovid19complications>
- Partinen, M. (2021). Sleep research in 2020: COVID-19-related sleep disorders. *Lancet Neurology*, *20*(1), 15–17. [https://doi.org/10.1016/S1474-4422\(20\)30456-7](https://doi.org/10.1016/S1474-4422(20)30456-7)
- Paul, E., Steptoe, A., & Fancourt, D. (2020). Anti-vaccine attitudes and risk factors for not agreeing to vaccination against COVID-19 amongst 32,361 UK adults: Implications for public health communications. *medRxiv*, 2020.2010.2021.20216218. <https://doi.org/10.1101/2020.10.21.20216218>
- Pierce, M., Hope, H., Ford, T., Hatch, S., Hotopf, M., John, A., . . . Abel, K. M. (2020). Mental health before and during the COVID-19 pandemic: A longitudinal probability sample survey of the UK population. *Lancet Psychiatry*, *7*, 883–892. [https://doi.org/10.1016/S2215-0366\(20\)30308-4](https://doi.org/10.1016/S2215-0366(20)30308-4)
- Power, K. (2020). The COVID-19 pandemic has increased the care burden of women and families. *Sustainability: Science, Practice and Policy*, *16*(1), 67–73. <https://doi.org/10.1080/15487733.2020.1776561>
- Puterman, E., Weiss, J., Hives, B. A., Gemmill, A., Karasek, D., Mendes, W. B., & Rehkopf, D. H. (2020). Predicting mortality from 57 economic, behavioral, social, and psychological factors. *Proceedings of the National Academy of Sciences*, *117*, 16273–16282. <https://doi.org/10.1073/pnas.1918455117>
- Raisi-Estabragh, Z., McCracken, C., Bethell, M. S., Cooper, J., Cooper, C., Caulfield, M. J., . . . Petersen, S. E. (2020). Greater risk of severe COVID-19 in Black, Asian and Minority Ethnic populations is not explained by cardiometabolic, socioeconomic or behavioural factors, or by 25(OH)-vitamin D status: Study of 1326 cases from the UK Biobank. *Journal of Public Health*, *42*, 451–460. <https://doi.org/10.1093/pubmed/fdaa095>
- Rees, C. J., Rutter, M. D., Sharp, L., Hayee, B., East, J. E., Bhandari, P., & Penman, I. (2020). COVID-19 as a barrier to attending for gastrointestinal endoscopy: Weighing up the risks. *The Lancet Gastroenterology & Hepatology*, *5*, 960–962. [https://doi.org/10.1016/S2468-1253\(20\)30268-5](https://doi.org/10.1016/S2468-1253(20)30268-5)
- Rhodes, R. E., Boudreau, P., Josefsson, K. W., & Ivarsson, A. (2020). Mediators of physical activity behaviour change interventions among adults: A systematic review and meta-analysis. *Health Psychology Review*, 1–15. <https://doi.org/10.1080/17437199.2019.1706614>
- Roberts, N. J., Kerr, S. M., & Smith, S. M. S. (2013). Behavioral interventions associated with smoking cessation in the treatment of tobacco use. *Health Services Insights*, *6*, HSI.S11092. <https://doi.org/10.4137/HSI.S11092>
- Robertson, E., Reeve, K. S., Niedzwiedz, C. L., Moore, J., Blake, M., Green, M., Benzeval, M. J. (2021). Predictors of COVID-19 vaccine hesitancy in the UK Household Longitudinal Study. *medRxiv*, 2020.2012.2027.20248899. <https://doi.org/10.1101/2020.12.27.20248899>

- Robinson, E., Boyland, E., Chisholm, A., Harrold, J., Maloney, N. G., Marty, L., . . . Hardman, C. A. (2021). Obesity, eating behavior and physical activity during COVID-19 lockdown: A study of UK adults. *Appetite*, *156*, 104853. <https://doi.org/10.1016/j.appet.2020.104853>
- Robinson, E., & Daly, M. (2020). Explaining the rise and fall of psychological distress during the COVID-19 crisis in the United States: Longitudinal evidence from the Understanding America Study. *The British Journal of Health Psychology*. <https://doi.org/10.1111/bjhp.12493>
- Rogers, N. T., Waterlow, N. R., Brindle, H., Enria, L., Eggo, R. M., Lees, S., & Roberts, C. H. (2020). Behavioral change towards reduced intensity physical activity is disproportionately prevalent among adults with serious health issues or self-perception of high risk during the UK COVID-19 lockdown. *Frontiers in Public Health*, *8*. <https://doi.org/10.3389/fpubh.2020.575091>
- Sabatino, S. A., Lawrence, B., Elder, R., Mercer, S. L., Wilson, K. M., DeVinney, B., . . . Glanz, K. (2012). Effectiveness of interventions to increase screening for breast, cervical, and colorectal cancers: Nine updated systematic reviews for the guide to community preventive services. *American Journal of Preventive Medicine*, *43*(1), 97–118. <https://doi.org/10.1016/j.amepre.2012.04.009>
- Society for the Study of Addiction. (2020). How is the COVID-19 pandemic changing our use of illegal drugs? An overview of ongoing research. Retrieved from <https://www.addiction-ssa.org/how-is-the-covid-19-pandemic-changing-our-use-of-illegal-drugs-an-overview-of-ongoing-research/>
- Spence, J. C., Rhodes, R. E., McCurdy, A., Mangan, A., Hopkins, D., & Mummery, W. K. (2021). Determinants of physical activity among adults in the United Kingdom during the COVID-19 pandemic: The DUK-COVID study. *The British Journal of Health Psychology*. <https://doi.org/10.1111/bjhp.12497>
- The King's Fund. (2012). Long-term conditions and mental health: The cost of co-morbidities. Retrieved from https://www.kingsfund.org.uk/sites/default/files/field/field_publication_file/long-term-conditions-mental-health-cost-comorbidities-naylor-feb12.pdf
- Thornton, J. (2020). Covid-19: A&E visits in England fall by 25% in week after lockdown. *BMJ*, *369*, m1401. <https://doi.org/10.1136/bmj.m1401>
- Trial Forge. The INCLUDE Ethnicity Framework. Retrieved from <https://www.trialforge.org/trial-forge-centre/include/>
- Tsipi, A., O'Connor, D. B., Branley-Bell, D., Day, F., Hall, L. H., Sykes-Muskett, B., . . . Conner, M. (2020). Promoting colorectal cancer screening: A systematic review and meta-analysis of randomised controlled trials of interventions to increase uptake. *Health Psychology Review*, *1*–24. <https://doi.org/10.1080/17437199.2020.1760726>
- UK Government. (2020). *Analysis of the health, economic and social effects of COVID-19 and the approach to tiering*. Retrieved from https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/944823/Analysis_of_the_health_economic_and_social_effects_of_COVID-19_and_the_approach_to_tiering_FINAL_-_accessible_v2.pdf
- Valtorta, N. K., Kanaan, M., Gilbody, S., Ronzi, S., & Hanratty, B. (2016). Loneliness and social isolation as risk factors for coronary heart disease and stroke: Systematic review and meta-analysis of longitudinal observational studies. *Heart*, *102*, 1009. <https://doi.org/10.1136/heartjnl-2015-308790>
- Wedig, I. J., Duelge, T. A., & Elmer, S. J. (2020). Infographic. Stay physically active during COVID-19 with exercise as medicine. *British Journal of Sports Medicine*, *55*, 346–347. <https://doi.org/10.1136/bjsports-2020-103282>
- Whisman, M. A. (2010). Loneliness and the metabolic syndrome in a population-based sample of middle-aged and older adults. *Health Psychology*, *29*, 550–554. <https://doi.org/10.1037/a0020760>
- Williams, L., Gallant, A. J., Rasmussen, S., Brown Nicholls, L. A., Cogan, N., Deakin, K., . . . Flowers, P. (2020). Towards intervention development to increase the uptake of COVID-19 vaccination among those at high risk: Outlining evidence-based and theoretically informed future intervention content. *The British Journal of Health Psychology*, *25*, 1039–1054. <https://doi.org/10.1111/bjhp.12468>

- Witham, M. D., Anderson, E., Carroll, C., Dark, P. M., Down, K., Hall, A. S. . . . On behalf of the, I.w. g. (2020). Developing a roadmap to improve trial delivery for under-served groups: Results from a UK multi-stakeholder process. *Trials*, *21*, 694. <https://doi.org/10.1186/s13063-020-04613-7>
- Wright, L., Steptoe, A., & Fancourt, D. (2020). Are adversities and worries during the COVID-19 pandemic related to sleep quality? Longitudinal analyses of 45,000 UK adults. *medRxiv*, 2020.2006.2002.20120311. <https://doi.org/10.1101/2020.06.02.20120311>

Received 8 March 2021