

Nutritional Considerations for Female Rock Climbers

LESLIE-WUJASTYK, Mina and GIBSON-SMITH, Edward
<<http://orcid.org/0009-0002-3984-7228>>

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/33142/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

LESLIE-WUJASTYK, Mina and GIBSON-SMITH, Edward (2024). Nutritional Considerations for Female Rock Climbers. *Journal of Science in Sport and Exercise*.

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>



Nutritional Considerations for Female Rock Climbers

Mina Leslie-Wujastyk¹ · Edward Gibson-Smith²

Received: 17 July 2023 / Accepted: 14 November 2023
© The Author(s) 2024

Abstract

Despite growing participation, there is limited research into the nutritional needs of climbers and none specific to female climbers. Female athletes in general are still significantly under-represented in sport and nutritional science research. The physiological requirements of climbing are extensive and variable, demanding both highly developed anaerobic and aerobic energy systems. Finger strength, upper body power and training hours have been highlighted as the key determinants of climbing performance in females. Nutritional implications of this include adequate energy and carbohydrate availability, appropriate protein intake and timing for recovery, and use of specific supplements. As a weight sensitive sport, energy intake and availability are key areas of consideration for female climbers' health and performance. Consideration of macronutrient intake and timing with an understanding of sex hormone interaction across the menstrual cycle confers some considerations to nutritional guidelines. Micronutrients of particular interest to the female climber include iron, vitamin D and calcium. Appropriate supplement use may be beneficial, however more research is needed to provide any female specific dosing strategies. It may be premature to prescribe generalised female specific nutrition recommendations for climbers. A personalised approach that considers the individual's menstrual cycle and experience is recommended. Further research into nutrition for the female climber is warranted.

Keywords Menstrual cycle · Sex hormones · Bouldering · Climbing · Female athlete

Introduction

Climbing debuted at the 2020 Tokyo Olympics and with this acceleration, there is a need for more research into the physiological demands, training practices, and nutritional needs of climbers. There is a significant paucity of literature when it comes to the female climbing population. Climbing includes multiple disciplines; this review will consider the nutritional needs of the female single pitch lead climber incorporating research that covers bouldering due to the similarities within these disciplines.

Although no nutritional guidelines specific to female climbers currently exist, there have been recent studies into the dietary intake of high performing female climbers [19, 46] and their anthropometric characteristics [22]. Previous research into dietary intakes of adult climbers either

exclusively studied males or did not state the gender of participants [35, 75]. To date, there are two review papers that discuss the nutritional demands for climbing, however neither of these have any specific considerations for female climbers beyond highlighting an increased risk for negative consequences of low energy availability (LEA) [44, 62].

Female athletes are still significantly under-represented in sport and nutritional science research and application from male only studies may not be appropriate [31, 47, 65, 74]. However, research into specific female athlete nutrition is emerging and will be considered in the context of the physiological demands and current nutritional guidelines for climbing. Table 1 lists a glossary of rock-climbing specific terminology for the benefit of the reader.

Method

Search terms including “nutrition”, “female athlete”, “climbing”, “bouldering”, “supplements”, “hydration”, “performance”, “carbohydrate”, “protein”, “fat” and “menstrual cycle” were used in various combinations in PubMed and Google Scholar. Papers were then selected by relevance and their reference lists examined for use.

✉ Edward Gibson-Smith
edward.smith@shu.ac.uk

¹ Lattice Training Ltd, Chesterfield, UK

² Academy of Sport and Physical Activity, College of Health, Wellbeing and Lifestyle, Sheffield Hallam University, Sheffield, UK

Table 1 Glossary of rock-climbing specific terminology

Term	Definition
Finger-boarding	A training modality that involves hanging from the fingers on a training tool called a “fingerboard” at prescribed intensities by adding or removing weight
Pump	The sensation of increasing fatigue and pressure in the forearm associated with build-up of metabolites and increased anaerobic contribution
Power out	The sensation of fatigue that inhibits further efforts at a high intensity
Crux	The most physically or technically demanding section of a climbing route
Single pitch lead climbing	A discipline of rock climbing that involves a single effort of climbing that can vary from approximately 10–50 m where the climber brings the rope with them, attaching to safety points (quickdraws) along the way but never deriving any assistance from the rope or other equipment. It has mixed physiological needs but is associated with endurance
Bouldering	This is a discipline of rock climbing where the climber climbs on a shorter boulder or rock face above crash pads for approximately 2–12 moves or 2–8 m. It has mixed physiological needs but is associated with strength, power, and anaerobic capacity

Physiological Demands of Rock Climbing

To consider the nutritional needs of female climbers, the physiological demands of the sport must be realised. Lead climbing routes vary in length, with a maximum height of 15 m on artificial walls used for competition and exceeding 50 m on natural rock. Routes may have a “crux” section akin to a high intensity boulder problem interspersed with opportunities to rest, or they may be continuous. Therefore, athletes are required to perform using the phosphocreatine system (PCr), anaerobic glycolysis, and aerobic metabolism interchangeably as the dominant source of energy provision. Climbing training should include optimisation of the PCr and the anaerobic glycolysis systems as well as upper body aerobic power and sport specific muscular strength, notably the finger flexors located in the forearm [62]. The forearm is often the limiting factor in a route attempt, with climbers experiencing “pump” or “power out” sensations indicative of the anaerobic contribution to the activity via repeated bouts of isometric contraction [21]. When considering sport climbing, optimising aerobic capacity is also paramount as rock climbing requires a large fraction of the climber’s peak oxygen uptake with a congruent high heart rate response [59]. This is best measured and trained in the upper body to ensure specificity to sport [45]. To date, we have no data that outlines any variance in the physiological response of climbing in female athletes.

Physiology of Female Athletes

This review will discuss the adult pre-menopausal female climber. For a detailed understanding of the hormonal changes across menstrual cycle phases the review by Davis

et al. [14] is recommended. Changes in nutritional requirements induced by the use of oral contraception is also described elsewhere in the literature [52].

Due to conflicting findings, large variation between studies, small effect sizes and poor quality of research, there is currently no scientific consensus on whether hormonal changes in the menstrual cycle affect a female athlete’s substrate metabolism during exercise and whether this relates to a meaningful performance effect [26]. Small performance changes across the menstrual cycle have been demonstrated, however, based on current evidence, standardised sex-specific guidelines for training and nutrition may be premature [41, 47]. Nevertheless, female athletes do report training and performance changes across the menstrual cycle [63] and a small effect can be meaningful for performance.

The optimal approach at present is to adapt non-female-specific guidelines to the individual in the context of potential hormone related differences and, most importantly, their individual response to hormonal changes [47].

Anthropometrics of Female Rock Climbers

Climbing is perceived as a weight sensitive sport, however, only one study to date has reported a correlation between performance and body fat levels in females [22]. A more consistent outcome could be masked by a degree of homogeneity in the athletic profile of the athletes, the limited span of ability level, and the method of performance assessment (i.e., self-reported ability). Although more data is needed, current recommendations focus on the development of absolute strength and healthy weight management to optimise strength to weight ratio, with finger strength and upper body power highlighted as the key determinants of climbing

performance [22]. This supports previous data that, rather than focusing on body fat levels, trainable variables are key determinants of performance in female climbers [43] and nutritional guidelines should support this.

Nutritional Strategies for Female Rock Climbers

Energy Intake

Appropriate energy intake is a pillar of sports nutrition for optimal athlete performance and health. Energy balance affects body composition, which is relevant to sports like climbing, affected by strength to weight ratio [47]. Current understanding of energy expenditure in climbing suggests MET values of 5.8–8, depending on the difficulty level [1] in addition to the suggestion of 10–11 kcal/min of active climbing time, although a variety of parameters, such as angle and efficiency, affect this estimation [71]. Resting metabolic rate may be higher in the luteal phase compared with the follicular phase although larger studies show less effect [5] and this slight increase may be mediated by increased spontaneous energy intake in the luteal phase [3].

A recent survey suggests female lead climbers are at higher risk for disordered eating, especially at elite level [34]. Gibson-Smith et al. [19] found a mean energy intake in elite female climbers of 2039 kcal/day, with 77.5% of athletes failing to meet predicted energy requirements based on a “moderate” training regime. This value is lower than the energy expenditure recorded for many elite female athletes [47] and similar findings arise in other weight sensitive sports [54]. At least two further studies have confirmed inadequate energy availability in female climbers [12, 46] highlighting the need for education among female climbers around optimal energy intake to avoid the negative consequences to health and performance associated with LEA [48].

Due to the paucity of data in climbing, the huge variety in movement, style, energy system uses and efficiency, it is hard to predict energy cost accurately, particularly as currently used RMR calculations are inaccurate in calculating energy needs of female climbers [11]. Individualised energy targets should be based on the athlete’s individual energy demands and periodized to match their training volume and body composition goals, with a minimum intake of 30–45 cal/kg fat free mass (FFM) to ensure energy availability [48], and where possible, the assessment of endocrine, haematological, metabolic, and bone health, to more accurately monitor and screen for LEA [28].

Macronutrients

Carbohydrate

Daily Intake

Gibson-Smith et al. [19] reported a mean carbohydrate intake of 3.7 g/kg/day and Monedero et al. [46] reported intakes of 3.4 ± 0.7 g/kg/day in female climbers both of which sit at the lower end of the suggested intake of 3–7 g/kg/day for climbers [44] and below the recommendation for boulderers [62]. While not always the case [11], outside of climbing female athletes have been reported to consume 30% less carbohydrate per kg body mass (BM) than their male counterparts, adding to data that suggest female athlete carbohydrate consumption is a key area for nutrition focus [6, 54].

Carbohydrate is essential for both performance and health in female athletes, playing a role in hormonal balance, aerobic and anaerobic metabolism, as well as brain and central nervous system function [58]. Whether female athletes need specific carbohydrate guidelines is unresolved; issues include the use of oral contraceptives in studies that may have otherwise given more insight [33, 47], and issues surrounding research quality including methodological control of ovarian hormones [37]. Female athletes would benefit from a focus on periodised carbohydrate availability according to body mass and training volume to ensure adequate fuel for the work required [58].

Adopting a low carbohydrate, high fat diet has been anecdotally popular amongst the climbing community, contrary to evidence which suggests this approach may have deleterious effects on intermittent high intensity performance [9], primarily due to the downregulation of glycogenolysis [66].

Pre-Exercise Intake

In the early follicular phase, when menstruating, exercise performance may be trivially impaired and women report negative symptoms [41]. Due to lower glycogen storage in this phase, an increase in carbohydrate intake to 6–8 g/kg/day has been suggested to maintain carbohydrate availability for performance [47, 58]. Pre-climbing carbohydrate intake of at least 1 g/kgBM is particularly important in the follicular phase to ensure carbohydrate availability when oxidation rates are higher [50, 74]. For the female climber, this must be considered on an individual basis to correlate with overall energy needs and the balance of other macronutrient intakes.

During Exercise

The benefits of intra-exercise carbohydrate fuelling are well established [7]. Although mostly based on male data, results from studies including female athletes show similar metabolic responses as well as an ergogenic effect across all phases of the menstrual cycle [40, 70]. A single attempt on a lead route is usually < 60 min and intermittent, therefore, intake of small amounts of carbohydrate (~ 20–30 g/hour) are likely sufficient providing the athlete's pre-attempt fuelling is adequate [44, 62]. In the luteal phase, when oestrogen concentrations are high, lower carbohydrate oxidation and greater fat oxidation is seen, with higher glycogen storage at rest and sparing during moderate intensity exercise [26, 50]. This warrants extra focus on ensuring exogenous carbohydrate availability in-sessions so that female climbers can access carbohydrates for higher intensity work [31]. Research suggests that women naturally consume more carbohydrate in the luteal phase [10], however, appetite suppression is seen around exercise [32].

Recovery

Much of the existing data on glycogen resynthesis in female athletes, while similar to males, is confounded by oral contraceptive use [47]. Recovery rates appear to be affected by menstrual cycle phase, with greater markers of muscle damage and inflammation in the follicular compared to luteal phase [25]. Female climbers should ensure recovery carbohydrate intakes in the range of at least 1 g/kg/hour, 0–4 h following training or performance [8, 44] with rapid ingestion of at least 0.75 g/kg post-session [74].

Carbohydrate Loading

Glycogen depletion is a proposed mechanism in the onset of performance decrement during repeated or sustained bouts of high-force muscle contractions; an effect which appears to adversely impact the intramyofibrillar stores within type II muscle fibres to a greater degree when the resistance is > 75% 1RM [30]. As force production of this magnitude is relied upon during on-limit climbing, and greater whole body glycogen stores may be used during long sessions with multiple route attempts, carbohydrate loading may be a relevant protocol for climbers before a redpointing attempt or competitive event, with a low-residue approach favoured to ameliorate impact on body mass [18].

Initially it was thought that female athletes had reduced capacity to store glycogen following a loading protocol, however, later work suggests it is not sex dependent provided sufficient absolute carbohydrate intake of 8–12 g/kg/

day [47]. Although glycogen storage may be reduced in the follicular phase, this difference was not apparent in ranges indicative of a loading protocol and, therefore, loading may be particularly advantageous in this phase to overcome lower resting muscle glycogen [40, 47, 56]. More research investigating carbohydrate loading in the context of climbing is warranted.

'Train Low' Protocols

In male athletes, manipulation of carbohydrate intake to include some restricted states can stimulate cellular adaptation, but this must be balanced with carbohydrate availability for training quality [7]. More female-specific research is needed to see whether these techniques are beneficial, neutral or maladaptive [47]. In the luteal phase, oestrogen causes a reduction in available carbohydrate which can be mediated by eating before exercise [56, 58], therefore, it may be prudent to avoid intentional fasted or carbohydrate restricted training in this phase. If applied, these must be managed carefully, especially in the case of a female climber who, as discussed previously, may be at a higher risk for LEA. Training with low muscle glycogen stores may also negatively influence cellular growth and attenuate strength adaptation [13] and increase inflammation and consequently hepcidin levels which can negatively impact iron absorption [61], a key micronutrient for the female climber [19].

Protein

Daily Intake

A study of 20 female climbers [19], found a mean protein intake of 1.6 g/kg/day which sits within the current suggestions of 1.4–2 g/kg/day for boulderers and 1.2–1.8 g/kg/day for lead climbers [44, 62] and the broader athletic range of 1.2–2 g/kg/day [67]. Interestingly, there was a significant positive correlation in female climbers between climbing ability and protein intake. While rates of muscle protein synthesis do not differ between sexes or across menstrual cycle phases [73], protein catabolism is higher in the luteal phase compared to follicular therefore increased protein intake in this phase is recommended [42, 47, 56, 74].

Post-Exercise Intake and Frequency

Post-exercise protein intakes of ~ 0.3 g/kg have been suggested for female athletes with particular benefit to upper

body strength development; a key determinant in a female climber's performance [22, 42, 53]. This dosing protocol is in line with the current climbing guidelines [44, 62]. Due to higher amino acid oxidation and attenuated muscle protein synthesis in sports with endurance components, such as climbing, female athletes may benefit from a post-exercise protein target of ~0.4 g/kg protein with ~0.3 g/kg consumed every 3–4 h thereafter in the ~24 h recovery period, across all menstrual phases [47]. Use of a high-quality protein powder, such as whey, may be practically advantageous for meeting these targets (Maughan et al., 2018).

Fat

Dietary fats are essential for absorption of fat-soluble vitamins and hormone production while also a fuel source for lower intensity exercise [69]. Female athletes demonstrate a lower respiratory exchange rate during exercise than men indicative of using more fat as fuel [58]. This may be due to female athletes having higher intramuscular triglyceride stores and a greater proportion of type 1 muscle fibres which have more fatty acid transport proteins [38]. Oestrogen increases lipid oxidation creating an increase in free fatty acids available for use in female athletes, evident in the high hormone phase [38, 51]. Higher fat stores and more efficient use of fat as fuel may make female athletes more suited to endurance exercise which would be advantageous for climbing routes ≥ 30 m where climbing duration can exceed 60 min.

There is no specific data for fat intake in climbers with Michael et al. [44] suggesting fat intake in the region of 20%–35% of energy intake. In female athletes, adequate fat intake of no less than 20% of energy intake, especially in the higher hormone phase, is necessary to support the increased use of fat as fuel [74]. Fat intake, provided it does not drop too low, can be manipulated to meet energy intake targets after appropriate carbohydrate and protein levels are determined.

Omega-3 fatty acids may optimise muscle recovery and injury rehabilitation in males, [55], however we do not know if this data transfers to female athletes. Daily supplementation of a DHA rich fish oil (1.75 g DHA + 155 mg EPA) for 4-weeks has been demonstrated to improve neurocognitive function in female athletes [24] which could be a benefit to performance in climbers who need to make rapid, complex decisions while climbing.

Hydration

Current sports nutrition evidence suggests avoiding dehydration of > 2% body mass, however, climbers embarking on shorter routes or bouldering may manage a limit of 3%–5%

before any performance deficit due to the high intensity nature, especially in cold conditions [62, 67].

Regulation of fluid balance differs between sexes; women sweat 18%–34% less, have lower plasma volumes, and lower potassium and sodium sweat losses [58, 72]. Evidence suggests that female hormones affect systems that control thirst, fluid and sodium regulation, although more research is needed [20, 57].

Oestrogen dominance (late follicular phase) promotes heat dissipation while progesterone dominance (luteal phase) promotes heat conservation, and women typically present with a higher core temperature in this phase [57]. Most research indicates that these differences in core temperature do not appear to affect fluid regulation, sweat rates or health risks during exercise [20, 57]. More research is needed to understand the potential hormone related fluid retention in the luteal phase of the menstrual cycle [57], and any associated performance effects.

Estimating fluid needs for the female climber is complex due to large inter-individual variances in hormonal fluctuations across the menstrual cycle, an incomplete understanding of the interactions between female sex hormones and fluid balance, and the variety in the demands and conditions in climbing. It would be prudent to advise on an individual basis by monitoring sweat rates, fluid needs, and changes in body water over the menstrual cycle. Female climbers should aim to start climbing well hydrated and keep urine colour pale by drinking 5–10 mL/kg body weight approximately 2 h before exercise followed by 250 mL/hour with adjustments for conditions such as heat and humidity [44, 67].

Micronutrients

Iron

Iron plays a role in oxygen transport, energy production, and affects exercise capacity and cognitive function [39] with deficiency in female athletes typically reported in the range of ~15%–35% [61]. In climbers, Gibson-Smith et al. [19] reported that 80% of females failed to meet the daily recommended intake (DRI) for iron, 45% had sub-optimal ferritin levels, with 30% meeting the criteria for iron deficiency. In a more recent study, the mean iron intake of female climbers was ~50% of the target DRI (7.6 ± 3.1 mg/day vs. 14.8 mg/day; [46]). In addition, it has been suggested that the DRIs may not be sufficient for an athlete population [61].

Low energy availability can also affect iron status with higher risks in those with heavy menstrual bleeding or those following vegetarian diets [29, 48, 61]. Iron status is associated with reduced training capacity and endurance performance in female rowers [16], which, like climbing, demands a high level of upper-body conditioning.

Exercise induced inflammation stimulates the release of hepcidin, the main iron regulatory protein, which impairs absorption of dietary iron intake for 3–6 h, therefore, a high frequency training schedule can make optimal iron absorption problematic [39]. Hepcidin also displays diurnal variation, with concentrations lower in the morning [68]. To combat deficiency in those at risk, intake of iron should be planned away from training sessions, with a preference for the morning [61]. Menstruation affects iron status through blood loss, moreover, inflammation at this point in the cycle can also increase hepcidin levels, compounding the issue [60]. The potential interaction between female sex hormones, iron metabolism, and post exercise hepcidin response remains unclear and is an area for future research, although maintaining oestrogen levels (through adequate energy availability), may be a positive factor in modulating the hepcidin response and preserving iron status [60, 61].

Due to evidence of a high prevalence of sub-optimal iron levels in female climbers, iron status should be screened and monitored on an individual basis and managed accordingly. Including regular iron-rich, bioavailable foods in the diet such as red meat, seafood or legumes combined with foods that contain vitamin C, along with avoidance of iron inhibitors (e.g. tannins, calcium) in the same meal, should be sufficient to maintain a positive iron status. For those deficient, supplementation may be necessary [39].

Vitamin D and Calcium

Preliminary research suggests vitamin D and calcium intake is inadequate in female climbers [12, 46]. Vitamin D is synthesised in the skin from exposure to sunlight, with only 10%–20% derived from dietary sources. It plays a role in immune health, exercise performance, injury recovery, bone health, muscular strength, and neuromuscular function [67]. Calcium is used by the body for bone metabolism, muscle contraction, and nerve conduction [39].

Oestrogen plays a role in the movement of calcium into the bone, therefore appropriate energy intake and availability to maintain hormonal function, as well as calcium intake, is crucial for bone health [64]. Vitamin D also regulates calcium via stimulation of intestinal calcium absorption and deficiency is associated with stress fractures [36]. Multidirectional load bearing exercise, such as climbing, has an independent positive effect on bone health [23].

Inadequate calcium intake, which is especially prevalent in female athletes [2], is associated with disordered eating, restricted intake, and avoidance of dairy/calcium-rich foods [67]. Vitamin D deficiency is also common in female athletes, especially in those with low energy availability or those who predominantly train indoors [17, 58]. Increasing levels of vitamin D may ameliorate negative symptoms during the premenstrual phase in women who are deficient, which could aid training and performance [27]. Rock climbing is predominantly an outdoor pursuit which may positively affect vitamin D status though increased sunlight exposure, depending on season and latitudes.

An athlete with a varied diet who is not presenting as deficient, does not typically require micronutrient supplementation [4], however this may be too simplistic, especially for vitamin D [15]. Dietary vitamin D has limited availability and absorption and supplemental dosing in the general population, and especially athletes, is controversial [15]. In the UK, the general population is advised to take 400 IU during the winter months, but it has been suggested that this is too low [49] and, although further research is needed, a dose of 1000–2000 IU in female athletes may be more appropriate [15, 31]. Calcium can be managed through dietary intake with a focus on regular intake of calcium rich foods such as dairy [23].

Table 2 outlines nutritional considerations and practical advice for the female climber. Figure 1 provides a visual summary of the nutritional considerations across the menstrual cycle.

Conclusion

It may be premature to prescribe female-specific nutrition recommendations for climbers, however, a personalised approach with an understanding of the influence of female hormones and an athlete's individual experience across the menstrual cycle is recommended. Based on the available evidence, specific areas for consideration for the female climber include energy intake and availability management, macronutrient intake and timing, micronutrients including iron, vitamin d, and calcium, and appropriate supplement use. Further research into nutrition for the female climber is warranted, as described in Table 3.

Table 2 Summary of nutrition guidelines for female rock climbers

Nutrition	Current Advice for Rock Climbers	Female Specific Considerations	Practical Recommendations
Energy Intake	5.8–8 MET value, 10–11 kcal/min of active climbing Minimum intake of 30–45 kcal/kgFFM	RMR may be slightly higher in the luteal phase Caution advised around ensuring adequate energy intake to avoid low energy availability	Monitor energy intake/expenditure to ensure adequate intake Education around the negative consequences of low energy availability and RED-S
Carbohydrate			Use of blood markers for LEA screening where appropriate
Daily Intake	3–7 g/kg BM/day	Overall CHO availability is particularly important in FP	Potentially increase CHO intake in FP to 6–8 g/kg/BM depending on individual training demands
During Exercise	Pre-exercise intake of 1 g/kg BM, 1–4 h before ~20–30 g/hour CHO during training/climbing	Exogenous CHO availability is important in LP	Special attention to intra-exercise CHO intake during LP
Recovery	Recovery intake of 1 g/kg/hour, 0–4 h post session	Increased markers of muscle damage and inflammation in the FP compared to LP	Recovery intake of 1 g/kg/hour, 0–4 h post session with rapid ingestion of 0.75 g/kg post-training
CHO Loading	8–12 g/kg BM/day	Possibly advantageous in FP to overcome lower resting muscle glycogen	8–12 g/kg/BM/day, possible benefit for performance days or heavy training sessions
Train Low Protocols	Potential benefit to endurance adaptation via enhanced gene signalling	More female specific research needed	Particular caution advised due to risk of RED-S
Protein			Avoid in LP
Daily Intake	1.4–2 g/kg BM boulderers 1.2–1.8 g/kg BM sport climbers	Protein catabolism is higher in the LP No sex difference in MPS rates	1.6–2 g/kg/day, with particular care in the LP
Intake Distribution	~0.3 g/kg BM, every 3–4 h	MPS is consistent across MC phases	Initial ~0.4 g/kg protein dose with ~0.3 g/kg every 3–4 h thereafter in the ~24-h recovery period
Fats	20%–35% of energy intake	Increased use of fat as fuel, especially in LP	No less than 20% of energy intake Include sources of omega-3 fats, e.g. oily fish or supplements
Vitamins & Minerals			
Iron	No existing guidelines specific to climbers	High risk for iron deficiency. Added risk for heavy menstruation and vegetarians Menstruation increases hepcidin release	Screen and monitor on an individual basis Include regular iron rich foods in diet. Intake away from training sessions, with a preference for the morning if possible Supplement if deficient

Table 2 (continued)

Nutrition	Current Advice for Rock Climbers	Female Specific Considerations	Practical Recommendations
Calcium + Vitamin D	<p>No existing guidelines specific to climbers</p> <p>General athlete recommendation of 1000–2000 IU, vitamin D, adjust for individual status if known</p> <p>1000 mg/day calcium for females, 1500 mg/day if postmenopausal or presenting with LEA</p>	Oestrogen affects calcium transport into bone	<p>Consider timing of iron intake</p> <p>Ensure adequate overall energy availability</p> <p>1000–2000 IU/day during winter months, adjust for individual status if known</p>
Hydration	<p>Monitor and advise on an individual basis</p> <p>Start climbing well hydrated and keep urine colour pale</p> <p>Drink 5–10 mL/kg body weight 2 h pre-exercise</p> <p>Drink 250 mL/hour adjusting for conditions</p>	<p>Core temperature changes due to female specific hormone fluctuations do not appear to affect fluid needs</p> <p>Women have lower sweat rates, lower plasma volumes and lower potassium and sodium sweat losses</p> <p>More research needed to understand fluid retention seen in LP</p>	<p>Include calcium rich foods in diet, e.g. dairy</p> <p>Monitor and advise on an individual basis</p> <p>Start climbing well hydrated and keep urine colour pale</p> <p>Drink 5–10 mL/kg body weight 2 h pre-exercise</p> <p>Drink 250 mL/hour adjusting for conditions</p>

CHO Carbohydrate, FP Follicular Phase, LP Luteal Phase




Fig. 1 Visual representation of Nutritional Considerations across the Menstrual Cycle

Table 3 Summary of recommendations for future research

Recommendations for Future Research

- The effect of ergogenic supplements such as creatine, beta alanine, caffeine and nitrate on female climbing performance
- Energy requirements of outdoor sport climbing in a range of conditions
- Performance variation in female climbers across the menstrual cycle
- Nutritional considerations for the post-menopausal female climber
- Nutritional considerations for pregnant and breastfeeding female climbers
- Research into the fluid retention reported in the luteal phase in female athletes
- Nutritional considerations for the female competition climber
- The interaction between female sex hormones, iron metabolism, and post exercise hepcidin response

Acknowledgements Mina would like to acknowledge her husband, David, for his support throughout this project.

Author Contributions The concept of the review was developed by ML-W; data interpretation, manuscript preparation, and editing were undertaken by ML-W and EG-S. All authors approved the final version of the paper.

Funding This review received no funding.

Data Availability No datasets were generated or analysed during the current study.

Declarations

Conflict of Interest The authors declare that the review was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated

otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bassett DR Jr, Tudor-Locke C, Greer JL, Vezina J, Whitt-Glover MC, Leon AS. 2011 compendium of physical activities: a second update of codes and MET values. *Med Sci Sports Exerc.* 2011;43(8):1575–81. <https://doi.org/10.1249/MSS.0b013e31821ece12>.
- Baranauskas M, Stukas R, Tubelis L, Žagminas K, Šurkienė G, Švedas E, Giedraitis VR, Dobrovolskij V, Abaravičius JA. Nutritional habits among high-performance endurance athletes. *Medicina (Kaunas).* 2015;51(6):351–62. <https://doi.org/10.1016/j.medic.2015.11.004>.
- Barr SI, Janelle KC, Prior JC. Energy intakes are higher during the luteal phase of ovulatory menstrual cycles. *Am J Clin Nutr.* 1995;61(1):39–43. <https://doi.org/10.1093/ajcn/61.1.39>.
- Beck KL, von Hurst PR, O'Brien WJ, Badenhorst CE. Micronutrients and athletic performance: a review. *Food Chem Toxicol: Int J.* 2021;158:112618. <https://doi.org/10.1016/j.fct.2021.112618>.
- Benton MJ, Hutchins AM, Dawes JJ. Effect of menstrual cycle on resting metabolism: a systematic review and meta-analysis. *PLoS ONE.* 2020;15(7): e0236025. <https://doi.org/10.1371/journal.pone.0236025>.
- Burke LM, Cox GR, Culmings NK, Desbrow B. Guidelines for daily carbohydrate intake: do athletes achieve them? *Sports Med.* 2001;31(4):267–99. <https://doi.org/10.2165/00007256-200131040-00003>.
- Burke LM, Hawley JA, Jeukendrup A, Morton JP, Stellingwerff T, Maughan RJ. Toward a common understanding of diet-exercise strategies to manipulate fuel availability for training and competition preparation in endurance sport. *Int J Sport Nutr Exerc Metab.* 2018;28(5):451–63. <https://doi.org/10.1123/ijsnem.2018-0289>.
- Burke LM, van Loon LJC, Hawley JA. Post exercise muscle glycogen resynthesis in humans. *J Appl Physiol.* 2017;122(5):1055–67. <https://doi.org/10.1152/jappphysiol.00860.2016>.
- Burke L, M. Ketogenic low-CHO, high-fat diet: the future of elite endurance sport? *J Physiol.* 2021;599:819–43. <https://doi.org/10.1113/JP278928>.
- Chappell S, Hackney AC. Associations between menstrual cycle phase, physical activity level and dietary macronutrient intake. *Biol Sport.* 1997;14(4):251–8.
- Chmielewska A, Kujawa K, Regulska-Ilow B. Accuracy of resting metabolic rate prediction equations in sport climbers. *Int J Environ Res Public Health.* 2023;20(5):4216. <https://doi.org/10.3390/ijerph20054216>.
- Chmielewska A, Regulska-Ilow B. The evaluation of energy availability and dietary nutrient intake of sport climbers at different climbing levels. *Int J Environ Res Public Health.* 2023;20(6):5176. <https://doi.org/10.3390/ijerph20065176>.
- Creer A, Gallagher P, Slivka D, Jemiolo B, Fink W, Trappe S. Influence of muscle glycogen availability on ERK1/2 and Akt signaling after resistance exercise in human skeletal muscle. *J Appl Physiol.* 2005;99(3):950–6.
- Davis HC, Hackney AC. The hypothalamic–pituitary–ovarian axis and oral contraceptives: regulation and function. In: Hackney AC, editor. *Sex hormones, exercise and women: scientific and clinical aspects.* London: Springer; 2017. pp. 1–17.
- De La Puente Yagüe M, Collado Yurrita L, Ciudad Cabañas MJ, Cuadrado Cenual MA. Role of Vitamin D in athletes and their performance: current concepts and new trends. *Nutrients.* 2020;12(2):579. <https://doi.org/10.3390/nu12020579>.
- Dellavalle DM, Haas JD. Iron status is associated with endurance performance and training in female rowers. *Med Sci Sports Exerc.* 2012;44(8):1552–9. <https://doi.org/10.1249/MSS.0b013e3182517ceb>.
- Farrokhyar F, Tabasinejad R, Dao D, Peterson D, Ayeni OR, Hadi-oonzadeh R, Bhandari M. Prevalence of vitamin D inadequacy in athletes: a systematic-review and meta-analysis. *Sports Med.* 2015;45(3):365–78. <https://doi.org/10.1007/s40279-014-0267-6>.
- Foo WL, Harrison JD, Mhizha FT, Langan-Evans C, Morton JP, Pugh JN, Areta JL. A short-term low-fiber diet reduces body mass in healthy young men: implications for weight-sensitive sports. *Int J Sport Nutr Exerc Metab.* 2022;32(4):256–64. <https://doi.org/10.1123/ijsnem.2021-0324>.
- Gibson-Smith E, Storey R, Ranchordas M. Dietary intake, body composition and iron status in experienced and elite climbers. *Front Nutr.* 2020;7:122. <https://doi.org/10.3389/fnut.2020.00122>.
- Giersch G, Charkoudian N, Stearns RL, Casa DJ. Fluid balance and hydration considerations for women: review and future directions. *Sports Med.* 2020;50(2):253–61. <https://doi.org/10.1007/s40279-019-01206-6>.
- Giles LV, Rhodes EC, Taunton JE. The physiology of rock climbing. *Sports Med.* 2006;36(6):529–45. <https://doi.org/10.2165/00007256-200636060-00006>.
- Giles D, Barnes K, Taylor N, Chidley C, Chidley J, Mitchell J, España-Romero V. Anthropometry and performance characteristics of recreational advanced to elite female rock climbers. *J Sports Sci.* 2021;39(1):48–56.
- Goolsby MA, Boniquit N. Bone health in athletes. *Sports Health.* 2017;9(2):108–17. <https://doi.org/10.1177/1941738116677732>.
- Guzman JF, Esteve H, Pablos C, Pablos A, Blasco C, Villegas JA. DHA-rich fish oil improves complex reaction time in female elite soccer players. *J Sports Sci Med.* 2011;10(2):301–5.
- Hackney AC, Kallman AL, Åggön E. Female sex hormones and the recovery from exercise: Menstrual cycle phase affects responses. *Biomed Hum Kinet.* 2019;11(1):87–9. <https://doi.org/10.2478/bhk-2019-0011>.
- Hackney AC. Menstrual cycle hormonal changes and energy substrate metabolism in exercising women: a perspective. *Int J Environ Res Public Health.* 2021;18(19):10024. <https://doi.org/10.3390/ijerph181910024>.
- Heidari H, Amani R, Feizi A, Askari G, Kohan S, Tavasoli P. Vitamin D supplementation for premenstrual syndrome-related inflammation and antioxidant markers in students with vitamin D deficient: a randomized clinical trial. *Sci Rep.* 2019;9(1):1–8.
- Heikura IA, Stellingwerff T, Areta JL. Low energy availability in female athletes: from the lab to the field. *Eu r J Sport Sci.* 2022;22(5):709–19. <https://doi.org/10.1080/17461391.2021.1915391>.
- Hinton P. Trace minerals of concern for female athletes: iron and zinc. In: Beals K, editor. *Nutrition for the female athlete.* London: CRC Press; 2013. pp. 75–92.
- Hokken R, Laugesen S, Aagaard P, Suetta C, Frandsen U, Ørtenblad N, Nielsen J. Subcellular localization- and fibre type-dependent utilization of muscle glycogen during heavy resistance exercise in elite power and olympic weightlifters. *Acta Physiol (Oxf).* 2021;231(2): e13561. <https://doi.org/10.1111/apha.13561>.
- Holtzman B, Ackerman KE. Recommendations and nutritional considerations for female athletes: health and performance. *Sports Med.* 2021;51(Suppl 1):43–57. <https://doi.org/10.1007/s40279-021-01508-8>.
- Howe SM, Hand TM, Larson-Meyer DE, Austin KJ, Alexander BM, Manore MM. No effect of exercise intensity on appetite

- in highly-trained endurance women. *Nutrients*. 2016;8(4):223. <https://doi.org/10.3390/nu8040223>.
33. Impey SG, Jevons EFP, Mees G, Cocks MS, Strauss JA, Chester N, Laurie I, Target D, Hodgson A, Shepherd SO, Morton JP. Glycogen utilization during running: intensity, sex, and muscle-specific responses. *Med Sci Sports Exerc*. 2020;52(9):1966–75. <https://doi.org/10.1249/MSS.0000000000002332>.
 34. Joubert LM, Gonzalez GB, Larson AJ. Prevalence of disordered eating among international sport lead rock climbers. *Front Sports Act Living*. 2020;2:86. <https://doi.org/10.3389/fspor.2020.00086>.
 35. Kemmler W, Roloff I, Baumann H, Schöfl V, Weineck J, Kalender W, Engelke K. Effect of exercise, body composition, and nutritional intake on bone parameters in male elite rock climbers. *Int J Sports Med*. 2006;27(8):653–9. <https://doi.org/10.1055/s-2005-872828>.
 36. Knechtel B, Jastrzębski Z, Hill L, Nikolaidis PT. Vitamin D and stress fractures in sport: preventive and therapeutic measures—a narrative review. *Medicina (Kaunas)*. 2021;57(3):223. <https://doi.org/10.3390/medicina57030223>.
 37. Kuikman MA, McKay AKA, Smith ES, Ackerman KE, Harris R, Elliott-Sale KJ, Stellingwerff T, Burke LM. Female athlete representation and dietary control methods among studies assessing chronic carbohydrate approaches to support training. *Int J Sport Nutr Exerc Metab*. 2023;33(4):198–208. <https://doi.org/10.1123/ijnsnem.2022-0214>.
 38. Maher AC, Tarnopolsky MA. Substrate utilization in female athletes: implications for fuel selection and macronutrient requirements. In: Beals K, editor. *Nutrition for the female athlete*. London: CRC Press; 2013. pp. 1–24.
 39. McClung JP, Gaffney-Stomberg E, Lee JJ. Female athletes: a population at risk of vitamin and mineral deficiencies affecting health and performance. *J Trace Elem Med Biol: Organ Soc Miner Trace Elem (GMS)*. 2014;28(4):388–92. <https://doi.org/10.1016/j.jtemb.2014.06.022>.
 40. McLay RT, Thomson CD, Williams SM, Rehrer NJ. Carbohydrate loading and female endurance athletes: effect of menstrual-cycle phase. *Int J Sport Nutr Exerc Metab*. 2007;17(2):189–205. <https://doi.org/10.1123/ijnsnem.17.2.189>.
 41. McNulty KL, Elliott-Sale KJ, Dolan E, Swinton PA, Ansdell P, Goodall S, Thomas K, Hicks KM. The effects of menstrual cycle phase on exercise performance in eumenorrheic women: a systematic review and meta-analysis. *Sports Med*. 2020;50(10):1813–27. <https://doi.org/10.1007/s40279-020-01319-3>.
 42. Mercer D, Convit L, Condo D, Carr AJ, Hamilton DL, Slater G, Snipe RMJ. Protein requirements of pre-menopausal female athletes: systematic literature review. *Nutrients*. 2020;12(11):3527. <https://doi.org/10.3390/nu12113527>.
 43. Mermier CM, Janot JM, Parker DL, Swan JG. Physiological and anthropometric determinants of sport climbing performance. *Br J Sports Med*. 2000;34(5):359–66. <https://doi.org/10.1136/bjism.34.5.359>.
 44. Michael MK, Witard OC, Joubert L. Physiological demands and nutritional considerations for Olympic-style competitive rock climbing. *Cogent Med*. 2019;6:1.
 45. Michailov ML, Morrison A, Ketenliev MM, Pentcheva BP. A sport-specific upper-body ergometer test for evaluating submaximal and maximal parameters in elite rock climbers. *Int J Sports Physiol Perform*. 2015;10(3):374–80. <https://doi.org/10.1123/ijsp.2014-0160>.
 46. Monedero J, Duff C, Egan B. Dietary intakes and the risk of low energy availability in male and female advanced and elite rock climbers. *J Strength Cond Res*. 2023;37(3):e8–15. <https://doi.org/10.1519/JSC.0000000000004317>.
 47. Moore DR, Sygo J, Morton JP. Fuelling the female athlete: carbohydrate and protein recommendations. *Eur J Sport Sci*. 2022;22(5):684–96. <https://doi.org/10.1080/17461391.2021.1922508>.
 48. Mountjoy M, Sundgot-Borgen JK, Burke LM, Ackerman KE, Blauwet C, Constantini N, Lebrun C, Lundy B, Melin AK, Meyer NL, Sherman RT, Tenforde AS, Klungland Torstveit M, Budgett R. IOC consensus statement on relative energy deficiency in sport (RED-S): 2018 update. *Br J Sports Med*. 2018;52(11):687–97. <https://doi.org/10.1136/bjsports-2018-099193>.
 49. Ogan D, Pritchett K. Vitamin D and the athlete: risks, recommendations, and benefits. *Nutrients*. 2013;5(6):1856–68. <https://doi.org/10.3390/nu5061856>.
 50. Oosthuysen T, Bosch AN. The effect of the menstrual cycle on exercise metabolism: implications for exercise performance in eumenorrheic women. *Sports Med*. 2010;40(3):207–27. <https://doi.org/10.2165/11317090-000000000-00000>.
 51. Oosthuysen T, Bosch A. Oestrogen's regulation of fat metabolism during exercise and gender specific effects. *Curr Opin Pharmacol*. 2012;12(3):363–71. <https://doi.org/10.1016/j.coph.2012.02.008>.
 52. Palmery M, Saraceno A, Vaiarelli A, Carlomagno G. Oral contraceptives and changes in nutritional requirements. *Eur Rev Med Pharmacol Sci*. 2013;17(13):1804–13.
 53. Pihoker AA, Peterjohn AM, Trexler ET, Hirsch KR, Blue M, Anderson KC, Ryan ED, Smith-Ryan AE. The effects of nutrient timing on training adaptations in resistance-trained females. *J Sci Med Sport*. 2019;22(4):472–7. <https://doi.org/10.1016/j.jsams.2018.09.236>.
 54. Pilis K, Stec K, Pilis A, Mroczek A, Michalski C, Pilis W. Body composition and nutrition of female athletes. *Rocz Panstw Zakl Hig*. 2019;70(3):243–51. <https://doi.org/10.32394/rpzh.2019.0074>.
 55. Rawson ES, Miles MP, Larson-Meyer DE. Dietary supplements for health, adaptation, and recovery in athletes. *Int J Sport Nutr Exerc Metab*. 2018;28(2):188–99. <https://doi.org/10.1123/ijnsnem.2017-0340>.
 56. Rehrer NJ, McLay-Cooke RT, Sims ST. Nutritional strategies and sex hormone interactions in women. In: Hackney A, editor. *Sex hormones, exercise and women*. Cham: Springer; 2017.
 57. Rodriguez-Giustiniani P, Rodriguez-Sanchez N, Galloway SDR. Fluid and electrolyte balance considerations for female athletes. *Eur J Sport Sci*. 2022;22(5):697–708.
 58. Rossi KA. Nutritional aspects of the female athlete. *Clin Sports Med*. 2017;36(4):627–53.
 59. Sheel AW, Seddon N, Knight A, McKenzie DC, Warburton RDE. Physiological responses to indoor rock-climbing and their relationship to maximal cycle ergometry. *Med Sci Sports Exerc*. 2003;35(7):1225–31. <https://doi.org/10.1249/01.MSS.0000074443.17247.05>.
 60. Sim M, Dawson B, Landers G, Trinder D, Peeling P. Iron regulation in athletes: exploring the menstrual cycle and effects of different exercise modalities on hepcidin production. *Int J Sport Nutr Exerc Metab*. 2014;24(2):177–87. <https://doi.org/10.1123/ijnsnem.2013-0067>.
 61. Sim M, Garvican-Lewis LA, Cox GR, Govus A, McKay AKA, Stellingwerff T, Peeling P. Iron considerations for the athlete: a narrative review. *Eur J Appl Physiol*. 2019;119(7):1463–78.
 62. Smith E, Storey R, Ranchordas M. Nutritional considerations for bouldering. *Int J Sport Nutr Exerc Metab*. 2017;27:1–28. <https://doi.org/10.1123/ijnsnem.2017-0043>.
 63. Solli GS, Sandbakk SB, Noordhof DA, Ihalainen JK, Sandbakk Ø. Changes in self-reported physical fitness, performance, and side effects across the phases of the menstrual cycle among competitive endurance athletes. *Int J Sports Physiol Perform*. 2020;15:1324–33. <https://doi.org/10.1123/ijsp.2019-0616>.
 64. Spence K. Nutrients needed for optimal bone health in the female athlete. In: Beals K, editor. *Nutrition for the female athlete*. London: CRC Press; 2013. pp. 117–38.

65. Stachenfeld NS. Including women in research. It's necessary, and really not so hard to do. *Exp Physiol*. 2018;103(10):1296–7. <https://doi.org/10.1113/EP087261>.
66. Stellingwerff T, Spriet LL, Watt MJ, Kimber NE, Hargreaves M, Hawley JA, Burke LM. Decreased PDH activation and glycogenolysis during exercise following fat adaptation with carbohydrate restoration. *Am J Physiol Endocrinol Metab*. 2006;290(2):E380–8. <https://doi.org/10.1152/ajpendo.00268.2005>.
67. Thomas DT, Erdman KA, Burke LM. American college of sports medicine joint position statement. Nutrition and athletic performance. *Med Sci Sports Exerc*. 2016;48(3):543–68. <https://doi.org/10.1249/mss.0000000000000852>.
68. Troutt JS, Rudling M, Persson L, Ståhle L, Angelin B, Butterfield AM, Schade AE, Cao G, Konrad RJ. Circulating human hepcidin-25 concentrations display a diurnal rhythm, increase with prolonged fasting, and are reduced by growth hormone administration. *Clin Chem*. 2012;58(8):1225–32. <https://doi.org/10.1373/clinchem.2012.186866>.
69. Volek JS, Forsythe CE, Kraemer WJ. Nutritional aspects of women strength athletes. *Br J Sports Med*. 2006;40(9):742–8. <https://doi.org/10.1136/bjism.2004.016709>.
70. Wallis GA, Dawson R, Achten J, Webber J, Jeukendrup AE. Metabolic response to carbohydrate ingestion during exercise in males and females. *Am J Physiol-Endocrinol Metab*. 2006;290(4):E708–15. <https://doi.org/10.1152/ajpendo.00357.2005>.
71. Watts PB. Physiology of difficult rock climbing. *Eur J Appl Physiol*. 2004;91(4):361–72. <https://doi.org/10.1007/s00421-003-1036-7>.
72. Wenner M, Stachenfeld N. Fluid and electrolyte requirements for female athletes. In: Beals K, editor. *Nutrition for the female athlete*. London: CRC Press; 2013. pp. 75–92.
73. West DWD, Burd NA, Churchward-Venne TA, Camera DM, Mitchell CJ, Baker SK, Phillips SM. Sex-based comparisons of myofibrillar protein synthesis after resistance exercise in the fed state. *J Appl Physiol* (1985). 2012;112(11):1805–13. <https://doi.org/10.1152/jappphysiol.00170.2012>.
74. Wohlgemuth KJ, Arieta LR, Brewer GJ, Hoselton AL, Gould LM, Smith-Ryan AE. Sex differences and considerations for female specific nutritional strategies: a narrative review. *J Int Soc Sports Nutr*. 2021;18(1):27. <https://doi.org/10.1186/s12970-021-00422-8>.
75. Zapf J, Fichtl B, Wielgoss S, Schmidt W. Macronutrient intake and eating habits in elite rock climbers. *Med Sci Sports Exerc*. 2001;33(5):S72.