

**Positively charged amino acids are essential for electron transfer and protein-protein interactions in the soluble methane monooxygenase complex from methylococcus capsulatus (Bath)**

BALENDRA, S., LESIEUR, C., SMITH, T. J. <<http://orcid.org/0000-0002-4246-5020>> and DALTON, H.

Available from Sheffield Hallam University Research Archive (SHURA) at:  
<http://shura.shu.ac.uk/368/>

---

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

**Published version**

BALENDRA, S., LESIEUR, C., SMITH, T. J. and DALTON, H. (2001). Positively charged amino acids are essential for electron transfer and protein-protein interactions in the soluble methane monooxygenase complex from methylococcus capsulatus (Bath). *Biochemistry*, 41 (8), 2571-2579.

---

**Copyright and re-use policy**

See <http://shura.shu.ac.uk/information.html>

## FIGURE LEGENDS

**Fig. 1. Cross-linking and neutralisation of primary amino groups associated with the hydroxylase.** The hydroxylase was reacted with the cross-linker (BS<sup>3</sup>) or the lysine-neutralising reagent (sulfo-NHS-acetate) and excess reagent was removed by ultrafiltration. Samples from the cross-linking experiment were analysed by means of (A) SDS-7.5% PAGE and (B) nondenaturing, native PAGE. Samples from the lysine neutralisation experiment were analysed by using nondenaturing PAGE (C). Samples analysed before measurement of propene oxygenation activity: lanes 1 and 2, unreacted and reacted hydroxylase, respectively. Samples analysed after measurement of propene oxygenation activity in the sMMO complex: lane 3, unreacted hydroxylase; lane 4, control in which the probe was removed by ultrafiltration before addition of the hydroxylase; lane 5, reacted hydroxylase. Samples analysed after measurement of propene oxygenation activity via the peroxide shunt: lanes 6 and 7, unreacted and reacted hydroxylase, respectively. The proteins were visualised with Coomassie blue; molecular masses of standards in lane M of part A are indicated in kDa. The cross-linked species observed by means of SDS-PAGE (A) have been tentatively assigned based on the known molecular masses of the  $\alpha$ ,  $\beta$  and  $\gamma$  subunits of the hydroxylase.

**Fig. 2. Surface plasmon resonance analysis** of the binding between the GST-protein B fusion protein and (A) native, (B) primary amine-modified and (C) arginine-modified hydroxylase. The points at which the hydroxylase was added to and removed from the buffer flowing across the sensor chip are indicated.

**Fig. 3. Distribution of charged groups on the surface of the hydroxylase**, based on the X-ray crystal structure (9). Nitrogen atoms of primary amino and arginyl side-chain guanidinium groups are shown in dark and light blue, respectively. Side-chain carboxylate oxygen atoms are shown in red. The uncharged parts of helices  $\alpha$ E and  $\alpha$ F of the  $\alpha$  subunit, proposed to form the hydrophobic binding site for protein B, are coloured light green.

Fig. 1.






Fig. 2.

A



B



C




Fig. 3.

