

## **An ionic-gelling alginate drink attenuates postprandial glycaemia in males**

HARDEN, Charlotte, RICHARDSON, J Craig, DETTMAR, Peter W, CORFE, Bernard M and PAXMAN, Jenny <<http://orcid.org/0000-0003-3596-489X>>

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/5722/>

---

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

### **Published version**

HARDEN, Charlotte, RICHARDSON, J Craig, DETTMAR, Peter W, CORFE, Bernard M and PAXMAN, Jenny (2012). An ionic-gelling alginate drink attenuates postprandial glycaemia in males. *Journal of functional foods*, 4 (1), 122-128.

---

### **Copyright and re-use policy**

See <http://shura.shu.ac.uk/information.html>

1 **AN IONIC-GELLING ALGINATE DRINK ATTENUATES**  
2 **POSTPRANDIAL GLYCAEMIA IN MALES**

3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

4 **Charlotte J. Harden<sup>a</sup>, J. Craig Richardson<sup>b</sup>, Peter W.**  
5 **Dettmar<sup>b</sup>, Bernard M. Corfe<sup>c\*</sup>, Jenny R Paxman<sup>a</sup>**

6

7 <sup>a</sup> Centre for Food Innovation, Sheffield Hallam University,  
8 Howard St, Sheffield, S1 1WB, UK.

9 <sup>b</sup> Technostics Limited, The Deep Business Centre, Tower  
10 Street, Kingston Upon Hull, HU1 4BG, UK.

11 <sup>c</sup> Department of Oncology, The University of Sheffield,  
12 The Medical School, Beech Hill Road, Sheffield, S10 2RX,  
13 UK.

14

---

\* Corresponding author: Dr Bernard Corfe, Department of  
Oncology, The University of Sheffield, The Medical  
School, Beech Hill Road, Sheffield, S10 2RX, UK,

*E-mail address:* b.m.corfe@sheffield.ac.uk

*Telephone:* +44 (0)114 271 3004

*Fax:* +44 (0)114 271 3314

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

15    **ABSTRACT**

16    Obese individuals are at increased risk of type 2 diabetes  
17    compared to their healthy weight counterparts. Dietary  
18    fibre, such as alginate, could attenuate glycaemic  
19    disturbances associated with obesity when included in  
20    the diet.

21    Forty self-reported, healthy males completed this  
22    randomised, single-blinded, controlled, parallel trial to  
23    determine the glycaemic response to a controlled test-  
24    lunch of mixed composition following an ionic-gelling  
25    alginate preload drink compared to an acidic-gelling  
26    control.

27    Individual baseline area under the curve was 52% lower  
28    ( $P=0.010$ ) and peak glycaemia was 14% lower ( $P<$   
29     $0.0005$ ) after the ionic-gelling alginate drink compared  
30    with the control. Body fatness was a predictor of  
31    postprandial glycaemia however there was no interaction  
32    effect between body fat % and treatment type.

33    We have shown ionic-gelling alginate can attenuate  
34    glycaemic response to set lunch of mixed composition.  
35    Functional foods that include ionic-gelling alginates may  
36    benefit those with elevated postprandial blood glucose.

37    **KEY WORDS**

38 Alginate; glucose; glycemia; gel; body fat

39

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 40 1.0 INTRODUCTION

41 As obesity increases, the incidence of associated co-  
42 morbidities rises concomitantly, most dramatically in  
43 relation to body mass index-related diabetes (McPherson  
44 et al., 2007). Abdominal fatness has been linked with  
45 elevated fasting blood glucose (Rezende et al., 2006).  
46 Pascot et al. (1999) showed visceral adipose tissue  
47 accumulation was accompanied by increased plasma  
48 glucose in the fasted state and after a 75g oral glucose  
49 load in young and middle aged women. In a six year  
50 prospective study Kriketos et al. (2003) showed baseline  
51 body fatness and increasing fatness over time to be  
52 strong predictors of elevated fasting plasma glucose in  
53 individuals 'at-risk' of type 2 diabetes

54 Epidemiological evidence suggests dietary fibres may  
55 have a preventive role in the development of type 2  
56 diabetes (Meyer et al., 2000). Several mechanisms by  
57 which soluble fibres may modulate glycaemic response  
58 have been proposed (Augustin et al., 2000). Soluble fibre  
59 ingestion reduces carbohydrate digestion rates, therefore  
60 aiding regulation of postprandial glycaemia (Augustin et  
61 al., 2000; Kimura et al., 1996; Welch, 1994).

62

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63 Soluble fibres have been shown to have beneficial effects  
64 in controlling glycaemia following carbohydrate ingestion  
65 in healthy volunteers (Goñi et al., 2000; Rigaud et al.,  
66 1998; Lavin and Read, 1995). Similarly, fibre-rich foods  
67 (Flammang et al., 2006) and fibre supplementation  
68 (Sierra et al., 2002) have been shown to help attenuate  
69 postprandial glycaemic responses in type 2 diabetic  
70 adults. Kaline et al. (2007) reviewed the potential  
71 mechanisms by which diets rich in dietary fibre can be  
72 useful in diabetes prevention.

73 Alginate is an algal polysaccharide found in the cell walls  
74 of certain brown seaweed species. This fibre has been  
75 used in several relevant human intervention studies. 5.0g  
76 of sodium alginate added to a meal significantly  
77 attenuated postprandial glycaemic response in type 2  
78 diabetics by 31% compared to the control meal  
79 (Torsdottir et al., 1991). Wolf and colleagues (2002)  
80 demonstrated that 1.5g of sodium alginate, incorporated  
81 into a 100g glucose-based preload drink with an acid-  
82 soluble calcium source (to produce an acid-induced  
83 viscosity complex), elicited a non-significant drop in peak  
84 glycaemia and a significant attenuation of incremental  
85 change from baseline area under the curve (AUC) in  
86 healthy, non-diabetic adults compared to a soluble fibre-  
87 based control. Williams et al. (2004) fed a "crispy bar"

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

88 containing 5.5g guar gum and 1.6g sodium alginate to  
89 healthy adults and measured the resultant glycaemic  
90 response compared to an alginate-free bar. Postprandial  
91 blood glucose excursions were significantly lower at 15,  
92 30, 45, and 120 minutes and the positive incremental  
93 AUC was significantly reduced (by 33%) after  
94 consumption of the enriched “crispy bar” compared to the  
95 alginate-free bar. Paxman et al. (2008a) reported a  
96 strong positive correlation between change from  
97 individual baseline AUC glycaemia and body fat % when  
98 a hypromellose control preload was ingested prior to a  
99 test lunch. This positive correlation was not apparent  
100 following an ionic gelling sodium alginate preload,  
101 providing preliminary evidence to suggest that the  
102 enhanced glycaemic response to a meal at higher body  
103 fat could be normalised following ingestion of an alginate  
104 preload identical to the one used in the present study.

105 Hoad et al. (2004) fed volunteers a strong gelling (high-G)  
106 and a weaker gelling (low-G) alginate meal, a guar-based  
107 meal or a control (without added fibre) and examined the  
108 resultant gastric emptying rates. *In vitro*, both alginate  
109 meals formed intragastric gel 'lumps', and in the case of  
110 the strong-gelling alginate, this was reportedly associated  
111 with a feeling of fullness and a reduction in hunger. Hoad  
112 and colleagues (2004) purport that acid-gelling agents

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

113 such as alginate may be usefully incorporated into  
114 weight-reducing diets/ foods in order to enhance antrum  
115 distension and/ or manipulate nutrient uptake from the  
116 ileum.

117 Alginate is widely used in the food industry as a thickener,  
118 stabiliser and gelling agent (Brownlee et al., 2005). Its  
119 constituent sugar residues are D-mannuronic (M) and L-  
120 guluronic acid (G). Homopolymeric G blocks (comprising  
121 diaxial linkages in the  ${}^1C_4$  conformation) can react with  
122  $Ca^{2+}$  and  $H^+$  ions to yield a strong, cross-linked gel  
123 (Brownlee et al., 2005; Seal and Mathers, 2001; Kimura  
124 et al, 1996). Consequently, the gel strength of alginate  
125 and its consequent biochemical and biophysical  
126 properties are determined by its chemical structure.

127 Specific alginates and specific alginate formulations are  
128 therefore likely to react differently within the  
129 gastrointestinal milieu.

130 The primary objective of the present study was to  
131 examine the effect of alginate gelled ionically compared  
132 to acidically (control) on glycaemic response to a  
133 standard meal of mixed composition. Secondary to this,  
134 we investigated how body fatness affects the postprandial  
135 glycaemic response when subjects ingest the ionic-  
136 gelling formulation compared to the acid-gelling control.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

137

138

## 139 **2.0 MATERIALS AND METHODS**

### 140 **2.1 Subjects**

141 41 male subjects participated in the study. Only one  
142 subject was excluded, due to unusually low fasting  
143 glucose levels, leaving complete datasets for 40  
144 participants. Subjects aged 18 to 65 years were eligible  
145 to take part providing they did not meet any of the criteria  
146 for exclusion which were; type 1 or 2 diabetes, history of,  
147 or current cardiovascular complaints ( or if they had been  
148 fitted with a pacemaker or other implantable electronic  
149 device) or gastrointestinal complaints (such as irritable  
150 bowel syndrome or inflammatory bowel disorder,  
151 dumping syndrome or Cushing's syndrome), current fibre  
152 supplement use, use of constipation-causing drugs such  
153 as codeine or morphine, bowel blockage, bowel muscle  
154 weakness or recent food poisoning. In addition, anyone  
155 with a known allergy to, or intolerance of, the foods or  
156 ingredients used in the experiment was excluded from  
157 taking part, as were vegans (due to the nature of the  
158 foods used).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

159 Baseline pre-screening took place less than one week  
160 prior to the experimental phase, in which subjects  
161 completed a general health questionnaire and various  
162 anthropometric measures were made. Height and weight  
163 were recorded (SECA 709 mechanical column scales  
164 with SECA 220 telescopic measuring rod; SECA United  
165 Kingdom, Birmingham) and body mass index (BMI) was  
166 calculated. Bioelectrical impedance analysis was  
167 undertaken following 5 minutes of supine rest on non-  
168 conducting foam matting using a BodyStat 1500  
169 (BodyStat Ltd., Isle of Man, British Isles). Body fat % was  
170 recorded. Subjects completed a 51-item Three Factor  
171 Eating Questionnaire (TFEQ; Stunkard and Messick,  
172 1985) to determine eating behaviour across three pre-  
173 defined factors. Mean values for all three factors;  
174 restraint, disinhibition and hunger, were low for the group  
175 as a whole (Stunkard and Messick, 1985). Subject  
176 characteristics are reported in Table 1. This study was  
177 approved by the relevant University Ethics Committee  
178 (Ref: FIRC/2006/RE21). All subjects gave informed  
179 consent to participate.

## 51 52 **2.2 Study Design**

53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

181 In this randomised, single-blinded, controlled parallel trial  
182 subjects ( $n = 40$ ) were split equally either side of the

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

183 median into haptiles by body fatness (lower body fat  
184 group: <16.10%, upper body fat group: ≥16.10%).

185 Following a 12 hour overnight fast, all subjects consumed  
186 a controlled breakfast at 9am (60g Kellogg's® Hint of  
187 Honey Corn Flakes; Kellogg's Company GB Limited,  
188 Manchester, 125ml semi-skimmed milk and 200ml 'Drink  
189 Fresh' orange juice; DCB Foodservice, Herts). After  
190 breakfast subjects were asked to travel to the laboratory  
191 using motorised transport to minimise energy expenditure.  
192 From breakfast until 11am subjects consumed only  
193 bottled spring water (Highland Spring still natural mineral  
194 water with a sports cap, 2 x 500ml; Highland Spring Ltd,  
195 Perthshire, Scotland) to a maximum volume of 1 litre.  
196 Water consumption was *ad libitum* but the bottles were  
197 weighed prior to the experiment and at 11am in order to  
198 determine the exact amount consumed before the test-  
199 lunch. Upon arrival at the facility for the experimental day,  
200 subjects were randomly allocated to one of two preload  
201 treatments; an ionic-gelling sodium alginate formulation  
202 (SA) or an acid-gelling excipient free control (EF).

### 203 **2.3 Preload Formulations and Glycaemia**

204 The SA formulation contained sodium alginate, calcium  
205 carbonate (CaCO<sub>3</sub>) and buffering agents. It was  
206 specifically formulated to undergo enhanced ionic

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

207 intragastric gelation upon ingestion. This is achieved by  
208 mixing sodium alginate with an acid soluble calcium salt.  
209 Post-ingestion solubilisation of calcium salt in acidic  
210 gastric fluid liberates free calcium ions which are then  
211 available to cross-link with the sodium alginate. The SA  
212 formulation has been described in detail by Paxman et al.  
213 (2008b). The EF control is identical in composition to SA  
214 with the omission of the CaCO<sub>3</sub>. This formulation yields a  
215 gel via acid gelation (in the absence of calcium), resulting  
216 in weaker intra-molecular hydrogen bonded mass.

217 Prior to preload ingestion, baseline glycaemia (11:45am,  
218 0 minutes) was determined using capillary blood taken  
219 from the finger. A single use Accu-check® Softclix® Pro  
220 lancing device was used to obtain a single droplet sample  
221 via OneTouch® Ultra® Test Strips with FastDraw™  
222 design. The OneTouch® Ultra® Blood Glucose  
223 Monitoring System was used to determine glycaemia  
224 (reference range 1.1 to 33.3mmol/l; Lifescan Inc., Bucks).

225 Each preload was served at 12:00pm (15 minutes after  
226 baseline glycaemia measurements) in an opaque non-  
227 descript plastic cup in standard feeding booths in green  
228 light. The coloured light masked a very slight colour  
229 difference between preload drinks. The drinks were  
230 flavoured with vanilla to yield an orosensory match.  
231 Subjects were instructed to drink the entire product. All

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

232 preloads were consumed within 5 minutes of their initial  
233 hydration with 100ml bottled water.  
234 Following ingestion of the product (12:15pm, 30 minutes  
235 from baseline), glycaemia was again determined  
236 following identical protocol.

#### 237 **2.4 Test-lunch**

238 Volunteers ingested a controlled test-lunch of mixed  
239 composition thirty minutes after consuming the preload  
240 drink (12:30pm, 45 minutes from baseline) in standard  
241 feeding booths in natural light. The test-lunch consisted  
242 of 300g pre-cooked then chilled penne pasta (Don Mario  
243 100% durum wheat semolina pasta quills', manufactured  
244 by Abbey Foods Ltd, PO BOX 178, Liverpool) and 100g  
245 Sacla Italia <sup>TM</sup> vine-ripened tomato and mascarpone stir  
246 through sauce (F.lli Sacla S.p.A. Asti Italy; Sacla UK LTD,  
247 Basil House, 21 London End, Bucks). This test-lunch  
248 was heated to a temperature of at least 72°C in a  
249 microwave and was served at a temperature of between  
250 60-65°C. Subjects were instructed to consume the entire  
251 test-lunch and all subjects adhered to protocol.

252

253 The meal provided 57%, 13% and 30% of total energy  
254 from carbohydrate, protein and fat respectively, as

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

255 analysed by NetWISP (version 3.0 for Windows, Tinuviel  
256 Software, Anglesey, UK). The test-lunch protocol used  
257 here has been described previously (Paxman et al.,  
258 2008a).

## 259 **2.5 Protocol Postprandially**

260 Further measures of capillary glucose were obtained at  
261 90, 120, 150, 180, 210, 240, 270 and 330 minutes from  
262 baseline. In total, ten capillary blood samples were taken  
263 to determine glycaemia up to 330 minutes from baseline  
264 (270 minutes postprandially).

## 265 **2.6 Statistical analysis**

266 Blood glucose measures were converted to delta area  
267 under the curve (AUC) using the trapezoid rule with  
268 subtraction of basal values (NCSS; Hintze, 2004, NCSS  
269 and PASS Number Cruncher Statistical Systems,  
270 Kaysville, Utah). Two-way between groups ANOVAs  
271 were performed in order to identify the main effects of  
272 treatment and body fat haptile and any interaction effects  
273 on glycaemia at each time point, change from individual  
274 baseline AUC glycaemia and peak postprandial  
275 glycaemia (SPSS; version 15.0 for Windows, SPSS Inc.,  
276 Chicago, IL, USA). Graphical presentations were  
277 produced using SPSS (version 15.0 for Windows, SPSS  
278 Inc., Chicago, IL, USA) and Microsoft Excel 2003

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

279 (Microsoft Office, Microsoft Corporation). Significance  
280 was set at  $p < 0.05$ . Data are presented as mean  $\pm$  1 SD.

281

## 282 **3.0 RESULTS**

283 Forty self-reported healthy male subjects (equal numbers  
284 in each treatment arm) successfully completed the  
285 experiment with no deviation from protocol.

### 286 **3.1 Ionic gelling sodium alginate attenuates the** 287 **glycaemic response to a meal**

288 Two-way between groups ANOVAs showed a significant  
289 effect of treatment type on glycaemia at 90 ( $p < .0005$ ),  
290 150 ( $p = .003$ ), 180 ( $p = .021$ ) and 210 ( $p = .013$ ) minutes  
291 (see Figure 1). Overall, ingestion of SA compared to EF  
292 resulted in a significant reduction in a mean change from  
293 individual baseline AUC glycaemia ( $\underline{M} = 148.43 \pm 148.65$   
294 vs.  $\underline{M} = 312.53 \pm 253.60$ ;  $p = .010$ ) of 52.5% (see Figure 1).  
295 Irrespective of treatment type, subjects in the lower  
296 haptile for body fat % had a reduced mean change from  
297 individual baseline AUC glycaemia ( $177.68 \pm 255.44$ )  
298 compared to those in the upper haptile for body fat %  
299 ( $283.28 \pm 171.66$ ;  $p = .065$ ; data not shown) however, this  
300 was not significant and there was no interaction effect  
301 between treatment type and body fat % grouping.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

302 **3.2 Ionic gelling sodium alginate reduces peak**

303 **postprandial glycaemia**

304 Preload type failed to affect the timing of peak glycaemia  
305 as shown in Figure 1. However, Figure 2 shows the  
306 significant 14% lower mean peak postprandial glycaemia  
307 at 90 minutes following SA versus EF ( $\underline{M}$ = 6.06 ± .59  
308 mmol/L vs.  $\underline{M}$ = 6.92 ± .70mmol/L;  $p$ < .0005) for the study  
309 group as a whole. Subjects in the lower body fat haptile  
310 had a lower peak postprandial glycaemia (6.39  
311 ± .85mmol/L) than those in the upper body fat haptile,  
312 irrespective of treatment type (6.59 ± .70mmol/L;  $p$ = .170;  
313 data not shown) however this was not significant and  
314 there was no interaction effect between treatment type  
315 and body fat % grouping.

316 **3.3 Body fat classification determines the**

317 **postprandial glycaemic response to a meal but the**

318 **beneficial effects of alginate remain**

319 Irrespective of treatment type, subjects in the upper body  
320 fat haptile had non-significantly elevated peak  
321 postprandial glycaemia and non-significantly greater  
322 mean change from individual baseline AUC glycaemia  
323 compared to those in the lower body fat haptile. In  
324 addition, body fat % grouping had a significant effect on  
325 delta glycaemia at 120 ( $p$ = .005), 150 ( $p$ = .012), 180



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

326 ( $p= .049$ ) and 210 minutes ( $p= .046$ ) from baseline, with  
327 subjects in the upper body fat haptile having higher mean  
328 glycaemia than those in the lower body fat haptile at  
329 these time points, irrespective of preload treatment  
330 (Figure 3).

331 For glycaemia at each time point, change from individual  
332 baseline AUC glycaemia and peak postprandial  
333 glycaemia however, the two-way between-groups  
334 ANOVA showed no interaction effect between treatment  
335 type and body fat % grouping in each case. Subjects  
336 appeared to respond to the ionic-gelling sodium alginate  
337 (SA) treatment in a similar fashion irrespective of body  
338 fat %.

339 Examination of the response to treatment type by body  
340 fat % grouping showed the lower body fat haptile on SA  
341 reduced their change from individual baseline AUC  
342 glycaemia by 68.3%, and their peak postprandial  
343 glycaemia by 16.2% compared to the lower body fat  
344 haptile on EF. A slightly weaker effect was apparent in  
345 the upper body fat haptile on SA who reduced their  
346 change from individual baseline AUC glycaemia by  
347 46.6%, and their peak postprandial glycaemia by 9.7%  
348 compared to those in the upper body fat haptile on the EF  
349 treatment type (Figure 3). This finding supports our

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

350 previous suggestions relating to altered glycaemic  
351 response and body fatness (Paxman et al., 2008a).  
352  
353 In summary, glycaemic response to the test-meal was  
354 reduced following ingestion of the ionic-gelling sodium  
355 alginate drink (SA) compared to the acid-gelling  
356 excipient-free formulation (EF) throughout the 330 minute  
357 measurement period. Body fatness influenced  
358 postprandial glycaemic response but the effect of the  
359 ionic-gelling alginate drink was maintained.

360

#### 361 **4.0 DISCUSSION**

362 The literature suggests soluble fibre can alter subjective  
363 hunger and fullness ratings (Peters et al., 2011), gastric  
364 emptying rate and intestinal nutrient absorption, though  
365 the extent and subsequent effect on glycaemia is poorly  
366 established (Wolf et al., 2002; Delargy et al., 1997;  
367 Fairchild et al., 1996). Contradictory reports are most  
368 likely explained by the type, dose, homogeneity and  
369 physicochemical properties of fibres used, and differing  
370 participant characteristics between studies.

371 The physicochemical properties of alginate have particular  
372 potential in terms of attenuating postprandial glycaemic

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

373 response and improving diabetic control (Williams et al.,  
374 2004; Wolf et al., 2002; Torsdottir et al., 1991). Highly  
375 viscous solutions are unpalatable; solutions which form  
376 solid gel particles in the gastric lumen may provide a  
377 more feasible alternative for controlling gastric emptying  
378 and nutrient uptake. In order to establish an optimum  
379 formulation for delivery of a glycaemia-modulating  
380 alginate, the physiological response to ionic- and acid-  
381 gelling alginates were compared in males of differing  
382 body fatness. Physiologic data show greater glucose  
383 intolerance among the obese and numerous prospective  
384 studies support such associations between measures of  
385 obesity and type 2 diabetes risk (Carey et al., 1997).  
386 Such differences are postulated to be connected with  
387 body fatness.

388 Our data show that the ionic-gelling sodium alginate drink  
389 (SA) reduced early-phase and peak postprandial  
390 glycaemia and flattened the postprandial glycaemic curve  
391 in comparison to the acid-gelling control (EF). From  
392 baseline to 30 minutes, the EF preload drink resulted in a  
393 slight elevation of blood glucose, most likely due to the 7g  
394 fructose contained within the formulation. In comparison,  
395 the SA preload treatment elicited no change in glycaemia  
396 during this period despite containing the same amount of  
397 fructose. The difference between these responses can

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

398 most probably be attributed to the addition of calcium  
399 carbonate in the SA formulation. The acid-soluble  
400 calcium salt was expected to facilitate intra-gastric ionic  
401 gelation of the drink (Kimura et al., 1996). When alginate  
402 formulations are pH dependent there is a known time lag  
403 of 25-40 minutes before gelation occurs (Mattes, 2007).  
404 The inhibition of a glycaemic response to the SA  
405 formulation could have resulted from immediate fructose  
406 entrapment, delayed gastric emptying or both. Torsdottir  
407 et al. (1991) reported delayed glucose delivery and  
408 reduced glycaemic peak in type 2 diabetics by the  
409 addition of alginate to meals. They attributed this  
410 response solely to delayed gastric emptying, measured  
411 by aspirated radioactive stomach contents. There is  
412 evidence to suggest alginate ingestion results in 'gel  
413 lump' formation, which alters nutrient transport to the  
414 small intestine (Hoad et al., 2004). In this study the  
415 glycaemic response to a test-lunch of mixed composition  
416 following the SA drink was consistently lower throughout  
417 the investigation, thus it seems likely that nutrients were  
418 captured within the gel matrix to some degree.

419 Hoad et al. (2004) used serial magnetic resonance  
420 imaging (MRI) to gather *in vivo* measurements of guar  
421 gum and weak and strong gelling alginates that had been  
422 incorporated into milk-based drinks. MRI images showed

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

423 heterogeneous distribution of alginate formulations in the  
424 stomach with the formation of 'lumps', compared to the  
425 homogenous distribution of guar gum. Initial 'gel lump'  
426 formation was observed 10 minutes postprandially, other  
427 'lumps' developed over time, compatible with the pH  
428 decrease normally observed following ingestion of a meal.  
429 In addition, the strong gelling alginate resulted in  
430 significantly increased intragastric gel 'lump' production  
431 compared to the weak gelling. Data from 'lump'  
432 classification showed liquid filled 'lumps' were formed  
433 predominantly with the strong gelling alginate; the  
434 researchers hypothesise this gel strength is sufficient to  
435 allow layer formation which resist break forces caused by  
436 stomach motion.

437 There is a prevailing assumption that BMI measurement  
438 is strongly associated with body fatness and consequent  
439 morbidity and mortality (Gallagher et al., 2000).

440 Increased postprandial blood glucose is independently  
441 related to the risk of cardiovascular disease and all-cause  
442 mortality in newly diagnosed type 2 diabetics. Some  
443 individuals classified overweight by BMI do not have  
444 high % body fat. Conversely, others who have normal or  
445 healthy BMIs have a relatively high body fat %.

446 Individuals who are misclassified by BMI are reportedly  
447 uncommon relative to the UK population as a whole but

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

448 since body fatness is a stronger predictor of increased  
449 fasting glucose than BMI (Kriketos et al., 2003) it is more  
450 appropriate and meaningful to divide subjects in the  
451 present study by body fat %. In support of this, the  
452 present study clearly shows subjects in the upper body  
453 fat haptile had comparatively elevated early-phase  
454 glycaemic excursion to those in the lower body fat haptile.

455

## 456 **5.0 CONCLUSIONS**

457 We conclude that an ionic-gelling sodium alginate drink  
458 can significantly attenuate postprandial glycaemic  
459 response in self-reported healthy males in comparison to  
460 an acid-gelling control. This effect persisted in subjects in  
461 both the lower and upper haptiles of body fatness. The  
462 benefits of optimising glycaemic control through the use  
463 of ionic-gelling sodium alginate products in patients with  
464 morbidity related to body fatness (including type 2  
465 diabetic and metabolic syndrome patients) warrant further  
466 investigation.

467

## 468 **ROLE OF THE FUNDING SOURCE**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

469 This work was financially supported by Technostics Ltd.,  
470 UK, who also had input into the overall design, write up  
471 and submission of this work.

472

473 **REFERENCES**

474 Augustin, L.S.A., Axelsen, M., Jenkins, D.J.A., Kendall,  
475 C.W.C., Smith, U., and Vuksan, V. Dietary fibre,  
476 carbohydrates, and insulin resistant diseases, *British*  
477 *Journal of Nutrition* 83 (2000), 157.

478 Brownlee, I.A., Allen, A., Pearson, J.P., Dettmar, P.W.,  
479 Havler, M.E., Atherton, M.R., and Onsøyen, E. Alginate  
480 as a source of dietary fibre, *Critical Reviews in Food*  
481 *Sciences and Nutrition* 45 (2005), pp. 497-510.

482 Carey, V.J., Walters, E.E., Colditz, G.A., Soloman, C.G.,  
483 Willett, W.C., Rosner, B.A., Speizer, F.E., and Manson,  
484 J.E. Body fat distribution and risk of non-insulin  
485 dependent diabetes mellitus in women, *American Journal*  
486 *of Epidemiology* 145 (1997), pp. 614-619.

487 Delargy, H.J., O'Sullivan, K.R., Fletcher, R.J., and  
488 Blundell, J.E. Effects of amount and type of dietary fibre  
489 (soluble and insoluble) on short-term control of appetite,  
490 *International Journal of Food Science and Nutrition* 48  
491 (1997), pp. 67-77.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

492 Fairchild, R.M., Ellis, P.R., Byrne, A.J., Luzio, S.D., and  
493 Mir, M.A. A new breakfast cereal containing guar gum  
494 reduces postprandial plasma glucose and insulin  
495 concentrations in normal-weight human subjects, *British*  
496 *Journal of Nutrition* 76 (1996), pp. 63-73.

497 Flammang, A.M., Kendall, D.M., Baumgartner, C.J.,  
498 Slagle, T.D., and Choe, Y.S. Effect of a viscous fiber bar  
499 on postprandial glycemia in subjects with type 2 diabetes,  
500 *Journal of the American College of Nutrition* 25 (2006), pp.  
501 409-414.

502 Gallagher, D., Heymsfield, S.B., Heo, M., Jebb, S.A.,  
503 Murgatroyd, P.R., and Sakamoto, Y. Healthy percentage  
504 body fat ranges: an approach for developing guidelines  
505 based on body mass index, *American Journal of Clinical*  
506 *Nutrition* 72 (2000), pp. 694-701.

507 Goñi, I., Valdivieso, L., and Garcia-Alonso, A. Nori  
508 seaweed consumption modifies glycemic response in  
509 healthy volunteers, *Nutrition Research* 20 (2000), pp.  
510 1367-1375.

511 Hoad, C.L., Rayment, P., Spiller, R.C., Marciani, L., de  
512 Celis Alonso, B., Traynor, C., Mela, D.J., Peters, H.P.F.,  
513 and Gowland, P.A. In vivo imaging of intragastric gelation  
514 and its effect on satiety in humans, *Journal of Nutrition*  
515 134 (2004), pp. 2293-2300.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

516 Jenkins, D.J.A., Axelsen, M., Kendall, C.W.C., Augustin,  
517 L.S.A., Vuksan, V., and Smith, U. Dietary fibre, lente  
518 carbohydrates and the insulin-resistant diseases, *British*  
519 *Journal of Nutrition* 83 (2000), pp. S157-S163.

520 Kaline, K., Bornstein, S.R., Bergmann, A., Hauner, H.,  
521 and Schwarz, P.E.H. The importance and effect of  
522 dietary fiber in diabetes prevention with particular  
523 consideration of whole grain products, *Hormone and*  
524 *Metabolic Research* 39 (2007), pp 687-693.

525 Kimura, Y., Watanabe, K., and Okuda, H. Effects of  
526 soluble sodium alginate on cholesterol excretion and  
527 glucose tolerance in rats, *Journal of Ethnopharmacology*  
528 54 (1996), pp. 47-54.

529 Kriketos, A.D., Carey, D.G., Jenkins, A.B., Chisholm, D.J.,  
530 Furler, S.M., and Campbell, L.V. Central fat predicts  
531 deterioration of insulin secretion index and fasting  
532 glycaemia: 6-year follow-up of subjects at varying risk of  
533 type 2 diabetes mellitus, *Diabetic Medicine* 20 (2003), pp.  
534 294-300.

535 Lavin, J.H., and Read, N.W. The effect on hunger and  
536 satiety of slowing the absorption of glucose: relationship  
537 with gastric emptying and postprandial blood glucose and  
538 insulin responses, *Appetite* 25 (1995), pp. 89-96.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

539 Mattes, R.D. Effects of a combination fibre system on  
540 appetite and energy intake in overweight humans,  
541 *Physiology and Behavior* 90 (2007), pp. 705-711.

542 McPherson, K., Marsh, T., and Brown, M. Modelling  
543 Future Trends in Obesity and the Impact on Health  
544 Foresight Tackling Obesities: Future Choices (2nd  
545 Edition, 2007) Government Office for Science, UK.

546 Meyer, K.A., Kushi, L.H., Jacobs Jr., D.R., Slavin, J.,  
547 Sellers, T.A., and Folsom, A.R. Carbohydrates, dietary  
548 fibre, and incident type 2 diabetes in older women.  
549 *American Journal of Clinical Nutrition* 71 (2000), pp. 921-  
550 930.

551 Pascot, A., Lemieux, S., Lemieux, I., Prud'homme, D.,  
552 Tremblay, A., Bouchard, C., Nadeau, A., Couillard, C.,  
553 Tchernof, A., Bergeron, J., and Després, J.P. Age-related  
554 increase in visceral adipose tissue and body fat and the  
555 metabolic risk profile of premenopausal women, *Diabetes*  
556 *Care* 22 (1999), pp. 1471-1478.

557 Paxman, J.R., Richardson, J.C., Dettmar, P.W., and  
558 Corfe, B.M. Alginate reduces the increased uptake of  
559 cholesterol and glucose in overweight male subjects: a  
560 pilot study, *Nutrition Research* 28 (2008a), pp. 501-505.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

561 Paxman, J.R., Richardson, J.C., Dettmar, P.W., and  
562 Corfe, B.M. Daily ingestion of alginate reduces energy  
563 intake in free-living subjects, *Appetite* 51 (2008b), pp.  
564 713-719.

565 Peters, H.P.F., Koppert, R.J., Boers, H.M., Ström, A.,  
566 Melnikov, S.M., Haddeman, E., Schuring, E.A.H., Mela,  
567 D.J. and Wiseman, S.A. Dose-dependent suppression of  
568 hunger by a specific alginate in a low-viscosity drink  
569 formulation, *Obesity* (2011), doi: 10.1038/oby.2011.63.

570 Pinhas-Hamiel, O., and Zeitler, P. The global spread of  
571 type 2 diabetes mellitus in children and adolescents, *The*  
572 *Journal of Pediatrics* 146 (2005), pp. 693-700.

573 Prentice, A., and Jebb, S. Energy intake/physical activity  
574 interactions in the homeostasis of body weight regulation,  
575 *Nutrition Reviews* 62 (2004), pp. S98-S104.

576 Rezende, F.A.C., Rosado, L.E.F.P.L., Ribeiro, R.C.L.,  
577 Vidigal, F.C., Vasques, A.C.J., Bonard, I.S., and de  
578 Carvalho, C.R. Body mass index and waist circumference:  
579 association with cardiovascular risk factors, *Arquivos*  
580 *Brasileiros de Cardiologia* 87 (2006), pp. 728-734.

581 Rigaud, D., Paycha, F., Meulemans, A., Merrouche, M.,  
582 and Mignon, M. Effect of psyllium on gastric emptying,  
583 hunger feeling and food intake in normal volunteers: a

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

584 double blind study. *European Journal of Clinical Nutrition*  
585 52 (1998), pp. 239-245.

586 Seal, C.J., and Mathers, J.C. Comparative  
587 gastrointestinal and plasma cholesterol responses of rats  
588 fed on cholesterol-free diet supplemented with guar gum  
589 and sodium alginate, *British Journal of Nutrition* 85 (2001),  
590 pp. 317-324.

591 Sierra, M., Garcia, J.J., Fernández, N., Diez, M.J., Calle,  
592 A.P., and Farmafibra Group. Therapeutic effects of  
593 psyllium in type 2 diabetic patients, *European Journal of*  
594 *Clinical Nutrition* 56 (2002), pp. 830-842.

595 Strugala, V., Kennington, E.J., Skjåk-Bræk, G., and  
596 Dettmar, P.W. Bioactive properties of epimerised  
597 alginates. In Williams, P.A., and Phillips, G.O. (Eds.),  
598 *Gums and Stabilisers for the Food Industry 12* Cambridge,  
599 UK, The Royal Society of Chemistry (2004), pp.262-271.

600 Stunkard, A.J., and Messick, S. The three-factor eating  
601 questionnaire to measure dietary restraint disinhibition  
602 and hunger, *Journal of Psychosomatic Research* 29  
603 (1985), pp. 71-83.

604 Torsdottir, I., Alpsten, M., Holm, G., Sandberg, A.S., and  
605 Tolli, J. A small dose of soluble alginate fiber affects  
606 postprandial glycemia and gastric emptying in humans

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

607 with diabetes. *Journal of Nutrition* 121 (1991), pp. 795-  
608 799.

609 Welch, R.W. Can dietary oats promote health? *British*  
610 *Journal of Biomedical Science* 51 (1994), pp. 260-270.

611 Williams, J.A., Lai, C., Corwin, H., Ma, Y., Maki, K.C.,  
612 Garleb, K.A., and Wolf, B.W. Inclusion of guar gum and  
613 alginate into a crispy bar improves postprandial glycemia  
614 in humans. *Journal of Nutrition* 134 (2004), pp. 886-889.

615 Wolf, B.W., Lai, C., Kipnes, M.S., Ataya, D.G., Wheeler,  
616 K.B., Zinker, B.A., Garleb, K.A., and Firkins, J.L.  
617 Glycemic and insulinemic responses of nondiabetic  
618 healthy adult subjects to an experimental acid-induced  
619 viscosity complex incorporated into a glucose beverage.  
620 *Nutrition* 18 (2002), pp. 621-626.

621 Zaninotto, P., Wardle, H., and Stamatakis, E. (on behalf  
622 of the Department of Health, UK). *Forecasting Obesity to*  
623 *2010* (2006) [online]  
624 [http://www.dh.gov.uk/en/Publicationsandstatistics/Publica](http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsStatistics/DH_4138630)  
625 [tions/PublicationsStatistics/DH\\_4138630](http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsStatistics/DH_4138630) (last accessed;  
626 May 2011).

627

628

629 **TABLES**

630 Table 1

631 Subject characteristics

		<i>n</i>	Range	Mean ±	SD
<b>Age (y)</b>		<b>40</b>	<b>18 – 55</b>	<b>30.03 ±</b>	<b>11.21</b>
<b>Sodium Alginate</b>	Lower BF%	9	20 – 31	23.89 ±	4.05
	Upper BF%	11	18 – 51	34.55 ±	10.47
	TOTAL	20	18 – 51	29.75 ±	9.71
<b>Excipient Free</b>	Lower BF%	11	21 – 32	24.00 ±	3.19
	Upper BF%	9	19 – 55	38.00 ±	15.94
	TOTAL	20	19 – 55	30.30 ±	12.78
<b>BMI (kg/m<sup>2</sup>)</b>		<b>40</b>	<b>18.6 – 39.4</b>	<b>26.02 ±</b>	<b>4.41</b>
<b>Sodium Alginate</b>	Lower BF%	9	21.7 – 24.7	23.34 ±	1.07
	Upper BF%	11	22.7 – 35.2	29.07 ±	3.03
	TOTAL	20	21.7 – 35.2	26.50 ±	3.72
<b>Excipient Free</b>	Lower BF%	11	18.6 – 26.0	22.32 ±	2.33
	Upper BF%	9	23.0 – 35.6	29.47 ±	4.72
	TOTAL	20	18.6 – 39.4	25.54 ±	5.06
<b>Body Fat (BF) %</b>		<b>40</b>	<b>7.1 - 35.6</b>	<b>17.54 ±</b>	<b>7.05</b>
<b>Sodium Alginate</b>	Lower BF%	9	7.1 – 11.9	10.31 ±	1.56
	Upper BF%	11	16.8 – 31.7	22.58 ±	4.94
	TOTAL	20	7.1 – 31.7	17.06 ±	7.29
<b>Excipient Free</b>	Lower BF%	11	9.2 – 15.4	12.76 ±	1.82
	Upper BF%	9	18.1 – 35.6	24.47 ±	5.09
	TOTAL	20	9.2 – 35.6	18.03 ±	6.96

632

633

634 **FIGURE CAPTIONS**

635 Figure 1

636 Mean delta AUC glycaemia ( $\pm 1$ SD)

637 Following ingestion of the SA preload (solid line, filled  
638 diamonds), mean delta AUC glycaemia was reduced by  
639 52.5% when compared to the EF preload (broken line,  
640 open squares). There was a significant effect of preload  
641 treatment type on mean delta AUC ( $p = .010$ ). In addition,  
642 preload treatment type had a significant effect (\*) on  
643 mean delta glycaemia at 90 minutes ( $p < .0005$ ), 150  
644 minutes ( $p = .003$ ), 180 minutes ( $p = .021$ ) and 210  
645 minutes ( $p = .013$ ).

646

647 Figure 2

648 Mean peak postprandial glycaemia ( $\pm 1$ SD)

649 There was a significant effect of preload treatment type  
650 on mean peak postprandial glycaemia (SA solid bars;  $\underline{M} =$   
651  $6.06 \pm .59$  mmol/L compared to EF shaded bars;  $\underline{M} =$   
652  $6.92 \pm .70$  mmol/L; \* $p < .0005$ ).

653

654 Figure 3

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

655 Mean delta AUC glycaemia by body fat haptile

656 When subjects were split by haptiles of body fat % (solid

657 line = upper body fat haptile, broken line = lower body fat

658 haptile) there was a significant effect of body fat %

659 classification (§) on mean delta glycaemia at 120 minutes

660 ( $p = .005$ ) 150 minutes ( $p = .012$ ) 180 minutes ( $p = .049$ )


661 and 210 minutes ( $p = .046$ ), irrespective of treatment type

662 (solid diamonds = sodium alginate, open squares =


663 excipient free).



# Figure 1



**Figure 2**



**Figure 3**

