

**Attitudes and preferences towards self-help treatments for depression in comparison to psychotherapy and antidepressant medication**

HANSON, Katie, WEBB, Thomas L, SHEERAN, Paschal and TURPIN, Graham

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/9558/>

---

This document is the Accepted Version [AM]

**Citation:**

HANSON, Katie, WEBB, Thomas L, SHEERAN, Paschal and TURPIN, Graham (2015). Attitudes and preferences towards self-help treatments for depression in comparison to psychotherapy and antidepressant medication. *Behavioural and Cognitive Psychotherapy*, 44 (02), 129-139. [Article]

---

**Copyright and re-use policy**

See <http://shura.shu.ac.uk/information.html>

**BEHAVIOURAL AND  
COGNITIVE PSYCHOTHERAPY**



**Attitudes and preferences towards self-help treatments for  
depression in comparison to psychotherapy and  
antidepressant medication**

Journal:	<i>Behavioural and Cognitive Psychotherapy</i>
Manuscript ID:	BCP-01026-14.R1
Manuscript Type:	Brief Clinical Report
Keywords:	depression, self-help, acceptance

SCHOLARONE™  
Manuscripts

**Abstract**

*Background:* Self-help is an effective treatment for depression. Less is known, however, about how acceptable people find different self-help treatments for depression.

*Aims:* To investigate preferences and attitudes toward different self-help treatments for depression in comparison to psychotherapy and antidepressants.

*Methods:*  $N = 536$  people who were not actively seeking treatment for depression were randomly assigned to read about one of five treatment options (bibliotherapy, Internet-based self-help, guided self-help, antidepressants, or psychotherapy) before rating how acceptable they found the treatment. Participants also ranked the treatments in order of preference.

*Results:* Psychotherapy and guided self-help were found to be the most acceptable and preferred treatment options. Antidepressants and bibliotherapy were found to be the least acceptable treatments, with antidepressants rated as the most likely to have side effects. Preference data reflected the above findings – psychotherapy and guided self-help were the most preferred treatment options.

*Conclusions:* The findings highlight differences in attitudes and preferences between guided and unguided self-help interventions; and between self-help interventions and psychotherapy. Future research should focus on understanding why unguided self-help interventions are deemed to be less acceptable than guided self-help interventions for treating depression.

Word count: 186 words

*Keywords:* Depression, Acceptability, Preference, Self-help

Running head: HOW ACCEPTABLE IS SELF HELP?

2

1  
2  
3 Attitudes and preferences towards self-help treatments for depression in comparison  
4  
5 to psychotherapy and antidepressant medication  
6

7 Marris (1995) defines self-help as “the use of written materials or computer  
8 programs ... for the purpose of gaining understanding or solving problems relevant to  
9 a person’s developmental or therapeutic needs” (p. 846). Self-help materials typically  
10 (1) provide the user with the means to identify their problem by offering information  
11 about the symptoms commonly experienced, and (2) offer advice on how to overcome  
12 problems, along with techniques for alleviating symptoms, and examples of how to  
13 use these techniques. Self-help can be delivered in many formats including books  
14 (termed ‘bibliotherapy’) or via the Internet. Self-help can also be offered as either a  
15 guided or unguided intervention, where guided self-help involves the patient helping  
16 themselves with some form of support from another person (Lucock, Barber, Jones, &  
17 Lovell, 2007). Self-help treatments are currently recommended by the National  
18 Institute for Health and Clinical Excellence (NICE, 2009) for depression and meta-  
19 analyses show that self-help interventions for depression are more effective than no-  
20 treatment and comparable to psychotherapies and antidepressants (Cuijpers et al.,  
21 2013).  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

40 Although the evidence suggests that self-help treatments for depression are  
41 relatively effective, less is known about peoples’ attitudes toward self-help  
42 treatments; in particular, whether people deem self-help interventions to be an  
43 acceptable treatment approach and the extent to which self-help interventions are  
44 preferred to other treatment options. Research suggests that patients with depression  
45 show a preference for psychotherapy over antidepressants (Raue & Schulberg, 2007)  
46 and that patients may benefit more from treatments that they show a preference for  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56 (e.g., Kocsis et al., 2009; Kwan, Dimidjian, & Rizvi, 2010; Lin, et al., 2005; Mergl  
57  
58  
59  
60

1  
2  
3 et al., 2011; Moradveisi, Huibers, Renner, & Arntz, 2014). Other studies, however,  
4  
5 have found no impact of patient preference on outcomes (e.g., Leykin et al., 2007;  
6  
7 Moradveisi et al., 2014; Raue, Schulberg, Heo, Klimstra, & Bruce, 2009) and these  
8  
9 discrepancies have led researchers to explore variables, such as beliefs about the  
10  
11 cause of depression (Dunlop et al., 2012; Khalsa, McCarthy, Sharpless, Barrett, &  
12  
13 Barber, 2011; Steidtmann et al., 2012), which may moderate the link between  
14  
15 preference and treatment outcome. Preference has also been linked to engagement  
16  
17 with treatment. Specifically, there is evidence that treatment preference influences  
18  
19 initiation of treatment (King et al., 2005; Raue et al., 2009; Raue & Schulberg, 2007),  
20  
21 adherence (Elkin et al., 1999; Raue et al., 2009), attrition (Kwan et al., 2010) and  
22  
23 therapeutic alliance (Iacoviello et al., 2007; Kwan et al., 2010). In short, attitudes  
24  
25 toward treatment are likely to influence treatment outcomes.  
26  
27  
28

29  
30 Although we know much about preferences for psychotherapy versus  
31  
32 antidepressants little research has examined preferences towards self-help treatments  
33  
34 and how they fare in relation to psychotherapy or antidepressants (Cooper-Patrick et  
35  
36 al., 1997). There are, however, some studies that can provide indicative evidence.  
37  
38 Landreville et al. (2001) investigated attitudes towards treatments for depression.  
39  
40 Participants aged 65 years and over were asked to read one of two descriptions of  
41  
42 depression (either mild to moderate or severe depression) before reading descriptions  
43  
44 of psychotherapy, bibliotherapy, and antidepressant treatments. Participants rated how  
45  
46 acceptable they believed that they would find each of the treatments using the  
47  
48 modified Treatment Evaluation Inventory (Landreville & Guérette, 1998).  
49  
50 Psychotherapy and bibliotherapy were both rated as more acceptable than  
51  
52 antidepressants for treating mild to moderate levels of depression (but not for severe  
53  
54 depression).  
55  
56  
57  
58  
59  
60

Running head: HOW ACCEPTABLE IS SELF HELP?

4

1  
2  
3 Mitchell and Gordon (2007) explored attitudes towards computerised  
4  
5 cognitive behavioural therapy (CCBT) amongst 122 university students, 65% of  
6  
7 whom had prior or current experience of depression or anxiety. Participants were  
8  
9 asked to read a brief description of CCBT before rating the treatment in terms of its  
10  
11 credibility, the expectancy that its use would improve the symptoms of depression and  
12  
13 the perceived likelihood of using this form of treatment. The findings suggested that  
14  
15 the sample rated CCBT as only ‘somewhat credible’, with moderately low  
16  
17 expectations for improvement reported. In terms of the participants rating the  
18  
19 likelihood of using the treatment, only 10% said that they would be likely to choose  
20  
21 this form of treatment as their first choice, with nearly 55% of the sample saying they  
22  
23 would prefer counselling.  
24  
25  
26

27 Schneider, Foroushani, Grime, and Thornicof (2014) explored how  
28  
29 acceptable self-help intervention for depression was deemed to be.  $N = 637$   
30  
31 employees, with symptoms of depression, took part in an online CCBT intervention  
32  
33 for 5 weeks. Prior to the intervention, participants were asked to rate how acceptable  
34  
35 they would find using CCBT over going to see a GP or psychologist. At the end of the  
36  
37 intervention they were also asked to rate how acceptable they found the treatment.  
38  
39 Schneider et al. found that, at baseline, 65% of the sample rated CCBT to be equally  
40  
41 acceptable to seeing a psychologist and 80% of the sample found CCBT as acceptable  
42  
43 as seeing a GP. There were no significant changes in how acceptable participants  
44  
45 found the treatments at the end of the study, suggesting that attitudes expressed in  
46  
47 response to hypothetical scenarios (e.g., “How do you think you would feel...?”)  
48  
49 reflect how people actually feel if they experience the treatment.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## The Present Research

Although the studies described above provide insight into how acceptable people find different self-help treatments for depression, a number of important questions remain unanswered. First, no study to date has compared how acceptable people find different types of self-help. The present research will examine attitudes toward and preferences for guided self-help, unguided bibliotherapy, and unguided Internet-based self-help. The research will also investigate how acceptable people find traditional treatments (namely, psychotherapy and antidepressants), in order to provide a comparison. Second, research to date has focused on how acceptable people find different treatments, but has not yet explored treatment preferences. Specifically, if peoples' first choice of treatment is unavailable (e.g., there is a long waiting list for psychotherapy), then it is currently unclear what treatment they might prefer instead. Pressures on health services mean that this question is significant. The present research, therefore, also asked participants to rank treatments in order of preference. We also measured current levels of depression and previous treatment experience to investigate whether they influence attitudes and preferences.

## Method

### Sample

Staff and students at a large University in the UK were emailed an invitation to take part in a study examining attitudes toward treatments for depression. As we were interested in attitudes towards treatments that are not clouded by actual help-seeking behaviour, we sought to recruit an analogue sample who were not actively seeking treatment for depression. No inclusion/exclusion criteria were set in terms of level of depression or diagnosis.  $N = 536$  participants responded. Participants were aged between 17 and 76 years ( $M = 29.90$ ,  $SD = 12.57$ ) and 65.11% were female,

Running head: HOW ACCEPTABLE IS SELF HELP?

6

53.73% were students, and 57.46% were White British. Participants' mean score on the Center for Epidemiologic Studies Depression Scale (CES-D; Radloff, 1977) was 26.75 ( $SD = 13.64$ ), indicating relatively high levels of depression (Radloff, 1991).

### Procedure

Participants who agreed to take part in the study were asked to read a brief description of depression and a personal account of how it feels to be depressed.<sup>1</sup> Participants were then randomly allocated to read a detailed description of one of five treatment for depression: psychological therapy, antidepressants, guided self-help, bibliotherapy or Internet-based self-help. Each description contained information regarding what the treatment involved, what the different treatment subtypes were (e.g., examples of the different types of psychotherapy available) and how the treatment could be accessed.<sup>2</sup>

Once participants had read the detailed treatment description, they rated how acceptable they found the treatment using a modified version of the Treatment Evaluation Inventory (TEI; Kazdin, 1980; Landreville & Guérette, 1998). The TEI was modified to measure how acceptable people find different treatments for depression and consisted of nine questions (e.g. "*How acceptable would you find this treatment for treating your depression?*" and "*To what extent do you think there might be risks in undergoing this kind of treatment?*"). In line with the findings of

---

<sup>1</sup> The description and personal account were 276 words in length and were taken from the website of the mental health charity, Mind (Stewart, 2010). Pilot research suggested that the account brought to life the experience of depression and accurately reflected how it feels to be depressed. Further details of this pilot research, along with the materials used are available in the supplementary materials.

<sup>2</sup> The descriptions of psychological therapy and antidepressants were taken from the UK mental health charity, Rethink (Rethink, 2012a, 2012b). These documents were edited to make them shorter and they were used as a template for the descriptions of the self-help treatments. Pilot research suggested that the treatment descriptions portrayed what the treatment involved and reflected what receiving the treatment would be like. Further details of this pilot research, along with the descriptions used are available in the supplementary materials.



Running head: HOW ACCEPTABLE IS SELF HELP?

7

1  
2  
3 Landreville and Guérette (1998) principle components analysis with oblimin rotation,  
4 identified two components that accounted for 69.36% of the variance. The two factors  
5 were labelled “acceptability” (e.g., “*How consistent is this treatment with your*  
6 *common sense or everyday notions about what a treatment for depression should*  
7 *be?*”) ( $\alpha = 0.92$ ) and “side effects” (e.g., “*To what extent do you think undesirable*  
8 *side effects are likely to result from this treatment?*”) ( $\alpha = 0.66$ ). Factor scores were  
9 computed for each component. Landreville and Guérette (1998) noted good  
10 concurrent validity, internal consistency and test-retest reliability when using the scale  
11 to assess treatment acceptability and side effects in relation to treatments for  
12 depression.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

25 All participants were then asked to read brief descriptions of all five  
26 treatments, which were developed by shortening the detailed treatment descriptions.  
27 Participants were asked to rank the five treatments in order of preference. Finally,  
28 participants completed a questionnaire, which measured current levels of depression  
29 (using the CES-D, Radloff, 1977) and treatment experience (e.g. “If you have  
30 suffered from depression, which treatments have you used?”), as well as demographic  
31 information (gender, age, ethnic origin, and occupation).  
32  
33  
34  
35  
36  
37  
38  
39  
40

#### 41 Analysis strategy

42 One-way between-groups multivariate analyses of variance (MANOVA) was  
43 used to investigate differences in ratings of acceptability and side effects between the  
44 five treatment descriptions, and to investigate the impact of current levels of  
45 depression and treatment experience on ratings of acceptability and side effects. A  
46 Friedman test was used to investigate differences in preference ratings, with  
47 Wilcoxon sign-ranks tests used for post-hoc comparison.  
48  
49  
50  
51  
52  
53  
54  
55

#### 56 Results

57  
58  
59  
60

Running head: HOW ACCEPTABLE IS SELF HELP?

8

### **How Acceptable are Treatments for Depression?**

Table 1 shows the average levels of acceptability and side effects for each of the five treatment options. Perceptions of both acceptability,  $F(4, 531) = 18.97, p < 0.01, \eta^2 = 0.13$ , and side effects,  $F(4, 531) = 18.19, p < 0.01, \eta^2 = 0.12$ , differed between treatments. Pairwise comparisons with Bonferroni adjustment revealed that psychotherapy and guided self-help were rated as the most acceptable treatments.

There was no significant difference in how acceptable participants rated psychotherapy and guided self-help ( $p = 0.30$ ). Psychotherapy and guided self-help were, in turn, rated as significantly more acceptable than antidepressants, bibliotherapy, and Internet-based self-help ( $p < 0.01$ ).

In terms of perceived side effects, pairwise comparisons with Bonferroni adjustment revealed that antidepressants were rated as significantly ( $p < 0.01$ ) more likely to have side effects than psychotherapy that, in turn, was deemed to have significantly more side effects than bibliotherapy, guided self-help and Internet-based self-help. There were no differences in perceived side effects between any of the other self-help interventions ( $ps < 0.05$ ).

### **Does Current Depression or Treatment Experience Influence how Acceptable People Find Treatments?**

Radloff (1991) proposed that scores of 16 or higher on the CES-D scale indicate the presence of depression symptoms. In the present sample 64.74% of participants scored above this cut off point. Table 1 shows how acceptable participants found each of the five treatments separately for those with and without symptoms of depression. There was only a statistically significant difference in ratings between those with and without symptoms of depression for guided self-help,  $F(1, 85) = 7.72, p = .01, \eta^2 = 0.08$ . Depressed participants rated guided self-help as

Running head: HOW ACCEPTABLE IS SELF HELP?

9

1  
2  
3 being significantly less acceptable than did participants without symptoms of  
4 depression. There were no differences in acceptability or side effects between  
5 participants with and without symptoms of depression for the remaining treatments  
6  
7  
8  
9  
10 ( $F_s < 2.99, ns$ ).

11  
12 Table 1 also shows levels of acceptability and side effects associated with  
13 each of the five treatments for participants who had previous experience of the  
14 treatments versus those who did not. A series of one-way between-groups  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
MANOVAs revealed no statistically significant differences between those with and  
without treatment experience on the combined dependent variables ( $F_s < 2.61, ns$ ).

### **Which Treatments for Depression do Participants Prefer?**

Table 2 shows participants preferences for the five different types of  
treatment. There were significant differences between the mean rank scores for the  
five brief treatment descriptions ( $X^2 = 853.34, p < 0.001$ ). Post-hoc comparisons  
showed that psychotherapy was preferred to all other treatments; guided self-help ( $z =$   
 $-14.23, p < 0.01$ , antidepressants ( $z = -16.79, p < 0.01$ ), bibliotherapy ( $z = -18.55, p <$   
 $0.01$ ), and Internet-based self-help ( $z = -18.99, p < 0.01$ ). Guided self-help was  
preferred to antidepressants ( $z = -4.53, p < 0.01$ ), bibliotherapy ( $z = -10.79, p < 0.01$ ),  
and Internet-based self-help ( $z = -14.77, p < 0.01$ ). Antidepressants were preferred to  
bibliotherapy ( $z = -4.38, p < 0.01$ ) and Internet-based self-help ( $z = -8.31, p < 0.01$ ).  
Finally, bibliotherapy was preferred to Internet-based self-help ( $z = -6.09, p < 0.01$ ).

### **Discussion**

To investigate peoples' attitudes toward self-help treatments for depression,  
the present research compared perceptions of three types of self-help with  
psychotherapy and antidepressants. Consistent with the findings of other research  
(e.g., Raue & Schulberg, 2007), psychotherapy was rated as more acceptable and

Running head: HOW ACCEPTABLE IS SELF HELP?

10

1  
2  
3 preferable to antidepressants. Extant research had not, however, explored how  
4  
5 acceptable people find different forms of self-help as an alternative to psychotherapy  
6  
7 and antidepressants. Our findings suggest that psychotherapy remained the most  
8  
9 preferred and most acceptable treatment option. However, guided self-help was  
10  
11 deemed to be equally acceptable, with the caveat that participants with depression  
12  
13 rated guided self-help as being less acceptable than non-depressed participants.  
14  
15 Across the sample as a whole, psychotherapy and guided self-help were rated as more  
16  
17 acceptable than bibliotherapy and Internet-based self-help.  
18  
19

20  
21 The preference for guided over unguided forms of self-help is consistent with  
22  
23 the findings of Mohr, Siddique, Ho, Duffecy, Jin, and Fokuo (2010) who found that  
24  
25 greater interest in receiving mental health treatment was associated with greater  
26  
27 interest in receiving face-to-face contact. The findings are also consistent with  
28  
29 findings in relation to anxiety. For example, Sharp, Power, and Swanson (2004) found  
30  
31 that the majority of people on a waiting list for treatment for anxiety disorders chose  
32  
33 to undertake individual therapy over unguided self-help. Antidepressants and  
34  
35 bibliotherapy were found to be the least acceptable treatments, with antidepressants  
36  
37 rated as the most likely to have side effects. This latter finding is consistent with  
38  
39 previous research suggesting that antidepressants are an unpopular treatment option  
40  
41 (Bedi et al., 2000), possibly due to associated side effects (Khawam, Laurencic, &  
42  
43 Malone, 2006).  
44  
45  
46

#### 47 **Limitations and Future Directions**

48  
49 **One potential** drawback to the present research is the use of a between sample  
50  
51 design, where participants read just one of five detailed treatment descriptions before  
52  
53 rating how acceptable they would find that treatment. Arguably, it may have been  
54  
55 preferable to have participants read detailed descriptions of all treatments. However,  
56  
57  
58  
59  
60

Running head: HOW ACCEPTABLE IS SELF HELP?

11

1  
2  
3 this was deemed to be overly onerous and not an accurate reflection of how  
4  
5 treatments are typically presented to people with depression. The other advantage of  
6  
7 randomly allocating participants to treatment over, for example, examining how  
8  
9 acceptable actual patients find a treatment that they have been offered, is that  
10  
11 potential confounds such as past experience or demographic factors are controlled for.  
12  
13 Moreover, the design enabled us to carefully control the amount and nature of  
14  
15 information that participants received about each treatment. The present research did,  
16  
17 however, also capitalise on a within sample design, where participants read brief  
18  
19 descriptions of each treatment and then ranked them in order of preference. The  
20  
21 preference data matched the acceptability data, in that both psychotherapy and guided  
22  
23 self-help were viewed as the most acceptable and most preferred treatment options. It  
24  
25 is, however, worth noting that the information provided in the brief treatment  
26  
27 descriptions may not have been detailed enough to provide sufficient information for  
28  
29 participants to make an informed decision on preference. In addition, the present  
30  
31 research did not consider preferences for the use of combined treatments (e.g.,  
32  
33 antidepressant medication and psychotherapy) or the preference for no-treatment or  
34  
35 watchful waiting (Dwight Johnson, Apesoa-Varano, Hay, Unutzer, & Hinton, 2013).  
36  
37 These might be useful issues to explore in future research.  
38  
39  
40  
41  
42

43 A second potential limitation is the use of an analogue design, recruiting  
44  
45 participants who were not actively seeking treatment for depression. The advantage of  
46  
47 this design is that attitudes towards treatments are not clouded by actual help-seeking  
48  
49 behaviour. Indeed, no differences were found in ratings of acceptability and perceived  
50  
51 side effects between participants with previous treatment experience and participants  
52  
53 without. Furthermore, there were few differences between those who had current  
54  
55 symptoms of depression and those who did not. Both these findings suggest that our  
56  
57  
58  
59  
60

Running head: HOW ACCEPTABLE IS SELF HELP?

12

1  
2  
3 analogue sample is likely to closely approximate the beliefs of a clinical sample,  
4  
5 which is often the case in the literature that compares clinical and analogue attitudes  
6  
7 towards treatments for mental health disorders (e.g., Feeny & Zoellner, 2004;  
8  
9 McHugh, Whitton, Peckham, Welge & Otto, 2013). Having said this, further research  
10  
11 could aim to replicate the present approach in a **treatment-seeking sample**.  
12  
13

#### 14 **Implications for research and clinical practice**

15  
16 One of the cornerstones of the stepped-care model is the assumption that the  
17  
18 treatments that are offered are acceptable to patients (Bower & Gilbody, 2005). As  
19  
20 such, researchers have begun to explore treatment attitudes and preferences for a  
21  
22 range of disorders (e.g., Sumner et al., 2014). **Our findings suggest that unguided**  
23  
24 **interventions are less acceptable and less preferable to interventions that contain an**  
25  
26 **element of personal contact, such as psychotherapy or guided self-help. Researchers**  
27  
28 **now need to further explore why interventions that contain personal contact are**  
29  
30 **preferred to unguided interventions. Macdonald, Mead, Bower, Richards and Lovell**  
31  
32 **(2007) interviewed participants who had received guided self-help for depression and**  
33  
34 **found that participants reported difficulties engaging with the intervention due to the**  
35  
36 **symptoms of depression, such as low motivation, or poor concentration. It is possible**  
37  
38 **that these issues are even more salient for those receiving unguided self-help as they**  
39  
40 **have no-one to help them to overcome these barriers. In addition, treatments that**  
41  
42 **incorporate personal contact may be perceived to provide more helpful and specific**  
43  
44 **guidance/coaching around the implementation of self-help techniques.**  
45  
46  
47  
48

49  
50 Finally, given that research suggests that patients allocated their preferred  
51  
52 treatment (out of psychotherapy or antidepressants) are more likely to engage with  
53  
54 that treatment, potentially improving efficacy (e.g., Kwan et al., 2010), future  
55  
56 research might usefully assess whether this is also the case for unguided self-help  
57  
58  
59  
60

1  
2  
3 interventions. Although less effective than guided self-help (Gellatly et al., 2007),  
4  
5 unguided interventions have been found to be effective for depression (e.g., Cuijpers,  
6  
7 1997), however there are often problems with poor engagement (e.g., Christensen,  
8  
9 Griffiths & Farrer, 2009). Future research needs to assess whether this is due to the  
10  
11 patient feeling that the treatment is unacceptable and/or having a preference for  
12  
13 another treatment. If this is the case, then possible solutions include; (1) providing  
14  
15 extra funding to increase the availability of acceptable treatment options, namely  
16  
17 psychotherapy and guided self-help, (2) investigating which forms of support are  
18  
19 acceptable, as some forms of support are less costly to administer and equally  
20  
21 effective (in comparison to face-to-face support) such as telephone support (Farrand  
22  
23 & Woodford, 2013), or (3) implementing protocols to boost the acceptability of  
24  
25 unguided interventions. For example, a large-scale publicity campaign to educate the  
26  
27 general public in the efficacy of such treatment approaches. The Department of  
28  
29 Health (2013) announced £16 million pounds worth of funding over the next four  
30  
31 years for a campaign against mental health stigma and within this campaign there  
32  
33 could be scope to promote the use of unguided interventions.  
34  
35  
36  
37

### 38 References

- 39  
40 Bedi, N., Lee, A., Harrison, G., Chilvers, C., Dewey, M., Fielding, K., ... Duggan, C.  
41  
42 (2000). Assessing effectiveness of treatment of depression in primary care:  
43  
44 Partially randomized preference trial. *The British Journal of Psychiatry*, 177,  
45  
46 312-318. doi: 10.1192/bjp.177.4.312  
47  
48  
49 Bower, P., & Gilbody, S., (2005). Stepped care in psychological therapies: Access,  
50  
51 effectiveness and efficacy. *British Journal of Psychiatry*, 186, 11-17. doi:  
52  
53 10.1192/bjp.186.1.11  
54  
55  
56  
57  
58  
59  
60

Running head: HOW ACCEPTABLE IS SELF HELP?

14

- 1  
2  
3 Christensen, H., Griffiths, K. M., & Farrer, L. (2009). Adherence in Internet  
4  
5 interventions for anxiety and depression: Systematic review. *Journal of*  
6  
7 *Medical Internet Research, 11*, e13. doi: 10.2196/jmir.1194  
8  
9  
10 Cooper-Patrick, L., Powe, N. R., Jenckes, M. W., Gonzales, J. J., Levine, D. M., &  
11  
12 Ford, D. E. (1997). Identification of patient attitudes and preferences  
13  
14 regarding treatment of depression. *Journal of General Internal Medicine, 12*,  
15  
16 431-438. doi: 10.1046/j.1525-1497.1997.00075.x  
17  
18 Cuijpers, P. (1997). Bibliotherapy in unipolar depression: A meta-analysis. *Journal of*  
19  
20 *Behavioral Therapy and Experimental Psychiatry, 28*, 139-147. doi:  
21  
22 10.1016/S0005-7916(97)00005-0  
23  
24 Cuijpers, P., Berking, M., Andersson, G., Quigley, L., Kleiboer, A., & Dobson, K. S.  
25  
26 (2013). A meta-analysis of cognitive-behavioural therapy for adult depression,  
27  
28 alone and in comparison with other treatments. *Canadian Journal of*  
29  
30 *Psychiatry, 58*, 376–385.  
31  
32  
33  
34 Dunlop, B. W., Kelley, M. E., Mletzko, T. C., Velasquez, C. M., Craighead, W. E., &  
35  
36 Mayberg, H. S. (2012). Depression beliefs, treatment preference, and  
37  
38 outcomes in a randomized trial for major depressive disorder. *Journal of*  
39  
40 *Psychiatric Research, 46*, 375-381. doi: 10.1016/j.jpsychires.2011.11.003  
41  
42  
43 Dwight Johnson, M., Apesoa-Varano, C., Hay, J., Unutzer, J., & Hinton, L. (2013).  
44  
45 Depression treatment preferences of older white and Mexican origin men.  
46  
47 *General Hospital Psychiatry, 35*, 59-65. doi:  
48  
49 10.1016/j.genhosppsy.2012.08.003  
50  
51  
52 Elkin, I., Yamaguchi, J. L., Arnkoff, D. B., Glass, C. R., Sotsky, S. M., & Krupnick,  
53  
54 J. L. (1999). "Patient-Treatment Fit" and early engagement in therapy.  
55  
56 *Psychotherapy Research, 9*, 437 - 451. doi: 10.1080/10503309912331332851  
57  
58  
59  
60



- 1  
2  
3 Farrand, P., & Woodford, J. (2013). Impact of support on the effectiveness of written  
4  
5 cognitive behavioural self-help: A systematic review and meta-analysis of  
6  
7 randomised controlled trials. *Clinical Psychology Review, 33*, 182 - 195. doi:  
8  
9 10.1016/j.cpr.2012.11.001  
10  
11  
12 Feeny, N. C., & Zoellner, L. A. (November, 2004). *Prolonged exposure versus*  
13  
14 *sertraline: Primary outcomes for the treatment of chronic PTSD*. Paper  
15  
16 presented at the International Society for the Study of Traumatic Stress, New  
17  
18 Orleans, LA.  
19  
20  
21 Gellatly, J., Bower, P., Hennessy, S., Richards, D., Gilbody, S., & Lovell, K. (2007).  
22  
23 What makes self-help interventions effective in the management of depressive  
24  
25 symptoms? Meta-analysis and meta-regression. *Psychological Medicine, 37*,  
26  
27 1217-1228. doi: 10.1017/S0033291707000062  
28  
29  
30 Iacoviello, B. M., McCarthy, K. S., Barrett, M. S., Rynn, M., Gallop, R., & Barber, J.  
31  
32 P. (2007). Treatment preferences affect the therapeutic alliance: Implications  
33  
34 for randomized controlled trials. *Journal of Consulting and Clinical*  
35  
36 *Psychology, 75*, 194-198. doi: 10.1037/0022-006X.75.1.194  
37  
38  
39 Kazdin, A. E. (1980). Acceptability of alternative treatments for deviant child  
40  
41 behavior. *Journal of Applied Behavior Analysis, 13*, 259–273.  
42  
43 doi:10.1901/jaba.1980.13-259  
44  
45  
46 Kocsis, J. H., Leon, A. C., Markowitz, J. C., Manber, R., Arnow, B., Klein, D. N., &  
47  
48 Thase, M. E. (2009). Patient preference as a moderator of outcome for chronic  
49  
50 forms of major depressive disorder treated with nefazodone, cognitive  
51  
52 behavioral analysis system of psychotherapy, or their combination. *Journal of*  
53  
54 *Clinical Psychiatry, 70*, 354-361. doi: 10.4088/JCP.08m04371  
55  
56  
57  
58  
59  
60

Running head: HOW ACCEPTABLE IS SELF HELP?

16

1  
2  
3 Khalsa, S. R., McCarthy, K. S., Sharpless, B. A., Barrett, M. S., & Barber, J. P.

4  
5 (2011). Beliefs about the causes of depression and treatment preferences.

6  
7 *Journal of Clinical Psychology, 67*, 539-549. doi: 10.1002/jclp.20785

8  
9 Khawam, E. A., Laurencic, G. & Malone, D. A. (2006). Side effects of

10  
11 antidepressants: An overview. *Cleveland Clinic Journal of Medicine, 73*, 351-

12  
13 353. doi: 10.3949/ccjm.73.4.351

14  
15 King, M., Nazareth, I., Lampe, F., Bower, P., Chandler, M., Morou, M., ... & Lai, R.

16  
17 (2005). Impact of participant and physician intervention preferences on

18  
19 randomized trials: A systematic review. *Journal of the American Medical*

20  
21 *Association, 293*, 1089-1099. doi: 10.1001/jama.293.9.1089

22  
23 Kwan, B. M., Dimidjian, S., & Rizvi, S. L. (2010). Treatment preference,

24  
25 engagement, and clinical improvement in pharmacotherapy versus

26  
27 psychotherapy for depression. *Behaviour Research and Therapy, 48*, 799-804.

28  
29 doi: 10.1016/j.brat.2010.04.003

30  
31 Landreville, P., & Guérette, A. (1998). Psychometric properties of a modified version

32  
33 of the treatment evaluation inventory for assessing the acceptability of

34  
35 treatments for geriatric depression. *Canadian Journal on Aging, 17*, 414-424.

36  
37 doi: 10.1017/S071498080001268X

38  
39 Landreville, P., Landry, J., Baillargeon, L., Guérette, A., & Matteau, E. (2001). Older

40  
41 adults acceptance of psychological treatments for depression. *Journal of*

42  
43 *Gerontology: Psychological Sciences, 56B*, 285-291. doi:

44  
45 10.1093/geronb/56.5.P285

46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Leykin, Y., DeRubeis, R. J., Gallop, R., Amsterdam, J. D., Shelton, R. C., & Hollon,  
4  
5 S. D. (2007). The relation of patients' treatment preferences to outcome in a  
6  
7 randomized clinical trial. *Behavior Therapy*, 38, 209-217. doi:  
8

9  
10 10.1016/j.beth.2006.08.002

11  
12 Lin, P., Campbell, D. G., Chaney, E. F., Liu, C. F., Heagerty, P., Felker, B. L., &  
13  
14 Hedrick, S. C. (2005). The influence of patient preference on depression  
15  
16 treatment in primary care. *Annals of Behavioral Medicine*, 30, 164-173. doi:  
17

18  
19 10.1207/s15324796abm3002\_9

20  
21 Lucock, M. P., Barber, R., Jones., A., & Lovell, J. (2007). Service users' views of  
22  
23 self-help strategies and research in the UK. *Journal of Mental Health*, 16,  
24  
25 795–805.  
26

27  
28 Macdonald, W., Mead, N., Bower, P., Richards, D., & Lovell, K. (2007). A  
29  
30 qualitative study of patients perceptions of a 'minimal' psychological therapy.  
31  
32 *International Journal of Social Psychiatry*, 53, 23-35. doi:  
33

34  
35 10.1177/0020764006066841

36  
37 Marrs, R. W. (1995). A meta-analysis of bibliotherapy studies. *American Journal of*  
38  
39 *Community Psychology*, 23, 843-870. doi: 10.1007/BF02507018

40  
41 McHugh, K. R., Whitton, S. W., Peckham, A. D., Welge, J. A., & Otto, M. W.  
42  
43 (2013). Patient preference for psychological vs pharmacologic treatment of  
44  
45 psychiatric disorders: A meta-analytic review. *Journal of Clinical Psychiatry*,  
46  
47 74, 595 - 602. doi: 10.4088/JCP.12r07757  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Running head: HOW ACCEPTABLE IS SELF HELP? 18

1  
2  
3 Mergl, R., Henkel, V., Allgaier, A. K., Kramer, D., Hautzinger, M., Kohnen, R., ... &  
4  
5 Hegerl, U. (2010). Are treatment preferences relevant in response to  
6  
7 serotonergic antidepressants and cognitive-behavioral therapy in depressed  
8  
9 primary care patients? Results from a randomized controlled trial including a  
10  
11 patients' choice arm. *Psychotherapy and Psychosomatics*, *80*, 39-47. doi:  
12  
13 10.1159/000318772  
14  
15

16 Mitchell, N., & Gordon, P. K. (2007). Attitudes towards computerized CBT for  
17  
18 depression amongst a student population. *Behavioural and Cognitive*  
19  
20 *Psychotherapy*, *35*, 421-430. doi: 10.1017/S1352465807003700  
21  
22

23 Mohr, D. C., Siddique, P. H., Ho, J., Duffecy, J., Jin, L., & Fokuo, J. K (2010).  
24  
25 Interest in behavioral and psychological treatments delivered face-to-face, by  
26  
27 telephone, and by internet. *Annals of Behavioral Medicine*, *40*, 89-98. doi:  
28  
29 10.1007/s12160-010-9203-7  
30  
31

32 Moradveisi, L., Huibers, M., Renner, F., & Arntz, A. (2014). The influence of  
33  
34 patients' preference / attitude towards psychotherapy and antidepressant  
35  
36 medication on the treatment of major depressive disorder. *Journal of Behavior*  
37  
38 *Therapy and Experimental Psychiatry*, *45*, 170-177. doi:  
39  
40 10.1016/j.jbtep.2013.10.003  
41  
42

43 NICE (2009). *Depression in adults: The treatment and management of depression in*  
44  
45 *adults*. CG90. London: NICE.  
46

47 Radloff, L. S. (1977). The CES-D scale: A self-report depression scale for research  
48  
49 in the general population. *Applied Psychological Measurement*, *1*, 385-401.  
50  
51 doi: 10.1177/014662167700100306  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 Radloff, L. S. (1991). The use of the center for epidemiologic studies depression  
4 scale in adolescents and young adults. *Journal of Youth and Adolescence*, 20,  
5 149-166. doi: 10.1007/BF01537606  
6  
7  
8  
9  
10 Raue, P., & Schulberg, H. C. (2007). Psychotherapy and patient preferences for the  
11 treatment of major depression in primary care. In: M. J. Henri (Ed.), *Trends in*  
12 *depression research* (pp. 31 - 51). Hauppauge, NY: Nova Science Publishers,  
13 Inc.  
14  
15  
16  
17  
18 Raue, P., Schulberg, H., Heo, M., Klimstra, S., & Bruce, M. (2009). Patients'  
19 depression treatment preferences and initiation, adherence, and outcome: A  
20 randomized primary care study. *Psychiatric Services*, 60(3), 337-343. doi:  
21 10.1176/appi.ps.60.3.337  
22  
23  
24  
25  
26  
27 Rethink mental illness. (2012a). *Medication for mental illness*. Retrieved from:  
28 [http://www.rethink.org/living\\_with\\_mental\\_illness/treatment\\_and\\_therapy/me](http://www.rethink.org/living_with_mental_illness/treatment_and_therapy/medication/index.html)  
29 [dication/index.html](http://www.rethink.org/living_with_mental_illness/treatment_and_therapy/medication/index.html)  
30  
31  
32  
33  
34 Rethink mental illness (2012b). *Talking therapies factsheet*. Retrieved from:  
35 [http://www.rethink.org/living\\_with\\_mental\\_illness/treatment\\_and\\_therapy/tal](http://www.rethink.org/living_with_mental_illness/treatment_and_therapy/talking_treatments/index.html)  
36 [king\\_treatments/index.html](http://www.rethink.org/living_with_mental_illness/treatment_and_therapy/talking_treatments/index.html)  
37  
38  
39  
40  
41 Schneider, J., Foroushani, P. S., Grime, P., & Thornicroft, G. (2014). Acceptability  
42 of online self-help to people with depression: users' views of MoodGYM  
43 versus informational websites. *Journal of Medical Internet Research*, 16, e90.  
44 doi:10.2196/jmir.2871  
45  
46  
47  
48  
49 Sharp, D. M., Power, K. G., & Swanson, V. (2004). A comparison of the efficacy  
50 and acceptability of group versus individual cognitive behavioral therapy in  
51 the treatment of panic disorder and agoraphobia in primary care. *Clinical*  
52 *Psychology and Psychotherapy*, 11, 73-82. doi: 10.1002/cpp.393  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 Steidtmann, D., Manber, R., Arnow, B. A., Klein, D. N., Markowitz, J. C.,  
4  
5 Rothbaum, B. O., . . . Kocsis, J. H. (2012). Patient treatment preference as  
6  
7 predictor of response and attrition in treatment for chronic depression.  
8  
9 *Depression and Anxiety*, 29, 896-905. doi: 10.1002/da.21977  
10  
11 Stewart, G. (2010). *Understanding depression*. Retrieved from:  
12  
13 [http://www.mind.org.uk/help/diagnoses\\_and\\_conditions/depression](http://www.mind.org.uk/help/diagnoses_and_conditions/depression)  
14  
15 Sumner, K., Haddock, G., Hartley, Kilbride, M., McCusker, M., Pitt, L . . .  
16  
17 Barrowclough, C. (2014). Preferences for psychological therapy in psychosis:  
18  
19 trail participation, mode of treatment and willingness to be randomised.  
20  
21 *Journal of Mental Health*, 23, 67–71. doi: 10.3109/09638237.2013.841865  
22  
23 The Department of Health, (2013). *Making mental health services more*  
24  
25 *effective and accessible*. London: Department of Health.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1**

*Mean Levels of Treatment Acceptability and Side Effects by Treatment Condition for the Whole Sample and by Symptoms of Depression and Treatment Experience*

Treatment	Acceptability			Side Effects		
	<i>N</i>	Mean	<i>SD</i>	<i>N</i>	Mean	<i>SD</i>
Psychotherapy	96	0.59	0.87	96	0.10	0.94
No depression symptoms	23	0.77	0.89	23	-0.06	0.97
Depression symptoms	60	0.61	0.87	60	0.15	0.96
No experience of treatment	25	0.66	0.84	25	0.18	1.05
Experience of treatment	43	0.62	0.84	43	0.00	0.92
Guided self-help	99	0.33	0.95	99	-0.07	0.92
No depression symptoms	27	0.76	0.93	27	-0.18	0.84
Depression symptoms	60	0.17	0.91	60	-0.05	0.95
No experience of treatment	2	0.26	1.03	2	0.03	0.81
Experience of treatment	71	0.01	0.99	71	-0.31	1.01
Bibliotherapy	104	-0.27	1.03	104	0.37	1.01
No depression symptoms	25	-0.49	0.93	25	-0.46	1.13
Depression symptoms	70	-0.09	1.03	70	-0.40	0.99
No experience of treatment	13	-0.18	1.15	13	-0.05	1.04
Experience of treatment	61	0.26	0.96	61	-0.40	0.88
Internet-based self-help	111	-0.26	0.95	111	-0.16	0.97
No depression symptoms	29	-0.24	0.98	29	-0.16	0.89
Depression symptoms	71	-0.32	0.92	71	-0.16	1.04
No experience of treatment	7	-0.41	1.08	7	-0.02	1.18
Experience of treatment	63	-0.30	0.86	63	-0.09	0.74

Antidepressants	112	-0.27	0.92	112	0.65	0.87
No depression symptoms	17	-0.24	0.87	17	0.42	0.89
Depression symptoms	77	-0.21	0.95	77	0.61	0.87
No experience of treatment	25	-0.16	1.04	25	0.54	0.98
Experience of treatment	48	-0.34	0.89	48	0.71	0.91

---

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 2**

*Mean Preference Ratings for the Whole Sample*

Treatment	Mean	SD
Psychotherapy	1.77	1.25
Guided self-help	3.25	1.40
Antidepressants	3.76	1.86
Bibliotherapy	4.23	1.27
Internet-based self-help	4.68	1.32

**Supplementary materials 1: Description of depression**

Depression is a serious mental illness that is characterized by a number of unpleasant symptoms. These symptoms include a low depressed mood which is constantly present every day, a lack of interest and pleasure in activities that used to be enjoyed, changes in weight and/or appetite, insomnia or hypersomnia (excess sleep), psychomotor agitation (which is a series of unintentional and purposeless motions that stem from mental tension and anxiety including pacing around a room, wringing one's hands and other similar actions), tiredness and a lack of energy, feelings of worthlessness or inappropriate guilt, an impaired ability to concentrate or indecisiveness and, most seriously, recurrent thoughts of death. A person who is depressed does not necessarily suffer from all of these symptoms, but generally 5 or more symptoms are present for more than two weeks.

**Supplementary materials 2: Personal account of depression**

“I felt detached from the world around me. All emotions – love, affection, anger – were gone. Actually, I can't say I had no emotions, I did, but they all seemed desperately negative. Most involved fear. Fear that I would never escape the condition. I was so scared of being alone with my thoughts. At night, everything seemed so bleak. I couldn't concentrate on anything; I couldn't read or watch TV. I couldn't relax or unwind. Sleep seemed impossible – so many thoughts were racing through my mind. I would spend hours fantasizing about ways of killing myself. Everything to do with everyday life seemed like such hard work. I simply didn't have the energy to go to work, to see friends, to shop, cook or clean. It all seemed pointless! What was the point in eating, when I didn't even want to be alive?”

### Supplementary materials 3: Detailed Treatment descriptions

#### *TREATMENT DESCRIPTION 1: Psychological therapy*

##### *What is psychological therapy?*

Many people find that psychological therapies, sometimes referred to as 'talking treatments', 'counseling or psychotherapy', are useful for the treatment of depression. With the help of a trained professional, clients are encouraged to talk through the problems associated with their depression. Therapy can also allow you to explore the factors involved in making you ill in the first place and what keeps you from recovering. Therapy can also help you to deal with a specific traumatic experience such as bereavement. Therapy is usually undertaken through one-to-one sessions with a trained therapist or can be held as a group. There are also non-talking based psychological therapies such as art therapy. The number of sessions required depends on the type of therapy, the nature of the illness and what is available if the therapy is accessed on the NHS. Therapy usually last around 1-2 hours per session for an average of 16 weeks, although the duration of therapy required depends on the type of therapy sought and the extent of the depression.

##### *Are there different types of psychological therapies?*

Yes, there are a number of different types of psychological therapies available, including:

- Psychodynamic psychotherapy is one of the longest established therapies and is a term that covers therapy of an analytical nature. It is a form of in-depth therapy that focuses on the unconscious and past experiences and the effects they have on current behavior and thinking. The client is encouraged to talk about childhood relationships and experiences during the sessions. The aim of the therapy is to help

1  
2  
3 a person to understand how experiences in the past can unconsciously affect their  
4  
5 behavior and thinking.

6  
7 • Cognitive Behavioral Therapy (CBT) can help you to change how you think  
8  
9 ("cognitive") and what you do ("behavior)". These changes can help you to feel  
10  
11 better. CBT focuses on the problems and difficulties in the "here and now" instead of  
12  
13 addressing the causes of your distress or symptoms. CBT looks for ways to improve  
14  
15 your state of mind by exploring how you currently think about yourself, the world and  
16  
17 other people and how that affects your reaction to situations.

18  
19 • Counseling is sometimes a term used generally to describe all types of therapy,  
20  
21 however it also refers to a specific type of therapy. Counseling allows a client to talk  
22  
23 to a trained counselor about a specific problem with the aim of helping the client to  
24  
25 understand their problem more clearly and come up with their own solutions to deal  
26  
27 with these difficulties. The role of the counselor is to listen and be non judgmental  
28  
29 towards the client, providing them with a safe and confidential environment to  
30  
31 discuss their difficulties.

32  
33  
34  
35  
36 *How do I access these treatments?*

37  
38 The first step is to discuss how you are feeling with your GP. He or she will then be  
39  
40 able to refer you to the most appropriate service. You may express a preference for  
41  
42 the type of talking therapy that you would like to receive. This will be taken into  
43  
44 account when a referral is made, however there may be times when the talking  
45  
46 therapy that you would like is not available or not the right one for the problems you  
47  
48 are experiencing.

49  
50  
51  
52  
53  
54 *TREATMENT DESCRIPTION 2: Internet administered self-help*

55  
56 *What is Internet administered self-help?*  
57  
58  
59  
60

1  
2  
3 In 2010 60% of the UK adult population accessed the Internet every day and many  
4  
5 of us are using the Internet as a means of understanding and treating our own  
6  
7 health. The NHS Direct website is one of the major sources of health information  
8  
9 online and it attracts more than 1.5 million visits each month.  
10

11 As a result of so many of us using the internet as a source of health information and  
12  
13 the problems the NHS face in treating depression in convention ways the internet is  
14  
15 being used to provide self-help treatments for depression.  
16  
17

18 *Are there different types of Internet administered self-help treatments?*  
19

20 Yes, there are many different websites providing self-help for depression, a few are  
21  
22 outlined below:  
23

24  
25 • Living life to the full ([www.livinglifetothefull.com](http://www.livinglifetothefull.com)) is an online course that aims to  
26  
27 provide access to high quality, practical and user-friendly training in practical  
28  
29 approaches you can use in your own life. The course content teaches key  
30  
31 knowledge in how to tackle and respond to issues/demands that we meet in our  
32  
33 lives. The course contains modules on issues such as understanding why we feel as  
34  
35 we do, practical problem solving skills, anxiety control, relaxation, overcoming  
36  
37 reduced activity, helpful and unhelpful behaviors, noticing and changing unhelpful  
38  
39 thoughts, healthy living (e.g. sleep, diet and exercise) and staying well. The course  
40  
41 delivers sound, text and video clips as well as short handouts and longer detailed  
42  
43 practical workbooks that develop and build upon the course. The website also offers  
44  
45 moderated discussion forums that allow course users to swap ideas, information and  
46  
47 provide support. Nothing on the website is compulsory - you are in control. After  
48  
49 completing the initial registration process and Session 1, you can choose to  
50  
51 complete as many or as few of the self-help life skills modules as you wish.  
52  
53

54  
55 • MoodGym (<http://www.moodgym.anu.edu.au/>) is an online interactive program  
56  
57  
58  
59  
60

1  
2  
3 designed to help you identify whether you are having problems with emotions like  
4  
5 anxiety and depression and learn skills that can help you cope with these emotions.  
6  
7 MoodGYM is based on two programs that are successful in preventing and treating  
8  
9 depression and anxiety. These are: Cognitive Behavior Therapy and Interpersonal  
10  
11 Therapy. MoodGym consists of interactive modules that are delivered to you in a  
12  
13 specific order. The modules are: feelings, thoughts, de-stressing, and relationships.  
14  
15 At the end of each module you can apply the material to your own circumstances  
16  
17 through a series of activities. As you move through the program, you are presented  
18  
19 with information, animated demonstrations, quizzes and “homework” exercises. Your  
20  
21 answers to the exercises are recorded in your own workbook, which keeps track of  
22  
23 your progress. At the end there is opportunity for you to view this progress and your  
24  
25 results.  
26  
27  
28

29  
30 • The Centre for Clinical Interventions (<http://www.cci.health.wa.gov.au/>) website  
31  
32 offers free downloadable workbooks that provide self-help guidance for depression.  
33  
34 The depression workbook contains nine modules that cover everything from the  
35  
36 symptoms and causes of depression, right through to how to maintain the progress  
37  
38 made once you have completed the modules. Specific self-help activities include,  
39  
40 behavioral strategies to increase activity levels, how to challenge automatic thoughts  
41  
42 that negatively impact upon feelings and finally core beliefs are examined and advice  
43  
44 & techniques are given to help the patient confront the core beliefs they hold that  
45  
46 may be leading to depression (for example: I’m unlovable).  
47  
48

49  
50 *How do I access these treatments?*

51  
52 Treatment is accessed via the Internet and thus requires a computer or other device  
53  
54 that is able to go online.  
55  
56  
57  
58  
59  
60

### *TREATMENT DESCRIPTION 3: Bibliotherapy*

#### *What is bibliotherapy?*

Bibliotherapy is basically books to help people solve problems. The use of literature can be used to help people cope with emotional problems, mental illnesses, or changes that have occurred in their lives. As a result of the change, it promotes personality and developmental growth. Bibliotherapy provides a sensitive way for a practitioner to guide reading to help an individual understand themselves and the environment, learn from others, and possibly find solutions to their problems.

#### *Are there different types of bibliotherapy?*

Yes, there are numerous amounts of self-help books for depression, three of the best-regarded are:

- *Overcoming Depression: A self-help guide using cognitive behavioral techniques* by Paul Gilbert. This book, written by an expert in the field of depression, uses cognitive-behavioral principles to provide a structured self-help treatment. Part One of the book helps the reader to gain an understanding of depression and its causes. Part Two provides the reader with guidance on managing their depression. It focuses on increasing activity levels, breaking problems into manageable steps and looks briefly at sleep management, diet, exercise and alcohol/drugs. It moves on to an exploration of the role of thoughts and feelings, and teaches the reader to identify and challenge thoughts that may contribute to low mood, giving particular focus to overcoming self-blaming and self-critical thoughts. The importance of developing inner compassion is highlighted, as a means of reducing signals to the body to produce a depressed response. Part Three looks at particular problems associated with depression, such as the need for approval, guilt, shame, anger, assertiveness, disappointment and perfectionism.



1  
2  
3 • Overcoming Depression: A five areas approach by Chris Williams. This book  
4 contains is a series of structured self-help workbooks for use by people experiencing  
5 depression. The course allows access to the cognitive behavior therapy (CBT)  
6 approach to treatment. Part one of the book helps the reader to understand why they  
7 feel as you do. Part two of the book is all about making changes to the readers life  
8 that will help them to overcome their depression. Topics covered include, making  
9 changes to do with people and events (such as practical problem solving, being  
10 assertive and building relationships with family/friends) making changes to behaviors  
11 and activity levels (such as using exercise to boost positive feelings and recognizing  
12 helpful and unhelpful behaviors) making changes to negative and upsetting thinking,  
13 making changes to things that affect bodily well-being (such as overcoming sleep  
14 problems) and finally making changes for the future, which is all about planning for a  
15 future that is healthy and free of depression.  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

32 • Feeling good: The New Mood Therapy by David Burns. This book describes an  
33 approach to mood modification that has a self-help component for those who suffer  
34 from depression. Research data is cited that suggests people can learn to control  
35 their mood swings and self-defeating behaviors with principles and techniques that  
36 the book describes in detail. The author describes dysfunctional thinking that can  
37 lead a person into a low mood or prevent them from healing. His theory involves  
38 helping the reader develop awareness of the connection between thoughts, feelings  
39 and behaviors and then taking active responsibility for identifying automatic thoughts,  
40 their distortions and replacing them with a rational response. Throughout the  
41 chapters, the author teaches techniques for building self-esteem, handling criticism  
42 anger and guilt as well as depression. He also focuses on prevention and personal  
43 growth work, skills for coping with stresses of daily living and finally offers advice for  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 handling suicide.

4  
5 *How do I access these treatments?*

6  
7 The reader can purchase self-help books or they are also readily available from  
8  
9  
10 libraries across the country.

11  
12  
13  
14 *TREATMENT DESCRIPTION 4: Guided self-help*

15  
16 *What is guided self-help*

17  
18 Guided self-help is provided when a therapist introduces a patient to a range of self-  
19  
20 help tools that they can work through independently, to gain a better understanding  
21  
22 of the issues that are affecting them. Guided self-help differs from pure self-help as  
23  
24 there is always a trained therapist on hand to provide support to the patient whilst  
25  
26 they are using the tools. In the UK, guided self-help for depression is provided by  
27  
28 therapists, who work within the NHS.  
29

30  
31  
32 *Are there different types of guided self-help?*

33  
34 Yes, there are a number of guided self-help interventions used by therapists, the  
35  
36 most commonly used are:

37  
38 • The most basic method by which guided self-help is provided, is for the therapist to  
39  
40 recommend resources for the patient to use or activities for the patient to do.

41  
42 Resources can include books, interactive websites, dvd's, cd-roms etc. All of which  
43  
44 provide information and self-help techniques for the patient to use. Activities that the  
45  
46 therapist may recommend include, changing your diet, exercising, socializing etc.

47  
48 The therapist is on hand to provide support to the patient but they are mainly there to  
49  
50 guide the patient on their self-help journey by providing the tools they need to help  
51  
52 themselves recover.  
53

54  
55  
56 • Cognitive restructuring is a guided self-help intervention that seeks to change  
57  
58  
59  
60

1  
2  
3 unhelpful thoughts by identifying, examining and challenging them. A major  
4  
5 component of depression is negative thinking. Most of the negative thoughts we  
6  
7 have when depressed are automatic and unhelpful, although they feel believable and  
8  
9 real at the time. Patients keep a diary in which they identify their unhelpful thoughts.  
10  
11 Patients then work with their therapist to identify which of their unhelpful thoughts is  
12  
13 responsible for negative emotions. Once the patient has identified which unhelpful  
14  
15 thought they wish to work on, they must examine the evidence for and against the  
16  
17 unhelpful thought. This can be difficult, especially when trying to come up with  
18  
19 evidence against but the therapist will help the patient by asking them questions like;  
20  
21 if you had a friend who had this thought what would you say to them to disprove it?  
22  
23 Once patients have gathered enough evidence for and against the unhelpful thought  
24  
25 they are able to revise their original thoughts and hopefully come up with an  
26  
27 alternate thought that is less damaging to their emotional state.  
28  
29

30  
31 • Problem solving is another guided self-help intervention that helps depressed  
32  
33 patients when their problems initially seem too big to solve. It helps patients consider  
34  
35 what solutions may exist if they take a systematic and step by step approach to  
36  
37 solving their problem. Patients begin by outlining the exact problem and breaking it  
38  
39 down into components. Potential solutions of all kinds, even apparently ridiculous  
40  
41 ones, are noted down and then analyzed in terms of their strengths and  
42  
43 weaknesses. The patient then, with the help of their therapist, can choose a solution  
44  
45 and begin to make a plan of how they will implement this solution. Patients will be  
46  
47 asked to keep a diary as they begin to put the plan into action. The patient will  
48  
49 continue to meet with their therapist, who will monitor progress and be on hand to  
50  
51 choose a new solution should the initial plan fail to solve the problem.  
52  
53  
54

55  
56 *How do I access these treatments?*  
57  
58  
59  
60

1  
2  
3 The first step is to discuss how you are feeling with your GP. He or she will then be  
4  
5 able to refer you to a therapist who specializes in providing guided self-help.  
6  
7

8  
9  
10 *TREATMENT DESCRIPTION 5: Antidepressant medication*

11 *What are antidepressants?*

12  
13 Antidepressant medication is a treatment for depression. Antidepressant medications  
14 are taken in tablet form and they work by increasing the amount of naturally  
15 occurring chemical messengers in the brain. The brain has many naturally occurring  
16 chemical messengers. Two of these are called serotonin and noradrenaline. Both  
17 are important in the areas of the brain that control or regulate mood and thinking. It is  
18 known that these two chemical messengers are not as effective or active as normal  
19 in the brain of someone who is depressed. Antidepressants increase the amount of  
20 these chemical messengers, thus helping to correct the lack of action of the  
21 messengers and improving mood.  
22  
23

24 *Are there different types of antidepressant medications?*

25  
26 Yes, there are a number of different types of antidepressants that may be prescribed  
27 to you, including:  
28  
29

30 • Tricyclic antidepressants were among the earliest antidepressants developed.  
31  
32 Tricyclic antidepressants are effective, but they have generally been replaced by  
33 other antidepressants that cause fewer side effects. Other antidepressants are  
34 prescribed more often, but Tricyclic antidepressants are still a good option for some  
35 people. In certain cases, Tricyclic antidepressants relieve depression when other  
36 treatments have failed. Tricyclic antidepressants ease depression by affecting  
37 chemical messengers in the brain (neurotransmitters). These naturally occurring  
38 brain chemicals are used to communicate between brain cells. Tricyclic  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 antidepressants block the absorption (reuptake) of the neurotransmitters serotonin  
4  
5 and norepinephrine making more of these chemicals available in the brain. This  
6  
7 seems to help brain cells send and receive messages, which in turn boosts mood.

8  
9  
10 • Selective serotonin re-uptake inhibitors (SSRIs) are the most commonly prescribed  
11  
12 antidepressants. SSRIs can ease symptoms of moderate to severe depression, are  
13  
14 relatively safe and generally cause fewer side effects than other types of  
15  
16 antidepressants. SSRIs ease depression by affecting chemical messengers  
17  
18 (neurotransmitters) used to communicate between brain cells. SSRIs block the  
19  
20 reabsorption (reuptake) of the neurotransmitter serotonin in the brain. Changing the  
21  
22 balance of serotonin seems to help brain cells send and receive chemical messages,  
23  
24 which in turn boosts mood. SSRIs are called selective because they seem to  
25  
26 primarily affect serotonin, not other neurotransmitters.  
27  
28

29  
30 • Mono-amine oxidase inhibitors (MAOIs) were the first type of antidepressant  
31  
32 developed. MAOIs are effective, but have generally been replaced by other  
33  
34 antidepressants that are safer and cause fewer side effects. MAOIs generally require  
35  
36 diet restrictions because they can cause dangerously high blood pressure when  
37  
38 taken with certain foods. In spite of side effects, MAOIs are still a good option for  
39  
40 some people. In certain cases, MAOIs relieve depression when other treatments  
41  
42 have failed. The enzyme monoamine oxidase is involved in removing the  
43  
44 neurotransmitters norepinephrine, serotonin and dopamine from the brain. MAOIs  
45  
46 prevent this from happening, which makes more of these brain chemicals available.  
47  
48 This is thought to boost mood by improving brain cell communication.  
49  
50

51  
52 *How do I access these treatments?*

53  
54 Antidepressant medications are only available by prescription from a medical  
55  
56 profession such as your GP, or Psychiatrist. You will be monitored throughout the  
57  
58  
59  
60

1  
2  
3 course of your treatment as often the dose or medication will need to be altered to  
4  
5 treat your depression or ease side effects.  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Supplementary materials 5: Brief treatment descriptions

### *Psychological therapy*

With the help of a trained therapist, clients are encouraged to talk through the problems associated with their depression. Therapy can also allow the client to explore the factors involved in making them ill in the first place and what keeps them from recovering. Therapy sessions usually last between 1-2 hours for an average of 16 weeks, although this varies depending on the severity of the depression.

### *Antidepressant medications*

Are taken in tablet form and work by increasing the amount of naturally occurring chemical messengers in the brain. It is known that chemical messengers are not as effective or active as normal in the brain of someone who is depressed.

Antidepressants increase the amount of these chemical messengers, thus helping to correct the lack of action of the messengers and improving mood.

### *Bibliotherapy*

The definition of bibliotherapy is simplified to basically books that help people solve problems. The use of literature can be used to help people cope with emotional problems, mental illnesses, or changes that have occurred in their lives.

Bibliotherapy provides a sensitive way for a practitioner to guide reading to help an individual understand themselves and the environment, learn from others, and possibly find solutions to their problems.

### *Internet administered self-help*

1  
2  
3 The Internet is a vast source of knowledge and many of use the Internet as a source  
4  
5 of health information. As a result of this, the Internet has been used as a way of  
6  
7 delivering self-help treatments for depression. Many websites provide free  
8  
9 information on depression and access to self-help techniques. These websites often  
10  
11 have downloadable workbooks, tutorials and advice on treating depression.  
12  
13

### 14 15 16 *Guided self-help*

17  
18 In the UK, therapists who work within the NHS often provide guided self-help for  
19  
20 patients suffering from depression. Guided self-help is often the first step of care  
21  
22 provided to patients who have decided to seek treatment for their depression.  
23  
24 Therapists provide the patient with self-help materials (such as books, interactive  
25  
26 CD-ROMs, worksheets and online resources), which they work though  
27  
28 independently. The therapist is on hand to offer guidance and support to the patient,  
29  
30 but the majority of the work is down to the patient.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



## Supplementary materials 6: Description of pilot research

A pilot study was conducted to develop materials to be used in the main study. Specifically, we wanted to ensure that the personal account of depression would enable the participants to understand how it feels to be depressed. In addition, we wanted to ensure that the treatment descriptions were comparable in terms of readability, understandability, and perceived bias. Finally, we wanted to ensure that the brief treatment descriptions were reflective of their full-length counterparts.

### *Participants and Procedure*

The pilot study recruited 54 postgraduate students studying in the Psychology department at the University of Sheffield. The sample was made up of 42 females (77.8%) and 12 males (22.2%). Participants were emailed a link to an online questionnaire, which took around 20 minutes to complete. Ethical approval for the preliminary study was granted alongside the main study from the University of Sheffield.

Participants read two different accounts of depression (one referring to 'Helen', the other referring to 'David') in a counterbalanced order. David's account is presented above (see Supplementary materials 2), while Helen's account was as follows:

"I was tired all the time and not normal fatigue but bone-weary exhaustion. I slept as if I had been knocked unconscious and struggle to wake in the morning, dragging my leaden limbs through the day. I was always cold; my fingers white and numb even during the summer, when I kept a heater going full blast. If I got too cold, I would find it almost impossible to get warm again and have to resort to lying in a bath with the hot water running. My arms and legs ached constantly, so painfully that, at times, I take painkillers every four hours. I felt constantly low and depressed;

1  
2  
3 I couldn't throw off the mood I have developed, a low feeling that seems always to  
4 envelop me like a cold, grey blanket. The crying grows worse and by now I am  
5 scarcely sleeping. I have started to cry in unexpected places, at inconvenient times.  
6  
7 One day, I cried at work. I was mortified. I never cry at work. I decided that I must be  
8 exhausted, and take a week off. It is the end of June. I spent the days walking  
9 around the streets, wearing dark glasses, with tears streaming down my face. I  
10 walked for hours every day".  
11  
12  
13  
14  
15  
16  
17

18 Participants were asked seven questions about each of the accounts of  
19 depression (see Table 1) and asked which of the two accounts they felt best  
20 captured the essence of how it feels to be depressed. Participants were then asked  
21 to read the treatment descriptions before being asked to answer four questions  
22 about each of the longer treatment descriptions (see Table 2) and five questions  
23 about each of the brief treatment descriptions (see Table 3).  
24  
25  
26  
27  
28  
29  
30  
31

### 32 *Results*

33  
34 Table 1 shows that participants rated each of the accounts of depression  
35 relatively positively (in the sense that they were easy to read, brought to life the  
36 experience of depression and so on). Scores were summed to create an overall  
37 score for Helen's account ( $\alpha = 0.86$ ) and David's account ( $\alpha = 0.85$ ). In order to  
38 establish whether there were significant differences between the overall mean  
39 scores for the two accounts of depression, a repeated measures ANOVA was used.  
40  
41 Significant differences were observed between the two personal accounts,  $F(1,48) =$   
42 8.47,  $p < .01$ , with participants showing a preference for David's account ( $M_s = 5.52$   
43 and 5.01,  $SDs = 1.04$  and 1.14, for David and Helen's account, respectively). When  
44 asked to choose a personal account of depression that best captured the essence of  
45 how it feels to be depressed, the majority of participants voted for David's account  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 (32 participants or 59.3%) rather than Helen's account (17 or 31.5%). Five  
4  
5 participants (9.3%) did not vote. We therefore used David's account of depression in  
6  
7 the main study.  
8

9  
10 Table 2 shows that the participants rated the detailed treatment descriptions  
11  
12 as easy to read and understand, and allowing them to imagine what receiving the  
13  
14 treatment would be like (i.e., all of the means were below the midpoint of the scale).  
15  
16 Participants felt that all of the treatment descriptions were slightly biased (i.e., all  
17  
18 means were above the midpoint of the scale). In order to investigate whether there  
19  
20 were significant differences between ratings of the different treatments, a series of  
21  
22 repeated measures ANOVAs were performed, using Bonferroni adjustment to  
23  
24 correct for multiple tests ( $p = 0.01$ ). No significant differences were observed  
25  
26 between the five treatment descriptions for any of the four questions ( $p > 0.05$  in all  
27  
28 cases).  
29  
30

31  
32 Table 3 shows that the participants rated the brief treatment descriptions as  
33  
34 easy to read, easy to understand and allowed them to imagine what receiving the  
35  
36 treatment would be like (i.e., all means were below the midpoint of the scale). As  
37  
38 with the detailed treatment descriptions, participants tended to feel that the brief  
39  
40 treatment descriptions were slightly biased (i.e., all means were above the midpoint  
41  
42 of the scale). Participants did, however, agree that the brief treatment descriptions  
43  
44 captured the essence of the longer treatment descriptions (i.e., all means were  
45  
46 above the midpoint of the scale). In order to investigate whether there were  
47  
48 significant differences in ratings of the five brief treatment descriptions, a series of  
49  
50 repeated measures ANOVAs were performed, with Bonferroni adjustment as before.  
51  
52 Significant differences were observed between the five brief treatment descriptions  
53  
54 for question 3 only ("How easy was it to imagine what receiving this type of treatment  
55  
56  
57  
58  
59  
60

1  
2  
3 would be like"?)  $F(4, 34) = 3.56, p < 0.01$  (all other  $F_s < 2.53, p > .06$ . Post hoc  
4  
5 analysis showed that the brief treatment description for guided self-help ( $M = 3.11,$   
6  
7  $SD = 1.56$ ) enabled participants to more easily imagine what the treatment would be  
8  
9 like than the brief treatment description for Internet self-help ( $M = 2.37, SD = 1.10$ ).  
10

### 11 *Conclusion*

12  
13  
14 The pilot research was able to fulfill its three main aims, which were to  
15  
16 establish (a) which personal account of depression best captured the essence of  
17  
18 how it feels to be depressed, (b) whether the detailed and brief treatment  
19  
20 descriptions were comparable and (c) whether the brief treatment descriptions  
21  
22 accurately portrayed their longer counterparts. In terms of the personal accounts of  
23  
24 depression, participants favored David's account and so that is the one that was  
25  
26 used in the main study. In terms of the full length and brief treatment descriptions  
27  
28 only one difference was noted, with participants rating the brief treatment description  
29  
30 for guided self-help as more easily enabling them imagine what the treatment would  
31  
32 be like than the brief treatment description for Internet self-help.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 1**

*Descriptive Statistics for the 7 Questions assessing the Personal Accounts of Depression*

Question	Account 1 (Helen)	Account 2 (David)
	Mean ( <i>SD</i> )	Mean ( <i>SD</i> )
The account was easy to read	4.90 (1.81)	5.57 (1.91)
The account brought to life the experience of depression	5.25 (1.33)	5.62 (1.33)
The account fitted my understanding of how it might feel to be depressed	5.22 (1.38)	5.78 (1.14)
The account made me realise the severity of depression	4.96 (1.48)	5.60 (1.36)
The account made it easy for me to imagine that I was depressed	4.35 (1.81)	4.61 (1.80)
The account highlighted a number of the symptoms of depression	5.02 (1.59)	5.67 (1.05)
The account made it easy for me to imagine how depression would affect my day-to-day life	5.13 (1.50)	5.56 (1.29)

*Note.* All questions were answered using a 7-point Likert scale where 1 = strongly disagree and 7 = strongly agree

**Table 2***Descriptive Statistics for the Five Treatment Descriptions*

Question	Psychotherapy Antidepressants	Bibliotherapy	Internet	Guided
	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)
1. How easy was the treatment description to read?	2.02 (1.16)	2.76 (1.69)	2.57 (1.36)	2.38 (1.23)
2. How easy was it to understand what this treatment involved?	2.42 (1.22)	2.66 (1.36)	2.61 (1.15)	2.30 (1.03)
3. How easy was it to imagine what receiving this type of treatment would be like?	2.64 (1.38)	3.30 (1.62)	2.98 (1.34)	3.02 (1.37)
4. The treatment description was neutral and unbiased	3.89 (2.13)	3.93 (1.99)	3.83 (1.90)	3.63 (1.93)

Note. All questions were answered using 7-point Likert scales. For questions 1 – 3, 1 = extremely easy and 7 = extremely difficult, while for question 4, 1 = strongly disagree and 7 = strongly agree

**Table 3***Descriptive Statistics for the Five Brief Treatment Descriptions*

Question	Psychotherapy Antidepressants	Bibliotherapy	Internet	Guided
	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)
1. How easy was the description to read?	1.66 (1.30)	1.74 (1.25)	2.03 (1.53)	1.71 (1.35)
2. How easy was it to understand what this treatment involved?	2.24 (1.26)	2.18 (1.52)	2.53 (1.45)	2.03 (1.33)
3. How easy was it to imagine what receiving this type of treatment would be like?	2.55 (1.35)	2.79 (1.56)	3.11 (1.49)	2.37 (1.10)
4. The description was neutral and unbiased	3.74 (2.06)	4.05 (2.05)	4.03 (2.17)	3.61 (2.52)
5. The description captured the essence of the longer description	4.76 (1.91)	4.75 (1.70)	4.76 (1.76)	5.46 (1.48)

*Note.* The response scales for questions 1 – 4 were as above and for question 5, 1 = strongly disagree and 7 = strongly agree