

Compassion in healthcare: a concept analysis

TAYLOR, Amy <<http://orcid.org/0000-0002-7720-6651>>, HODGSON, Denyse, GEE, Melanie <<http://orcid.org/0000-0001-9149-4314>> and COLLINS, Karen <<http://orcid.org/0000-0002-4317-142X>>

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/16040/>

This document is the author deposited version.

Published version

TAYLOR, Amy, HODGSON, Denyse, GEE, Melanie and COLLINS, Karen (2017). Compassion in healthcare: a concept analysis. *Journal of Radiotherapy in Practice*, 16 (4), 350-360. [Article]

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

Compassion in healthcare: a concept analysis

A. Taylor (MSc)^{1, 2}, D. Hodgson (EdD)², M. Gee (MSc)², K. Collins (Phd)²

¹. Weston Park Hospital, Sheffield. S10 2SJ

². Sheffield Hallam University, Sheffield. S10 2BP.

Correspondence: Amy Taylor, Weston Park Hospital, Sheffield. S10 2SJ
amy.taylor@sth.nhs.uk

There are no conflicts of interests

Abstract

Background

Compassion and compassionate care are central to radiographers' professional policy and practice and are congruent with the core values of the National Health Service (NHS) Constitution. The term compassion however is over-used, ambiguous and vague. This work sought to explore and provide contextual understanding to the term, compassion in healthcare.

Method

Walker and Avant's Eight-step model was used as the framework for the concept analysis. Data collection utilised a number of resources including online databases: Medline, CINAHL complete, Scopus, PubMed, PsycINFO, Science Direct, Cochrane and DARE; dictionaries, social media, internet sources, books and doctoral theses. 296 resources were included in the review.

Results

The concept analysis distinguishes the defining characteristics of compassion within a healthcare context, allowing for associated meanings and behaviours to be outlined aiding understanding of compassion. Compassion in healthcare requires five defining attributes to be present: Recognition, Connection, Altruistic desire, Humanistic response and Action.

Conclusion

The findings identify the complexity of the term and subjective nature in which it is displayed and in turn perceived. The concept analysis forms the basis of further research aiming to develop a healthcare explicit definition of compassion within healthcare, specifically cancer care and radiography practices. Lucidity will enhance understanding; facilitating active engagement and implementation into practice.

Introduction

The necessity to develop and subsequently utilise caring and compassionate behaviour within the healthcare workforce is central to radiographers' professional policy and practice [1] and in the United Kingdom (UK) is congruent with the core values of the National Health Service (NHS) Constitution [2]. Increases in patient complaints authenticated by Care Quality Commission reports of substandard care have raised concerns of malpractices, poor care and neglect. Compassion is a key recommendation of health legislation [3-6], further amplified following a number of high profile incidents in the UK where inadequate care and compassion reduced the quality of life of patients, with some instances resulting in death [7-11]. Numerous recommendations were formulated following these transgressions, the most pertinent being that the NHS needs to put the patient first, ensuring they receive services from caring, compassionate and committed staff [7].

The NHS has been criticised for promoting a culture amongst its staff which is more focused on doing the system's business rather than the patients in its care [7]. Tackling this compassion deficit has been placed at the centre of government initiatives [3,6, 7,12-14]. Despite reactive policy there continues to be a surfeit of failings which are all too often considered to be the inevitable [15]. This continuance raises questions regarding the ability of NHS trusts, service delivery managers and individual employees to understand, interpret and implement policy recommendations regarding compassionate practice at a local level. Thus, subsequently hindering their ability to design and develop infrastructures which support the implementation of policy.

The policies themselves provide limited definition and explanation of compassion and its meaning, leading some to consider whether the complex construct of compassion is fully understood [16,17] and, if not, can it therefore be successfully promoted in practice [18-21]. Concepts, such as 'compassion', are subjective, and are traditionally argued to be shaped and influenced by the environment and the objects in which they are situated [22]. Exploration and a contextual understanding was sought to refine the ambiguous, vague and often over-used term, compassion in healthcare.

Method

Walker and Avant's model [23] was used as the framework for this concept analysis. Their 8-step process (Table 1) was modified and simplified from Wilson's [24] original model and provides a more pragmatic and procedural format than other models which often place more emphasis on the philosophical and conceptual approaches [25, 26]. This pragmatic approach was favoured as the work focuses on healthcare within a physical environment and therefore requires understanding of the practical components of compassion. Analysis of the literature provides understanding of the defining attributes of the concept [23], allowing for the synthesis of existing views, enabling it to be distinguished from other similar and associated concepts thus resolving gaps or inconsistencies in the knowledge base of the discipline [27].

Step	Description	Method
One: Select a concept	Identification of an area of interest.	Compassion is the focus of the lead authors PhD and selected in accordance.
Two: Determine the purpose of the analysis	Outline why the analysis is being undertaken.	Compassion is the focus of the lead authors PhD and selected in accordance.
Three: Identify all uses of the concept	Identify all uses of the term both within and external to the chosen context, allowing for validation of defining attributes.	Resources incorporated into the concept analysis were reviewed.
Four: Determine the defining attributes	Defining attributes are characteristics of the concept identified repeatedly through the literature. These enable the classification of the occurrence of a specific phenomenon (concept) as differentiated from similar concepts.	Resources incorporated into the concept analysis were reviewed.
Five: Identify a model case	A model case demonstrates all the defining attributes, providing insight into the internal structure of the concept, allowing clarification of its meaning and the context.	A model case was created by the authors using a clinical example within a healthcare context. Behaviours, actions and responses associated with compassion identified in the literature were incorporated into the case to

		provide a practical example of compassion, where all five attributes of compassion were being displayed.
Six: Identify borderline*, related cases and contrary cases	Borderline and related cases contain many of the defining attributes of compassion, but not all of them. A contrary case demonstrates compassion is clearly not being displayed	Additional cases were created by the authors using clinical examples within a healthcare context. Behaviours, actions and responses associated with compassion and non-compassion identified in the literature were incorporated into the case to provide practical examples. The authors omitted behaviours and responses which represented three of the five defining attributes in the related case and all five within the contrary case.
Seven: Identify antecedents and consequences	Antecedents are events or incidents that must occur prior to the occurrence of the concept. Consequences are the events or incidents that occur as a result of occurrence of the concept.	Resources incorporated into the concept analysis were reviewed.
Eight: Define empirical referents	Empirical referents are categories of actual phenomena whose presence demonstrate the defining attributes and help determine the occurrence of the concept.	Resources incorporated into the concept analysis were reviewed.

Table 1: Eight steps of Walker and Avant's concept analysis [23] *Borderline case not included within this article

Data collection utilised a number of resources (Table 2).

Source category	Details
Dictionaries	Online English-language Nursing, medical, and psychology
Social Media	Twitter
Databases and Journals	Medline, CINAHL complete, Scopus, PubMed, PsycINFO, Science Direct, Cochrane and DARE Journal of Radiotherapy in Practice (JRP), Patient Education & Counselling, Journal of Medical Imaging and Radiation Oncology and the Journal of Medical Internet Research (JMIR)
Internet	Google and Google Scholar Websites including the Department of Health (DoH-GOV),

	INVOLVE, The Kings Fund
Books	Book chapters on human values and characteristics
Doctoral thesis	http://ethos.bl.uk , Database of over 40,000 theses

Table 2: Resources categories used in concept analysis

The databases were searched from October 2015 to June 2016 using a comprehensive search strategy and inclusion/exclusion criteria were applied to ensure only papers relevant to healthcare and with a focus on compassion were included (Appendix 1). The Twitter search used #compassion to capture current discussion over an 11 day period, providing a broader meaning to the term outside the published healthcare context by collating accounts and posts from individuals, newspapers, charities and organisations. Figure 1 displays the stages of the review process.

Relevant results from each literature source category were strategically examined; key points were documented, collated and compared across the categories allowing for the development of key themes and generation of a thematic map. The author produced a Wordle (Figure 2) documenting key words and phrases used in conjunction with the concept of compassion. This contributed to the development of behaviours and attributes associated with compassion, forming the basis of the antecedents. Saturation was reached once the literature sources generated no new themes.



Figure 2: Wordle denoting key terms from the literature associated with compassion.

Findings

Step 1: Concept

Compassion in healthcare

Step 2: Purpose of the analysis

To understand what compassion is and how it is displayed within a healthcare context by health professionals.

Step 3: Uses of the concept

Eleven terms about the concept of compassion were identified within the literature. Six of them were included to form this concept analysis, being considered by the authors to reflect accepted use of the concept within healthcare: Compassion, Compassionate, Compassionate-care, Compassion satisfaction, Compassionate practice and compassionate caring. The other five terms were excluded from this concept analysis. Compassion-fatigue and Self-Compassion (inner-compassion) are heavily discussed in relation to compassion and compassionate care, but their focus is on the self rather than towards another. The terms Compassionate-love, Compassionate-leave, and Compassionate-use, although possessing a focus on another rather than the self, represent the love of a partner or spouse, a policy of authorised leave from work or the prescription of non-licensed drugs respectively.

Step 4: Defining attributes:

The literature examined identified compassion in healthcare as comprising of five defining attributes (Figure 3 and Table 3). Although these occur sequentially and each attribute needs to occur, the individual who is to display compassion may need to move between the attributes depending on the situation.




Figure 3: Five sequential attributes of compassion and compassionate response

Attribute	Description
(1) Recognition	Cognitive recognition of another's adverse circumstances, physical, psychological or emotional wellbeing
(2) Connection	Personal connection with another based on automatic, authentic and genuine thought
(3) Altruistic desire	Altruistic desire to aid another
(4) Humanistic response	Humanistic, person-to-person, understanding of what it is to be human
(5) Action	Undertaking of an act or responsive behaviour

Table 3: Defining attributes of compassion.

Step 5: Model case

Kate was deteriorating; she had been diagnosed with advanced cancer and had been admitted to hospital, very ill with a serious infection.

Kate was distressed and felt very poorly. The nurse who was looking after her could see (1) how vulnerable and frightened Kate was, so he gently placed his hand on her arm, knelt down beside her bed (2) and said to Kate we're going to look after you (3). Kate relayed her experiences to her friends and followers on twitter, tweeting about a number of similar events over the course of her stay as an in-patient. She recalled her nurse ran her bath it was a hot, deep, bubbly bath, just the way she liked it (4, 5). On another evening she mentioned to the nurse that she had been struggling to sleep to which the nurse had replied don't sit awake in the night, just buzz me and I'll come and sit with you until you fall back to sleep (4,5).

Stage 6: Additional cases

Related case

Clive a therapy radiographer for 20 years, was chatting with a 19 year old patient called John who was about to start radical radiotherapy for testicular cancer. John was telling him that he'd already had the tumour and his left testicle removed and how he'd had to make the difficult decision as to whether to have his sperm frozen or not before he started radiotherapy. John told Clive that it has sparked him and his girlfriend to think about their futures together and decided to get married. He confessed he was worried and kept on stressing over whether the sperm banking process may not have worked and that he was worried how his fiancé would take the news if this was the case. Clive sat and listened to the young man's fears, ensuring that he knew he was there for him to talk to (2). Clive fully understood how John was feeling (1), not only had he treated many patients like John over his professional career, but when he was 22 he had received the same diagnosis so knew what John must have been going through.

Contrary case

Frank was attending his weekly review with the specialist radiographer; he had a review every week as part of his radiotherapy treatment for prostate cancer. He'd been worried for some time about problems he was experiencing with his erection and had wanted to ask someone for a while but he had felt too embarrassed. His wife had told him that he must speak to someone at his next appointment as it was becoming a problem for them as a couple. Towards the end of his review the radiographer asked him if there was anything else she could help him with. Frank told him about the problems he and his wife were experiencing when they tried to have intercourse, to which she replied it's just an effect of his diagnosis and the radiotherapy treatment. Frank expressed that he knew this but his original consultant had said there were some options and maybe some medication he could take. Reluctantly the radiographer nodded and replied this wasn't her area and so would go and ask a colleague for some advice. The radiographer exited the room into the main waiting area; leaving Frank sat alone with the door open. Whilst Frank was waiting he could hear laughter,

listening in he heard his specialist radiographer saying, (laughing)...I know tell me about it and it his age it....(laughter), yeh good point I'll tell him to google it there be plenty of stuff on the internet for that kind of thing.....well yeh, I suppose whatever floats your boat (laughter).

Step 7: Antecedents and Consequences

Figure 4 identifies antecedents for each of the five attributes of compassion identified in Figure 2.




Figure 4: Compassion's defining attributes and their antecedents

Figure 5 details the consequences which transpire when the five components (defining attributes) of compassion occur.




Figure 5: Consequences of the five components (defining attributes) of compassion.

Step 8: Empirical referents

The concept analysis ascertains the empirical referents of compassion can be structured into three categories: non-verbal, verbal and professional practice.

Non-verbal display

Compassionate behaviours included: eye contact [28], engaged body language [29], listening with full attention [30-32] and facial expressions which matched the subject of conversation [33]. All of these were deemed to display commitment and devotion by the health professional to what was being said, the significance of the topic and therefore demonstrated they were vested in them [34,35].

Verbal display

Compassionate behaviours included: being provided with clear, non-jargon, individual-patient-tailored information, being given time and the opportunity to ask questions, not being spoken over, the professionals asking questions of them about their preferences and level of understanding [33, 35-43]. The tone of voice and the language used was of particular

importance in a professional's ability to portray compassion due to the effects of an imbalance of power between the health professional and the patient [32, 44]. Belittling, judgemental attitude, oversimplifying and not taking the patient seriously were all perceived as uncompassionate behaviours [45,46].

Cutting across both verbal and non-verbal display was a display of warmth through touch, tone of voice and body language. Warmth made patients feel comfortable and gave the feeling of being 'cared for' even if the health professional wasn't actively involved in their treatment [38,47]. Also underlying throughout was a desire for the health professional to show respect to the patient [48]. Patients wanted to feel like their opinions, beliefs and preferences were not only known but valued by those responsible for their care [29,31,37,46,47,49]. Patients wanted interactions in all formats to be non-judgemental and be both understanding and accepting of the circumstances surrounding their needs [50].

Professional practice

Compassionate behaviours included undertaking and completing any required standard tasks as part of the patient's pathway and treatment [43], because ensuring that all process components have been completed prior to appointments prevents the patient facing additional distress of waiting, worrying about delays and being fearful of the consequences such delays could have on their prognosis. Patients want the health professionals to understand and appreciate the impact their current issue (diagnosis, bereavement, treatment etc.) is having on the physical, emotional and social rudiments of their life [30,31,33,51,52].

Furthermore, the practice of elementary tasks is deemed to be compassionate when delivered in such a way where a patient's dignity is maintained and considered as paramount [31, 38].

One of the five defining attributes of compassion was a personal connection between the two parties [37, 44, 53, 54]. Often this is on the basis of shared experience, knowledge or understanding of the current situation. There are caveats however within this, as some level of professional boundary is required in order for health professionals to make balanced, informed decisions about the care they provide which are not based on previous emotive events, beliefs or feelings [49].

Limitations

Despite the implementation of a rigorous search and review strategy, consideration needs to be given to omission of possible resources which may have limited the rigour of the findings. However, a saturation of themes occurred across all of resource categories denoting that further resources would not have brought additional themes to the analysis. Furthermore it may be argued that the use of Twitter did not align with the original aim to review compassion within a healthcare context. Twitter however provided a broader meaning to the term outside the published healthcare context as the twitter hashtag collated accounts and posts from a wide range of individuals, newspapers, charities and organisations.

Both Twitter and the dictionary definitions underpinned the review, clarified compassion outside of the healthcare environment by identifying patterns of use [22] and as such provided a boundary for understanding the differences between the concepts of compassion when used in different environments.

Previous concept analyses have focused on different uses of the topic which do not denote the practice or presentation of compassion within healthcare, for example compassion fatigue [55] and self-compassion [56]. Where compassion has been the focus of the analysis, defining attributes have failed to be identified [57]. Classification of these not only enables the concept to be distinguished from other similar concepts [23], but by inclusion provides a model to identify if each of the five attributes of compassion is being displayed. Furthermore this work has provided defining attributes, antecedents and consequences which are derived from and relevant to health care practice giving real world contextual meaning to the findings. Understanding and defining compassion is not just of relevance or only applicable to UK practice, the findings of this concept analysis didn't focus purely on one specific discipline as reported in other cases [58]. Therefore findings will be transferable to all health professions where treatment and patient-centred care is an essential requirement of the role.

Conclusion

Employment of a concept analysis has distinguished compassion in healthcare from other contexts and identifies it as composed of five attributes: Recognition, Connection, Altruistic desire, Humanistic response and Action. Associated meanings and behaviours have been outlined aiding an understanding of compassion. The findings however identify the complexity of the term and subjective nature in which it is displayed and in turn perceived.

The literature utilised within this process has identified there is no one agreed definition of compassion. Those which have been developed have been given an assigned meaning based on previous literature and dictionary definitions as opposed to being developed within an appropriate healthcare context. Further research needs to be conducted based upon these findings to develop a healthcare explicit definition of compassion within healthcare which will enhance understanding, thus facilitating active engagement and implementation into practice.

Acknowledgments

There are no acknowledgements

References

1. Society and College of Radiographers. Code of Professional Conduct. London, 5th July 2013.
2. Department of Health. The NHS Constitution for England. 26th March 2013.
3. Department of Health. Delivering high quality, effective, compassionate care: developing the right people with the right skills and the right values. England, Williams Lea for the Department of Health. 2013.
4. Department of Health. Education Outcomes Framework. England. 28 March 2013
5. NHS England. Putting Patients First: the NHS England business plan for 2014/15 – 2016/17. London..
6. Department of Health. Compassion in Practice; Nursing, Midwifery and Care staff, Our Vision and Strategy. London, December 2012
7. Report of the Mid Staffordshire NHS Foundation Trust Public Inquiry. Chaired by Robert Francis QC. London. 2013
8. Department of Health. Transforming care: A national response to Winterbourne View Hospital: Department of Health Review Final Report .December 2012
9. NHS. Review into the quality of care and treatment provided by 14 hospital trusts in England: overview report. Professor Sir Bruce Keogh. Chaired by Professor Sir Bruce Keogh KBE. July 2013
10. Care Quality Commission. Inspection Report; Ashcroft Care Home Limited . 10th January 2013
11. Department of Health. Government response to the House of Commons Health Select Committee Fourth Report of session 2014–15 Complaints and Raising Concerns. March 2015
12. Department of Health. High Quality Care for All NHS. Next Stage Review Final Report. June 2008
13. Health Education England. Clinical academic careers. <https://www.hee.nhs.uk/our-work/developing-our-workforce/clinical-academic-careers>
14. Royal College of Nursing. Quality with Compassion: the future of nursing education. Report of the Willis Commission on Nursing Education, 2012.
15. Traynor, M. (2014) Caring after Francis: moral failure in nursing reconsidered. Journal of Research in Nursing. 19(7–8) 546–556.

16. Crawford, P., et al. (2013). The language of compassion in acute mental health care. *Qualitative health research*. 23 (6) 719-727.
17. Bleiker, J., et al (2016) Compassionate care in radiography recruitment, education and training: A post-Francis report review of the current literature and patient perspectives. *Radiography*. 22, 257-262.
18. Dewar, B. (2013). Cultivating compassionate care. *Nursing standard*, 27 (34) 48-55.
19. Dewar, B. Christley, Y. (2013). A critical analysis of Compassion in Practice. *Nursing standard*: 28 (10), 46-50.
20. Dewar, B. Nolan, M. (2013). Caring about caring: Developing a model to implement compassionate relationship centred care in an older people care setting. *International journal of nursing studies*, 50 (9) 1247-1258.
21. Dewar, B. (2012). Using creative methods in practice development to understand and develop compassionate care. *International Practice Development Journal*. 2 (1) 1-11 11p.
22. Cronin, P. Ryan, F. and Coughlan, M., (2010) Concept analysis in healthcare research, *International Journal of Therapy and Rehabilitation*, 17 (2) 62-68.
23. Walker L and Avant K. *Strategies for theory construction in nursing*. Pearson Prentice Hall. Pearson Education. 2005
24. Wilson, J. *Thinking with concepts*. New York: Cambridge University Press. 1963
25. Carper, B. (1978). Fundamental patterns of knowing in nursing. *Advances in Nursing Science*, 1(1) 13-23.
26. Chinn, P. Kramer, M. *Integrated knowledge development in nursing (6th ed.)*. St. Louis, MO: Mosby. 2004
27. Knafl, K., Deatrick, J. (2000) Knowledge Synthesis and Concept Development in Nursing. In Rodgers, B., Knafl, K. (eds.) *Concept Development in Nursing. Foundations, Techniques and Applications*. 2nd ed. WB Saunders, Philadelphia: (39-54).
28. Saunders, J (2015). Compassion. *Clinical medicine (London, England)*, 15 (2) 121-124.
29. Marchuck, A. (2014). A personal nursing philosophy in practice. *Journal of neonatal nursing*, 20 (6), 266-273.
30. Van Der Cingel, M. (2011). Compassion in care: A qualitative study of older people with a chronic disease and nurses. *Nursing ethics*. 18 (5) 672-685.
31. Proctor, S. (2007) Can nurses show compassion? *Nursing Management*. 14 (8) 10.

32. Penson, R., et al. (2001). Caring for colleagues. *The oncologist*, 6 (2) 197-204.
33. Priddis, H., et al. (2014). " A patchwork of services" - caring for women who sustain severe perineal trauma in New South Wales - from the perspective of women and midwives. *BMC pregnancy and childbirth*, 14 (1).
34. Marcum, J. (2011). Care and competence in medical practice: Francis Peabody confronts Jason Posner. *Medicine, Health care, and Philosophy*. 14 (2) 143-153.
35. Bynum, W. (2014). Why physicians need to be more than automated medical kiosks. *Academic medicine: Journal of the association of American medical colleges*, 89 (2) 212-214.
36. Haq, C. (2014). Compassion in medicine. *Family medicine*, 46 (7) 549-550.
37. Thorne, S., et al. (2005). 'Being known': patients' perspectives of the dynamics of human connection in cancer care. [online]. *psycho-oncology*. 14 (10) 887-898.
38. Horsburgh, D., Ross, J. (2013). Care and Compassion: The experiences of newly qualified staff nurses. *Journal of clinical nursing*, 22 (1124-1132).
39. Schneider, M, Smith, C, Pomidor, M.. (2015). Compassionate Care for Patients With Complex Regional Pain Syndrome. *The journal of neuroscience nursing: Journal of the American association of neuroscience nurses*, 47 (4), 204-210.
40. Owen, R and Jeffrey, D (2008). Communication: Common challenging scenarios in cancer care. *European journal of cancer*, 44 (8) 1163-1168.
41. Perry, B (2009) Conveying compassion through attention to the essential ordinary.. *Nursing Older People*, 21 (6).
42. Halifax, J. (2011). The precious necessity of compassion. *Journal of pain and symptom management*, 41 (1) 146-153.
43. Steele, A. C., et al. (2013). Bereaved parents and siblings offer advice to health care providers and researchers. *Journal of paediatric Haematology/Oncology*. 35 (4) 253-259.
44. YouTube - Dr Kate Granger talks about compassionate care and #hellomynameis. Aug 8, 2014 https://www.youtube.com/watch?v=Be_nIItj8bs
45. Smith, J., et al. (2013). Clinical care for sexual assault survivors multimedia training: a mixed-methods study of effect on healthcare providers' attitudes, knowledge, confidence, and practice in humanitarian settings. *Conflict and health*, 7, 14.
46. Santen, S., Hemphill, R. (2011). A window on professionalism in the emergency department through medical student narratives. *Annals of emergency medicine*, 58 (3) 288-294.

47. Goetz, J, Keltner, D, Simon-Thomas, E (2010). Compassion: An Evolutionary Analysis and Empirical Review. [online]. *Psychological bulletin*. 136 (3) 351-374.
48. Rohan, E. Bausch, J. (2009). Climbing Everest: oncology work as an expedition in caring. *Journal of psychosocial oncology*. 27 (1) 84-118 35p.
49. Schapira, Lidia, et al. (2014). Caring for one of our own. *The oncologist*, 19 (5) 545-549.
50. Puchalski, C. Jafari, N (2015). Acknowledging the person in the clinical encounter: Whole person care for patients and clinicians alike. Commentary on Chochinov et al. *Journal of pain and symptom management*, 49 (6) 973.
51. Mannix, R. (2012). A compassionate care checklist. *Academic emergency medicine*, 19 (8), 992.
52. Straughair, C. (2012). Exploring compassion: Implications for contemporary nursing. Part 1. *British journal of nursing*, 21 (3) 160-164.
53. Price, B. (2013). Promoting compassionate care through learning journeys. *Nursing standard (royal college of nursing (Great Britain) : 1987)*, 27 (48) 51-57.
54. Bolderston, A., Lewis, D, Chai, M J. (2010). The concept of caring: Perceptions of radiation therapists. *Radiography*. 16 (3) 198-208.
55. Coetzee, S., K Klopper, H. (2010) Compassion fatigue within nursing practice: A concept analysis. *Nursing and Health Sciences*. 12, 235–243.
56. Reyes, D. Self-Compassion: A Concept Analysis. (2012) *Journal of Holistic Nursing*, 30 (2) 81-89.
57. Schantz, M. (2007) Compassion: A Concept Analysis. *Nursing Forum*, 42, (2) 48-55.
58. Menage, D, et al. (2016) A concept analysis of compassionate midwifery. *The Journal of Advanced Nursing*. 73 (3), 558-573.

Appendix 1: Search strategy

An initial scoping search identified a number of fundamental papers whose key words have been used to develop search terms in conjunction with database specific indexed control vocabulary (including MeSH terms).

A building block approach as advocated by Booth (2008) provided a foundation to the search with the numerous facets connected or eliminated by the use of BOOLEAN operators.

Truncations were used to widen the search for example: compassion* retrieved compassion and compassionate. Similarly wildcards allowed for variances in spelling of common words, for example behavio#r or behavio?r.

Key terms

compassion* , "compassion* car*"

"Professional behav*" "practitioner-patient relation*", "nurse-patient relation*", "patient car*", "person-centred car*", "professional-patient relation*", "relational practice", "staff-client relation*", "relationship-centred car*", "professional issues", "patient-centred car*"

AND

healthcare, "health profession*", medic*, "clinical medic*", "Medical car*", "nurs* practice", "allied health profession*", "multidisciplinary team", hospital*, "professional carer*", "health service*" "healthcare organisation", "health person*".

AND

Behav*, behavio#r, attribute*, trait*, relation*, attitude*.

Inclusion Criteria

Resources which:

- Were written in English
- Published between 1995 and 2015.
- Reviewed compassion and its concepts within healthcare
- Discussed and/or examined compassionate behaviour in healthcare

NB Evidence based material was not exclusively searched as the concept of compassion is fluid and open to individual interpretation. The literature review therefore wanted to capture this discussion and as such included non-research based material including discussion papers, commentary's, letters to editors, book chapters etc. These provided background and rationale for the project in addition to helping to understand what is meant by compassion in healthcare.

Exclusion Criteria

Resources which:

- Reviewed compassion outside of healthcare practice
- Addressed self-compassion
- Investigated/reported approaches to reduce compassion fatigue
- Solely addressed the management of patients receiving End of Life/Palliative Medicine
- Related to professional practice issues outside of compassion for example, medical errors which are not due to poor compassionate care.
- Addressed other behaviours for example medical skill levels which are not attributable to compassion.

Google and Google Scholar were used as method of searching grey literature and as a supplementary tool in conjunction with reference searching from seminal papers respectively.

Reference

Booth A. (2008) Using Evidence in Practice. Unpacking your literature search toolbox: on search styles and tactics. *Health Information and Libraries journal*. 25, 313-317.