

Exercise fidelity and progression in a supervised exercise programme for adults with venous leg ulcers

KLONIZAKIS, Markos <<http://orcid.org/0000-0002-8864-4403>>, GUMBER, Anil <<http://orcid.org/0000-0002-8621-6966>>, MCINTOSH, Emma, KING, Brenda, MIDDLETON, Geoff, MICHAELS, Jonathan A and TEW, Garry

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/20908/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

KLONIZAKIS, Markos, GUMBER, Anil, MCINTOSH, Emma, KING, Brenda, MIDDLETON, Geoff, MICHAELS, Jonathan A and TEW, Garry (2018). Exercise fidelity and progression in a supervised exercise programme for adults with venous leg ulcers. *International Wound Journal*, 15 (5), 822-828.

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

1 Introduction

2 Venous leg ulceration is a chronic and devastating condition that affects approximately 1% of the
3 adult population in the Western world (1). It costs up to 198 million sterling pounds in national
4 healthcare expenditure in the U.K. alone (2), affecting significantly, in a negative manner patients'
5 quality of life (3). Moreover, venous leg ulcers tend to recur quite frequently, with recurrence
6 rates reaching 70% within a year of healing (4).

7 With such costs involved and the considerable devastation in patients' lives, it is no surprise that
8 adjunct and alternative therapies to compression therapy (which is considered as the golden
9 standard) (5) have been pursued (e.g. ultrasound (6), larval therapy (7), biomaterials (8)), with
10 exercise and physical activity promotion being considered as well (e.g., walking (9), increased
11 physical activity (10), resistance exercise (11)).

12 The concept of using exercise as an adjunct therapy to compression isn't new and indeed exercise
13 is included as a recommendation in the NICE Clinical Knowledge Summary for venous leg ulcers'
14 management (e.g., "regular walking", "exercising to improve calf muscle pump function") (12).

15 Nevertheless, there is a fear shared by both clinicians and the patients that exercise may be either
16 inappropriate or harmful and actually delay rather than promote healing (13,14). This notion
17 together with the mixed results of previous studies (13,15-16), has limited the exploration of
18 regimes that could potentially benefit patients and improve clinical outcomes. Overall, there is
19 little published data on the ability of this patient group to undertake different types of exercise
20 training and on rates of exercise progression. The data has the potential to inform practitioners
21 and researchers involved in prescribing and supervising exercise with venous ulcer patients. .

22 FISCU (17) is a recently-completed, two-center study exploring the feasibility of using exercise as
23 an adjunct therapy to compression in patients with venous leg ulcers. This trial represents an

24 attempt to implement a supervised exercise programme with this patient population, in a manner
25 similar to what has been promoted successfully in other clinical populations in the UK (e.g.
26 peripheral arterial disease (18), chronic obstructive pulmonary disease (19), cardiac diseases (20)).
27 Central to the internal validity of all intervention trials is intervention fidelity, which refers to the
28 extent an experimental manipulation has been implemented in a comparable manner to all
29 participants, as intended (21). Furthermore, it is important to present all in-session exercise safety
30 data to better inform clinicians, policy makers and patients with venous ulcers. As such, having
31 published the main study findings, which supported the feasibility of conducting a future full-scale
32 trial (17), our aim here was to present a detailed appraisal of exercise data collected during the
33 FISCU trial, focussing on treatment fidelity and exercise progression.

34

35 **Methods**

36 FISCU was a two-arm, parallel-group, randomised feasibility trial that received ethical clearance
37 from the NHS National Research Ethics Committee for Yorkshire and the Humber (14/YH/0091),
38 and was prospectively registered (ISRCTN09433624). Thirty-eight adults who were receiving
39 lower-limb compression for a new venous leg ulcer of greater than 1 cm diameter were recruited
40 from tissue viability clinics and newspaper advertisement in Sheffield, United Kingdom. Following
41 provision of consent and baseline assessment, participants were randomly assigned to receive
42 usual care (n=20) or usual care plus a 12-week supervised exercise programme (n=18). A full
43 description of the protocol is available elsewhere (22); however, for the purpose of this article the
44 exercise training protocol is described below.

45

46 **Exercise protocol**

47 Following study enrolment and randomisation, exercise group participants were referred for a 12-
48 week exercise intervention, undertaken 3 times per week (typically being delivered on Mondays,
49 Wednesdays and Fridays to allow sufficient recovery between sessions). A maximum of an
50 additional 2 weeks was allowed for the participants to complete the 36 sessions in case sessions
51 were missed because of illness, family/work commitments or holiday. The sessions were
52 supervised by an exercise physiologist and were typically undertaken in a group form (no more
53 than 4 patients per session, to ensure proper supervision and adequate progression monitoring).
54 Each exercise session lasted approximately 60 minutes and comprised a combination of aerobic,
55 resistance and flexibility exercises. Each session began and ended with 5 minutes of low-intensity
56 treadmill walking or cycling for a warm-up and cool-down, respectively. The aerobic component
57 was aimed to last approximately 30 minutes, with the exercise mode being treadmill walking,
58 cycling, or a combination of both, with the mode being determined by the physical function and
59 preference of participants.

60 Resistance and flexibility exercises were performed for approximately 20 minutes in order to
61 improve calf muscle pump function, leg (predominantly calf) muscle strength, and joint
62 (predominantly ankle) mobility. Resistance exercises mainly involved dynamic body-weight
63 exercises with or without the use of dumbbells and stability balls (e.g., calf raises and partial
64 squats). Exercise was aimed to be performed for two or three sets of 10 to 15 repetitions to the
65 point of moderate muscle fatigue (23). For flexibility, static stretches were performed for all of the
66 major muscle groups of the legs, for a total of 60 seconds per muscle group (comprising 3 × 20-
67 second stretches), held at the point of mild discomfort (23).

68

69 **Exercise intensity: prescription and measurement**

70 The intensity of aerobic and resistance exercises was guided using Borg’s 6-20 ratings of
71 perceived exertion (RPE) scale (24), aiming for an exertion level of 12 to 14 (“moderate” to
72 “somewhat hard”) on the 6-20 scale, which equates to the ventilatory threshold (24). Each patient
73 was familiarized with the scale and the recommended researcher instructions for scale
74 administration were used (25). Perceived exertion, heart rate (via telemetry; Polar RS400,
75 Kempele, Finland), and aerobic and resistance exercise indices (e.g., treadmill speed and gradient)
76 were recorded at regular intervals during the whole session to allow accurate quantification of the
77 exercise stimulus and to facilitate progression of the programme over time.

78

79 **Exercise safety**

80 Compression garments (stockings/bandages) were monitored throughout each exercise session.
81 The exercise supervisor was instructed to terminate the session if these were affected by exercise,
82 with participants being referred to the tissue-viability nursing team for re-application, and
83 additional visits were to be noted for the health-economics analyses. Our safety monitoring
84 procedure indicated that all serious adverse events, as well as all non-serious adverse events that
85 are deemed to be related to participation in the research (e.g., exercise strains or injuries,
86 excessive wound discharge, in-session exercise bandage slipping) were to be recorded during the
87 period between provision of informed consent through to 12 months after randomisation.
88 Participants were asked to contact the study team to inform them about adverse events if and
89 when they occur. Study investigators also questioned participants about the occurrence of
90 adverse events during each participant study visit.

91 **Statistical analysis**

92 Descriptive statistics were used to calculate the session attendance data, completion rates as per
93 protocol for aerobic (duration, intensity, combination of duration and intensity), resistance
94 (intensity, number of exercises, sets, repetitions) and flexibility exercises (number of exercises,
95 duration, intensity), and present baseline demographics. The Shapiro-Wilk test was used to assess
96 data normality and Mauchly's Test of Sphericity was used to indicate data sphericity (the assumption
97 of sphericity was not violated in any case). Exercise progression was assessed by comparing Session
98 1 (baseline), with Sessions 18 (midpoint) and 36 (intervention completion) using Analysis of
99 Variance (ANOVA) for Repeated Measures (SPSS v.23, Armonk, NY: IBM Corp). Post-hoc analysis
100 was undertaken using Bonferroni corrected t-tests. To calculate the effect sizes we used eta-
101 square for ANOVA assessments and Cohen's d for post-hoc analysis, using the magnitudes
102 determined by Cohen (26): For η^2 0.01 is considered a small effect, 0.06 is considered a medium
103 effect and 0.14 is considered a large effect. Similarly for Cohen's d: 0.2 is considered a small effect,
104 0.5 is considered a medium effect and 0.8 is considered a large effect. Data are described as
105 means (SD), unless otherwise stated. Significance set at $p < 0.05$ and for post hoc analysis at
106 $p < 0.0167$.

107

108 **Results**

109 *Participants*

110 Characteristics of the 18 exercise-group participants are shown in Table 1. Ten of these
111 participants were female and the mean \pm SD age, stature and body mass were 66.9 ± 13.9 years,
112 171.1 ± 11.9 cm and 102.1 ± 29.4 kg, respectively. Median ulcer size was 4.9 cm^2 .

113

114 *Attendance*

115 The overall exercise attendance rate was 79% (512/648), with 13 of the 18 participants (72%)
116 attending all exercise sessions. Amongst those who completed the study, 411/468 sessions were
117 completed within the 12-week period, with the rest (57/468) being completed within the
118 additional 2-week period. Of the five participants who did not complete all sessions, one withdrew
119 fully from the trial before the 3-month follow-up assessment due to non-ulcer-related health
120 problems, and four withdrew from treatment (i.e. stopped attending before the end of the 12-
121 week intervention period) but remained in the study (one due to ulcer-related problems, three
122 due to non-ulcer-related health problems). These five participants had completed 2, 4, 6, 15 and
123 17 exercise sessions, respectively, before withdrawing. Reasons for not attendance included lack
124 of transportation (n=34), non-ulcer related health reasons (n=74) and ulcer-related health reasons
125 (n=32), with more than one reasons given on some occasions.

126

127 *Exercise Safety*

128 No serious, in-session adverse events were experienced and the bandaging was also not disrupted
129 during any exercise session. Two incidents of excessive fluid discharge were detected the day after
130 exercise sessions, possibly or probably related to exercise. Following consultations with healthcare
131 personnel, these were dealt by postponing the exercise session following the incident reporting

132 (incident 1) and temporarily removing the resistance element from the training programme

133 (incident 2).

134

135 *Exercise choices*

136 The majority of the participants (72%) chose treadmill as their main aerobic mode of training at

137 baseline, with the rest preferring cycling due to frailty and lack of confidence with exercising on

138 the treadmill. One participant changed briefly from treadmill to exercise cycle, before reverting to

139 treadmill again. Only one of the participants was training via exercise cycle at the end of the 12-

140 week intervention, with the rest of the participants that completed the intervention using the

141 treadmill instead.

142

143 In regards to the resistance element of the intervention, four participants started the programme

144 stating that they were unable to do squats, step-ups or calf raises. This number was reduced to

145 two at the end of their participation (they were however, able to complete the rest of the regime).

146 Finally, one of the participants could not do squats on Session 18, due to a pre-existing pain

147 unrelated to exercise, completing however, the rest of the session without issues. The participant

148 completed his programme as well.

149 *Exercise Intensity*

150 All of the aerobic and 91% of the resistance training components, across all participants, were

151 performed at the desired moderate intensity, as determined using RPE responses in the 12-14

152 range (Tables 2 and 4).

153

154 *Exercise Progression*

155 Table 3 presents data on changes in the duration of the aerobic component and the number of
156 repetitions completed for four lower-limb resistance exercises. The number of minutes spent on
157 aerobic exercise increased through the 12-week period (Baseline: 19 min (8), Mid-point: 26 min
158 (5), End-point: 29 min (3)).

159 Performance of the participants in the resistance exercise indices was also improved (Table 3): For
160 example, calf raises increased from 19 (13) at baseline, to 36 (13) at mid-point, reaching 42 (14) at
161 the end of the intervention.

162

163 *Exercise Fidelity*

164 For the aerobic exercise element all completed sessions were completed according to the
165 prescribed intensity. For resistance this was the case in 466/512 (91%) completed sessions.
166 Duration of the exercise elements was close to the prescribed duration as well (413/512 = 81% for
167 aerobic, 474/512 = 93% for resistance). The majority of those not completing the prescribed
168 duration were at the beginning of their programme, and was due to lack of physical fitness (n=4),
169 discomfort (n=2) and unfamiliarity with the training equipment/exercises (n=4) – with more than
170 one reason being given by some participants.

171 Similarly, the main reason for resistance components not being completed according to protocol
172 was lack of physical fitness. This, however, became less of an issue as the programme progressed,
173 reaching almost 100% completion in the last sessions.

174 Finally, flexibility exercises were completed as per protocol in regards to duration and number of
175 exercises.

176 **Discussion**

177 Using a supervised exercise regime as an adjunct therapy to reduce venous leg ulcer healing time,
178 represents a plausible, yet largely unassessed therapeutic strategy (16). The lack of appropriately
179 designed studies, which would substantiate its use and the fear of healthcare professionals and
180 the patients themselves about the safety and applicability of exercise are two main reasons, why
181 the advice of a more "active lifestyle" is not being taken up more widely within this patient
182 population (13,14).

183 We have recently presented data supporting the feasibility of a full-scale trial of adjunctive
184 exercise therapy for venous leg ulceration (17). The aim of the current paper was to undertake a
185 detailed evaluation of the exercise session data. When adhering to pre-determined safety criteria,
186 our results show primarily a very high fidelity of our proposed programme. It is evident from our
187 data that not only is it possible to exercise this primarily-older and largely-frail, patient population
188 at moderate intensities, but it is also possible to see a positive exercise progression over the
189 duration of a medium-term training programme. This is the first study to report in-session data on
190 this patient group and this acts as a comparator for researchers and practitioners embarking on
191 similar trials with exercise as a therapy with this patient group.

192 *Attendance, compliance, and safety*

193 Our overall session attendance (79%) for the 18 participants across the 12-week exercise
194 intervention compares well with an attendance range of 58–77% for other exercising, clinical
195 populations (27,28). Our attendance results can be interpreted even more favourably to those

196 achieved in other exercise studies, considering the fact that ours was a time-demanding (e.g. 3
197 times per week), 3-month intervention, focusing on a group which is older, sedentary and without
198 an exercising culture; the large majority of our participants have not previously followed an
199 exercise programme. Consequently, it can be postulated that our participants were keen to
200 embrace such an intervention and participated whole-heartedly. Our results also show that most
201 missed sessions can be accounted to reasons unrelated to the exercise programme (e.g. illnesses
202 and family commitments) rather than the exercise programme itself. This knowledge, combined
203 with the very good safety record (e.g., no participants had their compression garments affected
204 during the exercise sessions), is a sign of trust of moving the intervention into the next stage, that
205 of the definitive trial. Nevertheless, much more data is required to evaluate the safety of the
206 intervention properly in this patient group.

207 When evaluating the fidelity of exercise training interventions researchers should ideally consider
208 both session attendance and meeting the prescribed exercise intensity, as this interaction
209 constitutes the dose of the intervention and influences the physiological response to exercise
210 training (21). Although in our case this might have been considered as a relatively easy task (as our
211 aim was to have participants exercising at "moderate" intensity, e.g. 12-14 in the 6-20 RPE Borg
212 scale), which is considerably lower to that sought by high intensity training (e.g., 85–95% of peak
213 heart rate) (29) exercise regimes, results should not be overlooked: our participants' unfamiliarity
214 with exercise interventions and in some cases frailty, meant that even the intended moderate
215 intensity could potentially be difficult to achieve in practice. For the aerobic exercise element of
216 our intervention this was achieved and maintained throughout the duration of the intervention,
217 matching the performance of other regimes, conducted in clinical settings, in older clinical
218 populations (e.g. Alzheimer's Disease) (30). Results differ in regards to resistance and flexibility, as

219 certain participants found difficulty to complete all resistance exercises to the required level
220 (Table 3) or intensity (Table 4). This was mainly due to frailty and lack of physical fitness (number
221 of sets/repetitions for resistance) or patients finding the exercises easier than expected (intensity
222 for flexibility). This can only be considered as part of our learning process to introduce more
223 challenging exercises (for flexibility) and a varying introductory pace (for resistance), in the future
224 study stages.

225

226 *Exercise progression*

227 The main aim of this article was to present our findings on attendance, compliance and safety.
228 Nevertheless, our detailed collection and analysis of exercise training data permits the objective
229 appraisal of our regime in regards to exercise progression as well: To facilitate a positive
230 adaptation to training, the prescription of exercise needs to advance over time (27). Many
231 programmes have failed to achieve this, presenting a need to re-define targets, following an in-
232 programme assessment (31) (which can be costly and resource-intensive). In the study presented
233 in this article, we used relative measures of exercise intensity to assess adherence to the
234 prescribed intensity. The fact that our aim was achieved was reflected in all of our exercise
235 indices, which show a statistically-significant increase in most measures, as well as a moderate-to-
236 large effect sizes: this demonstrates a clear exercise progression. Although it is difficult to
237 compare our findings with that of other trials in clinical or older populations (as in-session data is
238 not usually reported), our data is equally- or more favourably- comparable to similar interventions
239 in other clinical populations where physical functioning indices appear to be reduced (e.g. chronic
240 kidney disease) (32) or improved (e.g. older people living in retirement communities (33), multiple

241 sclerosis (34)) when compared with normative values. It remains to see whether this exercise
242 progression will be achievable in the definitive trial as well, nevertheless, the indicators are
243 encouraging, suggesting that participants with venous ulcers can benefit in multiple ways (e.g.
244 improved cardiorespiratory endurance (35) and better physical function (27), which are related to
245 high exercise session attendance) by taking part in such an intervention combining medium-
246 intensity aerobic, flexibility and resistance exercise, as previous studies in clinical populations have
247 shown (36,37).

248 *Limitations*

249 With this study exploring the feasibility of the intervention, the number of participants was
250 relatively small to what the definitive trial is expected to include. With that in mind findings should
251 be treated as indicative. Additionally, an in-depth assessment of fidelity in a definitive, multi-
252 centre exercise intervention will examine the consistency of the exercise dose across the different
253 sites, something that was not possible on this occasion. Finally, we cannot rule out the possibility
254 of underreported RPE scores due to the influence of observer sex as it has been suggested that
255 male participants report lower RPE values when a female observer, as opposed to male, is in the
256 room (38). Nevertheless, our sessions were delivered by both male and female physiologists and
257 our findings appear to be consistent throughout the intervention, hence the likelihood of that is
258 small.

259 *Conclusions*

260 This is the first study to provide a detailed quantification of the exercise sessions performed across
261 an exercise intervention combining aerobic, resistance and flexibility exercises for patients with
262 venous ulcers. The data will act as a comparator for researchers embarking on similar trials and

263 advocating exercise to this patient group in their practice. Our findings showed that our
264 participants trained at the intended exercise intensity, improving their performance amongst all
265 exercise domains in which they trained (e.g., number of minutes in aerobic exercise, number of
266 squats and calf raises etc), without having their safety compromised. We conclude that it is
267 possible to exercise this patient population at moderate exercise intensities. This is purposeful for
268 further studies which consider deploying similar supervised exercise regimes as an adjunct
269 therapy to compression, in an attempt to reduce healing times in patients with venous ulcers.

270 **Conflicts of Interest**

271 None to declare.

272 **References**

-
- 273 1. Donnelly R, London NJ. The Ulcerated lower limb. In: Donnelly R, London N. NJ eds., *ABC of*
274 *Arterial and Venous Disease*. 2nd ed. Chichester, United Kingdom; Wiley-Blackwell; 2009:77-83.
- 275 2. Posnett J, Franks PJ. The burden of chronic wounds in the UK. *Nurs Times*. 2008;**104**:44-45.
- 276 3. Carradice D, Mazari FA, Samuel N, Allgar V, Hatfield J, Chetter IC. Modelling the effect of
277 venous disease on quality of life. *Br J Surg*. 2011;**98**:1089-1098.
- 278 4. Barwell JR, Davies CE, Deacon J, Harvey K, Minor J, Sassano A, Taylor M, Usher J, Wakely C,
279 Earnshaw JJ, Heather BP. Comparison of surgery and compression with compression alone in
280 chronic venous ulceration (ESCHAR study): randomised controlled trial. *Lancet*. 2004;**363**:1854-
281 1859.
- 282 5. O'Meara S, Cullum N, Nelson EA, Dumville JC. Compression for venous leg ulcers. *Cochrane*
283 *Database Syst Rev*. 2012;**11**:CD000265.

- 284 6. Watson JM, Kang'ombe AR, Soares MO, et al. Use of weekly, low dose, high frequency
285 ultrasound for hard to heal venous leg ulcers: the VenUS III randomised controlled trial. *BMJ*.
286 2011;**342**:d1092-1101.
- 287 7. Dumville JC, Worthy G, Bland JM, Cullum N, Dowson C, Iglesias C, Mitchell JL, Nelson EA, Soares
288 MO, Torgerson DJ. Larval therapy for leg ulcers (VenUS II): randomised controlled trial. *BMJ*.
289 2009;**338**:b773-780.
- 290 8. Mostow EN, Haraway GD, Dalsing M, Hodde JP, King D, OASIS Venus Ulcer Study Group.
291 Effectiveness of an extracellular matrix graft (OASIS Wound Matrix) in the treatment of chronic leg
292 ulcers: a randomized clinical trial. *Journal of vascular surgery. J Vasc Surg.* 2005;**41**:837-843.
- 293 9. Meagher H, Ryan D, Clarke-Moloney M, O'Laighin G, Grace PA. An experimental study of
294 prescribed walking in the management of venous leg ulcers. *J Wound Care.* 2012;**21**:421-430.
- 295 10. Heinen MM, Evers AW, Van Uden CJ, Van der Vleuten CJ, van de Kerkhof PC, Van Achterberg
296 T. Sedentary patients with venous or mixed leg ulcers: determinants of physical activity. *J Adv
297 Nurs.* 2007;**60**:50-57.
- 298 11. O'Brien J, Finlayson K, Kerr G, Edwards H. Evaluating the effectiveness of a self-management
299 exercise intervention on wound healing, functional ability and health-related quality of life
300 outcomes in adults with venous leg ulcers: a randomised controlled trial. *Int Wound J.*
301 2017;**14**:130-137.
- 302 12. Clinical Knowledge Summaries: Leg ulcer – venous, National Institute for Health and Clinical
303 Excellence <https://cks.nice.org.uk/leg-ulcer-venous#!topicsummary>. Accessed Jan 25, 2018.
- 304 13. O'Brien J, Finlayson K, Kerr G, Edwards H. The perspectives of adults with venous leg ulcers on
305 exercise: an exploratory study. *J Wound Care.* 2014;**23**:496-509.
- 306 14. McCulloch J, Mahoney E, McCallon S. Enhancing the role of physical therapy in venous leg
307 ulcer management. *JAMA Dermatol.* 2015;**151**:327-327. doi: 10.1001/jamadermatol.2014.4042

- 308 15. Jull A, Parag V, Walker N, Maddison R, Kerse N, Johns T. The prepare pilot RCT of home-based
309 progressive resistance exercises for venous leg ulcers. *J Wound Care*. 2009;**18**:497-503.
- 310 16. Yim E, Kirsner RS, Gailey RS, Mandel DW, Chen SC, Tomic-Canic M. Effect of physical therapy
311 on wound healing and quality of life in patients with venous leg ulcers: a systematic review. *JAMA*
312 *Dermatol*. 2015;**151**:320-327.
- 313 17. Klonizakis M, Tew GA, Gumber A, Crank H, King B, Middleton G, Michaels JA. Supervised
314 exercise training as an adjunct therapy for venous leg ulcers: a randomised controlled feasibility
315 trial. *Br J Dermatol*. 2017;**27**: In Press. doi: 10.1111/bjd.16089
- 316 18. Layden J, Michaels J, Bermingham S, Higgins B, Guideline Development Group. Diagnosis and
317 management of lower limb peripheral arterial disease: summary of NICE guidance. *BMJ*.
318 2012;**345**:e4947. doi: 10.1136/bmj.e4947
- 319 19. Liddell F, Webber J. Pulmonary rehabilitation for chronic obstructive pulmonary disease: a
320 pilot study evaluating a once-weekly versus twice-weekly supervised programme. *Physiotherapy*.
321 2010;**96**:68-74.
- 322 20. Sandercock GR, Cardoso F, Almodhy M, Pepera G. Cardiorespiratory fitness changes in patients
323 receiving comprehensive outpatient cardiac rehabilitation in the UK: a multicentre study. *Heart*.
324 2013;**99**:785-90.
- 325 21. Taylor KL, Weston M, Batterham A. Evaluating intervention fidelity: an example from a high-
326 intensity interval training study. *PLoS One*. 2015;**10**:e0125166. doi: 10.1371/journal.pone.0125166
- 327 22. Tew GA, Michaels J, Crank H, Middleton G, Gumber A, Klonizakis M. Supervised exercise
328 training as an adjunctive therapy for venous leg ulcers: study protocol for a randomised controlled
329 trial. *Trials*. 2015;**6**:443-453.
- 330 23. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, Nieman DC, Swain
331 DP. American College of Sports Medicine position stand. Quantity and quality of exercise for

332 developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in
333 apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc.* 2011;**43**:1334-
334 1359.

335 24. Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc.* 1982;**14**:377-381.

336 25. Purvis JW, Cureton KJ. Ratings of perceived exertion at the anaerobic threshold. *Ergonomics.*
337 1981;**24**:295-300.

338 26. Cohen J. *Statistical power analysis for the behavioral sciences.* 2nd ed. Hillsdale, NJ: Lawrence
339 Earlbaum Associates; 1988.

340 27. Picorelli AM, Pereira LS, Pereira DS, Felício D, Sherrington C. Adherence to exercise programs
341 for older people is influenced by program characteristics and personal factors: a systematic
342 review. *J Physiother.* 2014;**60**:151-6.

343 28. Weston M, Batterham AM, Tew GA, Kothmann E, Kerr K, Nawaz S, Yates D, Danjoux G.
344 Patients Awaiting Surgical Repair for Large Abdominal Aortic Aneurysms Can Exercise at Moderate
345 to Hard Intensities with a Low Risk of Adverse Events. *Front Physiol.* 2017; **9**:684.

346 29. Weston KS, Wisløff U, Coombes JS. High-intensity interval training in patients with lifestyle-
347 induced cardiometabolic disease: a systematic review and meta-analysis. *Br J Sports Med.*
348 2014;**48**:1227-1234.

349 30. Watson E, Yu F. Monitoring exercise delivery to increase participation adherence in older
350 adults with Alzheimer's disease. *J Gerontol Nurs.* 2013;**39**:11-4.

351 31. West MA, Loughney L, Lythgoe D, Barben CP, Sriadam R, Kemp GJ, Grocott MP, Jack S. Effect
352 of rehabilitation on objectively measured physical fitness after neoadjuvant treatment in
353 preoperative rectal cancer patients: a blinded interventional pilot study. *Br J Anaesth.*
354 2015;**114**:244-251.

- 355 32. Padilla J, Krasnoff J, Da Silva M, Hsu CY, Frassetto L, Johansen KL, Painter P. Physical
356 functioning in patients with chronic kidney disease. *J Nephrol.* 2008;**21**:550-559.
- 357 33. Baker MK, Kennedy DJ, Bohle PL, Campbell DS, Knapman L, Grady J, Wiltshire J, McNamara M,
358 Evans WJ, Atlantis E, Fiatarone Singh MA. Efficacy and feasibility of a novel tri-modal robust
359 exercise prescription in a retirement community: a randomized, controlled trial. *J Am Geriatr Soc.*
360 2007;**55**:1-10.
- 361 34. Carter AM, Daley AJ, Kesterton SW, Woodrooffe NM, Saxton JM, Sharrack B. Pragmatic
362 exercise intervention in people with mild to moderate multiple sclerosis: a randomised controlled
363 feasibility study. *Contemp Clin Trials.* 2013;**35**:40-47.
- 364 35. Rhodes RE, Martin AD, Taunton JE, Rhodes EC, Donnelly M, Elliot J. Factors associated with
365 exercise adherence among older adults. An individual perspective. *Sports Med.* 1999; **28**:397-411.
- 366 36. Pothirat C, Chaiwong W, Phetsuk N. Efficacy of a simple and inexpensive exercise training
367 program for advanced chronic obstructive pulmonary disease patients in community hospitals. *J*
368 *Thorac Dis.* 2015;**7**:637-643.
- 369 37. Taaffe DR, Newton RU, Spry N, Joseph D, Chambers SK, Gardiner RA, Wall BA, Cormie P,
370 Bolam KA, Galvão DA. Effects of Different Exercise Modalities on Fatigue in Prostate Cancer
371 Patients Undergoing Androgen Deprivation Therapy: A Year-long Randomised Controlled Trial. *Eur*
372 *Urol.* 2017;**72**:293-299.
- 373 38. Winchester R, Turner LA, Thomas K, Ansley L, Thompson KG, Micklewright D, Gibson AS.
374 Observer effects on the rating of perceived exertion and affect during exercise in recreationally
375 active males. *Percept Mot Skills.* 2012;**115**:213-227.
- 376
- 377
- 378

Variable	Exercise group (n=18)
Age, years	66.9 (13.9)
Gender, number male/female	8/10
Stature, cm	171.1 (11.9)
Body mass, kg	102.1 (29.4)
Ulcer size, cm ² , median (range)	4.9 (1.9 to 136.4)
Duration of ulcer, months, median (range)	5 (1 to 72)
Ankle-brachial index	1.05 (0.14)
Ankle circumference, cm	27.1 (5.5)
Calf circumference, cm	37.3 (7.6)
Comorbidities, n (%)	
Hypertension	7 (39)
History of other CVD	1 (6)
Non-insulin-dependent diabetes	4 (22)
History of cancer	2 (11)
Hypercholesterolemia	1 (6)

Table 1: Exercise group participant characteristics (Values are mean (SD) unless otherwise stated; CVD, cardiovascular disease).

380
381
382
383
384
385
386
387
388
389
390

Variable	Estimated Range <i>(Moderate Intensity 60% - 80%)</i>	Base-Line (n= 18)	Mid-point (n= 13)	Intervention End (n=13)	P value; η^2	Post Hoc (Baseline-Midpoint); Cohen's d	Post Hoc (Baseline-Intervention End); Cohen's d	Post Hoc (Midpoint-Intervention End); Cohen's d
Aerobic Training HR	91-121 (10-13)	103 (14)	107 (12)	112 (18)	0.7; 0.01	0.38; 0.32	0.14; 0.35	0.45; 0.30
Aerobic Training RPE	12-14	12(0)	12(0)	12(0)	0.3; 0.05	0.71; 0.65	0.14; 0.26	0.13; 0.37
Resistance Training RPE	12-14	12(1)	12(0)	12(0)	0.3; 0.05	0.5; 0.25	0.14; 0.62	0.92; 0.38

391
392
393
394
395
396

397

398

399

400

401

402

Table 2: Changes in Exercise Intensity Indices between Exercise Sessions (p<0.05 for Repeated Measures ANOVA and p<0.0167 for post-hoc analysis).

	Base- Line (n=18)	Mid- point (n=13)	Intervention End (n=13)	P value; η^2	Post Hoc (Baseline- Midpoint); Cohen's d)	Post Hoc (Baseline- Intervention End); Cohen's d	Post Hoc (Midpoint- Intervention End); Cohen's d
Aerobic (Min)	19 (8)	26 (5)	29 (3)	<0.01; 0.35	<0.01; 1.10	<0.01; 1.82	0.11; 0.68
Squats	5 (12)	14 (18)	36 (18)	<0.01; 0.42	0.08; 0.64	<0.01; 2.10	<0.01; 1.21
Sit to Stand	12 (10)	29 (17)	36 (19)	<0.01; 0.32	<0.01; 1.21	<0.005; 1.68	0.28; 0.43
Step Ups	14 (15)	24 (18)	31 (22)	0.04; 0.14	0.13; 0.56	<0.01; 0.98	0.29; 0.42
Calf Raises	19 (13)	36 (13)	42 (14)	<0.01; 0.35	<0.01; 1.21	<0.01; 1.62	0.28; 0.43

403
404 Table 3: Changes in Aerobic and Resistance Exercise Indices between Exercise Sessions (p<0.05 for
405 Repeated Measures ANOVA and p<0.0167 for post-hoc analysis).
406
407
408
409
410
411
412
413
414
415
416
417
418
419

Element	Fidelity Element	Percentage of Completion According to Protocol
Aerobic	Duration	81%*
	Intensity	100%
	Duration and Intensity	81%
Resistance	Number of Exercises	62%
	Repetitions	78%
	Sets	73%
	Intensity	91%
Flexibility	Duration	93%
	Number of Exercises	93%

Table 4: Assessment of Exercise Fidelity (* ≥25 minutes of total duration of aerobic exercises).

420
421
422
423