

A simple new method for identifying performance characteristics associated with success in elite tennis

FITZPATRICK, Anna, STONE, Joseph <<http://orcid.org/0000-0002-9861-4443>>, CHOPPIN, Simon <<http://orcid.org/0000-0003-2111-7710>> and KELLEY, John <<http://orcid.org/0000-0001-5000-1763>>

Available from Sheffield Hallam University Research Archive (SHURA) at:
<http://shura.shu.ac.uk/22802/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

FITZPATRICK, Anna, STONE, Joseph, CHOPPIN, Simon and KELLEY, John (2018). A simple new method for identifying performance characteristics associated with success in elite tennis. *International Journal of Sports Science & Coaching*, 14 (1), 43-50.

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

1 **Article type:** Original Research Article

2 **Corresponding Author:**

3 Anna Fitzpatrick, S001 Chestnut Court, Sheffield Hallam University, Collegiate Crescent, Sheffield,
4 S10 2BP, UK. Tel: +44 114 225 2355; E-mail: anna.fitzpatrick@shu.ac.uk.

5 **Article Title:** A simple new method for identifying performance characteristics associated with
6 success in elite tennis.

7 **Authors**

8 Anna Fitzpatrick.¹ Joseph A. Stone.² Simon Choppin.¹ John Kelley.¹

9 ¹Centre for Sports Engineering Research, Sheffield Hallam University, UK

10 ²Academy of Sport and Physical Activity, Sheffield Hallam University, UK

11

12

1 **Abstract**

2 Performance analysis and identifying performance characteristics associated with success are of great
3 importance to players and coaches in any sport. However, while large amounts of data are available
4 within elite tennis, very few players employ an analyst or attempt to exploit the data to enhance their
5 performance; this is partly attributable to the considerable time and complex techniques required to
6 interpret these large datasets. Using data from the 2016 and 2017 French Open tournaments, we tested
7 the agreement between the results of a simple new method for identifying important performance
8 characteristics (the Percentage of matches in which the Winner Outscored the Loser, PWOL) and the
9 results of two standard statistical methods, to establish the validity of the simple method. Spearman's
10 rank-order correlations between the results of the three methods demonstrated excellent agreement,
11 with all methods identifying the same three performance characteristics (*points won of 0-4 rally*
12 *length, baseline points won and first serve points won*) as strongly associated with success.
13 Consequently, we propose that the PWOL method is valid for identifying performance characteristics
14 associated with success in tennis, and is therefore a suitable alternative to more complex statistical
15 methods, as it is simpler to calculate, interpret and contextualise.

16 **Keywords**

17 Elite tennis strategy; clay courts; match statistics; successful performance; winning in tennis

18

1 **Introduction**

2 Performance analysis is of great applied importance to players and coaches in any sport,¹ enabling
3 identification of strengths and areas for improvement,² assessing technical and tactical effectiveness,³
4 opposition analysis,¹ guiding the development of athletes' training programmes,^{4,5} injury
5 rehabilitation,⁶ identifying successful patterns of play^{7,8} and predicting future match outcomes.⁹
6 However, while it is a valued discipline and well-established tool in sports such as soccer¹⁰ and
7 cycling,¹¹ the comparative progress of performance analysis within tennis has been slow.¹² In a 2012
8 poll, tennis was ranked as the second least progressive sport with regards to its use of performance
9 data.¹³ Since then, tennis performance analysis research has advanced, however, with recent studies
10 examining the prediction accuracy of different types of tennis match forecasting models,⁹ comparing
11 the physical demands and performance characteristics of professional tennis to those of the junior
12 game,¹⁴ and developing a comprehensive tennis shot taxonomy based on spatiotemporal data.¹²

13 To continue the development of performance analysis research in tennis, theoretical match
14 investigations, whereby data from multiple matches are used to identify the typical performance
15 characteristics of a sport,¹⁵ could be employed. Theoretical match investigations are helpful for
16 identifying performance characteristics associated with success, monitoring the characteristics of a
17 sport over time, on different playing surfaces and for varying playing positions, as well as
18 investigating the influence of rule changes. For example, a theoretical match investigation in soccer
19 showed that the number of passes and pass success rates have increased over time, but that the
20 magnitudes of these increases depend on playing position; thus, providing benchmark requirements
21 for current elite players.¹⁶ Similarly, Murray et al.¹⁷ found that recent rule changes in elite squash (the
22 new 11 point-per-rally scoring system and reduced tin height) decreased mean match duration and
23 reduced the time players have to perform strokes. Murray et al.¹⁷ also identified that more attacking
24 strategies were adopted by players after these rule changes were implemented. Consequently,
25 implications for training and conditioning had to be reconsidered, to ensure that players' training
26 behaviours were representative of their match-play behaviours.¹⁸

1 Published investigations of elite tennis match-play have demonstrated that the characteristics of
2 performance differ depending on court surface and sex.⁵ Reid et al.⁵ revealed several sex-based
3 differences in match-play characteristics at the Australian Open (i.e. on hard court), and subsequently
4 endorsed sex-specific training designs. In this context, such investigations facilitate a better
5 understanding of the way tennis is played, allowing coaches to better prepare their players for
6 matches.¹⁹ In turn, the identification of performance characteristics associated with success on
7 different surfaces enables the periodisation of training according to court surface, whereby a player's
8 tournament schedule is built around sub-seasons (e.g. the clay court season, the grass court season),
9 which are characterised by surface-specific training methods.¹⁹ For example, if winning a higher
10 percentage of net-points is found to be more closely associated with success on grass courts than on
11 any other surface, this should be reflected in training sessions, with approach shots and net-play
12 afforded more practice time around the grass court season. The serve is generally considered to be the
13 most important stroke in tennis^{20,21} and several studies have described the serve and serve-return as
14 key factors to overall success,^{22,23} but few attempts have been made to objectively identify the
15 performance characteristics most strongly associated with success.

16 With the development of increasingly sophisticated methods for monitoring and recording aspects of
17 tennis match-play performance, data collection has become more prevalent.²⁰ Consequently, huge
18 amounts of data are readily (and freely) available to players. Typically, however, complex data
19 processing and analysis techniques are required to uncover more useful information,²⁴ particularly
20 when the data were recorded using motion tracking systems such as Hawk-Eye (Hawk-Eye
21 Innovations Ltd., Basingstoke, UK). For example, a recent study analysing "big Hawk-Eye data"
22 featured several stages of data cleaning and further processing before a magnus linear model was
23 produced to examine the relationship between serve ball trajectory and winning-point probabilities.²⁰
24 Other studies have also used point-based probabilistic modelling^{25,26} and "common opponent
25 models"²⁷ to predict match outcomes. Even analysing and interpreting relatively simple sports data,
26 such as notational analysis data, requires time and a degree of statistical knowledge. Notational
27 analysis-based studies aiming to identify performance characteristics associated with success can

1 incorporate data from 100+ matches and have employed methods including Pearson's correlation
2 coefficients (with additional log transformations) and stepwise regression procedures (tennis²⁸),
3 Student's *t*-tests (tennis,²⁹ basketball³⁰) discriminant analysis (rugby³¹) and Kruskal-Wallis H
4 (soccer³²).

5 It has become commonplace to employ full time performance analysts in many sports,³³ but analysts
6 are not cheap, so even at tennis' elite level, it is uncommon for players to work with an analyst or
7 attempt to exploit the vast amounts of data available to them.²⁴ Instead, coaches (before and after
8 matches) and players (during matches) typically attempt to fulfil the role of analyst themselves, but
9 this match-by-match approach lacks objectivity and risks neglecting long-term performance
10 development, in favour of identifying short-term solutions. Taking this and the complex analysis
11 methods typically required into account, coaches are not likely to have the time⁴ nor often the desire²⁴
12 to analyse the large quantities of data, or potentially the expertise to transform these large datasets
13 into meaningful interpretations with respect to tennis.³⁴ Therefore, a simple method of identifying
14 performance characteristics associated with success, that is easy to understand and contextualise, may
15 help make performance analysis more accessible and user-friendly for players and coaches. Before a
16 method can be considered appropriate for use within elite tennis, however, it must first be compared
17 to existing methods to assess its validity.

18 The aim of this study was to establish the validity of a simple new method for identifying
19 performance characteristics associated with success in tennis, by testing the agreement between the
20 results of the simple method and results of two standard statistical methods; 1) paired *t*-tests, as used
21 by O'Donoghue²⁹ and Scanlan et al.³⁰ to identify match-play characteristics associated with success in
22 tennis and basketball, respectively, and 2) point-biserial correlations, recently used by Cowden,³⁵ to
23 investigate the association between mental toughness and match outcome in tennis, and Scanlan et
24 al.³⁰ to assess the association between basketball match-play characteristics and match outcome.
25 Analysis was undertaken using sample data from men's and women's elite tennis match-play at the
26 French Open.

1 **Method**

2 *Matches*

3 With institutional ethics approval, performance characteristics for the 2016 and 2017 French Open
4 men's (n = 244) and women's (n = 250) singles matches were obtained from the Roland Garros
5 website.³⁶ All performance characteristics available on the website were included. Incomplete
6 matches (i.e. walkovers, retirements and defaults) were excluded from the study.

7 *Performance characteristics*

8 The following performance characteristics were collected for both players in each match:

- 9 • number of aces, number of double faults
- 10 • number of first serves in
- 11 • average (i.e. mean) first serve speed*
- 12 • number of first serve points won, number of second serve points won
- 13 • number of first serve-return points won, number of second serve-return points won
- 14 • number of baseline points won, number of net points won
- 15 • number of break points won
- 16 • number of winners, number of forced errors, number of unforced errors
- 17 • number of points won of 0-4, 5-8 and 9+ rally length, respectively* .

18 * Collected only for those matches where a serve speed radar was available.

19 Data were classified by match outcome (i.e. winning player or losing player) and normalised using the
20 equations in Table 1, before being reduced to mean values ($\pm sd$).

21 **Table 1. Normalised performance characteristic equations, derived from O'Donoghue and Ingram³⁷ and**
22 **O'Donoghue.³⁸**

Performance characteristic	Equation
Aces (%)	Number of aces/number of serves performed x 100
Double faults (%)	Number of double faults/number of points served x 100
Successful first serves (%)	Number of first serves in/number of first serves attempted x 100
First serve points won (%)	Number of first serve points won/number of first serve points played x 100
First serve-return points won (%)	Number of first serve-return points won/number of first serve-return points played x 100
Second serve points won (%)	Number of second serve points won/number of second serve points played x 100

Second serve-return points won (%)	Number of second serve-return points won/number of second serve-return points played x 100
Break points won (%)	Number of break points won as returner/number of break points played as returner x 100
Net points won (%)	Number of net points won/number of net points played x 100
Baseline points won (%)	Number of baseline points won/number of baseline points played x 100
Winners (%)	Number of winners/number of rally points played x 100
Forced errors (%)	Number of forced errors/number of rally points played x 100
Unforced errors (%)	Number of unforced errors/number of rally points played x 100
Points won of 0-4 rally length (%)	Number of points won of 0-4 rally length/number of points played of 0-4 rally length x 100
Points won of 5-8 rally length (%)	Number of points won of 5-8 rally length/number of points played of 5-8 rally length x 100
Points won of 9+ rally length (%)	Number of points won of 9+ rally length/number of points played of 9+ rally length x 100

1

2 *Data analysis*

3 *Statistical correlation-based method:* Normalised data were imported into in SPSS (v23.0, SPSS Inc,
4 USA). Point-biserial correlations between match outcome and each performance characteristic were
5 calculated, to identify which characteristics were associated with match outcome, for both sexes.

6 *Statistical paired t-test method:* For each performance characteristic, a paired *t*-test was used to
7 compare winning and losing players' data, for each sex, with the simplifying assumption of normality
8 of the winner-loser differences. The *t* values were used to identify the performance characteristics that
9 best distinguished between winning and losing players.²⁹

10 *Proposed method:* For each performance characteristic, the winning player's performance was
11 compared to that of their opponent (i.e. the losing player), to identify which player 'outscored' the
12 other. Then, the *Percentage of matches in which the Winner Outscored the Loser* (PWOL) was
13 calculated for each performance characteristic. For example, if the winning player hit *more* aces than
14 (i.e. outscored) the losing player in 200 out of 250 matches, the PWOL for aces would be 80.0%.
15 Similarly, if the winning player hit *more* unforced errors than (i.e. outscored) the losing player in 100
16 out of 250 matches, the PWOL for unforced errors would be 40%; this would mean that the losing
17 player hit *more* unforced errors than the winning player in 60% of matches. To the authors'
18 knowledge, the method has not previously been applied within sports performance analysis.

1 A PWOL value of 50% for a particular performance characteristic means that players who outscored
 2 their opponent won the match in 50% of cases, indicating no association with match outcome (i.e.
 3 success). As the PWOL value increases towards 100%, this indicates a stronger positive association
 4 with match outcome or success. Correspondingly, as the PWOL value decreases towards 0%, this
 5 indicates a stronger negative association with match outcome (i.e. a stronger association with losing).
 6 Therefore, performance characteristics with either a high PWOL value or a low PWOL value are
 7 considered important in terms of winning, whereas those with PWOL values close to 50% are
 8 considered less important. PWOL values are simple to calculate, with no need for statistical software
 9 packages, such as SPSS. Furthermore, users do not require a comprehensive understanding of data
 10 analysis techniques to apply the method or interpret the results, so it may be more suitable for coaches
 11 and other sports practitioners.

12 The results of each of the three methods were used to indicate the relative importance of each
 13 performance characteristic. To assess the agreement between the results of the methods (i.e. establish
 14 the validity of the PWOL method), pairwise comparisons between the PWOL values, t values and
 15 point-biserial correlation coefficients were performed using Spearman's rank-order correlations.

16 Results

17 Table 2 (men) and Table 3 (women) show the mean and standard deviation of the performance
 18 characteristics for winning and losing players. Table 2 and 3 also show the results of each statistical
 19 method; point-biserial correlations between each performance characteristic and match outcome (r_{pb}),
 20 t values from paired t -tests comparing winning and losing players (t) and the Percentage of matches in
 21 which the Winner Outscored the Loser (PWOL).

22 **Table 2. Men's performance characteristics (presented as mean \pm sd), point-biserial correlations with**
 23 **match outcome, t values and PWOL values (\pm 95% confidence intervals); sorted by r_{pb}**

Performance characteristic	Winning players	Losing players	r_{pb}	t_{dof}	PWOL
Points won of 0-4 rally length (%)	55.2 \pm 5.0%	44.8 \pm 5.0%	0.72	44.47 ₁₈₀	89.0 (\pm 4.6%)
Baseline points won (%)	53.1 \pm 6.5%	42.2 \pm 6.3%	0.65	48.43 ₂₄₃	82.4 (\pm 4.8%)

First serve points won (%)	74.7 ± 8.1%	65.1 ± 8.3%	0.51	45.64 ₂₄₃	85.2 (± 4.5%)
First serve-return points won (%) ⁺	34.9 ± 8.3%	25.3 ± 8.1%	0.51	45.64 ₂₄₃	85.2 (± 4.5%)
Points won of 5-8 rally length (%)	54.4 ± 8.1%	45.6 ± 8.1%	0.48	29.27 ₁₈₀	65.2 (± 6.9%)
Second serve points won (%)	56.5 ± 9.4%	46.2 ± 9.5%	0.47	42.18 ₂₄₃	76.6 (± 5.3%)
Second serve-return points won (%) ⁺	53.8 ± 9.5%	43.5 ± 9.4%	0.47	42.18 ₂₄₃	76.6 (± 5.3%)
Points won of 9+ rally length (%)	55.6 ± 14.7%	44.4 ± 14.7%	0.35	27.57 ₁₈₀	65.7 (± 6.9%)
Winners (%)	18.1 ± 4.7%	15.1 ± 4.4%	0.31	16.67 ₂₄₃	63.9 (± 6.0%)
Break points won (%)	46.1 ± 16.0%	33.7 ± 22.5%	0.30	35.83 ₂₃₀	70.9 (± 5.9%)
Net points won (%)	67.6 ± 13.1%	61.4 ± 13.4%	0.23	22.22 ₂₄₃	61.9 (± 6.1%)
Aces (%)	5.1 ± 4.0%	3.7 ± 2.8%	0.21	10.46 ₂₄₃	59.4 (± 6.2%)
Successful first serves (%)	61.7 ± 7.4%	60.3 ± 7.2%	0.09	6.71 ₂₄₃	55.7 (± 6.2%)
Average first serve speed (km/h)	181.8 ± 9.7 km/h	180.9 ± 10.7 km/h	0.04	3.21 ₁₈₀	50.8 (± 7.3%)
Double faults (%)	3.1 ± 2.1%	3.4 ± 2.2%	-0.08	-3.09 ₂₄₃	43.9 (± 6.2%)
Unforced errors (%)	13.8 ± 4.5%	17.2 ± 5.0%	-0.34	-19.39 ₂₄₃	32.8 (± 5.9%)
Forced errors (%)	16.4 ± 3.3%	19.5 ± 3.9%	-0.40	-22.80 ₂₄₃	22.1 (± 5.2%)

1 ⁺ A player's first (or second) serve-return points won (%) = 100 – opponent's first (or second) serve
2 points won (%), hence identical associations with match outcome.

3 **Note:** Degrees of freedom differ, as some performance characteristics (e.g. first serve speed and rally
4 lengths) are only recorded on some match courts.

5 **Table 3. Women's performance characteristics (presented as mean ± *sd*), point-biserial correlations with**
6 **match outcome, *t* values and PWOL values (± 95% confidence intervals); sorted by *r_{pb}***

Performance characteristic	Winning players	Losing players	<i>r_{pb}</i>	<i>t_{dof}</i>	PWOL
Baseline points won (%)	54.1 ± 7.0%	42.6 ± 7.0%	0.64	49.58 ₂₄₉	83.9 (± 4.6%)
Points won of 0-4 rally length (%)	55.5 ± 6.8%	44.5 ± 6.8%	0.63	39.57 ₁₇₂	84.5 (± 5.4%)
First serve points won (%)	67.5 ± 9.7%	56.1 ± 10.1%	0.50	46.83 ₂₄₉	82.8 (± 4.7%)
First serve-return points won (%) ⁺	43.9 ± 10.1%	32.5 ± 9.7%	0.50	46.83 ₂₄₉	82.8 (± 4.7%)
Points won of 5-8 rally length (%)	54.8 ± 10.8%	45.2 ± 10.8%	0.41	27.25 ₁₇₂	67.8 (± 7.0%)
Second serve points won (%)	50.2 ± 11.0%	40.5 ± 10.9%	0.41	38.78 ₂₄₉	76.4 (± 5.3%)
Second serve-return points won (%) ⁺	59.5 ± 10.9%	49.8 ± 11.0%	0.41	38.78 ₂₄₉	76.4 (± 5.3%)
Winners (%)	18.0 ± 6.1%	13.9 ± 5.2%	0.34	20.47 ₂₄₉	68.0 (± 5.8%)
Points won of 9+ rally length (%)	54.3 ± 16.4%	45.7 ± 16.4%	0.25	19.57 ₁₇₁	55.7 (± 7.4%)
Break points won (%)	53.7 ± 17.3%	43.0 ± 24.9%	0.24	29.64 ₂₄₁	66.0 (± 6.0%)
Aces (%)	2.7 ± 2.6%	1.7 ± 1.8%	0.23	9.47 ₂₄₉	57.2 (± 6.1%)
Net points won (%)	66.8 ± 17.5%	59.9 ± 19.1%	0.18	20.19 ₂₄₉	53.6 (± 6.2%)
Successful first serves (%)	64.5 ± 8.1%	62.8 ± 8.7%	0.10	7.83 ₂₄₉	58.0 (± 6.1%)
Average first serve speed (km/h)	155.3 ± 10.5 km/h	154.7 ± 9.9 km/h	0.03	2.13 ₁₇₂	51.7 (± 7.4%)
Double faults (%)	4.0 ± 3.2%	4.5 ± 3.4%	-0.08	-3.82 ₂₄₉	45.6 (± 6.2%)
Forced errors (%)	14.7 ± 4.1%	17.5 ± 4.0%	-0.26	-12.27 ₂₄₉	34.4 (± 5.9%)
Unforced errors (%)	15.7 ± 5.1%	19.6 ± 6.3%	-0.32	-19.81 ₂₄₉	33.6 (± 5.9%)

7 ⁺ A player's first (or second) serve-return points won (%) = 100 – opponent's first (or second) serve
8 points won (%), hence identical associations with match outcome.

1 **Note: Degrees of freedom differ, as some performance characteristics (e.g. first serve speed and rally**
2 **lengths) are only recorded on some match courts.**

3 Point-biserial correlations, *t* values and PWOL values all identified *points won of 0-4 rally length*,
4 *baseline points won* and *first serve points won* as the performance characteristics most strongly
5 associated with match outcome (i.e. success), for both sexes. *Forced errors* and *unforced errors* were
6 the performance characteristics most negatively associated with match outcome, i.e. associated with
7 losing. Serve-related performance characteristics including *aces*, *double faults*, *successful first serves*
8 and *average first serve speed* were least associated with match outcome.

9 *Agreement between methods*

10 Table 4 displays the Spearman's rank-order correlation coefficients for point-biserial correlation
11 coefficients, *t* values and PWOL values for men's and women's data, respectively.

12 **Table 4. Spearman's rank-order correlation coefficients for point-biserial correlation coefficients, *t***
13 **values and PWOL values, for both sexes.**

Pairwise comparison	Men	Women
r_{pb} and <i>t</i>	0.95	0.95
r_{pb} and PWOL	0.96	0.95
<i>t</i> and PWOL	0.98	0.94

14 **Note: all correlations were significant at $p < 0.001$.**

15 All Spearman's rank-order correlation coefficients demonstrated excellent agreement between the
16 results of the different methods.³⁹

17 **Discussion**

18 The aim of this paper was to establish the validity of a simple new method for identifying
19 performance characteristics associated with success in tennis, by testing the agreement between the
20 results of the simple method and two previously used statistical methods. Spearman's rank-order
21 correlations between results of the new PWOL method and those of two statistical methods (paired *t*-
22 tests and point-biserial correlations) demonstrated excellent agreement (r_s between 0.94 and 0.98),³⁹
23 for men's and women's datasets. These high correlations show that the PWOL method can identify

1 performance characteristics associated with success as effectively as more complex statistical
2 methods; and is therefore a valid method. Accordingly, we suggest elite coaches consider employing
3 the PWOL method, as it is simpler to calculate, interpret and contextualise than standard statistical
4 methods.

5 *Points won of 0-4 rally length* demonstrated the highest PWOL value (89.0%) for men; this
6 corresponded to a point-biserial correlation with match outcome of 0.722 and a t value of 44.47. In
7 simple terms, a PWOL value of 89.0% means that players who won more points of 0-4 rally length
8 than their opponent won the match in almost 9 out of 10 cases. Similarly, in the women's event,
9 *points won of 0-4 rally length* demonstrated the highest PWOL value (84.5%), a point-biserial
10 correlation with match outcome of 0.633 and t value of 39.57. So, players who won more points of 0-
11 4 rally length than their opponent won the match in 84.5% of cases. *Baseline points won* and *first*
12 *serve points won* were also associated with success for men and women, exhibiting PWOL values of
13 above 80%. These values corresponded to moderate point-biserial correlation coefficients and t
14 values. Collectively, these results imply that *points won of 0-4 rally length*, *baseline points won* and
15 *first serve points won* may be considered the three most important performance characteristics at the
16 French Open, with superior performance in these areas closely associated with success for men and
17 women. These three performance characteristics all pertain to 'points won' and often comprise a large
18 proportion of the total points played within a match, so in a tennis context, it may be considered
19 unsurprising that they demonstrated associations with success.

20 For both sexes, *forced errors* and *unforced errors* were the performance characteristics with the
21 lowest PWOL values (between 22.1% and 34.4%), corresponding to negative point-biserial
22 correlation coefficients and negative t values. For example, male players who hit more forced errors
23 than their opponent won the match in only 22.1% of cases and female players who hit more forced
24 errors than their opponent won the match in 34.4% of cases. These results indicate a negative
25 association with match outcome, i.e. an association with losing, and show that hitting fewer forced
26 and unforced errors is advantageous in terms of winning on clay.

1 Serve-related performance characteristics including *aces*, *double faults*, *average first serve speed* and
2 *successful first serves* were among the least associated with match outcome for both sexes, exhibiting
3 PWOL values between 43.9% and 59.4%. The PWOL values for *aces* and *successful first serve*
4 *percentage* were above 50%, demonstrating weak associations with success, whereas the PWOL
5 value for *double faults* was below 50%, demonstrating a weak association with losing. Together,
6 results of the serve-related performance characteristics indicate that a player's serve performance is
7 not closely linked with success on clay courts and may therefore be considered less important in terms
8 of winning.

9 The PWOL method has several advantages compared to standard statistical methods. Calculating the
10 PWOL value of a performance characteristic is straight forward and the process does not require a
11 comprehensive understanding of statistical methods or a software package. A pen, notepad and basic
12 mobile phone calculator (if necessary) are sufficient; to tally the number of matches in which the
13 winner outscored the loser on a relevant performance characteristic and calculate the tallied number of
14 matches as a percentage of total number of matches. Furthermore, the PWOL value of a single
15 performance characteristic can be interpreted in isolation, whereas the result of a paired *t*-test (for
16 example) is more difficult to contextualise, as a series of values are required to gauge relative
17 importance.

18 We propose that analysts adopt the PWOL method to investigate important performance
19 characteristics, as its ease of interpretation means that it can be effectively fed back to coaches
20 directly.⁴⁰ Clearly, representative tournaments should be chosen for analysis, i.e. coaches of elite
21 female players should only consider elite level women's tournaments in their sample, rather than
22 lower level or men's events, and the court surface should also be considered. The PWOL confidence
23 intervals displayed in Table 2 and Table 3 demonstrated that a smaller *n* resulted in a larger
24 confidence interval. In terms of the sample size required to draw valid conclusions using the PWOL
25 method, our results suggest that analysing data from 200 matches will give confidence intervals of
26 approximately $\pm 5\%$ for PWOL values of 80 - 90%. As such, Grand Slam events are ideal for this
27 type of analysis, as performance data from over 100 men's and women's matches are readily and

1 freely available to all players, coaches and analysts at every Grand Slam. Further work is needed to
2 establish whether PWOL can be used on a smaller scale (i.e. using data from fewer matches); if so,
3 the method is such that coaches could adopt it themselves if they so wished. The process would also
4 be efficient with an appropriately designed spreadsheet template.

5 In addition to its simple calculation and interpretation in a coaching context, the PWOL method offers
6 a further benefit compared to standard statistical methods. Paired *t*-tests and point-biserial correlations
7 both consider the magnitude of the differences between winning and losing players' values on a
8 particular performance characteristic, when establishing that characteristic's association with match
9 outcome. In contrast, the PWOL method simply acknowledges the fact that the winning player's value
10 was either higher or lower than the losing player's value, irrespective of the magnitude of the
11 difference. For this reason, the PWOL is more robust than paired *t*-tests and point-biserial correlations
12 in the case of extreme values or outliers. For example, in this study, men's *average first serve speed*
13 exhibited a *t* value of 3.21, which indicates a *significant* difference between winning and losing
14 players' values. However, if we were to remove from the dataset the *two* matches in which the lowest
15 average first serve speeds occurred, the *t* value would decrease to a *non-significant* 1.76, which
16 demonstrates the strong influence of extreme values in a paired *t*-test. While the PWOL may be more
17 robust to outliers (by disregarding the magnitude of the difference between two players' values), it
18 should be noted that in some instances, the magnitude of the difference can offer relevant information.

19 As we are proposing the PWOL method as a more user-friendly alternative for coaches, statistical
20 significance was not incorporated in the results here, as it is unlikely to be relevant to coaches. It is
21 worth highlighting, however, that if performance analysts or other users wish to calculate statistical
22 significance for PWOL values, this can be done using a binomial distribution, with parameters *n* and
23 *p*, where *n* is the sample size and *p* is the probability of, in this case, the winning player outscoring the
24 losing player in a single match.

25 In previous studies attempting to identify performance characteristics associated with success in
26 tennis,^{28,29} methodological differences such as the performance characteristics included in the analyses

1 and their respective calculations appear to have contributed to inconsistent results. For example, rally
2 length statistics have not previously been included, but here, *points won of 0-4 rally length* was most
3 closely associated with success. This study has shown that the PWOL method, correlation-based
4 method and paired *t*-test method demonstrate excellent pairwise agreement. Therefore, if performance
5 characteristics, operational definitions and calculations are consistent between studies in future, we
6 can assume that any differences in the characteristics identified as strongly associated with success are
7 attributable to differences in the context of the performances analysed (e.g. court surface, sex, time
8 etc.), and not differences in the data analysis methods. The list of performance characteristics
9 presented here comprises a more comprehensive selection than those in previous studies. Future
10 research aiming to identify performance characteristics associated with success in tennis should
11 endeavour to incorporate a comprehensive range of characteristics, including rally length statistics.
12 Accordingly, a standardised list of tennis strategy performance characteristics, calculations and
13 definitions would be beneficial. In this context, the PWOL method can be used to help prevent
14 ‘paralysis by analysis’, in that a performance analyst or future studies analysing the sport (e.g. on
15 different surfaces) may employ the method to narrow down or ‘filter’ a full list of performance
16 characteristics and highlight those most strongly associated with success on a particular surface.
17 Subsequently, the filtered list would provide a concise summary of relevant areas for coaches to focus
18 on during training sessions. Additionally, with more sophisticated methods of performance tracking
19 (e.g. Hawkeye Innovations Ltd., Basingstoke, UK) now commonplace in elite tennis,⁵ performance
20 characteristics not typically reported (e.g. distance travelled, average and maximum movement speed)
21 could become more accessible. Future work should investigate the potential use of the PWOL method
22 for new (and currently under-used) performance characteristics.

23 In conclusion, results of the PWOL method demonstrated excellent agreement with results of the
24 point-biserial correlation and paired *t*-test methods. As such, this study has shown that the PWOL
25 method is able to successfully identify performance characteristics associated with success in elite
26 tennis. The method is simple to calculate and does not require statistical software or the expertise of a
27 performance analyst to understand the results; this may encourage players and coaches to begin to

1 engage with performance analysis as a discipline and recognise its potential benefits. Furthermore, the
2 PWOL method is robust in the case of extreme values. We therefore propose the PWOL method as a
3 suitable, more user-friendly alternative to common statistical methods of data analysis in elite tennis.

4 **Acknowledgements**

5 The author(s) would like to acknowledge and thank Roland Garros and Brain Game Tennis for
6 facilitating this study.

7 **Declaration of Conflicting Interests**

8 The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or
9 publication of this article.

10 **Funding**

11 The author(s) received no financial support for the research, authorship, and/or publication of this
12 article.

13 **References**

- 14 1. Groom R, Cushion CJ and Nelson LJ. The delivery of video-based performance analysis by
15 England youth soccer coaches: towards a grounded theory. *J Appl Sport Psychol* 2011; 23: 16-32.
- 16 2. Smith DJ, Norris SR and Hogg JM. Performance evaluation of swimmers. *Sports Med* 2002; 32(9):
17 539-554.
- 18 3. Passos P, Araujo D and Volossovitch A. *Performance Analysis in Team Sports*. Routledge:
19 London, 2017.
- 20 4. Reilly T, Morris T and Whyte G. The specificity of training prescription and physiological
21 assessment: a review. *J Sport Sci* 2009; 27: 575-589.

- 1 5. Reid M, Morgan S and Whiteside D. Matchplay characteristics of Grand Slam tennis: implications
2 for training and conditioning. *J Sport Sci* 2016; 34(19): 1791-1798.
- 3 6. Blobel T, Pfab F, Wanner P, Haser C and Lames M. Healthy Reference Patterns (HRP) supporting
4 prevention and rehabilitation process in professional football. In: World conference on science and
5 soccer, Rennes, France, 31 May-2 June 2017, pp. 183-184.
- 6 7. Hughes M. Computerised notation of racket sports. In: Reilly T, Hughes M and Lees A (eds)
7 *Science and Racket Sports*. London: E & FN Spon, 1995, pp. 249-256.
- 8 8. Unierzyski P and Wieczorek A. Comparison of tactical solutions and game patterns in the finals of
9 two grand slam tournaments in tennis. In: Lees A, Kahn J-F and Maynard I (eds) *Science and Racket*
10 *Sports III*. Oxon: Routledge, 2004, pp. 169-174.
- 11 9. Kovalchik SA. Searching for the GOAT of tennis win prediction. *J Quant Anal Sports* 2016; 12(3):
12 127-138.
- 13 10. Thompson KG. *Pacing Individual Strategies for Optimal Performance*. Champaign: Human
14 Kinetics, 2015.
- 15 11. Trimble L. et al. *Sport in the UK*. Exeter: Learning Matters, 2010.
- 16 12. Kovalchik SA and Reid M. A shot taxonomy in the era of tracking data in professional tennis. *J*
17 *Sport Sci* 2018; 36(18): 2096-2104.
- 18 13. Martin T, Annacone P, O'Shannessy C and Stein M. Tennis Analytics.
19 <http://www.sloansportsconference.com/content/tennis-analytics/>. (2012, accessed 18 July 2018).
- 20 14. Kovalchik SA and Reid M. Comparing matchplay characteristics and physical demands of junior
21 and professional tennis athletes in the era of big data. *J Sport Sci Med* 2017; 16: 489-497.
- 22 15. O'Donoghue P. Match analysis in racket sports. In: Lees A, Kahn J-F and Maynard I (eds)
23 *Science and Racket Sports III*. Oxon: Routledge, 2004, pp.155-162.

- 1 16. Bush M, Barnes C, Archer DT, Hogg B and Bradley PS. Evolution of match performance
2 parameters for various playing positions in the English Premier League. *Hum Mov Sci* 2015; 39: 1-11.
- 3 17. Murray S, James N, Hughes MD, Pers J, Mandeljc R and Vuckovic G. Effects of rule changes on
4 physical demands and shot characteristics of elite-standard men's squash and implications for
5 training. *J Sport Sci* 2017; 34(23): 2170-2174.
- 6 18. Pinder RA, Davids K, Renshaw I and Araujo, D. Representative learning design and functionality
7 of research and practice in sport. *J Sport Exerc Psychol* 2011; 33: 146-155.
- 8 19. Over S and O'Donoghue P. What's the point – tennis analysis and why. *ITF Coach Sport Sci Rev*
9 2008; 15(45): 19-21.
- 10 20. Mecheri S, Rioult F, Mantel B, Kauffmann F and Benguigui N. The serve impact in tennis: first
11 large-scale study of big Hawk-Eye data. *The ASA Data Science Journal* 2016; 9(5): 310-325.
- 12 21. Bollettieri N. *Nick Bollettieri's Tennis Handbook*. 2nd ed. Champaign: Human Kinetics, 2015.
- 13 22. Gillet E, Leroy D, Thouvarecq R and Stein J-F. A notational analysis of elite tennis serve and
14 serve-return strategies on slow surface. *J Strength Cond Res* 2009; 23(2): 532-539.
- 15 23. Klaus A, Bradshaw R, Young W, O'Brien B and Zois J. Success in national level junior tennis:
16 tactical perspectives. *Int J Sports Sci Coach* 2017; 12(5): 618-622.
- 17 24. Briggs S. Why tennis is ready to embrace its inner geek. *The Telegraph*,
18 <https://www.telegraph.co.uk/tennis/2018/03/27/tennis-ready-embrace-inner-geek/> (2018, accessed 27
19 March 2018).
- 20 25. Barnett T and Clarke SR. Combining player statistics to predict outcomes of tennis matches. *IMA*
21 *J Manag Math* 2005; 16(2): 113-120.
- 22 26. Spanias D and Knottenbelt WJ. Predicting the outcomes of tennis matches using a low-level point
23 model. *IMA J Manag Math* 2012; 24(3): 311-320.

- 1 27. Knottenbelt WJ, Spanias D and Madurska AM. A common-opponent stochastic model for
2 predicting the outcome of professional tennis matches. *Comput Math Appl* 2012; 64(12): 3820-3827.
- 3 28. Reid M, McMurtrie D and Crespo M. The relationship between match statistics and top 100
4 ranking in professional men's tennis. *Int J Perform Anal Sport* 2010; 10: 131-138.
- 5 29. O'Donoghue P. Performance models of ladies' and men's singles tennis at the Australian Open.
6 *Int J Perform Anal Sport* 2002; 2(1): 73-84.
- 7 30. Scanlan TA, Teramoto M, Delforce M and Dalbo JV. Do better things come in smaller packages?
8 Reducing game duration slows game pace and alters statistics associated with winning in basketball.
9 *Int J Perform Anal Sport* 2017; 16: 157-170.
- 10 31. Ortega E, Villarejo D and Palao JM. Differences in game statistics between winning and losing
11 rugby teams in the Six Nations tournament. *J Sport Sci Med* 2009; 8(4): 523-527.
- 12 32. Lago-Penas C, Lago-Ballesteros J, Dellal A and Gomez M. Game-related statistics that
13 discriminated winning and losing teams from the Spanish soccer league. *J Sport Sci Med* 2010; 9(2):
14 288-293.
- 15 33. Hughes M and Bartlett R. What is performance analysis? In: Hughes M and Franks IM (eds)
16 *Essentials of Performance Analysis in Sport*. 2nd ed. London: Routledge, 2015, pp. 18-28.
- 17 34. Hughes M and Franks IM. *Notational Analysis of Sport: Systems for Better Coaching and*
18 *Performance in Sport*. 2nd ed. London: Routledge, 2004.
- 19 35. Cowden RG. Competitive performance correlates of mental toughness in tennis: a preliminary
20 analysis. *Percept Mot Skills* 2016; 123: 341-360.
- 21 36. Roland Garros. Completed Matches, www.rolandgarros.com (2017, accessed 30 June 2017).
- 22 37. O'Donoghue P and Ingram B. A notational analysis of elite tennis strategy. *J Sport Sci* 2001;
23 19(2): 107-115.

- 1 38. O'Donoghue P. Normative profiles of sports performance. *Int J Perform Anal Sport* 2005; 5: 104-
2 119.
- 3 39. Hahs-Vaughn DL and Lomax RG. *An Introduction to Statistical Concepts*. 3rd ed. New York:
4 Routledge, 2012.
- 5 40. Sigrist R, Rauter G, Reiner R and Wolf P. Augmented visual, auditory, haptic, and multimodal
6 feedback in motor learning: a review. *Psychon Bull Rev* 2013; 20: 21-53.