

Exercise Training as a Mediator for Enhancing Coronary Collateral Circulation: A Review of the Evidence

NICKOLAY, Thomas <<http://orcid.org/0000-0002-1481-7949>>, NICHOLS, Simon <<http://orcid.org/0000-0003-0377-6982>>, INGLE, Lee and HOYE, Angela

Available from Sheffield Hallam University Research Archive (SHURA) at:
<http://shura.shu.ac.uk/25036/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

NICKOLAY, Thomas, NICHOLS, Simon, INGLE, Lee and HOYE, Angela (2019). Exercise Training as a Mediator for Enhancing Coronary Collateral Circulation: A Review of the Evidence. *Current Cardiology Reviews*, 15.

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

REFERENCES

- [1]
Zimarino M, D'Andreamatteo M, Waksman R, Epstein SE, De Caterina R. The dynamics of the coronary collateral circulation. *Nat Rev Cardiol* 2014; 11(4): 191-7.
[\[http://dx.doi.org/10.1038/nrcardio.2013.207\]](http://dx.doi.org/10.1038/nrcardio.2013.207) [PMID: 24395049]
- [2]
Werner GS, Surber R, Ferrari M, Fritzenwanger M, Figulla HR. The functional reserve of collaterals supplying long-term chronic total coronary occlusions in patients without prior myocardial infarction. *Eur Heart J* 2006; 27(20): 2406-12.
[\[http://dx.doi.org/10.1093/eurheartj/ehi270\]](http://dx.doi.org/10.1093/eurheartj/ehi270) [PMID: 17003048]
- [3]
Cui K, Lyu S, Song X, *et al.* Effect of Coronary Collaterals on Prognosis in Patients Undergoing Primary Percutaneous Coronary Intervention for Acute ST-Segment Elevation Myocardial Infarction: A Meta-Analysis. *Angiology* 2018; 69(9): 803-11.
[\[http://dx.doi.org/10.1177/0003319718768399\]](http://dx.doi.org/10.1177/0003319718768399) [PMID: 29656656]
- [4]
Baroldi G, Mantero O, Scomazzoni G. The collaterals of the coronary arteries in normal and pathologic hearts. *Circ Res* 1956; 4(2): 223-9.
[\[http://dx.doi.org/10.1161/01.RES.4.2.223\]](http://dx.doi.org/10.1161/01.RES.4.2.223) [PMID: 13293825]
- [5]
Fulton WFM. The Time Factor in the Enlargement of Anastomoses in Coronary Artery Disease. *Scott Med J* 1964; 9(1): 18-23.
[\[http://dx.doi.org/10.1177/003693306400900104\]](http://dx.doi.org/10.1177/003693306400900104) [PMID: 14103801]
- [6]
Zoll PM, Wessler S, Schlesinger MJ. Interarterial coronary anastomoses in the human heart, with particular reference to anemia and relative cardiac anoxia. *Circulation* 1951; 4(6): 797-815.
[\[http://dx.doi.org/10.1161/01.CIR.4.6.797\]](http://dx.doi.org/10.1161/01.CIR.4.6.797) [PMID: 14879489]
- [7]
Meier P, Hemingway H, Lansky AJ, Knapp G, Pitt B, Seiler C. The impact of the coronary collateral circulation on mortality: a meta-analysis. *Eur Heart J* 2012; 33(5): 614-21.
[\[http://dx.doi.org/10.1093/eurheartj/ehr308\]](http://dx.doi.org/10.1093/eurheartj/ehr308) [PMID: 21969521]
- [8]
Koerselman, van der Graaf, de Jaegere, Grobbee. Coronary Collaterals. *Circulation* 2003; 107(19): 2507-11.
[PMID: 12756191]
- [9]
Pipp F, Boehm S, Cai WJ, *et al.* Elevated fluid shear stress enhances postocclusive collateral artery growth and gene expression in the pig hind limb. *Arterioscler Thromb Vasc Biol* 2004; 24(9): 1664-8.
[\[http://dx.doi.org/10.1161/01.ATV.0000138028.14390.e4\]](http://dx.doi.org/10.1161/01.ATV.0000138028.14390.e4) [PMID: 15242864]
- [10]
Van Royen N, Piek JJ, Schaper W, Bode C, Buschmann I. Arteriogenesis: mechanisms and modulation of collateral artery development. *J Nucl Cardiol* 2001; 8(6): 687-93.
[\[http://dx.doi.org/10.1067/mnc.2001.118924\]](http://dx.doi.org/10.1067/mnc.2001.118924) [PMID: 11725265]
- [11]
Baeyens N, Bandyopadhyay C, Coon BG, Yun S, Schwartz MA. Endothelial fluid shear stress sensing in vascular health and disease. *J Clin Invest* 2016; 126(3): 821-8.
[\[http://dx.doi.org/10.1172/JCI83083\]](http://dx.doi.org/10.1172/JCI83083) [PMID: 26928035]
- [12]
Schaper. Collateral circulation. *Basic Res Cardiol* 2009; 104(1): 5-21.
[\[http://dx.doi.org/10.1007/s00395-008-0760-x\]](http://dx.doi.org/10.1007/s00395-008-0760-x) [PMID: 19101749]
- [13]
Scholz D, Cai WJ, Schaper W. Arteriogenesis, a new concept of vascular adaptation in occlusive disease. *Angiogenesis* 2001; 4(4): 247-57.
[\[http://dx.doi.org/10.1023/A:1016094004084\]](http://dx.doi.org/10.1023/A:1016094004084) [PMID: 12197469]
- [14]
Seiler C, Stoller M, Pitt B, Meier P. The human coronary collateral circulation: development and clinical importance. *Eur Heart J* 2013; 34(34): 2674-82.
[\[http://dx.doi.org/10.1093/eurheartj/ehi195\]](http://dx.doi.org/10.1093/eurheartj/ehi195) [PMID: 23739241]
- [15]
Carmeliet P. Mechanisms of angiogenesis and arteriogenesis. *Nat Med* 2000; 6(4): 389-95.
[\[http://dx.doi.org/10.1038/74651\]](http://dx.doi.org/10.1038/74651) [PMID: 10742145]
- [16]
Degen A, Millenaar D, Schirmer SH. Therapeutic approaches in the stimulation of the coronary collateral circulation. *Curr Cardiol Rev* 2014; 10(1): 65-72.
[\[http://dx.doi.org/10.2174/1573403X113099990027\]](http://dx.doi.org/10.2174/1573403X113099990027) [PMID: 23721076]
- [17]
Eitenmüller I, Volger O, Kluge A, *et al.* The range of adaptation by collateral vessels after femoral artery occlusion. *Circ Res* 2006; 99(6): 656-62.
[\[http://dx.doi.org/10.1161/01.RES.0000242560.77512.dd\]](http://dx.doi.org/10.1161/01.RES.0000242560.77512.dd) [PMID: 16931799]
- [18]
Qin X, Deng Y, Wu D, Yu L, Huang R. Does Enhanced External Counterpulsation (EECP) Significantly Affect Myocardial Perfusion?: A Systematic Review & Meta-Analysis. *PLoS One* 2016; 11(4): e0151822
[\[http://dx.doi.org/10.1371/journal.pone.0151822\]](http://dx.doi.org/10.1371/journal.pone.0151822) [PMID: 27045935]
- [19]
Gloekler S, Meier P, de Marchi SF, *et al.* Coronary collateral growth by external counterpulsation: a randomised controlled trial. *Heart* 2010; 96(3): 202-7.
[\[http://dx.doi.org/10.1136/hrt.2009.184507\]](http://dx.doi.org/10.1136/hrt.2009.184507) [PMID: 19897461]
- [20]

- Duncker DJ, Bache RJ. Regulation of coronary blood flow during exercise. *Physiol Rev* 2008; 88(3): 1009-86. [<http://dx.doi.org/10.1152/physrev.00045.2006>] [PMID: [18626066](https://pubmed.ncbi.nlm.nih.gov/18626066/)]
- [21]
- Telkova IL, Tepliakov AT. [Relationships between changes of coronary blood flow, energy metabolism of the myocardium, and hyperinsulinemia in patients with ischemic heart disease]. *Kardiologija* 2005; 45(8): 61-8. [PMID: [16091647](https://pubmed.ncbi.nlm.nih.gov/16091647/)]
- [22]
- Magnani S, Roberto S, Sainas G, *et al.* Metaboreflex-mediated hemodynamic abnormalities in individuals with coronary artery disease without overt signs or symptoms of heart failure. *Am J Physiol Heart Circ Physiol* 2018; 314(3): H452-63. [<http://dx.doi.org/10.1152/ajpheart.00436.2017>] [PMID: [29127237](https://pubmed.ncbi.nlm.nih.gov/29127237/)]
- [23]
- Schirmer SH, Degen A, Baumhäkel M, *et al.* Heart-rate reduction by If-channel inhibition with ivabradine restores collateral artery growth in hypercholesterolemic atherosclerosis. *Eur Heart J* 2012; 33(10): 1223-31. [<http://dx.doi.org/10.1093/eurheartj/ehr255>] [PMID: [21821843](https://pubmed.ncbi.nlm.nih.gov/21821843/)]
- [24]
- Gloekler S, Traupe T, Stoller M, *et al.* The effect of heart rate reduction by ivabradine on collateral function in patients with chronic stable coronary artery disease. *Heart* 2014; 100(2): 160-6. [<http://dx.doi.org/10.1136/heartjnl-2013-304880>] [PMID: [24186565](https://pubmed.ncbi.nlm.nih.gov/24186565/)]
- [25]
- Eckstein RW. Effect of exercise and coronary artery narrowing on coronary collateral circulation. *Circ Res* 1957; 5(3): 230-5. [<http://dx.doi.org/10.1161/01.RES.5.3.230>] [PMID: [13427133](https://pubmed.ncbi.nlm.nih.gov/13427133/)]
- [26]
- Scheel KW, Ingram LA, Wilson JL. Effects of exercise on the coronary and collateral vasculature of beagles with and without coronary occlusion. *Circ Res* 1981; 48(4): 523-30. [<http://dx.doi.org/10.1161/01.RES.48.4.523>] [PMID: [7460222](https://pubmed.ncbi.nlm.nih.gov/7460222/)]
- [27]
- Heaps CL, Parker JL. Effects of exercise training on coronary collateralization and control of collateral resistance. *J Appl Physiol* 2011; 111(2): 587-98. [<http://dx.doi.org/10.1152/jappphysiol.00338.2011>] [PMID: [21565987](https://pubmed.ncbi.nlm.nih.gov/21565987/)]
- [28]
- Nolewajka AJ, Kostuk WJ, Rechnitzer PA, Cunningham DA. Exercise and human collateralization: an angiographic and scintigraphic assessment. *Circulation* 1979; 60(1): 114-21. [<http://dx.doi.org/10.1161/01.CIR.60.1.114>] [PMID: [445714](https://pubmed.ncbi.nlm.nih.gov/445714/)]
- [29]
- Niebauer J, Hambrecht R, Marburger C, *et al.* Impact of intensive physical exercise and low-fat diet on collateral vessel formation in stable angina pectoris and angiographically confirmed coronary artery disease. *Am J Cardiol* 1995; 76(11): 771-5. [[http://dx.doi.org/10.1016/S0002-9149\(99\)80224-0](http://dx.doi.org/10.1016/S0002-9149(99)80224-0)] [PMID: [7572652](https://pubmed.ncbi.nlm.nih.gov/7572652/)]
- [30]
- Fujita M, Sasayama S, Asanoi H, Nakajima H, Sakai O, Ohno A. Improvement of treadmill capacity and collateral circulation as a result of exercise with heparin pretreatment in patients with effort angina. *Circulation* 1988; 77(5): 1022-9. [<http://dx.doi.org/10.1161/01.CIR.77.5.1022>] [PMID: [2834115](https://pubmed.ncbi.nlm.nih.gov/2834115/)]
- [31]
- Belardinelli R, Georgiou D, Ginzton L, Cianci G, Purcaro A. Effects of moderate exercise training on thallium uptake and contractile response to low-dose dobutamine of dysfunctional myocardium in patients with ischemic cardiomyopathy. *Circulation* 1998; 97(6): 553-61. [<http://dx.doi.org/10.1161/01.CIR.97.6.553>] [PMID: [9494025](https://pubmed.ncbi.nlm.nih.gov/9494025/)]
- [32]
- Werner GS. The role of coronary collaterals in chronic total occlusions. *Curr Cardiol Rev* 2014; 10(1): 57-64. [<http://dx.doi.org/10.2174/1573403X10666140311123814>] [PMID: [24611646](https://pubmed.ncbi.nlm.nih.gov/24611646/)]
- [33]
- Zbinden R, Zbinden S, Windecker S, Meier B, Seiler C. Direct demonstration of coronary collateral growth by physical endurance exercise in a healthy marathon runner. *Heart* 2004; 90(11): 1350-1. [<http://dx.doi.org/10.1136/hrt.2003.023267>] [PMID: [15486146](https://pubmed.ncbi.nlm.nih.gov/15486146/)]
- [34]
- Zbinden R, Zbinden S, Meier P, *et al.* Coronary collateral flow in response to endurance exercise training. *Eur J Cardiovasc Prev Rehabil* 2007; 14(2): 250-7. [<http://dx.doi.org/10.1097/HJR.0b013e3280565dee>] [PMID: [17446804](https://pubmed.ncbi.nlm.nih.gov/17446804/)]
- [35]
- Möbius-Winkler S, Uhlemann M, Adams V, *et al.* Coronary Collateral Growth Induced by Physical Exercise: Results of the Impact of Intensive Exercise Training on Coronary Collateral Circulation in Patients With Stable Coronary Artery Disease (EXCITE) Trial. *Circulation* 2016; 133(15): 1438-48. [<http://dx.doi.org/10.1161/CIRCULATIONAHA.115.016442>] [PMID: [26979085](https://pubmed.ncbi.nlm.nih.gov/26979085/)]
- [36]
- Pijls NHJ, De Bruyne B, Peels K, *et al.* Measurement of fractional flow reserve to assess the functional severity of coronary-artery stenoses. *N Engl J Med* 1996; 334(26): 1703-8. [<http://dx.doi.org/10.1056/NEJM199606273342604>] [PMID: [8637515](https://pubmed.ncbi.nlm.nih.gov/8637515/)]
- [37]
- Laskey WK. Brief repetitive balloon occlusions enhance reperfusion during percutaneous coronary intervention for acute myocardial infarction: a pilot study. *Catheter Cardiovasc Interv* 2005; 65(3): 361-7. [<http://dx.doi.org/10.1002/ccd.20397>] [PMID: [15945105](https://pubmed.ncbi.nlm.nih.gov/15945105/)]
- [38]
- Crisafulli A, Melis F, Tocco F, *et al.* Exercise-induced and nitroglycerin-induced myocardial preconditioning improves hemodynamics in patients with angina. *Am J Physiol Heart Circ Physiol* 2004; 287(1): H235-42. [<http://dx.doi.org/10.1152/ajpheart.00989.2003>] [PMID: [15016635](https://pubmed.ncbi.nlm.nih.gov/15016635/)]
- [39]

Togni M, Gloekler S, Meier P, *et al.* Instantaneous coronary collateral function during supine bicycle exercise. *Eur Heart J* 2010; 31(17): 2148-55.

[\[http://dx.doi.org/10.1093/eurheartj/ehq202\]](http://dx.doi.org/10.1093/eurheartj/ehq202) [PMID: 20584776]

[40]

Lin S, Lu X, Chen S, *et al.* Human coronary collateral recruitment is facilitated by isometric exercise during acute coronary occlusion. *J Rehabil Med* 2012; 44(8): 691-5.

[\[http://dx.doi.org/10.2340/16501977-0989\]](http://dx.doi.org/10.2340/16501977-0989) [PMID: 22729798]

[41]

Shu W, jing J, Fu LC, *et al.* The relationship between diastolic pressure and coronary collateral circulation in patients with stable angina pectoris and chronic total occlusion. *Am J Hypertens* 2013; 26(5): 630-5.

[\[http://dx.doi.org/10.1093/ajh/hps096\]](http://dx.doi.org/10.1093/ajh/hps096) [PMID: 23391622]

[42]

MacDougall JD, Tuxen D, Sale DG, Moroz JR, Sutton JR. Arterial blood pressure response to heavy resistance exercise. *J Appl Physiol* 1985; 58(3): 785-90.

[\[http://dx.doi.org/10.1152/jappl.1985.58.3.785\]](http://dx.doi.org/10.1152/jappl.1985.58.3.785) [PMID: 3980383]

[43]

McCartney N. Acute responses to resistance training and safety. *Med Sci Sports Exerc* 1999; 31(1): 31-7.

[\[http://dx.doi.org/10.1097/00005768-199901000-00007\]](http://dx.doi.org/10.1097/00005768-199901000-00007) [PMID: 9927007]