

Early transient suppression of immune checkpoint proteins T-cell immunoglobulin mucin-3 and programmed cell death-1 in peripheral blood lymphocytes after blastocyst transfer is associated with successful implantation

ZHANG, Tao, ZHU, Wenyue, ZHAO, Yiwei, CHEUNG, Wing Ching, LIU, Yingyu, CHEN, Xiaoyan, DU, Yan, LEUNG, Kam Tong, CHAN, Yiu Leung, WANG, Chi Chiu, LAIRD, Susan <<http://orcid.org/0000-0003-4020-9020>> and LI, Tin Chiu

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/26006/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

ZHANG, Tao, ZHU, Wenyue, ZHAO, Yiwei, CHEUNG, Wing Ching, LIU, Yingyu, CHEN, Xiaoyan, DU, Yan, LEUNG, Kam Tong, CHAN, Yiu Leung, WANG, Chi Chiu, LAIRD, Susan and LI, Tin Chiu (2020). Early transient suppression of immune checkpoint proteins T-cell immunoglobulin mucin-3 and programmed cell death-1 in peripheral blood lymphocytes after blastocyst transfer is associated with successful implantation. *Fertility and Sterility*, 114 (2), 426-435.

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

1

F & S ____ style revision

2 **Running title: Tim-3 and PD-1 during peri-implantation**

3

4

5

6

7

8

9

10

11

12

13

14

15

1 **Early transient suppression of immune checkpoint proteins**
2 **T-cell immunoglobulin mucin-3 and programmed cell death-1**
3 **in peripheral blood lymphocytes after blastocyst transfer is**
4 **associated with successful implantation**

5 Tao Zhang Ph.D^a, Wenyue Zhu Mphil^a, Yiwei Zhao MBBS^a, Wing Ching
6 Cheung Ph.D^a, Yingyu Liu Ph.D^a, Xiaoyan Chen Ph.D^a, Yan Du DrPH^b, Kam
7 Tong Leung Ph. D^c, Yiu Leung Chan Ph.D^a, Chi Chiu Wang Ph.D^{a,d,e}, Susan
8 Laird Ph.D^f, Tin Chiu Li Ph.D^a

9 ^a Assisted Reproductive Technology Unit, Department of Obstetrics and
10 Gynaecology, Faculty of Medicine, The Chinese University of Hong Kong,
11 Hong Kong, China

12 ^b Office of Clinical Epidemiology, Obstetrics and Gynaecology Hospital of
13 Fudan University, Shanghai, China

14 ^c Department of Pediatrics, The Chinese University of Hong Kong, Hong Kong
15 China

16 ^d Reproduction and development laboratory, Li Ka Shing Institute of Health
17 Sciences, The Chinese University of Hong Kong, Hong Kong, China

18 ^e School of Biomedical Sciences, The Chinese University of Hong Kong, Hong
19 Kong, China

20 ^f Biomolecular Sciences Research Center, Sheffield Hallam University,
21 Sheffield, United Kingdom

22

23 Correspondence:

24 Tin Chiu Li

25 Assisted Reproductive Technology Unit, Department of Obstetrics and
26 Gynaecology, The Chinese University of Hong Kong, Hong Kong, China.

27 Email: tinchiu.li@cuhk.edu.hk

28

29

30

1

2 **Capsule:**

3 Successful implantation is associated with a transient reduction of Tim-3 and
4 PD-1 expression in peripheral lymphocytes on days 3 and 6, but not on day 9,
5 after ET

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

1 **ABSTRACT**

2 Objective: To compare the changing peripheral levels of immune checkpoint
3 proteins Tim-3/Galectin-9 (Gal-9) and PD-1/PL-1 over a nine-day period after
4 blastocyst transfer between women who did and did not conceive.

5 Design: Prospective observational study.

6 Setting: A university teaching hospital.

7 Patients(s): 51 infertile women who were undergoing day-5 blastocyst transfer.

8 Intervention(s): Serial blood samples were obtained on the day of embryo
9 transfer (ET), and 3, 6 and 9 days afterwards for measurement of (1)
10 membranous Tim-3 and PD-1 expression on various peripheral lymphocytes
11 by flow cytometry; and (2) serum concentrations of ligands Gal-9 and PD-L1
12 by ELISA.

13 Main Outcome Measure(s): Membranous Tim-3 and PD-1 expression on
14 lymphocytes and serum Gal-9 and PD-L1 concentrations and comparison of
15 results between pregnant and non-pregnant women.

16 Result(s): In women who conceived, the measurements exhibited three
17 different types of response: (1) a transient and significant reduction of
18 Tim-3⁺NK-like T cells, Tim-3⁺/PD-1⁺CD8⁺ T cells and Tim-3⁺/PD-1⁺CD4⁺ T cells
19 which returned back to baseline level 9 days after ET; (2) a reduction followed
20 by steady increase to above baseline level on day 9 (Tim-3⁺CD56^{dim}NK cells);
21 (3) a steady increase in expression after ET to reach a level significantly higher
22 than that of the baseline by day 9 (Tim-3⁺CD56^{bright}NK cells). Women who did
23 not conceive showed no significant fluctuation in all the parameters measured
24 across the 4 time points.

1

2 Conclusion(s): Successful blastocyst implantation is associated with a
3 reduction of Tim-3 and PD-1 expression in peripheral lymphocytes on days 3
4 and 6, which is no longer apparent on day 9.

5 Key Words: implantation, immune checkpoint, Tim-3, PD-1, natural killer (NK)
6 cells

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

1

2 **INTRODUCTION**

3 Immunological adaptation is a pre-requisite for endometrial preparation which
4 is necessary for successful implantation (1). The uterus becomes receptive
5 during the mid-secretory phase (days 19-23) of the menstrual cycle, commonly
6 known as the window of implantation (WOI). There are considerable changes
7 in endometrial immune cell numbers during WOI, especially the uterine natural
8 killer (uNK) cells, which increase from 40% of the total leukocyte population
9 during the proliferative phase to 60% during WOI and 70% during early
10 pregnancy (2). The exact role of uNK cells is unknown but they may play a role
11 in angiogenesis and decidualization of stromal cells to create a unique
12 immunological microenvironment suitable for the attachment and invasion of
13 embryo expressing paternal antigens (3, 4).

14 However, a limitation of immune cell studies at the maternal fetal interface
15 is that obtaining suitable tissue is invasive and cannot be repeated at regular
16 intervals to observe the dynamic changes which occur as the implantation
17 process progresses. In contrast, peripheral immune parameters may be
18 measured at repeated regular intervals to examine how peripheral immune
19 responses change during the implantation process. Recently, it has been
20 demonstrated that both peripheral and decidual NK cell cytotoxicity is inhibited
21 by T-cell immunoglobulin mucin-3 (Tim-3) which has higher expression in pNK
22 cells in the first trimester of pregnancy (5, 6). Tim-3 has been shown to be an
23 immune checkpoint marker of immune suppression, similar to programmed
24 death-1 (PD-1), the well-known target for immunotherapy in cancer (7).
25 Previous studies have demonstrated that together with Tim-3, PD-1 modulates
26 decidual CD4⁺T and CD8⁺T cells to produce Th2 type cytokines which may

1 lead to maternal-fetal immune tolerance after implantation (8). However, it is
2 uncertain to what extent PD-1 and Tim-3 are involved in immunological
3 adaptation at the very early stages of implantation.

4 The aim of this study was to compare the serial changes, in the
5 expression of Tim-3 and PD-1 in peripheral NK, T and NK-like T cells, and the
6 serum levels of their soluble ligands Galectin-9 (Gal-9) and programmed death
7 -ligand 1 (PD-L1), in the nine days after embryo transfer, between women who
8 did and did not conceive.

9 **MATERIALS AND METHODS**

10 **Subjects**

11 Women undergoing embryo transfer after in-vitro-fertilization (IVF) or
12 intracytoplasmic sperm injection (ICSI) treatment at the Assisted Reproductive
13 Technology Unit, Prince of Wales Hospital, Chinese University of Hong Kong
14 between November 2018 and August 2019 were recruited into the study.

15 The inclusion criteria were women (1) having one or two day 5 blastocysts
16 transferred in fresh or frozen cycles; (2) maternal age 20-40 years old. The
17 exclusion criteria include: (1) hydrosalpinx; (2) structural uterine abnormalities;
18 (3) antiphospholipid syndrome; (3) abnormal thyroid function; (4) polycystic
19 ovarian syndrome; (5) hormonal or metabolic disorders; (6) known clinical
20 autoimmune disease; (7) undergoing immunotherapy; (8) cancer or
21 post-cancer status; (9) significant medical complications.

22 **Patients recruitment**

23 All subjects included in this study were recruited on the day of ET following
24 informed, written consent. They had blood samples taken on the day of ET,

1 and 3, 6, 9 days afterwards. This study has been approved by our Institutional
2 Review Board (CREC Ref: 2014.637).

3 **IVF treatment**

4 Ovarian stimulation was initiated by human menopausal gonadotrophins
5 (HMG) (Pergonal, Serono, Switzerland) or recombinant follicle stimulating
6 hormone (rFSH) (Gonad-F, Serono, Switzerland). The ovulation trigger used
7 was 10,000 units HCG (Profasi, Serono, Switzerland) administered
8 intramuscularly when there were three or more leading follicles reaching 16
9 mm or more in diameter on transvaginal ultrasound. Transvaginal oocyte
10 retrieval was performed 36 h after hCG trigger. Luteal support was
11 commenced in the evening of oocyte retrieval in the form of vaginal
12 progesterone, either a 90 mg daily dose of Crinone (Merck, Germany) or
13 Endometrin (Ferring, Saint-Prex, Switzerland) 100 mg TDS. Blastocysts were
14 transferred on day 5.

15 In the cycle of frozen-thawed embryo transfer, natural cycle and hormonal
16 replacement treatment cycle were monitored with endometrial thickness,
17 ovarian activity and hormonal levels, as previously described (9). Blastocysts
18 were thawed and transferred five days following the estimated day of ovulation
19 or progesterone administration respectively.

20 **Culture system**

21 Cumulus cells were removed from oocytes with hyaluronidase (Vitrolife,
22 Goteborg, Sweden). After denudation, all mature oocytes were incubated in 60
23 µl G-IVF medium for at least an hour in culture, and then fertilized by
24 conventional IVF or inseminated by ICSI. Fertilized oocytes were cultured in
25 G-1 medium (Vitrolife, Goteborg, Sweden) supplemented with 10% serum

1 substitute supplement, (SSS (Irvine Scientific, Santa Ana, USA)). On the
2 morning of day 3 development, each embryo was cleaned and rinsed with G1
3 medium using a 170- μ m pipette (Flexipets, Cook Medical, Bloomington, USA).
4 Each embryo was individually cultured in G-2 medium (Vitrolife, Goteborg,
5 Sweden) supplemented with 10% SSS until blastocyst stage. All embryos in
6 this study were cultured using sequential culture media in 30- μ l microdrops
7 under oil, with incubation conditions of 37.0 °C and 6% CO₂/5% O₂ balanced in
8 N₂ as previously described by our group(10).

9 **Embryo cryopreservation**

10 All blastocysts included in the FET cycles of this study were cryopreserved by
11 vitrification on day 5, using a commercial vitrification kit (Vit Kit® Freeze,
12 90133-SO, Irvine Scientific, Santa Ana, CA, USA) and commercial warming kit
13 (Vit Kit® Thaw, 90137-SO, Irvine Scientific), following the manufacturer's
14 instructions.

15 **Confirmation of pregnancy**

16 All subjects included in this study had serum β -hCG measurement on the
17 blood samples obtained 9 days after ET, according to our routine clinical
18 practice to verify if pregnancy has occurred. Women who had serum β -hCG
19 (<10 mIU/L) were included into the non-conception group. Women who had
20 serum β -hCG (>10 mIU/L) were offered ultrasonography 23 days and 30 days
21 after ET to confirm viability and location of the pregnancy; those who had
22 ongoing intrauterine pregnancy beyond 12 weeks of gestation were included
23 into the conception group. For the purpose of this study, women with
24 biochemical pregnancy loss and miscarriage were excluded because the main
25 aim of this study was to examine the changes of peripheral Tim-3 and PD-1

1 levels in subjects with normal pregnancy as compared to subjects who did not
2 conceive. Biochemical pregnancy was defined as one in which the serum level
3 of β -hCG>10 mIU/L on 9th day after embryo transfer, but no ultrasound
4 evidence of intrauterine pregnancy at 6 weeks of gestation. Miscarriage was
5 defined as one in which there was ultrasound evidence of intrauterine
6 pregnancy but the pregnancy did not progress to beyond 12 weeks of
7 gestation.

8 **Blood sampling**

9 Peripheral blood was collected for serum and lymphocyte isolation. 3 ml whole
10 blood sample was obtained and placed into a tube with serum separator clot
11 activator and EDTA coated tube respectively on the day of blastocyst transfer,
12 and repeated 3, 6 and 9 days after, abbreviated as D0, D3, D6 and D9. Serum
13 was separated after centrifuged at 3000 rpm. at 4°C for 30 min. and stored at
14 -80°C for later analysis.

15 **Laboratory measurements**

16 **1. Peripheral blood mononuclear cell (PBMC) separation**

17 Peripheral blood was diluted 1:1 with 0.9% saline and lymphocytes were
18 isolated by Lymphopure Density Gradient Medium (Biolegend, USA) with
19 centrifuged at 800g for 30 min. After centrifugation, PBMCs formed a defined
20 cell layer at the plasma Lymphopure interface. The layer of PBMCs were
21 collected and washed in 1% BSA/PBS. The cells were centrifuged,
22 resuspended and counted for flow cytometry.

23 **2. Labeling of PBMCs and flow cytometric analysis**

24 1×10^6 PBMCs were resuspended in 100 μ l 1% BSA/PBS per tube and

1 incubated with anti-human CD56-BV605, CD3-FITC, CD4-APC/Cy7,
2 CD8-PE/Cy7 and PD-1-PE or Tim-3-PE purchased from BioLegend (San
3 Diego, CA, USA) for 30 min. in the dark at 4°C. In parallel, each sample was
4 incubated with matched isotype control antibodies. Then cells were washed
5 twice with cold 1% BSA/PBS. The expression of Tim-3 and PD-1 on
6 CD3⁺CD4⁺/CD8⁺T cells, NK-like T (CD3⁺CD56⁺) cells, and CD3⁻CD56^{dim/bright}
7 NK cells were assayed by FACS (BD LSRFortessa, USA) and analyzed by
8 FlowJo version 10 software (Tree Star). A minimum of 100 000 events per
9 sample were collected for phenotypic analysis. As regards flow cytometry
10 gating strategy, total lymphocytes were first gated on a forward scatter (FSC)
11 and side scatter (SSC) and then gated on the CD56^{bright/dim}CD3⁻ population as
12 NK cells, CD56⁺CD3⁺ population as NK-like T cells, CD56⁻CD3⁺ population as
13 T cells which was further gated on the CD8⁺ or CD4⁺ population as CD8⁺ T
14 cells and CD4⁺ T cells. These cells were further gated on Tim-3 and PD-1 for
15 the subsets of interest, namely, Tim3⁺NK cells, Tim-3⁺NK-like T cells, Tim-3⁺
16 CD8⁺ T cells, Tim-3⁺CD4⁺ T cells, PD-1⁺ NK-like T cells, PD-1⁺CD8⁺ T cells
17 and PD-1⁺CD4⁺ T cells.

18 **3. Quantification of PD-L1, Gal-9 in serum**

19 The levels of soluble ligands PD-L1 and Gal-9 in serum collected at the same
20 time points were measured by commercial ELISA kits (R and D Systems,
21 Minneapolis, USA) according to the manufacturer's protocols.

22 **Statistical analysis**

23 The observed immune parameters were expressed as mean±standard error of
24 the mean (SEM). Paired sample t tests or Wilcoxon signed-rank tests were
25 used to compare the change in percentage of lymphocyte subsets and Tim-3,

1 PD-1 levels in lymphocyte subsets at different time-points. The comparison of
2 results between the pregnant and nonpregnant groups was made by using
3 independent sample *t* tests for parametric data or Mann-Whitney *U* test for
4 nonparametric data. The difference in qualitative data was compared by
5 chi-square test. P-values < 0.05 were considered statistically significant in all
6 statistical tests. The statistical analyses were performed using SPSS (Version
7 22; SPSS Inc., New York, NK, USA).

8 **RESULTS**

9 **Demographics**

10 From November 2018 to August 2019, 60 subjects were recruited. Nine
11 subjects were excluded because of biochemical pregnancy loss, miscarriage
12 and ectopic pregnancy. In total, 51 subjects were included in the study, with 33
13 patients in conception group (all singleton) and 18 in non-conception group.
14 The demographic details of these two groups are compared in table 1. The
15 mean (\pm SD) age of women was 34.9 (\pm 3.2) in the conception group and 37.3
16 (\pm 1.8) years ($P < 0.05$) in the non-conception group. Otherwise, there was no
17 difference in baseline FSH, body mass index, duration of infertility, type of
18 infertility, endometrial thickness on the day of hCG trigger and the number of
19 blastocysts transferred between the two groups.

20 **Phenotype analysis of peripheral lymphocytes**

21 We investigated the relative proportion of NK cells ($CD3^-CD56^+$), T cells
22 ($CD3^+CD56^-$) and NK-like T cells ($CD3^+CD56^+$), and the proportion of two NK
23 cell subtypes ($CD56^{dim}$ NK cells and $CD56^{bright}$ NK cells) and two T cell subsets
24 ($CD4^+$ and $CD8^+$ T cells) in peripheral blood lymphocytes on the day of
25 blastocyst transfer, and three, six and nine days after blastocyst transfer in

1 women who did and did not conceive. The proportions of lymphocyte subsets
2 did not alter at any of the time points and there was no significant difference in
3 the results between women who did or did not conceive for all the time points.
4 (Table 2).

5 **Levels of Tim-3 expression after ET**

6 The changing levels of Tim-3 expression in women who conceived are shown
7 in Table 2. On the day of ET, Tim-3 was expressed in 34.82 % of pNK cell
8 ($CD56^{dim}$ NK cells (35.91%) and $CD56^{bright}$ NK cells (25.58%)), 3.37% of T cells
9 ($CD8^+$ T cells (4.14%) and $CD4^+$ T cells (2.91%)) and 3.96% of NK-like T cells.
10 On day 3 after ET, a significant ($p < 0.05$) drop in Tim-3 expression was
11 observed in 4 out of 5 lymphocyte subsets ($CD56^{dim}$ NK cells, $CD8^+$ T cells,
12 $CD4^+$ T cells and NK-like T cells). By day 6 after ET, there was a noticeable
13 reversal of the earlier trend, resulting in a minor increase of Tim-3 expression
14 in these 4 lymphocyte subsets, although at this stage the expression level was
15 still significantly different to the baseline (ET). By day 9 after ET, the increased
16 expression of Tim-3 had continued in all cell types, with Tim-3 expression
17 being significantly higher than the baseline in $CD56^{dim}$ NK cells and
18 $CD56^{bright}$ NK cells.

19 The changing levels of Tim-3 in women who did not conceive across the
20 four time points are also summarized in table 2. There was no significant
21 change in Tim-3 expression among all 5 lymphocyte subsets on all the time
22 points examined with the exception of $CD56^{dim}$ NK and $CD56^{bright}$ NK cells on
23 day 9 after ET, which were significantly (42.17 ± 3.24 , 28.11 ± 3.20) higher than
24 the baseline level (34.49 ± 1.97 , 22.02 ± 1.67).

25 **Comparison of Tim-3 expression between women who did and did not** 26 **conceive**

1 On the day of ET, the percentage of Tim-3⁺pNK cells, Tim-3⁺T cells and
2 Tim-3⁺NK-like T cells were not significantly different between women who did
3 and did not conceive. On day 3 after ET, Tim-3 expression in four of the 5
4 lymphocytes subsets (CD56^{dim}NK cells, CD8⁺T cells, CD4⁺T cells and NK-like
5 T cells) was significantly lower in women who conceived compared to women
6 who did not. There was no difference in CD56^{bright}NK cells. On day 6, a
7 significant decrease in Tim-3 positive cell numbers was seen in CD56^{dim}NK
8 cells, CD4⁺T cells and NK-like T cells, with no difference in CD8⁺T cells and
9 CD56^{bright}NK cells. On day 9, there was no significant difference in Tim-3
10 expression in all lymphocyte subsets (Figure 2A-E).

11

12 **Levels of PD-1 expression after ET**

13 The levels of PD-1 expression in women who conceived after ET are shown in
14 Table 2. On the day of ET, PD-1 was not expressed in either type of pNK cells,
15 but in 26.83% of T cells and 31.11% of NK-like T cells. There were similar
16 proportions of cells expressing PD-1 in the 2 T-cell subtypes: 26.53% of CD8⁺
17 cells and 26.88% of CD4⁺T cells. On day 3 after ET, a significant ($p < 0.05$) drop
18 in PD-1 expression was observed only in CD4⁺T cells. By day 6 after ET, PD-1
19 expression was significantly lower in both CD4⁺T cells and CD8⁺T cells. By day
20 9 after ET, there was a reversal of the decreasing trend, resulting in an
21 increase of PD-1 expression in both CD4⁺T cells and CD8⁺T cells to the
22 baseline level. The levels of PD-1 expression in NK-like T cells were not
23 significantly different at any time point.

24 In the non-pregnant group, there were no significant changes in the
25 percentage of PD-1⁺T cells and PD-1⁺ NK-like T cells at any of the time points
26 examined.

1 **Comparison of PD-1 expression between women who did and did not**
2 **conceive**

3 The percentages of PD-1⁺T cells and PD-1⁺NK-like T cells were not
4 significantly different between women who did and did not conceive at all time
5 points after ET (Figure 2F-H).

6 **Concentration of serum Gal-9 and PD-L1**

7 Figure 2I and 2G show the levels of Gal-9 and PD-L1 in serum of pregnant and
8 non-pregnant women on days 0-9 after ET. There was no difference in levels of
9 either ligand in women who conceived and those who did not.

10

11 **DISCUSSION**

12 **Main findings**

13 In the present study, we found that there is a transient and significant reduction
14 of Tim-3 and PD-1 expression on the cell surface of peripheral lymphocytes
15 within 9 days after blastocyst transfer. The findings suggest that the Tim-3 and
16 PD-1 may be useful immunological biomarkers of successful embryo
17 implantation. In addition, the study results may also help to elucidate
18 immunological mechanisms that facilitate implantation.

19 **Proportion of peripheral NK and T cells after ET**

20 We found that the relative proportion of subsets of peripheral NK and T cells
21 were not changed after ET, regardless of whether conception occurred or not.
22 Earlier studies in the first trimester of pregnancy also showed that the
23 proportion of peripheral NK subsets prior to conception is similar to those in the
24 first trimester of pregnancy (11). Our observation further suggested that there

1 is no alteration in numbers of pNK cell and T cell after ET. Taken together, it
2 suggests that the process of implantation does not significantly alter the
3 proportion of lymphocyte subsets in peripheral blood.

4 **Changing levels of Tim-3 after ET**

5 We observed a significant but transient decrease in Tim-3 expression in
6 CD56^{dim}NK cells, T cells and NK-like T cells in the nine days after ET.
7 Measurement beyond day 9 after ET were not carried out in our study, but
8 earlier reports suggested that the level of expression of Tim-3 in the peripheral
9 NK cells remained elevated in the first trimester of pregnancy (6). In addition,
10 Tim-3 was strongly expressed in decidual NK and T cells (5, 8). Tim-3 was
11 originally identified as a surface marker for interferon- γ (IFN- γ)-producing
12 CD4⁺ T helper 1 (Th1) specific cell and CD8⁺ T cytotoxic (Tc) T cell whose role
13 is to suppress inflammatory responses and to facilitate the development of
14 immunological tolerance through Gal-9 engagement (12, 13). Later studies
15 have shown that Tim-3 is expressed on other innate immune cells, such as NK
16 cells, monocyte-macrophages, and dendritic cells and may regulate their
17 differentiation and functions (14). The abundance of Tim-3 at the site of
18 embryo implantation has led to the hypothesis that the anti-inflammatory
19 environment mediated by Tim-3 is a pre-requisite for the establishment of a
20 successful pregnancy.

21 The unexpected finding in our study is a significant and transient reduction
22 in Tim-3 expression after ET. The exact biological mechanism for such a
23 biphasic response is not clear, but supports the hypothesis that the initial
24 phase of embryo implantation is associated with a pro-inflammatory response
25 (1, 15). The implantation site is characterized by the presence of inflammatory
26 cytokines interleukin-6 (IL-6), IL-8, IL-15, granulocyte-macrophage

1 colony-stimulating factor (GM-CSF), CXC chemokine ligand 1(CXCL1),
2 CC-chemokine ligand 4 (CCL4), osteopontin and tumor necrosis factor (TNF)
3 which are derived from both endometrial stromal cells and infiltrated immune
4 cells (15). Precisely coordinated interplay between immune cells, cytokines
5 and hormones is crucial for the tissue remodeling required for successful
6 embryo implantation (16). The transient reduction of Tim-3 may indicate that in
7 peripheral blood the pro-inflammatory response occurs after ET followed by
8 anti-inflammatory response associated with the increase in Tim-3.

9 **Changing levels of PD-1 after ET**

10 PD-1 expression was not seen in NK cell, but was expressed by T cells and
11 NK-like T cells. The changing pattern of PD-1 expression after ET in T-cell and
12 NK like T cells was similar to that of Tim-3, but with subtle differences. There
13 was a reduction in expression among all three lymphocytes subsets 3-days
14 after ET, but the reduction was significant in only CD4⁺T cells. On day 6 after
15 ET there was a significant reduction of PD-1 expressed in both CD4⁺T cells
16 and CD8⁺T cells. By day 9 after ET, the expression level had returned back to
17 baseline level in all three subsets. PD-1 in decidual T cells plays a critical role
18 in the maintenance of maternal tolerance by interacting with PD-L1 expressed
19 on stromal cells (17), trophoblast cells (18), and antigen presenting cells (19).
20 Moreover, PD-1 is involved in the T-cell homeostasis, namely the
21 Th1/Th2/Th17/Tregs immune balance at the maternal fetal interface (20, 21).
22 Given that both Tim-3 and PD-1 are recognized to be anti-inflammatory
23 mediators (13, 22); the similar changing pattern after ET supports the
24 hypothesis that a pro-inflammatory response may be a pre-requisite of early
25 embryo implantation. However, whether this is also reflected at the fetal
26 endometrial interface is unknown.

1 **Tim-3 and PD-1 results in women who did not conceive**

2 In contrast to women who conceived, the expression of Tim-3 and PD-1 in
3 various lymphocyte subsets in women who did not conceive did not show any
4 significant changes 3, 6 and 9 days after ET, with the exception of increased
5 Tim-3 expression on CD56^{dim}NK cells 9 days after ET. This observed increase
6 in CD56^{dim}NK cells may be a consequence of progesterone therapy used for
7 luteal support, as it has been reported that physiological concentrations of
8 progesterone induces Tim-3 expression on pNK cells via the IL-4-STAT6
9 pathway, either directly or indirectly (6).

10 There was no difference in expression of PD-1 between subjects who did
11 and did not conceive in all lymphocyte subsets on all the days examined. The
12 observation suggests the change in the expression of Tim-3 associated with
13 successful implantation is more pronounced than that of PD1.

14 **Levels of ligands (Gal-9 & PD-L1)**

15 In contrast to the changes in immune inhibitory receptors Tim-3 and PD-1
16 expression in lymphocyte subsets, there were no observable changes in the
17 serum concentration of their respective ligands Gal-9 and PD-L1 3, 6 and 9
18 days after ET, and no difference in the levels between women who did or did
19 not conceive. The findings suggest that the *initial phase* of immune modulation
20 during pregnancy is primarily, if not entirely, mediated via immune inhibitory
21 receptors but not their ligands. This is in contrast to the finding of an earlier
22 study which showed that maternal blood level of Gal-9 and PD-L1 were
23 significantly elevated after confirmation of pregnancy (equivalent to nine days
24 after ET), which might be caused by their secretion from trophoblasts (23).
25 The later observation represents the second phase of immunomodulation in
26 pregnancy. Gal-9 and PD-L1 are expressed by both haemopoietic and

1 non-haemopoietic cells (24, 25). The source of these ligands during pregnancy
2 might be primarily from trophoblasts since high level of both proteins are
3 expressed in the placenta (23). Based on animal studies, the function of these
4 two ligands are proposed to mediate maternal immune tolerance though
5 binding to immune cells expressing Tim-3 or PD-1(8). Prospectively planned
6 clinical studies are required to confirm their roles in humans.

7 **Strength of current study**

8 A particular strength of this study is that serial measurements were obtained at
9 several precisely timed points after ET in a population undergoing the same
10 treatment. Many earlier studies on immune-modulation of pregnancy involved
11 a single time point (26, 27), not as precisely timed, usually in terms of
12 gestational weeks rather than days (23, 28). In addition, the observations in
13 this study were made in the very early stage of embryo implantation, well
14 before it is known whether or not the implantation is successful, which enabled
15 us to study the initial, transient phase of immunomodulation of pregnancy.
16 Thirdly, we have included in our study two cohorts, one cohort with successful
17 implantation leading to ongoing pregnancy and the other cohort with failure of
18 implantation leading to non-conception. The latter cohort serves as a
19 comparison group with which the observations in the conception cohort could
20 be meaningfully compared. Finally, we have also obtained baseline
21 measurement prior to the replacement of embryos in both cohorts, with which
22 any changes following ET could be compared and quantified; moreover, the
23 observation that the relevant baseline immunological parameters are the same
24 between those with success or failure of implantation suggest that these
25 parameters are unlikely to be confounding factors.

26 **Limitations of current study**

1 Whilst we have found that successful embryo implantation is associated with a,
2 transient and significant response in the maternal peripheral blood immune
3 system, it is not possible to conclude in this study if the lack of a normal
4 maternal immune response is the cause or consequence of the implantation
5 failure; further well-designed prospective studies are required. A further
6 limitation of the study is that the measurements in peripheral blood may not
7 reflect what is happening at the site of implantation. Other limitations of this
8 study include the relatively small sample size, which did not enable as to
9 analyze the possible impact of various confounding variables such as patient
10 population (different underlying causes of infertility), the type of embryo
11 transfer cycle (fresh or frozen), the methods of endometrium preparation
12 (natural or artificial) and embryo quality.

13 **Clinical implication**

14 The significant and transient reduction of Tim-3 and PD-1 was detected on day
15 3 after ET, but the changes reverted back to normal or above baseline level by
16 the 9th day after ET. It implies that in order to ascertain if there is an underlying
17 immune cause of implantation failure, serum markers should be measured as
18 soon as an embryo is replaced; doing the measurement after the pregnancy
19 test result is known (9 days after ET) is already too late. The monitoring of any
20 proposed immune therapy to improve embryo implantation should also
21 commence prior to confirmation of a pregnancy test result.

22 **CONCLUSION**

23 To conclude, we have found that successful embryo implantation is
24 characterized by a biphasic immunological response in peripheral blood
25 lymphocytes, with a transient and significant reduction in Tim-3 and PD-1
26 expression, signifying a pro-inflammatory response 3-6 days after ET, followed

1 by a rebound increase in Tim-3 expression consistent with the beginning of a
2 second phase of anti-inflammatory response.

3 **Conflict of interest:**

4 The authors declare no financial or commercial conflict of interest.

5 **References**

- 6 1. Mor G, Aldo P, Alvero AB. The unique immunological and microbial aspects of pregnancy.
7 Nat Rev Immunol 2017;17:469-82.
- 8 2. Tang AW, Alfirevic Z, Quenby S. Natural killer cells and pregnancy outcomes in women
9 with recurrent miscarriage and infertility: a systematic review. Hum Reprod 2011;26:1971-80.
- 10 3. Chen X, Man GCW, Liu Y, Wu F, Huang J, Li TC *et al.* Physiological and pathological
11 angiogenesis in endometrium at the time of embryo implantation. Am J Reprod Immunol
12 2017;78:e12693.
- 13 4. Zhang Y, Wang Y, Wang XH, Zhou WJ, Jin LP, Li MQ. Crosstalk between human
14 endometrial stromal cells and decidual NK cells promotes decidualization in vitro by
15 upregulating IL25. Mol Med Rep 2018;17:2869-78.
- 16 5. Li YH, Zhou WH, Tao Y, Wang SC, Jiang YL, Zhang D *et al.* The Galectin-9/Tim-3
17 pathway is involved in the regulation of NK cell function at the maternal-fetal interface in early
18 pregnancy. Cell Mol Immunol 2016;13:73-81.

- 1 6. Li Y, Zhang J, Zhang D, Hong X, Tao Y, Wang S *et al.* Tim-3 signaling in peripheral NK
2 cells promotes maternal-fetal immune tolerance and alleviates pregnancy loss. *Sci Signal*
3 2017;10.
- 4 7. Koyama S, Akbay EA, Li YY, Herter-Sprie GS, Buczkowski KA, Richards WG *et al.*
5 Adaptive resistance to therapeutic PD-1 blockade is associated with upregulation of alternative
6 immune checkpoints. *Nat Commun* 2016;7:10501.
- 7 8. Wang S, Zhu X, Xu Y, Zhang D, Li Y, Tao Y *et al.* Programmed cell death-1 (PD-1) and
8 T-cell immunoglobulin mucin-3 (Tim-3) regulate CD4+ T cells to induce Type 2 helper T cell
9 (Th2) bias at the maternal-fetal interface. *Hum Reprod* 2016;31:700-11.
- 10 9. Law TSM, Cheung WC, Wu F, Zhang R, Chung JPW, Wang CC *et al.* Endometrial
11 Vascularization Characterized by Optical Coherence Tomography and Immunohistochemistry
12 in Women Undergoing In Vitro Fertilization-Embryo Transfer Treatment. *Medicina (Kaunas)*
13 2019;55:81.
- 14 10. Yeung QSY, Zhang YX, Chung JPW, Lui WT, Kwok YKY, Gui B *et al.* A prospective study
15 of non-invasive preimplantation genetic testing for aneuploidies (NiPGT-A) using
16 next-generation sequencing (NGS) on spent culture media (SCM). *J Assist Reprod Genet*
17 2019;36:1609-21.
- 18 11. Kwak JY, Kwak FM, Ainbinder SW, Ruiz AM, Beer AE. Elevated peripheral blood natural
19 killer cells are effectively downregulated by immunoglobulin G infusion in women with
20 recurrent spontaneous abortions. *Am J Reprod Immunol* 1996;35:363-9.

- 1 12. Monney L, Sabatos CA, Gaglia JL, Ryu A, Waldner H, Chernova T *et al.* Th1-specific cell
2 surface protein Tim-3 regulates macrophage activation and severity of an autoimmune
3 disease. *Nature* 2002;415:536-41.
- 4 13. Sanchez-Fueyo A, Tian J, Picarella D, Domenig C, Zheng XX, Sabatos CA *et al.* Tim-3
5 inhibits T helper type 1-mediated auto- and alloimmune responses and promotes
6 immunological tolerance. *Nat Immunol* 2003;4:1093-101.
- 7 14. Han G, Chen G, Shen B, Li Y. Tim-3: an activation marker and activation limiter of innate
8 immune cells. *Front Immunol* 2013;4:449.
- 9 15. Mor G, Cardenas I, Abrahams V, Guller S. Inflammation and pregnancy: the role of the
10 immune system at the implantation site. *Ann N Y Acad Sci* 2011;1221:80-7.
- 11 16. Zenclussen AC, Hammerling GJ. Cellular Regulation of the Uterine Microenvironment
12 That Enables Embryo Implantation. *Front Immunol* 2015;6:321.
- 13 17. Nagamatsu T, Schust DJ, Sugimoto J, Barrier BF. Human decidual stromal cells suppress
14 cytokine secretion by allogenic CD4+ T cells via PD-1 ligand interactions. *Hum Reprod*
15 2009;24:3160-71.
- 16 18. Holets LM, Hunt JS, Petroff MG. Trophoblast CD274 (B7-H1) is differentially expressed
17 across gestation: influence of oxygen concentration. *Biol Reprod* 2006;74:352-8.
- 18 19. Sayama S, Nagamatsu T, Schust DJ, Itaoka N, Ichikawa M, Kawana K *et al.* Human
19 decidual macrophages suppress IFN-gamma production by T cells through costimulatory
20 B7-H1:PD-1 signaling in early pregnancy. *J Reprod Immunol* 2013;100:109-17.

- 1 20. Guleria I, Khosroshahi A, Ansari MJ, Habicht A, Azuma M, Yagita H *et al.* A critical role for
2 the programmed death ligand 1 in fetomaternal tolerance. *J Exp Med* 2005;202:231-7.
- 3 21. Shepard MT, Bonney EA. PD-1 regulates T cell proliferation in a tissue and
4 subset-specific manner during normal mouse pregnancy. *Immunol Invest* 2013;42:385-408.
- 5 22. Francisco LM, Sage PT, Sharpe AH. The PD-1 pathway in tolerance and autoimmunity.
6 *Immunol Rev* 2010;236:219-42.
- 7 23. Enninga EAL, Harrington SM, Creedon DJ, Ruano R, Markovic SN, Dong H *et al.* Immune
8 checkpoint molecules soluble program death ligand 1 and galectin-9 are increased in
9 pregnancy. *Am J Reprod Immunol* 2018;79.
- 10 24. Freeman GJ, Casasnovas JM, Umetsu DT, DeKruyff RH. TIM genes: a family of cell
11 surface phosphatidylserine receptors that regulate innate and adaptive immunity. *Immunol*
12 *Rev* 2010;235:172-89.
- 13 25. Kowanz M, Zou W, Gettinger SN, Koeppen H, Kockx M, Schmid P *et al.* Differential
14 regulation of PD-L1 expression by immune and tumor cells in NSCLC and the response to
15 treatment with atezolizumab (anti-PD-L1). *Proc Natl Acad Sci USA* 2018;115:E10119-E26.
- 16 26. Wang Q, Li TC, Wu YP, Cocksedge KA, Fu YS, Kong QY *et al.* Reappraisal of peripheral
17 NK cells in women with recurrent miscarriage. *Reprod Biomed Online* 2008;17:814-9.
- 18 27. Katano K, Suzuki S, Ozaki Y, Suzumori N, Kitaori T, Sugiura-Ogasawara M. Peripheral
19 natural killer cell activity as a predictor of recurrent pregnancy loss: a large cohort study. *Fertil*
20 *Steril* 2013;100:1629-34.

1 28. Mosimann B, Wagner M, Shehata H, Poon LC, Ford B, Nicolaides KH *et al.* Natural killer
2 cells and their activation status in normal pregnancy. *Int J Reprod Med* 2013;2013:906813.

3

4

5 **Figure Legends**

6 Figure 1. Differential Tim-3 and PD-1 expression in pNK, T and NK-like T cells.
7 (A-C) Gating strategy of peripheral NK cells (CD3⁻CD56⁺), T cells (CD3⁺CD56⁻),
8 NK-like T cells (CD3⁺CD56⁺), CD8⁺T cells (CD3⁺CD56⁻CD8⁺ CD4⁻) and CD4⁺T
9 cells (CD3⁺CD56⁻CD4⁺CD8⁻). (D-G) Gating strategy and representative plots
10 of flow cytometric analysis of Tim-3⁺NK cells, Tim-3⁺NK-like T cells,
11 Tim-3⁺CD4⁺T cells and Tim-3⁺CD8⁺T cells. (H-K) Gating strategy and
12 representative plots of flow cytometric analysis of PD-1⁺NK cells, PD-1⁺NK-like
13 T cells, PD-1⁺CD4⁺T cells and PD-1⁺CD8⁺T cells.

14 Figure 2. Comparison of Tim-3 and PD-1 expression on peripheral NK subsets,
15 NK-like T and T cell subtypes between women who did and did not conceive
16 over a nine-day period after ET. (A) Tim-3⁺CD56^{dim}NK, (B) Tim-3⁺CD56^{bright}NK
17 cells, (C) Tim-3⁺CD8⁺T cells, (D) Tim-3⁺CD4⁺T cells, (E) Tim-3⁺NK-like T cells,
18 (F) PD-1⁺CD8⁺T cells, (G) PD-1⁺CD4⁺T cells, (H) PD-1⁺NK-like T cells.
19 Significant difference was found in (A), (D) and (E) at three and six days after
20 ET between two groups. The percentage of (C) was significantly lower in
21 pregnant women compared with non-pregnant women at three days after ET.
22 No significant difference was found in (B, F-H). Asterixis indicate comparison
23 which showed a significantly different result, whereas if the asterixis is not
24 shown, it means the results were not significantly different. The concentrations

1 of circulating Gal-9 (I) and PD-L1 (J) were compared between pregnant and
2 non-pregnant women at serial time points. No significant difference in
3 concentration of serum Gal-9 and PD-L1 was observed within groups and
4 between two groups.

5