

**Symptoms of Overtraining in Resistance Exercise:  
International Cross-Sectional Survey**

GRANDOU, Clementine, WALLACE, Lee, COUTTS, Aaron J, BELL, Lee  
<<http://orcid.org/0000-0003-0583-3522>> and IMPELLIZZERI, Franco M

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/26713/>

---

This document is the Accepted Version [AM]

**Citation:**

GRANDOU, Clementine, WALLACE, Lee, COUTTS, Aaron J, BELL, Lee and IMPELLIZZERI, Franco M (2020). Symptoms of Overtraining in Resistance Exercise: International Cross-Sectional Survey. *International Journal of Sports Physiology and Performance*, 1-10. [Article]

---

**Copyright and re-use policy**

See <http://shura.shu.ac.uk/information.html>



**Symptoms of Overtraining in Resistance Exercise:  
International Cross-Sectional Survey**

Journal:	<i>International Journal of Sports Physiology and Performance</i>
Manuscript ID	IJSPP.2019-0825.R2
Manuscript Type:	Original Investigation
Date Submitted by the Author:	19-Feb-2020
Complete List of Authors:	Grandou, Clementine; University of Technology Sydney, Sport & Exercise Discipline Group, Faculty of Health Wallace, Lee; University of Technology Sydney, Sport & Exercise Discipline Group, Faculty of Health Coutts, Aaron; University of Technology Sydney, Sport & Exercise Discipline Group, Faculty of Health Bell, Lee; Sheffield Hallam University, Academy for Sport and Physical Activity, Faculty of Health and Wellbeing Impellizzeri, Franco; University of Technology Sydney, Sport & Exercise Discipline Group, Faculty of Health
Keywords:	fatigue, overreaching, overtraining, resistance training, strength training

SCHOLARONE™  
Manuscripts

## 1 ABSTRACT

2

3 **Purpose:** To provide details on the nature and symptomatic profile of training maladaptation in  
4 competitive resistance-based athletes to examine whether there are symptoms that may be used as  
5 prognostic indicators of overtraining. Identifying prognostic tools to assess for training maladaptation  
6 is essential for avoiding severe overtraining conditions.

7

8 **Methods:** A web-based survey was distributed to a cross-sectional convenience sample of competitive  
9 athletes involved in sports with a significant resistance training component. The 46 item anonymous  
10 survey was distributed via industry experts and social media from July-August 2019.

11

12 **Results:** The final sample included 605 responses (completion rate: 84%). Seventy-one percent of  
13 respondents indicated that they had previously experienced an unexplained decrease in performance.  
14 Among those, the majority reported a performance decrement lasting from 1 week – 1 month (43.8%).  
15 General feelings of fatigue were the most frequent self-reported symptom of maladaptation. Acute  
16 training maladaptation, lasting <1 month, was also accompanied by symptoms of musculoskeletal aches  
17 and pain. In the majority of cases (92.5%) training maladaptation was accompanied by additional non-  
18 training stressors. A greater proportion of respondents with more severe maladaptation (>4 months)  
19 were training to muscular failure.

20

21 **Conclusion:** The results from this study support the multifactorial nature of training maladaptation. The  
22 multidimensional nature of fatigue and individual variability in symptomatic responses precludes  
23 definitive prognostic symptoms or differential diagnostic factors of functional/non-functional  
24 overreaching or the overtraining syndrome in resistance exercise.

25

26 **Keywords:** *fatigue, overreaching, overtraining, resistance training, strength training*

## 27 INTRODUCTION

28

29 Resistance training is often undertaken by athletes who strive to improve muscular strength,  
30 hypertrophy and power.<sup>1</sup> It is well-accepted that resistance training can enhance an athlete's ability to  
31 perform general sporting skills (e.g. jumping and sprinting) which may result in superior competition  
32 performance during sport specific tasks.<sup>2</sup> However, designing an optimal resistance training program is  
33 a complex process that involves the careful manipulation of several training variables (e.g. training load,  
34 volume, frequency, rest periods and exercise selection).<sup>3</sup> Intensified resistance training in combination  
35 with inadequate recovery can result in a decline in performance with or without related physiological  
36 and/or psychological signs and symptoms.<sup>4</sup> Resulting maladaptive conditions may include functional  
37 overreaching (FOR), non-functional overreaching (NFOR) or the overtraining syndrome (OTS). The  
38 well-accepted definitions of Meeusen et al<sup>5</sup> suggest that the differential diagnosis of these conditions is  
39 based on the time required for performance restoration. Additionally, it has been suggested that  
40 complete performance restoration may not ever be possible with the OTS.<sup>2</sup> However, as current  
41 diagnostic criteria has been established through the study of overtraining in endurance activities it is  
42 unknown whether these definitions are compatible with such conditions in resistance training.

43

44 The prevention and early diagnosis of maladaptive conditions is critical as there are no firmly  
45 established therapeutic agents (other than rest) capable of reversing the detrimental effects of  
46 overtraining.<sup>5</sup> Many previous studies have searched for prognostic and diagnostic markers that may be  
47 used to determine the onset or presence of overtraining (for review, see Meeusen et al<sup>5</sup>). It has  
48 previously been established that considerable variability exists in the way that individuals respond to  
49 resistance exercise stress.<sup>6</sup> Contextual factors and non-training stressors such as environmental,  
50 physical and/or emotional stressors, including insufficient calorie intake, pressure to perform,  
51 inadequate nutrition, insufficient sleep or excessive socio-economic concerns may result in large  
52 interindividual variability of stress symptoms.<sup>7, 8</sup> In a recent study designed to examine overtrained  
53 athletes, Cadejani et al<sup>9</sup> identified a unique combination of clinical and biochemical manifestations in  
54 each individual affected. Furthermore, this study demonstrated a relationship between additional non-  
55 training stressors or contextual factors and susceptibility to overtraining. Additionally, in a synopsis of  
56 previous literature Fry et al<sup>10</sup> listed 84 major symptoms and manifestations of overtraining in a variety  
57 of sports. Determining which signs and symptoms to monitor from such an extensive list continues to  
58 challenge both coaches and athletes.

59

60 Estimates of the prevalence of overtraining have varied widely depending on the authors' definition of  
61 overtraining, the population and the study methodology. It has previously been suggested that  
62 endurance- and resistance-trained athletes respond differently to training stress.<sup>4</sup> Early studies imply  
63 that athletes involved in primarily anaerobic activities may be more susceptible to overtraining than

64 endurance based athletes.<sup>11, 12</sup> However, much of the early overtraining literature in resistance exercise  
65 was limited to anecdotal accounts with inconsistent use of definitions making quantification of  
66 prevalence difficult.<sup>12</sup> Additionally, majority of previous overtraining studies have included only male  
67 participants.<sup>13</sup> At present, information concerning overtraining prevalence, potential mechanisms and  
68 symptomatology in resistance-trained athletes is scarce.

69

70 Many previous studies that have investigated the mechanisms and manifestation of overtraining in  
71 resistance exercise have failed to appropriately establish FOR, NFOR or OTS (for review, see Grandou  
72 et al<sup>13</sup>). Therefore, the objective of the present exploratory study is to identify possible prognostic  
73 symptoms of training maladaptation in resistance exercise. Determining the point at which training  
74 becomes maladaptive is of key practical significance for athletes and coaches. At present, the correct  
75 diagnosis of maladaptive conditions can only be made retrospectively. Therefore, if prognostic  
76 symptoms of overtraining can be identified, remedial reductions in training stress can be implemented  
77 and training maladaptation may be avoided.

78

## 79 **METHODS**

80

81 An open international survey was used to identify the symptoms of unexplained training maladaptation  
82 amongst competitive athletes in resistance-based sports. Detailed methods according to the Checklist  
83 for Reporting Results of Internet E-Surveys (CHERRIES)<sup>14</sup> are available in Table S1. This study was  
84 approved by the Human Research Ethics Committee of the University of Technology Sydney (ETH19-  
85 3898).

86

### 87 **Survey Development**

88

89 An anonymous survey was developed on REDCap (Research Electronic Data Capture software version  
90 8.11.3 – University of Technology Sydney), a secure web application for building and managing online  
91 surveys. The survey was created by the authors in conjunction with a multidisciplinary team of experts  
92 in overtraining and resistance training who provided feedback. In order to establish the content validity  
93 and reduce response bias the first draft of the survey was piloted with a convenience sample of 24  
94 athletes who participate in resistance-based sports. Based on the resulting feedback, the survey was  
95 modified to improve its content, clarity, readability and overall quality. The revised survey was further  
96 piloted on a focus group of 6 participants (industry experts and athletes). Based on feedback from the  
97 pilot testing the content and format of the survey was further refined. Finally, the authors completed a  
98 heuristic evaluation to establish the usability of the survey interface on various devices (PC, Macintosh,  
99 iPhone, Android).

100

101 The final survey consisted of 46 items distributed between 5 sections: (1) demographic information, (2)  
102 strength, (3) performance, (4) training, (5) symptoms and (6) recovery (Table 1). The survey concluded  
103 at stage (3) for subjects that indicated that they have never experienced an unexplained decline in  
104 performance. Both open-ended and dichotomous questions were included throughout the survey. In  
105 order to avoid acquiescence bias, respondents were not prompted by pre-loaded questions asking if they  
106 did/did not experience a particular symptom. Such questions increase the likelihood of participants  
107 falsely reporting the presence of a symptom that may not have experienced. Rather, respondents were  
108 required to self-report their symptoms in open text boxes. The survey was available in 4 languages  
109 (English, Italian, Portuguese and Spanish). Native speakers assessed the validity of each translation  
110 based on the original English survey.

111

112 INSERT TABLE 1 ABOUT HERE

113

#### 114 **Sample Selection & Administration**

115

116 A voluntary convenience sample of competitive athletes involved in sports with a resistance training  
117 component were recruited. Eligible sports were categorised into ‘resistance exercise only’ sports  
118 (powerlifting, bodybuilding, weightlifting and strongman) and ‘resistance exercise combined’ sports  
119 (CrossFit, rugby, sprinting, hurdles, long/triple/high jump, shot put, javelin, discus and pole vault).  
120 Collectively, respondents will be referred to as ‘resistance-based athletes’. Participants must have  
121 competed in their respective sports, however, no restriction was placed on the level of competition (i.e.  
122 club to international level athletes).

123

124 Respondents were recruited through various means from July to August 2019 to obtain a sample of  
125 approximately one thousand responses. The primary methods of recruitment were through emails  
126 distributed to industry experts in relevant sports/disciplines and by direct sharing of a survey recruitment  
127 flyer on social media (Figure S1). In order to avoid sampling bias, terms related to ‘overtraining’ and  
128 ‘overreaching’ were not used in the survey advertisement or until section (6) of the survey. Thus,  
129 reducing the likelihood that the resulting sample overrepresents individuals who have strong opinions  
130 or experiences with overreaching and overtraining.

131

#### 132 **Statistical Analysis**

133

134 Statistical analysis of the anonymous data set were conducted using IBM SPSS v25 (2019). Missing  
135 data checks were conducted to confirm data integrity. Frequencies were calculated for respondents  
136 demographic and training characteristics, respondents were categorised according to their respective

137 sport/discipline (resistance exercise only and resistance exercise combined). The proportion of athletes  
138 who reported that they had previously experienced an unexplained decrease in performance were  
139 tabulated. Bivariate statistics (chi-square analyses) were used to determine whether experiences of  
140 training maladaptation varied by training history and/or training style.

141

## 142 **RESULTS**

143

144 Among the 961 online survey views, 760 responded (84% completion rate) (Figure 1). Following the  
145 exclusion of recreational athletes and those that were involved in non-eligible sports the final sample  
146 size included 605 responses. Overall, 70.9% of participants indicated that they had previously  
147 experienced an unexplained decrease in competition performance (Table 2). The majority of  
148 respondents (76.5%) were involved in resistance exercise only sports (powerlifting, weightlifting and  
149 bodybuilding). Demographic characteristics of participants are summarised by sporting category in  
150 Table 3.

151

152

153 INSERT FIGURE 1 ABOUT HERE

154

155 INSERT TABLE 2 ABOUT HERE

156

157 INSERT TABLE 3 ABOUT HERE

158

159

160 Among the participants that indicated that they had previously experienced an unexplained decrease in  
161 performance, 70.9% were indicative of acute maladaptation indicating possible acute fatigue, FOR or  
162 NFOR (<1 week=26.8%, 1 week – 1 month=43.8%) (Figure 2). Only 17.8% of responses were  
163 reflective of chronic training maladaptation indicating possible NFOR or OTS (1-3 months=13.1%, >4  
164 months=4.7%). No interaction effect was found between how many years participants had been training  
165 and unexplained decreases in performance ( $p=0.259$ ). However, findings showed that a greater  
166 proportion of participants with more severe training maladaptation (>4 months) reported training to  
167 muscular failure (Figure 3). The majority of participants who had experienced a decline in performance  
168 also reported experiencing additional stress outside of training (92.5%). The most commonly reported  
169 stressor was work (25.5%) followed by personal life (23.2%) and external factors such as  
170 dieting/negative energy balance (22.9%) (Table 4).

171

172

INSERT FIGURE 2 ABOUT HERE

173

174

INSERT FIGURE 3 ABOUT HERE

175

176

INSERT TABLE 4 ABOUT HERE

177

178 The most reported symptoms for each duration of training maladaptation are displayed in Figure 4. A  
179 complete list of symptoms and frequency of reporting can be found in Table S2. General feelings of  
180 fatigue were the most common self-reported symptom overall (n=153, 35.7%) and in each time frame  
181 of training maladaptation (<1 week n=48, 28.9%; 1 week – 1 month n=74, 27.0%; 1-3 months n=25,  
182 25.5%; >4 months n=6, 22.2%). Musculoskeletal aches and pain were the second most frequent self-  
183 reported symptom for acute maladaptation to training (FOR: n=15, 9.0%; NFOR: n=25, 9.1%).

184

185

INSERT FIGURE 4 ABOUT HERE

186

## 187 **DISCUSSION**

188

189 Finding the optimal balance between training and recovery in order to enhance athletic performance  
190 and simultaneously avoid maladaptive training conditions continues to challenge both coaches and  
191 athletes. Despite the potentially serious implications of overtraining on athletic performance, to date, a  
192 decrease in sport specific performance, that cannot be explained by underlying conditions, is the only  
193 diagnostic marker.<sup>13</sup> The purpose of this study was to provide details on the nature and symptomatic  
194 profile of training maladaptation in competitive resistance-based athletes to examine whether there are  
195 symptoms that may be used as prognostic indicators of overtraining. A greater understanding of the  
196 symptomatic profile of maladaptation in resistance-based athletes may allow for the development of  
197 valid, reliable and objective prognostic and diagnostic tools that do not involve maximal performance  
198 tests. This is the first global survey of experiences and self-reported symptoms of unexplained  
199 underperformance in a large sample of competitive resistance-based athletes across 50 countries.

200

201 Previous studies have suggested that training maladaptation is a significant problem in resistance-based  
202 sports.<sup>11</sup> The purpose of the present study did not include the determination of the prevalence or  
203 incidence of overtraining conditions in resistance exercise. However, a significant number of  
204 respondents indicated that they had previously experienced an unexplained decline in performance  
205 (70.9%). The majority of athletes reported a performance impairment that lasted between 1 week and 1  
206 month (43.8%). A decline in performance of this duration may be reflective of FOR or NFOR, in  
207 accordance with the well-accepted presentation of stages of overtraining.<sup>5</sup> Based on this classification,  
208 in our sample, the proportions of respondents reporting a decline in performance of more than 4 months



209 (i.e. potentially indicative of severe OTS) was relatively low (4.7%). The low proportion of long-term  
210 underperformance, compared to overall prevalence of training maladaptation is consistent with previous  
211 literature which suggests that cases of the OTS are less common.<sup>4</sup> However, the differential diagnosis  
212 of FOR, NFOR or OTS based on survey responses is beyond the scope of this paper. Due to the  
213 continuous nature of overtraining defining the cut-off point between FOR, NFOR and OTS based on  
214 the proposed definitions of Meeusen et al<sup>5</sup> is difficult. Examination of the appropriateness of using  
215 similar diagnostic criteria (time-frame of performance decline) for both endurance- and resistance-  
216 based sports should be considered.

217

218 The detrimental effects of overtraining on athletic performance has led to the search for prognostic  
219 symptoms that can be used to prevent imminent overtraining. However, self-reported symptom  
220 prevalence has not been studied in competitive athletes involved in resistance-based sports.  
221 Determination of the most frequent self-reported symptoms of training maladaptation is of key practical  
222 significance for athletes and coaches in resistance-based sports as it may provide a simple tool, that  
223 does not require physical tests or biochemical analysis, to assess both training and competition  
224 performance. Subjective feelings of general fatigue were the most common self-reported symptom in  
225 this study regardless of the duration of training maladaptation. Athletes and coaches should attempt to  
226 monitor for subjective feelings of fatigue during periods of high training load to monitor for possible  
227 maladaptation. However, the symptom of fatigue is poorly defined and therefore careful inquiry is  
228 needed to distinguish lack of energy from exhaustion or sleepiness.<sup>15</sup> Furthermore, it is unclear whether  
229 the general symptom of fatigue is a physiological response, psychological perception or symptom of  
230 physical or psychiatric disorders.<sup>16</sup> The multidimensional nature of fatigue introduces difficulties in  
231 quantifying the subjective feeling. Future studies should test or develop appropriate instruments to  
232 differentiate acute from chronic fatigue.

233

234 It is difficult to separate the symptoms of overtraining with those of normal fatigue that may result from  
235 high intensity and/or volume training sessions which are necessary to promote physiological  
236 adaptations. Although the decline in performance associated with overtraining appears to be  
237 accompanied by symptoms of generalised fatigue, the presence of fatigue is not necessarily synonymous  
238 with forthcoming training maladaptation. A multidimensional approach to the assessment and  
239 monitoring of training maladaptation is likely required. Athletes and coaches should regularly monitor  
240 a combination of performance and symptomatic variables to identify athletes that may be approaching  
241 FOR, NFOR or OTS. However, the diagnostic criteria of maladaptive conditions remains a decline in  
242 maximal performance lasting 'days-weeks' for FOR, 'weeks-months' for NFOR and 'months...' for  
243 OTS.<sup>5</sup> Accordingly, an accurate diagnosis can only be made retrospectively, once confounding factors  
244 that may be attributed to the decline in performance have been excluded.<sup>5</sup>

245

246 Apart from fatigue, a variety of secondary psychological and physiological symptoms were reported by  
247 athletes, the expression of which varying depending on the individual, training methods and other  
248 contextual factors. Musculoskeletal aches and pains were a frequent complaint of respondents who  
249 reported acute durations of training maladaptation (<1 week and 1 week – 1 month). Although the  
250 etiology of musculoskeletal aches and pain is multifactorial, there is a general consensus that repeated  
251 loading of high forces may create the potential for acute joint or musculotendinous injury.<sup>17</sup> These  
252 results may suggest that an imbalance between training and recovery may result in acute  
253 musculoskeletal pain responses, such as swelling or micro trauma to tissues. However, these responses  
254 are likely to be transient as musculoskeletal aches and pain were not among the most reported symptoms  
255 in chronic maladaptive conditions (1 – 3 months and >4 months). These findings are supported by  
256 Cadegiani et al<sup>18</sup> who reported that impaired muscle recovery may be a characteristic of the OTS.  
257 During periods of high intensity resistance training, athletes and coaches should monitor subjective  
258 symptoms of musculoskeletal aches and pain as chronic maladaptation to training may be avoided if  
259 sufficient recovery is implemented.

260

261 Given the heterogeneity and variability of symptoms of overtraining in resistance exercise and the  
262 present state of research, a single symptom that can be used to predict ensuing overtraining is unlikely.  
263 The interindividual variability of the symptoms of overtraining as well as the variable nature of the  
264 stressors that may cause overtraining suggest the need for a variety of parameters as markers of training  
265 maladaptation. The results of this survey suggest that, at present, no particular symptomatic profile can  
266 be used to distinguish between maladaptive conditions (FOR, NFOR and OTS) in resistance-based  
267 athletes. This may be due to the variability of symptoms or the proposed continuum theory of training  
268 maladaptation. If overtraining exists on a continuum it may be difficult to differentiate between the  
269 symptoms of acute fatigue, overreaching and overtraining. Kuipers et al<sup>19</sup> suggest that symptoms of  
270 overtraining in endurance activities appear progressively as fatigue accumulates. Future studies should  
271 seek to evaluate the severity and progression of symptoms in resistance-based athletes at different stages  
272 of overreaching and overtraining.

273

274 The pathogenesis of maladaptation in resistance-based sports is not completely understood. However,  
275 it is well-established that causes of training maladaptation are multifactorial.<sup>5, 9</sup> In agreement with  
276 previous research in endurance activities,<sup>20, 21</sup> the present findings suggest that a combination of training  
277 and non-training stressors are present at the time of resistance training maladaptation. The vast majority  
278 (92.5%) of athletes experiencing an unexplained decline in performance also reported experiencing  
279 additional stress outside of training. Similarly, Cadegiani et al<sup>9</sup> identified a relationship between  
280 susceptibility to maladaptation and non-training stressors. These findings suggest that training  
281 maladaptation is not only the result of excessive physical training. Coaches and athletes should be aware  
282 of potential stressors outside of training including caloric and macronutrient intake as well as concurrent

283 cognitive demands. Additional social and environmental stressors must be considered and managed  
284 appropriately to prevent training maladaptation. Therefore, regular psychological assessment of athletes  
285 who complete intensive resistance training may provide benefits by detecting early symptoms of  
286 maladaptive training.

287

288 A large proportion of athletes with a decrease in performance lasting greater than 4 months reported  
289 training to muscular failure (65%). This finding suggests that severe maladaptation is associated with  
290 high intensity resistance training to muscular failure. Previous studies have suggested that high intensity  
291 training may increase susceptibility to overtraining in resistance exercise.<sup>13, 22</sup> However, at present no  
292 previous study has examined how training to muscular failure may impact the etiology of overtraining  
293 in resistance exercise. It is well-accepted that there is interindividual variability in the way that athletes  
294 tolerate increases in training load, training intensity and, competition and non-training stress.<sup>6</sup>  
295 Therefore, training load and intensity must be individualised depending on each athlete's response.

296

### 297 **Limitations**

298

299 This is the first global survey of maladaptation to training in resistance-based sports. However, there  
300 are limitations to this study that must be acknowledged. The main limitation of this study was the self-  
301 reported quantification of performance decrements. Possible inter-participant variability in the  
302 determination of an unexplained decrease in performance and recency bias should be considered when  
303 interpreting the findings of this study. For example, low energy availability resulting in relative energy  
304 deficiency in sport may have contributed to a decline in performance that participants unknowingly  
305 interpreted as 'unexplained underperformance'. Future studies are required to understand eating  
306 patterns as additional stressors in relation to overtraining in resistance exercise. Study participants were  
307 drawn from a convenience sample and restricted to competitive resistance-based athletes, therefore  
308 findings cannot be generalised to non-competitive athletes or endurance athletes. In addition, this study  
309 adopted a cross-sectional design precluding any causal inferences (causal link with symptoms).  
310 Furthermore, no precautionary measures were used to prevent multiple entries from the same individual.  
311 Caution should be used when comparing athletes undertaking resistance exercise only or combined  
312 training, it is unknown whether resistance exercise only participants were engaging in additional aerobic  
313 exercise at the time of maladaptation. The difficulties and ethical considerations of studying  
314 overtraining in athletic populations are appreciated, however, well-designed studies are required to gain  
315 a greater understanding of maladaptive training conditions in resistance-based athletes.

316

317

318

## 319 **Practical Applications**

320

321 The purpose of this study was to provide details on the nature and symptomatic profile of training  
322 maladaptation in competitive resistance-based athletes to examine whether there are symptoms that may  
323 be used as prognostic indicators of overtraining. Considering that overtraining is likely associated with  
324 the duration and severity of symptoms rather than a singular symptom, practitioners should routinely  
325 and systematically assess fatigue, musculoskeletal pain and non-training stressors during periods of  
326 high training load and/or intensity. The development of valid and reliable questionnaires or other  
327 diagnostic measures in athletic populations are required for assessing fatigue and additional symptoms  
328 related to training maladaptation. The persistence of such symptoms may be indicative of a negative  
329 progression in training. Given the multifactorial nature of the etiology and symptoms of overtraining,  
330 future studies are required to gain a greater understanding of the underlying mechanisms of training  
331 maladaptation. Well-designed studies with demonstrated decreases in performance and follow-up  
332 measures are required for the development of causal assumptions of overtraining for use in future  
333 prognostic studies.

334

## 335 **Conclusion**

336

337 Overtraining conditions are characterised by an imbalance between training as well as non-training  
338 stress and recovery. Maladaptive conditions may present with a wide range of clinically significant  
339 symptoms such as fatigue, musculoskeletal aches and pain, loss of motivation, insomnia and/or other  
340 physical or psychological symptoms. Subjective feelings of fatigue are likely to accompany training  
341 maladaptation in resistance exercise. However, the results from this exploratory study do not permit  
342 causal conclusions. It is evident that the accompanying symptoms of training maladaptation may be  
343 variable in nature. The multidimensional nature of fatigue and individual variability in symptomatic  
344 responses precludes the definitive prognosis or differential diagnosis of FOR, NFOR or OTS in  
345 resistance exercise. At present, the correct diagnosis of overreaching and overtraining in resistance  
346 exercise remains a demonstrated decrease in maximal performance capacity.

347

348 **Acknowledgments:** The authors wish to thank Brad Schoenfeld, Mike Israetel, Stuart McMillan, Carlo  
349 Buzzichelli, Greg Nuckols, Johann Bilsborough, David Nolan, Layne Norton, Bill Campbell, Michael  
350 Pang, Jackson Peos, Jared Feather, Luke Tulloch, Kristen Dunsmore, Patrick Umphrey, Brandon  
351 Roberts and all others who aided in the distribution of this survey.

352

353 **Funding:** No sources of funding were received in the preparation of this manuscript.

354

355 **Disclosure Statement:** Clementine Grandou, Lee Wallace, Franco M. Impellizzeri, Aaron J. Coutts  
356 and Lee Bell declare that they have no conflicts of interest relevant to the content of this study.

357

358 **Data Availability Statement:** The raw and coded data that support these findings of this study are  
359 available from <https://osf.io/je2rf> and <https://osf.io/bgxpa>, respectively.

For Peer Review

## 360 REFERENCES

361

- 362 1. Kraemer WJ, Ratamess NA. Fundamentals of resistance training: progression and exercise  
363 prescription. *J Med Sci Sports Exc.* 2004;36(4):674-88.
- 364 2. Suchomel TJ, Nimphius S, Stone MH. The importance of muscular strength in athletic  
365 performance. *Sports Med.* 2016;46(10):1419-49.
- 366 3. Bird SP, Tarpenning KM, Marino FE. Designing resistance training programmes to enhance  
367 muscular fitness. *Sports Med.* 2005;35(10):841-51.
- 368 4. Kreider RB, Fry AC, O'Toole ML. Overtraining in sport. Champaign, IL: Human Kinetics;  
369 1998.
- 370 5. Meeusen R, Duclos M, Foster C, et al. Prevention, diagnosis, and treatment of the overtraining  
371 syndrome: joint consensus statement of the European College of Sport Science and the American  
372 College of Sports Medicine. *Med Sci Sports Exerc.* 2013;45(1):186-205.
- 373 6. Erskine RM, Jones DA, Williams AG, et al. Inter-individual variability in the adaptation of  
374 human muscle specific tension to progressive resistance training. *Eur J Appl Physiol.*  
375 2010;110(6):1117-25.
- 376 7. McEwen BS. Stress, adaptation, and disease: Allostasis and allostatic load. *Ann N Y Acad Sci.*  
377 1998;840(1):33-44.
- 378 8. Selye H. The stress of life. New York: McGraw Hill; 1976.
- 379 9. Cadegiani FA, Kater CE. Novel insights of overtraining syndrome discovered from the EROS  
380 study. *BMJ Open Sport Exerc Med.* 2019;5(1):542.
- 381 10. Fry RW, Morton AR, Keast D. Overtraining in athletes. *Sports Med.* 1991;12(1):32-65.
- 382 11. Fry AC, Kraemer WJ. Resistance exercise overtraining and overreaching: neuroendocrine  
383 responses. *Sports Med.* 1997;23(2):106-29.
- 384 12. Wolf W. A contribution to the question of overtraining. A collection of papers presented at the  
385 Institute of Normal Human Anatomy and the Ministry of Foreign Affairs; Rome, Italy 1961;291.
- 386 13. Grandou C, Wallace L, Impellizzeri F, et al. Overtraining in resistance exercise: an exploratory  
387 systematic review and methodological appraisal of the literature. *Sports Med.* 2019.
- 388 14. Eysenbach G. Improving the quality of web surveys: the checklist for reporting results of  
389 internet e-surveys (CHERRIES). *J Med Internet Res.* 2004;6(3):e34.
- 390 15. Sharpe M, Wilks D. Fatigue. *BMJ.* 2002;325(7362):480-3.
- 391 16. Kennedy H. Fatigue and fatigability. *Br J Psychiatry.* 1988;153(1):1-5.
- 392 17. Pecina MM, Bojanic I. Overuse injuries of the musculoskeletal system: CRC press; 2003.
- 393 18. Cadegiani, F. A., Kater, C. E. Basal hormones and biochemical markers as predictors of  
394 overtraining syndrome in male athletes: the EROS-BASAL study. *J Athl Train.* 2019;54(8), 906-914.
- 395 19. Kuipers H, Keizer H. Overtraining in elite athletes. *Sports Med.* 1988;6(2):79-92.
- 396 20. Coutts AJ, Wallace L, Slattery K. Monitoring changes in performance, physiology,  
397 biochemistry, and psychology during overreaching and recovery in triathletes. *Int J Sports Med.*  
398 2007;28(02):125-34.
- 399 21. Morgan WP, Costill DL, Flynn MG, et al. Mood disturbance following increased training in  
400 swimmers. *Med Sci Sports Exerc.* 1988.
- 401 22. Fry AC, Kraemer WJ, Van Borselen F, et al. Performance decrements with high-intensity  
402 resistance exercise overtraining. *Med Sci Sports Exerc.* 1994;26(9):1165-73.

403

404

**Table 1.** Survey Instrument.

Field	Answer Choice
<b>1. Demographics</b>	
Country	Country dropdown
Gender	1, Male 2, Female 3, Other
Age	Numerical
Body weight (kg)	Numerical
Height (cm)	Numerical
What sport or discipline do you compete in?	1, Weightlifting 2, Powerlifting 3, Strongman 4, Bodybuilding 5, Sprinting 6, Hurdles 7, Long jump 8, Triple jump 9, High jump 10, Shot put 11, Javelin 12, Discus 13, CrossFit 14, Rugby Union 15, Rugby League 16, American Football 17, Pole Vault 18, Other
If "other", please specify	Open
How many years have you been training in your sport?	Numerical
What is the highest level you have competed at?	1, Club 2, Regional 3, State 4, National 5, International 6, Other
<b>2. Strength</b>	
Load SQUAT: What is the maximum weight you have lifted for a given amount of repetitions? (i.e. 100kg for 1RM or 75kg for 3RM etc.) (kg)	Numerical
Repetitions?	Numerical
Load BENCH PRESS: What is the maximum weight you have lifted for a given amount of repetitions? (i.e. 100kg for 1RM or 75kg for 3RM etc.) (kg)	Numerical
Repetitions?	Numerical
<b>3. Performance</b>	
Have you ever experienced an unexplained decrease in performance?	1, Yes 2, No (survey termination if "No")
<b>4. Training</b>	
How many times have you experienced this?	1, Once 2, Twice 3, Three or more times
Please answer the following questions in reference to your most severe case (if more than once)	

How was your performance affected?	1, Decreased strength 2, Decreased running speed 3, Blunted hypertrophy 4, Increased perception of effort while training 5, Decrease in sport performance (e.g. jump height, throwing, sprint) 6, Other
If "other", please specify	Open
How long did the decrease in performance last? (i.e. when did your performance return to normal?)	1, < 1 week 2, 1 week - 1 month 3, 1 - 3 months 4, > 4 months
Leading up to the decline in performance how many times per week were you performing resistance training?	Open
What was the average duration of each resistance training session?	1, < 1 hour 2, 1-2 hours 3, >2 hours
Leading up to the decline in performance were you also performing technical/skill training?	1, Yes 2, No
Leading up to the decline in performance were you also performing metabolic and/or conditioning training?	1, Yes 2, No
Which option best describes the INTENSITY of resistance training you were performing at the time you experienced the decline in performance?	1, heavy loads to muscular failure 2, heavy loads without muscular failure 3, light loads to muscular failure 4, light loads without muscular failure
Which option best describes the VOLUME of resistance training you were performing at the time you experienced the decline in performance?	1, high repetitions, high sets 2, high repetitions, low sets 3, low repetitions, high sets 4, low repetitions, low sets
How frequently were you training each muscle group?	1, 1 time per week 2, 2 times per week 3, 3 times per week 4, >3 times per week
Leading up to/during this period, were you experiencing any other non-training stress?	1, Dieting (calorie restriction) 2, Private life stress 3, Work-related stress 4, Other non-training stressors 5, None of the above 6, Viral infection 7, Mental health condition
<b>5. Symptoms</b>	
Symptoms	Open
Please rate the severity of the symptoms above	Insignificant   Extreme
Did you also experience any of these symptoms before you noticed the decline in performance?	Open
Did you experience any other symptoms during this period?	Open
Please describe the other symptoms you experienced	Open
<b>6. Recovery</b>	
Did you seek help/search for information when this occurred?	1, No 2, Yes – Doctor 3, Yes - Coach 4, Yes - Internet/web-sources 5, Yes – Other
What recovery/training strategies did you use in order to return performance to normal?	1, Decrease training intensity 2, Decrease training frequency




---

	3, Decrease training volume 4, Increase calorie intake 5, Eliminate stressful contextual (non-training) factors 6, Supplementation 7, Medicine/seek medical help 8, Other
If "other", please specify	Open
Would you define this period as	1, Overtraining 2, Overreaching 3, Acute fatigue 4, Other

---

For Peer Review



**Figure 1.** Sample selection.

Figure 1. Sample selection  
210x297mm (300 x 300 DPI)

**Table 2.** Frequency of self-reported unexplained decrease in performance.

Category	Sport/discipline	Total	Decrease in performance	No decrease in performance
RE only	Powerlifting	231	69.3%	30.7%
	Bodybuilding	138	76.0%	24.0%
	Weightlifting	71	76.0%	24.0%
	Strongman	23	47.8%	52.2%
	<i>Total</i>	<i>463</i>	<i>71.3%</i>	<i>28.7%</i>
RE combined	Rugby	54	63.0%	37.0%
	CrossFit	42	81.0%	19.0%
	Sprint	21	66.7%	33.3%
	American Football	10	60.0%	40.0%
	Hurdles	4	50.0%	50.0%
	Javelin	3	66.7%	33.3%
	Long jump	3	100.0%	0.0%
	Shot put	2	100.0%	0.0%
	Discus	1	0.0%	100.0%
	Pole vault	1	100.0%	0.0%
	Triple jump	1	100.0%	0.0%
	<i>Total</i>	<i>142</i>	<i>69.7%</i>	<i>33.3%</i>
	<b>Combined Total</b>		<b>605</b>	<b>70.9%</b>

*RE*, resistance exercise.

**Table 3.** Descriptive characteristics of 605 respondents.

<b>Characteristic</b>	<b>RE only n (<math>\pm</math>SD)</b>	<b>RE combined n (<math>\pm</math>SD)</b>	<b>Total n (<math>\pm</math>SD)</b>
Male	340	112	452
Female	123	30	153
Age (years)	29.7 (9.05)	28.0 (9.04)	29.27 (9.07)
Weight (kg)	82.37 (18.11)	84.06 (20.1)	83.53 (18.59)
Height (cm)	173.95 (8.79)	178.25 (12.45)	174.96 (9.93)
Training years (years)	4.45 (2.89)	6.54 (3.14)	4.94 (3.08)
Squat 1RM (kg)			
<i>Male</i>	178.62 (45.05) Range: 50-300	161.73 (44.29) Range: 30-290	174.37 (45.40) Range: 30-300
<i>Female</i>	116.48 (33.25) Range: 55-215	99.89 (27.35) Range: 60-174	113.54 (32.77) Range: 55-215
Bench press 1RM (kg)			
<i>Male</i>	125.69 (30.97) Range: 50-265	115.93 (28.64) Range: 20-200	123.28 (30.67) Range: 20-265
<i>Female</i>	69.97 (20.85) Range: 28-155	52.15 (13.41) Range: 25-85	66.85 (20.86) Range: 25-155
Competition level			
<i>Club</i>	166	33	149
<i>Regional</i>	90	22	112
<i>State</i>	54	12	66
<i>National</i>	141	50	191
<i>International</i>	62	25	87

RE, resistance exercise; 1RM, one repetition maximum.





Figure 2. Flow chart of maladaptation to training responses.

Figure 2. Flow chart of maladaptation to training responses.

210x297mm (300 x 300 DPI)



**Figure 3.** Difference between the proportions of participants training to muscular failure and participants training without muscular failure by the severity of training maladaptation.

Figure 3. Difference between the proportions of participants training to muscular failure and participants training without muscular failure by the severity of training maladaptation

293x217mm (96 x 96 DPI)

**Table 4.** Dichotomous frequency of additional stressors in respondents experiencing training maladaptation.

Additional stressor	‘Yes’ responses N(%)				Total
	<1 week	1 week – 1 month	1-3 months	>4 months	
Work	53 (48.1)	86 (48.5)	29 (54.7)	9 (45.0)	177 (25.5)
Life	48 (43.6)	76 (42.9)	28 (52.8)	9 (45.0)	161 (23.2)
Dieting	49 (44.5)	78 (44.0)	23 (43.4)	9 (45.0)	159 (22.9)
Other	21 (19.0)	36 (20.3)	9 (16.9)	4 (20.0)	70 (10.1)
Mental health	20 (18.1)	22 (12.4)	15 (28.3)	4 (20.0)	61 (8.8)
Virus	5 (4.5)	4 (2.2)	4 (7.5)	2 (10.0)	15 (2.2)
No additional stressors	20 (18.1)	23 (12.9)	5 (9.4)	4 (20.0)	52 (7.5)

For Peer Review




Figure 4. Most frequent self-reported symptoms of training maladaptation.

Figure 4. Most frequent self-reported symptoms of training maladaptation

210x297mm (300 x 300 DPI)





**Table S1.** Checklist for Reporting Results of Internet E-Surveys (CHERRIES) [7].

Item category	Checklist Item	Description
Design	Describe survey design	An open cross-sectional international survey was used to identify the prevalence, predicting factors and symptoms of overreaching and overtraining amongst competitive athletes in resistance based sports. The target population was competitive athletes involved in sports with a resistance training component. Eligible sports included powerlifting, bodybuilding, weightlifting, CrossFit, strongman, rugby, sprinting, hurdles, long/triple/high jump, shot put, javelin, discus and pole vault. Subjects must have competed in their respective sports, no restriction was placed on the level of competition (i.e. club to international level athletes). A voluntary non-randomised convenience sample was recruited. Due to the extremely large eligible population, sample randomisation was not possible.
Ethics	Ethics approval	Ethics approval was obtained from the Human Research Ethics Committee of the University of Technology Sydney (ETH19-3898).
	Informed consent	The survey was voluntary and anonymous. Therefore, consent was deemed to be given by beginning the online survey. The banner advertisement included information about the length of the survey and the voluntary/anonymous nature of the survey. The banner advertisement also have an email address that participants were encouraged to contact if they required further information regarding the purpose of the study, identity and qualifications of investigators etc.
	Data protection	REDCap (Research Electronic Data Capture Software Version 8.11.3 – University of Technology Sydney), a secure web application was used to develop and managing this survey. The survey was anonymous, no personal information was linked to the survey results. Data did require de-identification as the survey contained no identifiable questions. The dataset was kept on password protected computers.
Development and pre-testing	Development and testing	The survey was developed by the authors in conjunction with a multidisciplinary team of experts in overtraining and resistance training providing feedback. In order to establish the content validity and reduce response bias the first draft of the survey was piloted with a convenience sample of 24 athletes who participate in resistance-based sports. Based on the resulting feedback, the survey was modified to improve its content, clarity, readability and overall quality. The revised survey was further tested on a focus group of 6 people (industry experts and athletes). Based on feedback from the focus group the content and format of the survey was further refined. Finally, the authors completed a heuristic evaluation to establish the usability of the survey interface on various devices (PC, Macintosh, iPhone, Android). The final survey consisted of 46 items divided between 5 sections: (1) demographic information, (2) strength, (3) performance, (4) training practices, (5) symptoms, (6) recovery. The survey

Item category	Checklist Item	Description
		contained both open-ended and closed questions with response scales. The survey was translated by native speakers into 4 languages; English, Italian, Portuguese and Spanish.
Recruitment process and description of the sample having access to the questionnaire	Open vs closed survey	This was an open survey.
	Contact mode	Not applicable. Potential participants were not contacted.
	Advertising the survey	Survey recruitment was achieved in multiple ways, predominately through banner advertisements, contacting industry experts, sharing via social media and posting in online communities. A banner advertisement was posted with a link to the online survey on Twitter, Instagram and Facebook. The ad was reposted by industry contacts approximately 50 times collectively across all social media platforms. In order to avoid voluntary response bias, terms related to overtraining and overreaching were not used in the survey advertisement. Therefore, reducing the likelihood that the sample may over represent individuals who have strong opinions or experiences with overtraining.
Survey administration	Web/E-mail	This was a web-based survey, with respondents gathered through social media advertisement. Responses were collected through the secure online survey platform REDCap and stored on secure local servers. Responses included open-ended, multiple choice and response scales.
	Context	The survey advertisement was shared by industry experts (e.g. sports science researchers, coaches of eligible sports, athletes involved in eligible sports). The survey was also shared among online communities on Facebook (e.g. Weightlifting Australia, Bodybuilding forums). Therefore, the survey would have likely only captured individuals active on social media. However, this would likely not contribute to response bias. Wording of the survey advertisement was carefully selected to not include terms related to overtraining and overreaching to further reduce the likelihood of response bias.
	Mandatory/voluntary	Voluntary.
	Incentives	Respondents were not incentivised for their participation.
	Time/date	Responses were collected over two months from 24 July to 23 August, 2019.
	Randomisation of items	No randomisation of items was used.
	Adaptive questioning	Adaptive questioning (branched logic) was used throughout the survey to reduce the number and complexity of questions. Certain questions were only relevant for specific populations (e.g. those who indicated that they compete in bodybuilding were further prompted to indicate whether they competed in a natural bodybuilding federation or non-natural bodybuilding federation). Additionally, a stop action was in place on page three where those who indicated that they had never experienced an unexpected decline in performance were required to end the survey. Respondents who indicated that they had experienced an unexplained decrease in performance were required to continue on with the survey.

Item category	Checklist Item	Description
	Number of items	The full survey comprised a total of 46 items, although because of the adaptive nature of the survey, not all respondents answered all items.
	Number of screens	The entire survey was distributed over 6 pages.
	Completeness check	A completeness check was completed after responses were submitted. Page 3 was deemed mandatory <i>“Have you ever experienced an unexplained decrease in performance?”</i> Drop off was low at 16%. Most items, except demographic questions and those items required for adaptive questioning included an ‘other’ or ‘don’t know’ option.
	Review step	Respondents were able to change their responses on previous screens through a ‘Back’ button whilst they were completing the survey. However, respondents were unable to change their responses once submitted. The beginning of the survey was preceded by the sentence <i>“Please ensure that you have time to complete the entire survey before beginning as you cannot save your responses and return at a later time.”</i>
Response rates	Unique site visitor	Not applicable. Open survey.
	View rate	961 online survey views
	Participation rate	Not applicable. Open survey.
	Completion rate	Of the 906 respondents who commenced the survey, 760 completed it, giving a completion rate of 84%. 155 responses were excluded due to eligibility criteria (competitive athletes in resistance-based sports).
Preventing multiple entries from the same individual	Cookies used	Not used.
	IP check	Not used.
	Log file analysis	Not used.
	Registration	Not used.
Analysis	Handling of incomplete questionnaires	Only responses completed up until the third screen where a stop action was in place were included in the final dataset. Respondents who indicated they had experienced an unexplained decrease in performance were prompted to continue to answer questions about their experience. Respondents who indicated that they had never experienced an unexplained decrease in performance were prompted to finish the survey.
	Questionnaires with atypical timestamp	No respondents were removed from the data set for atypical completion times.
	Statistical correction	No methods such as weighting of items or propensity scores were used to adjust for the non-representative sample.



 **UTS** UNIVERSITY OF TECHNOLOGY SYDNEY

## **Demands of Sports with a Resistance Training Component**

### **About**

The purpose of this study is to investigate the training demands of athletes involved in sports with a resistance training component. These insights are important to further our understanding of the topic and provide information as to how practices can be improved.

### **Participants**

Competitive Athletes involved in:


**Bodybuilding, Weightlifting, Powerlifting**

**CrossFit, Rugby, Strongman, American Football, Sprinting, Hurdles,**

**Long/Triple/High Jump, Shot Put, Javelin, Discus, Pole Vault**

You are invited to **anonymously** contribute to this research study organised by the University of Technology Sydney.

**Link: [j.mp/30LU5Nh](https://j.mp/30LU5Nh)**

 This survey will take about 5 minutes

For further information please contact: [resistancetrainingonlinesurvey@gmail.com](mailto:resistancetrainingonlinesurvey@gmail.com)

**Figure S1.** Survey Advertisement.

**Table S2.** Symptoms.

<b>Duration of training maladaptation</b>	<b>Symptom</b>	<b>Frequency of reporting (n)</b>	<b>Frequency of reporting</b>
<b>&lt;1 week</b>	General feelings of fatigue	48	28.9%
	Musculoskeletal aches and pain	15	9.0%
	Decreased motivation	11	6.6%
	Lethargic	10	6.0%
	Perceived weakness	10	6.0%
	Increased perception of effort during training	9	5.4%
	Muscle soreness	8	4.8%
	Poor sleep quality	6	3.6%
	Depression	5	3.0%
	Stress	5	3.0%
	Emotional instability	5	3.0%
	Irritable	4	2.4%
	Anxiety	3	1.8%
	Decreased appetite	3	1.8%
	Poor concentration	3	1.8%
	Light headedness	3	1.8%
	Flu-like illness	2	1.2%
	Digestive discomfort	2	1.2%
	Injury	2	1.2%
	Excessive perspiration	2	1.2%
	Headaches	2	1.2%
	Inflammation	2	1.2%
	Decreased libido	1	0.6%
	Weight gain	1	0.6%
	Slowed recovery	1	0.6%
	Increased appetite	1	0.6%
Loss of coordination	1	0.6%	
Decreased hypertrophy	1	0.6%	
	<b>Total</b>	<b>166</b>	<b>100.0%</b>
<b>1 week – 1 month</b>	General feelings of fatigue	74	27.0%
	Musculoskeletal aches and pain	25	9.1%
	Decreased motivation	23	8.4%
	Muscle soreness	17	6.2%
	Lethargic	14	5.1%
	Perceived weakness	12	4.4%
	Emotional instability	11	4.0%
	Poor concentration	9	3.3%
	Poor sleep quality	8	2.9%
	Increased perception of effort during training	8	2.9%
	Irritable	8	2.9%
	Slowed recovery	6	2.2%
	Increased appetite	6	2.2%
	Decreased libido	5	1.8%
	Headaches	5	1.8%
	Light headedness	5	1.8%
	General apathy	5	1.8%

	Anxiety	4	1.5%
	Stress	4	1.5%
	Flu-like illness	4	1.5%
	Depression	3	1.1%
	Decreased appetite	3	1.1%
	Injury	3	1.1%
	Weight loss	3	1.1%
	Increased heart rate	2	0.7%
	Decreased hypertrophy	2	0.7%
	Weight gain	1	0.4%
	Digestive discomfort	1	0.4%
	Amenorrhea	1	0.4%
	Loss of coordination	1	0.4%
	Inflammation	1	0.4%
	<b>Total</b>	<b>274</b>	<b>100.0%</b>
<b>1 – 3 months</b>	General feelings of fatigue	25	25.5%
	Poor sleep quality	8	8.2%
	Lethargic	5	5.1%
	Decreased motivation	5	5.1%
	Poor concentration	5	5.1%
	Increased perception of effort during training	4	4.1%
	Decreased libido	4	4.1%
	Muscle soreness	4	4.1%
	Anxiety	3	3.1%
	Musculoskeletal aches and pain	3	3.1%
	Emotional instability	3	3.1%
	Flu-like illness	3	3.1%
	Increased appetite	3	3.1%
	Perceived weakness	3	3.1%
	Injury	3	3.1%
	Loss of coordination	3	3.1%
	Slowed recovery	2	2.0%
	Weight loss	2	2.0%
	Headaches	2	2.0%
	Depression	1	1.0%
	Increased heart rate	1	1.0%
	Decreased appetite	1	1.0%
	Stress	1	1.0%
	Irritable	1	1.0%
	Digestive discomfort	1	1.0%
	Light headedness	1	1.0%
	Decreased hypertrophy	1	1.0%
	<b>Total</b>	<b>98</b>	<b>100.0%</b>
<b>&gt;4 months</b>	General feelings of fatigue	6	22.2%
	Decreased motivation	4	14.8%
	Perceived weakness	3	11.1%
	Poor sleep quality	2	7.4%
	Irritable	2	7.4%
	Muscle soreness	2	7.4%
	Lethargic	1	3.7%
	Depression	1	3.7%

---

Anxiety	1	3.7%
Emotional instability	1	3.7%
Flu-like illness	1	3.7%
Injury	1	3.7%
Inflammation	1	3.7%
Decreased hypertrophy	1	3.7%
<b>Total</b>	<b>27</b>	<b>100.00%</b>

---

For Peer Review