

## **Current Insights into Exercise-based Cardiac Rehabilitation in Patients with Coronary Heart Disease and Chronic Heart Failure**

NICHOLS, Simon <<http://orcid.org/0000-0003-0377-6982>>, MCGREGOR, Gordon <<http://orcid.org/0000-0001-8963-9107>>, BRECKON, Jeff and INGLE, Lee

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/26911/>

---

This document is the Accepted Version [AM]

### **Citation:**

NICHOLS, Simon, MCGREGOR, Gordon, BRECKON, Jeff and INGLE, Lee (2020). Current Insights into Exercise-based Cardiac Rehabilitation in Patients with Coronary Heart Disease and Chronic Heart Failure. International Journal of Sports Medicine. [Article]

---

### **Copyright and re-use policy**

See <http://shura.shu.ac.uk/information.html>

## Abstract:

Cardiac rehabilitation is a package of lifestyle secondary prevention strategies designed for patients with coronary heart disease and chronic heart failure. A community-based cardiac rehabilitation programme provides patients with a structured exercise training intervention alongside educational support and psychological counselling. This review provides an update regarding the clinical benefits of community-based cardiac rehabilitation from a psycho-physiological perspective, and also focuses on the latest epidemiological evidence regarding potential survival benefits. Behaviour change is key to long-term adoption of a healthy and active lifestyle following a cardiac event. In order for lifestyle interventions such as structured exercise interventions to be adopted by patients, practitioners need to ensure that behaviour change programmes are mapped against patient's priorities and values, and adapted to their level of readiness and intention to engage with the target behaviour. We review the evidence regarding behaviour change strategies for cardiac patients and provide practitioners with the latest guidance. The "dose" of exercise training delivered to patients attending exercise-based cardiac rehabilitation is an important consideration because an improvement in peak oxygen uptake requires an adequate physiological stimulus to invoke positive physiological adaptation. We conclude by critically reviewing the latest evidence regarding exercise dose for cardiac patients including the role of traditional and more contemporary training interventions including high intensity interval training.

Word count: 200 words

Keywords: cardiac, behaviour change, high intensity interval training, exercise dose.

## **A brief history of cardiac rehabilitation**

Exercise training has been used to induce weight loss, increase muscular strength and mass, manage diabetes, and reduce the 'incidence of disease' for more than 2,500 years [1]. However, accounts of exercise training being used to improve the health of someone with coronary heart disease (CHD) did not emerge until the 18<sup>th</sup> century [2]. Despite this early observation, bedrest, rather than exercise, remained the most common treatment for patients with CHD until the 1940's when early mobilisation was shown to prevent complications caused by prolonged bedrest [3]. In 1964, the World Health Organisation acknowledged the important role of 'reconditioning' patients with heart disease [4] and, in 1993, recommended that exercise training become a key component of cardiac rehabilitation (CR) [5]. Exercise training is now considered a core component of a comprehensive CR programme [6] and is thought to improve patient survival [7] and quality of life [8]. However, despite the wide application of exercise training in CR, controversy relating to best practice remains.

## **Scope of the review**

The term 'heart disease' refers to all diseases of the coronary arteries, electrical system or and/or mechanical function of the heart [9]. The benefits of participating in exercise-based CR can vary depending on the type of heart disease being treated and for simplicity, this review will focus on the benefits of exercise-based CR in patients with CHD and chronic heart failure (CHF). The term 'exercise' will be applied in line with WHO definitions, where physical activity is 'any bodily movement produced by skeletal muscles that requires energy expenditure' and exercise is physical activity that is 'planned, structured, repetitive, and purposeful' [5].

Coronary heart disease is caused by atherosclerotic plaques that develop in one or more coronary arteries which can restrict the blood supply to the myocardium and cause angina, or rupture and cause a myocardial infarction (MI) [9]. Although CHD is the leading cause of CHF [10], CHF is fundamentally different, and has other causes which include cardiomyopathies and cardiac valve

disease [10]. Regardless of aetiology, CHF is characterised by structural and/or functional cardiac abnormalities, reduced cardiac output and/or elevated intracardiac pressures which cause patients to experience severe breathlessness, ankle swelling and/or fatigue [11]. Whilst patients with CHD and CHF can both benefit from exercise-based CR, the benefits of exercise-based CR may differ. This review was conducted ethically in accordance with the requirements of the journal [12].

### ***Clinical benefit of cardiac rehabilitation***

Although the effects of exercise training in patients with angina are under reported [13], systematic reviews and meta-analyses conducted in 2011 and 2016 have reported that exercise-based CR reduces cardiovascular mortality and hospital admissions by up to 26% and 18% respectively, over 12 months in patients with CHD [7, 14]. Nevertheless, the effect of exercise-based CR in the era of contemporary medical care has been brought in to question by a recent systematic review and meta-analysis of trials conducted after the year 2000 [15]. Powell et al. [14] found that exercise-based CR did not reduce all-cause (risk difference [RD] 0.00; 95% confidence interval [CI] -0.02 to 0.01,  $P=0.38$ ) or cardiovascular mortality (RD -0.01; 95% CI -0.02 to 0.01,  $P=0.25$ ) in patients with CHD. These findings are supported by data from sequential Cochrane reviews which report ever smaller reductions in all-cause and cardiovascular mortality following exercise-based CR for patients with CHD [7, 14, 16]. It is likely that this reflects the beneficial effect that contemporary revascularisation techniques [17, 18] and potent medical management have on shorter and long-term patient survival. In contrast to data recorded in patients with CHD however, exercise-based CR has not historically been considered to improve all-cause mortality in patients with CHF (relative risk [RR] 0.88; 95% CI 0.75 to 1.02) [8, 19]. In addition to the physiological impact, evidence suggests enhanced psychological functioning and emotional regulation reducing both stress and anxiety and increasing the perception of autonomy (self-determination) and control [20].

Despite exercise-based CR having limited impact on patient survival [8, 15], patients with CHD (RD  $-0.05$ ; 95% CI  $-0.10$  to  $-0.00$ ;  $P=0.05$ ) [15] and CHF may still benefit from fewer hospital admissions (RR 0.70; 95% CI 0.60 to 0.83) [8]. Furthermore, exercise-based CR may improve patients' health-related quality of life [8, 21] and improve cardiometabolic risk factors [22, 23] which could allow patients to reduce the number and/or dose of secondary prevention medications, although further research is needed to confirm this assertion. In light of these findings, practitioners should continue to explore ways of optimising exercise-based CR.

### **Modifiable cardiovascular risk factors**

Clinical trials and systematic reviews have shown that exercise-based CR can reduce the severity of cardiovascular risk factors associated with the progression of CHD or CHF (Table 1). These include a reduction in low-density lipoprotein (LDL) cholesterol [23, 24], triglycerides, fasting blood glucose, obesity [25], inflammatory markers such as C-reactive protein (CRP) [25], resting blood pressure [24], and end-diastolic volume [26]. Conversely, exercise training can increase high-density lipoprotein (HDL) cholesterol [22, 25] and improve endothelial function [27]. Limited evidence indicates that exercise training could limit myocardial remodelling [28] and attenuate the progression of atherosclerosis [29], although participation for between six and twelve months respectively, may be needed to achieve this outcome.

Clinical trials have shown that aerobic fitness, a key cardiovascular risk factor, may also be improved following exercise-based CR [22, 25, 26, 28-30]. However, data from 950 patients [31] indicated that aerobic fitness improvements following routine exercise-based CR may be small ( $\sim 0.5$  metabolic equivalents; METs) when compared to the effects noted in clinical trials ( $\sim 1.5$  METs) [32]. This may indicate that the dose of exercise delivered in routine healthcare is insufficient [33, 34] and recent data from 332 exercise sessions conducted in a routine exercise-based CR setting supports this

observation [35]. The mean exercise training intensity during a routine exercise-based CR session was reportedly  $37\% \pm 10\%$  of heart rate reserve [HRR], despite practitioners prescribing exercise at 40-70% [35]. This suggests that patients are not being adequately supported to engage in exercise-based CR, a finding emphasised by recent data from the pan-European research project, EUROASPIRE V [36]. EUROASPIRE V reported that only 34% ( $n=2,809$ ) of 8,261 patients with CHD were taking part in 30 minutes of exercise on five or more days per week, and that 42% ( $n=3,470$ ) of patients had no intention of changing their exercise behaviour [36]. Furthermore, 46% ( $n=3,800$ ) of patients did not receive advice to increase their exercise levels. A similar pattern was evident in relation to weight management [36], and recent findings indicate that obesity may increase following participation in some exercise-based CR programmes [37]. Increasing participation in exercise and reducing levels of obesity and overweight are core objectives of exercise-based CR. A failure to support patients to modify these risk factors could explain why exercise-based CR delivered in routine clinical practice may not improve survival [38]. Exercise-based CR can only be expected to improve cardiovascular risk factors and survival if patients meaningfully engage in the intervention. There is a clear need for researchers, policy makers and practitioners to re-examine the evidence for exercise-based CR to allow the translation of established benefits identified from clinical trials to be integrated into clinical practice. A focus of support from CR programme teams should be to assist patients in making positive lifestyle choices which help improve aerobic fitness and cardiometabolic health.

### **Behaviour change strategies**

While the benefits of structured exercise regimens are unequivocal for those CR patients, practitioners and rehabilitation teams cannot assume that the patient will initiate and maintain recommended rehabilitation programmes just because it is a clinical recommendation. Indeed, evidence suggests that those patients that do adopt lifestyle behaviour change (e.g. diet, PA,

smoking, alcohol and medication adherence) are highly likely to relapse, with programmes suffering from around 75% dropout within 12 weeks of their start [39]. For several years, evidence has highlighted the limits of the long-term effectiveness of exercise for sedentary individuals (which typically reflects these individuals' level of activity) [40] with an over-emphasis by practitioners on exercise initiation, neglecting long term-behaviour change strategies [41]. This action planning approach has typically relied on passive information exchange (e.g. leaflets and signposting to guidelines) which has demonstrated little impact on patient's long-term behaviour change and lifestyle modification. That is not to say that information and knowledge exchange does not have a role to play, but rather that it should not be the only tool in the practitioner's toolbox. To increase impact, an appreciation of effective exercise prescription and other lifestyle intervention components and strategies for the maintenance of such changes, are required [42].

In order for lifestyle interventions such as structured exercise interventions to be adopted by patients, practitioners need to ensure that behaviour change programmes are mapped against the individual's priorities and values [43] and adapted to their level of readiness [and intention] to engage with the target behaviour [e.g. exercise and dietary modification] [44]. An understanding of the patients' intention to change (an individual's desire to perform a given behaviour) is a facet of their likelihood to maintain change and has been highlighted as fundamental by numerous theories. These include: The Theory of Planned Behaviour (TPB) [45], Theory of Reasoned Action (TRA) [46], and Health Action Process Approach (HAPA) [47], all of which place intention as the proximal determinant of behaviour separating motivation and action. The support required for patients in exercise programming and lifestyle change is therefore multifaceted and requires any combination of resilience development, autonomy building, identifying flexible rather than rigid change goals, agenda mapping and environmental support in the form of peers and family members. Evidence has suggested that exercise interventions should also include clearly identifiable Behaviour Change Techniques (BCTs) to increase a patients' autonomy and intention toward change. Examples of BCT strategies for exercise adoption and maintenance include 'behavioural goal setting', 'self-monitoring

(behaviour)' or 'behavioural practice/rehearsal', all of which have been recommended as key ingredients to promoting long-term change [48]. These approaches increase an individual's perception of self-control and sense of autonomy toward their own change, fundamental aspects when trying to build and maintain motivation.

Patient lifestyle behaviour change is complex and often difficult to achieve. Practitioners should consider utilising the clients experience and raised awareness of their own resource in order to build their engagement in the planning, initiation and maintenance phases [44]. While BCTs have been cited as important ingredients toward supporting patient change, it is important that interventions are evidence-based and practitioners be aware of both the *what* to change as well as the *how* to facilitate this [49].

### **The role of aerobic fitness**

Peak oxygen uptake ( $\dot{V}O_{2peak}$ ) is determined during a maximal cardiopulmonary exercise test (CPET) and can be summarised by the Fick Equation [50]:

$$\dot{V}O_2 = \dot{Q} \times a-vO_{2\text{ diff}}$$

Peak oxygen uptake therefore reflects the functional reserve of the heart (cardiac output;  $\dot{Q}$ ) and muscles ( $a-vO_{2\text{ diff}}$ ) when responding to meet the metabolic demands of severe physical stress. The functional reserve of patients with CHD and CHF may be limited, as evidenced by a low  $\dot{V}O_{2peak}$  [51-53]. Multiple factors may contribute to a patient having a low  $\dot{V}O_{2peak}$ . For example, chronotropic incompetence caused by poor sympathetic and parasympathetic tone, may limit a rise in heart rate during exercise and thus limits  $\dot{Q}$  [54]. Similarly, exercise may induce left ventricular dysfunction which can also limit peak stroke volume and therefore,  $\dot{Q}$  [55]. These factors may impede the delivery of oxygenated blood to the exercising muscle [56,57] where adverse changes to skeletal muscle physiology including low muscle mass [58,59], poor perfusion and matching of the capillaries



and muscle fibres, as well as reduced muscle oxidative capacity [56], may reduce  $a-vO_{2\text{ diff}}$  and aerobic synthesis of adenosine triphosphate [55]. A low  $\dot{Q}$  and  $a-vO_{2\text{ diff}}$  causes a marked reduction in  $\dot{V}O_{2\text{ peak}}$  such that the  $\dot{V}O_{2\text{ peak}}$  of men ( $20.2\text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ ) [51] and women ( $15.1\text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ ) [52] with heart disease who are aged between 50 and 59 years is typically half of that found in healthy men ( $42.1\text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ ) and women ( $33.7\text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ ) of a similar age [60]. This is concerning because individuals with a low  $\dot{V}O_{2\text{ peak}}$  also have a higher risk of death [61] and disability [62] when compared to those with a higher  $\dot{V}O_{2\text{ peak}}$ .

The importance of maintaining aerobic fitness in to older age can be demonstrated by data which suggests that men and women with a  $\dot{V}O_{2\text{ peak}}$  below  $18$  and  $15\text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$  respectively, may struggle to maintain physical independence [62]. Furthermore, men with a  $\dot{V}O_{2\text{ peak}} < 14.9\text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$  have significantly greater risk of death (HR 15.15; 95% CI 7.68 to 29.88) when compared to men with a  $\dot{V}O_{2\text{ peak}} < 22.8\text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ . Similar, data are reported for women with a  $\dot{V}O_{2\text{ peak}} < 11.9\text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$  (HR 5.87; 95% CI 2.60 to 13.10) when compared to women with a  $\dot{V}O_{2\text{ peak}} < 16.6\text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$  [60]. The reasons for this are not fully understood, however patients with a low  $\dot{V}O_{2\text{ peak}}$  typically have more risk factors including advanced age, poor sympathetic and parasympathetic tone indicated by a slower post-exercise heart rate recovery [63], higher markers of cardiac stress (N-terminal pro-brain natriuretic peptide) and inflammation (hs-CRP), higher non-fasting blood glucose levels, more severe atherosclerosis, and lower haemoglobin and haematocrit [63]. Importantly however, increasing a patient's  $\dot{V}O_{2\text{ peak}}$  may improve survival and quality of life.

In 1995, Vanhees and colleagues [64] demonstrated that a 1% improvement in  $\dot{V}O_{2\text{ peak}}$  conferred a 2% reduction in mortality over 5-years, following exercise-based CR in patients with CHD. More recent evidence suggests that a  $3.5\text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$  improvement in  $\dot{V}O_{2\text{ peak}}$  following exercise training reduces mortality by 25%, if improvements are maintained for more than one year [65]. Exercise-based CR has the potential to increase a patients  $\dot{V}O_{2\text{ peak}}$  by  $5.4\text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$  (95% CI  $4.2\text{--}6.6\text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ ) [32]

suggesting that long-term adherence to accurately prescribed exercise-based CR should be capable of improving survival and quality of life.

We have recently investigated the effects of a routine, twice weekly, exercise-based CR programme for eight weeks (intervention group) compared with abstention from supervised exercise training [control group] in patients with coronary heart disease [66]. The primary outcome was  $\dot{V}O_{2peak}$  measured using criterion methods. In 70 patients (age  $63.1 \pm 10.0$  years; 86% male;  $n = 48$  intervention;  $n = 22$  controls),  $\dot{V}O_{2peak}$  was  $23.3 \text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$  at baseline, and there were no changes in  $\dot{V}O_{2peak}$  between groups at any time point. Based on our findings, routine CR does not lead to an increase in  $\dot{V}O_{2peak}$  and is unlikely to improve long-term physiological outcomes. One of the key reasons for this lack of improvement in peak oxygen uptake may relate to how we prescribe exercise in this patient group.

### **Exercise prescription strategies**

Exercise prescription recommendations for CR have been highlighted in national and international guidelines (Table 2) [6, 67]. Exercise prescription should first take account of the needs and goals of the individual, including their clinical status and readiness to change behaviour. Furthermore, exercise must be accessible to improve adherence and facilitate sustainable long-term change. Accordingly, exercise-based CR is delivered in numerous settings, either in supervised centre-based programmes, or at home, facilitated by manuals and online platforms. Every individual is different, thus a personalised programme taking account of these factors is likely to be the most successful approach. Unfortunately, in different healthcare settings around the world, it is common for prescription to be dictated by convention and resource, rather than preference or evidence-based best practice [68].

Supervised, centre-based CR is by far the most common delivery model, particularly in the UK [69]. Indeed, the accumulated evidence of the benefits of CR is based almost exclusively on studies adopting this approach either in the 'laboratory' or in pragmatic trials. However, recent evidence from the REACH HF study [21] has definitively demonstrated that a home-based CR intervention, that includes a progressive exercise training programme, can improve quality of life in patients with CHF. Patients in the intervention group reported a 5.7 point reduction in the Minnesota Living with Heart Failure Score after four months (95% confidence interval  $-10.6$  to  $-0.7$ ;  $P=0.025$ ), indicating a significant improvement in quality of life. This was maintained after 12 months (Minnesota Living with Heart Failure Score  $-3.2$ ; 95% CI  $-5.7$  to  $-0.6$ ;  $P=0.016$ ). Whilst the intervention did not increase estimated aerobic fitness, historical data from patients with coronary artery bypass grafts indicates that patients engaging in home-based CR are more likely to sustain improvements in aerobic fitness after 12 months, when compared to patients enrolling on centre-based CR [70]. The reasons for these conflicting findings are likely to be multifaceted, but could include factors such as patients with CHF typically experiencing faster disease progression than patients with coronary artery bypass graft, differences in intervention compliance, or exercise stimulus. Nonetheless, self-facilitated or home-based exercise-based CR is significantly underutilised and under researched, despite it showing great potential as an effective alternative to traditional centre-based CR. With countries currently in lockdown due to the outbreak of Covid-19, home-based exercise is, and should, benefit from increased interest amongst healthcare professionals and researchers. Whilst this will benefit patients, practitioners, and healthcare services, it would be easy to lose sight of the longer-term goal of widening access to exercise-based CR and improving the outcomes of patients who attend. Instead of exclusively focusing on the use of home-based CR during the Covid-19 crisis, researchers and practitioners should identify opportunities to incorporate high quality home-based CR in to future standard practice and research projects.

A common aim of exercise-based cardiac CR studies is to improve aerobic capacity, both as a means to enhance quality of life, and as a proxy for improved survival [71,72]. Likewise, improvements in

muscular strength, endurance and flexibility are considered important to improve the cardio-metabolic risk profile and maintain functional independence into older age [73]. To achieve this, guidelines promote comprehensive and holistic exercise assessment and prescription [74]. Perhaps the most contentious issue is that of exercise intensity. The lower and upper safe and effective limits are still debated, as is assessment and prescription in practice. Guidelines vary; Europe, North America and Canada, for example, advocate higher intensity exercise than the UK [67]. Higher intensity exercise results in greater improvements in cardiorespiratory fitness [75] but, for some patients, may be associated with an increased risk of cardiac events [76] or may not be personally acceptable and/or sustainable. With individually tailored exercise prescription, there is a delicate balance to be achieved between optimal intensity, acceptability and safety.

In the context of contemporary pharmacological and interventional management of CHD and CHF, evidence increasingly supports the safety and efficacy of higher intensity exercise-based CR. In meta-analyses, aerobic interval or high intensity interval training (HIIT) have been consistently shown to improve aerobic capacity more than conventional moderate intensity continuous training (MICT) [77-80]. Short bouts (1 to 4 min) of higher intensity ( $>85\% \text{VO}_{2\text{ peak}}$ ) exercise interspersed with low intensity recovery allow patients to accumulate a greater overall 'dose' of exercise. Benefits appear consistent, regardless of which HIIT protocol is adopted (i.e. 4 x 4 min or 10 x 1 min models) [81]. Historically, there has been a reluctance to deviate from conservative CR exercise intensity guidelines, but large datasets now confirm the safety of HIIT [82,83] and, importantly, patient acceptability [81]. However, two pragmatic multi-centre trials (SAINTEX-CAD [84] and SMARTEX-CHF [85]) did not report the superiority of HIIT using the 4 x 4 model in CHD and CHF. Both studies highlighted issues relating to compliance with the prescribed exercise intensity. In both cases, it was observed that there was little difference between the training intensities in the HIIT or MICT groups due to under achievement in the HIIT groups and over achievement in the MICT groups. This raises issues of compliance with HIIT which are likely multi-factorial, potentially relating to discomfort experienced when exercising above the ventilatory anaerobic threshold (VAT) in addition to logistics

and implementation [81]. Nevertheless, with appropriate medical approval, pre-screening, supervision and monitoring, HIIT can be prescribed safely and effectively in the CR setting for people with CHD and CHF [86, 87]. It should be considered an alternative to, but not replacement for MICT, rather an additional tool available to CR practitioners.

Exercise intensity in CR is commonly determined using predictive equations such as 40-70% heart rate reserve, derived from an estimated maximal heart rate with or without a sub-maximal exercise assessment [67]. This range equates to an RPE of 11-14, eliciting an exercise training intensity that straddles the VAT and is known to be effective at improving cardiorespiratory fitness [88]. In lieu of formal assessment of aerobic capacity using CPET, this approach provides a guide with which exercise can be prescribed and subsequently adjusted. However, recent studies have highlighted the inherent inaccuracy of such calculations, leading to exercise intensities that are consistently lower or higher than the intended optimal thresholds [88, 89]. This may contribute to reports of the diminishing efficacy of CR for some patients and of poor adherence and compliance for others [15, 69]. Whilst CPET can enable precision in exercise prescription, it is not routinely available, particularly in the UK, and is an expensive resource compared to commonly used submaximal 'field tests' such as the six-minute walk test, the incremental shuttle walk test (ISWT), and step tests. Field-tests may provide a useful indication of a patients' aerobic fitness, the presence of a 'normal response' to exercise, and provide information that can be used to inform exercise prescription. However, estimating  $\dot{V}O_{2peak}$  may be inaccurate, particularly when measuring *changes* in  $\dot{V}O_{2peak}$  after exercise-based CR [90]. Maximal cycle ergometry data from patients with CHD, collected by our group, showed that estimating changes in  $\dot{V}O_{2peak}$  resulted in significant measurement error, when compared to directly determined  $\dot{V}O_{2peak}$  (limits of agreement -4.7 to 5.9 ml·kg<sup>-1</sup>·min). Measurement error was such that estimated changes in  $\dot{V}O_{2peak}$  significantly increase following exercise-based CR (1.3 ml·kg<sup>-1</sup>·min; 95% CI 0.4 to 2.2 ml·kg<sup>-1</sup>·min;  $P=0.006$ ), whilst directly determined changes in  $\dot{V}O_{2peak}$  did not (0.5 ml·kg<sup>-1</sup>·min; 95% CI 0.6 to 1.8 ml·kg<sup>-1</sup>·min;  $P=0.332$ ). We recently observed similar findings in a study of patients attending CR where estimated changes in  $\dot{V}O_{2peak}$  following treadmill work

significantly increased, but directly determined  $\dot{V}O_{2peak}$  did not [66]. This may have important implications for the way we interpret data from studies that have only estimated  $\dot{V}O_{2peak}$ . At the very least, this should encourage caution when discussing whether individual patients had an increase in  $\dot{V}O_{2peak}$  after completing exercise-based CR. However, further research is needed to confirm whether other field tests have similar levels of measurement error. Whilst the use of CPET should be encouraged as widely as possible, it is likely that the adoption of CPET prescribed exercise intensity in CR will be dictated by resource implications rather than the potential to confer additional benefit to the patient with CHD or CHF.

One area of exercise prescription in CR that still requires investigation relates to the consistently observed inter-individual difference in training effects. Up to one third of CR patients appear to gain only minimal [or no] improvement in aerobic capacity despite apparent compliance and adherence with their exercise prescription [92]. The reasons for this are not well understood but are likely to include genetic, epigenetic, environmental, nutritional and medical factors, all of which may have a significant impact on exercise prescription for individuals with CHD and CHF [93]. Optimised prescription to maximise outcome may need to take account of these factors, as has personalised medicine in the treatment of disease. Identifying individuals who are the most or least likely to benefit from CR, and how best to optimise exercise prescription, will be the focus of ongoing translation research investigating the determinants and mechanisms of improved exercise capacity in CR [93].

In conclusion, CR is a key treatment vehicle for a broad spectrum of patients with heart disease. Its delivery format has evolved based on a robust body of evidence demonstrating improvements in psychological wellbeing and quality of life. However, CR appears to be underutilised by many clinicians which inevitably has a negative impact on patient outcomes. Further refinements to CR interventions may be required as current evidence indicates that there is a failure to improve peak oxygen uptake following CR, and longer-term survival benefits may be open to question. The “dose”

of exercise provided by community-based CR teams may not be of sufficient stimulus to drive physiological adaptation required to invoke positive changes in aerobic fitness and ultimately improve long-term mortality outcomes. Adopting effective behaviour change strategies are key to maintaining positive lifestyle changes in the longer-term, and strategies adopted must be individually tailored to meet specific patient requirements.

#### **Table legends**

**Table 1.** Favourable impact of exercise on physical and psychological aspects of health.

**Table 2.** Exercise prescription guidelines for the cardiac patient. Based on guidance from the BACPR [6].

## References

1. Tipton CM. The history of "Exercise Is Medicine" in ancient civilizations. *Advances in Phys Ed* 2014; 38:109-17.
2. Heberden W. *Commentaries on the History and Cure of Diseases*: T. Payne; 1802.
3. Mampuya WM. Cardiac rehabilitation past, present and future: an overview. *Cardiovasc Diagnos Ther* 2012; 2:38-49.
4. World Health Organisation. Expert committee on the rehabilitation of patients with cardiovascular disease. Geneva; 1964.
5. World Health Organisation. Rehabilitation after cardiovascular diseases, with special emphasis on developing countries: a report of a WHO Committee. World Health Organisation Technical Report Series 1993. p. 1-122.
6. BACPR. Standards and Core Components for Cardiovascular Disease Prevention and Rehabilitation 2017. Available from: <http://www.bacpr.com>. [accessed March 2020].
7. Anderson L, Oldridge N, Thompson DR, et al. Exercise-Based Cardiac Rehabilitation for Coronary Heart Disease Cochrane Systematic Review and Meta-Analysis. *J Am Coll Cardiol* 2016; 67: 1-12.
8. Long L, Mordi IR, Bridges C, et al. Exercise-based cardiac rehabilitation for adults with heart failure. *Cochrane Database Syst Rev* 2019; 1:Cd003331.
9. AACVPR. AACVPR Cardiac Rehabilitation Resource Manual. Leeds, UK: Human Kinetics; 2006.
10. Savarese G, Lund LH. Global Public Health Burden of Heart Failure. *Cardiac Fail Rev* 2017;3: 7-11.
11. Ponikowski P, Voors AA, Anker SD, et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology [ESC]. Developed with the special contribution of the Heart Failure Association [HFA] of the ESC. *Eur J Heart Fail* 2016; 18: 891-975.
12. Harriss DJ, Macsween A, Atkinson, G. Ethical Standards in Sport and Exercise Science Research: 2020 Update. *Int J Sports Med* 2019; 40: 813–817.



13. Long L, Anderson L, He j, et al. Exercise-based cardiac rehabilitation for stable angina: systematic review and meta-analysis. *Open Heart*. 2019;6[1]:e000989.
14. Heran BS, Chen J, Ebrahim S, et al. Exercise-based cardiac rehabilitation for coronary heart disease. *Cochrane Database Syst Rev*. 2011;7.
15. Powell R, McGregor G, Ennis S, et al. Is exercise-based cardiac rehabilitation effective? A systematic review and meta-analysis to re-examine the evidence. *BMJ Open*. 2018;8: e019656.
16. Jolliffe J, Rees K, Taylor RR, et al. Exercise-based rehabilitation for coronary heart disease. *Cochrane Library*. 2001.
17. D'Souza SP, Mamas MA, Fraser DG, et al. Routine early coronary angioplasty versus ischaemia-guided angioplasty after thrombolysis in acute ST-elevation myocardial infarction: a meta-analysis. *Eur Heart J*. 2011;32: 972-82.
18. Keeley EC, Boura JA, Grines CL. Primary angioplasty versus intravenous thrombolytic therapy for acute myocardial infarction: a quantitative review of 23 randomised trials. *Lancet* 2003; 361: 13-20.
19. Taylor RS, Sagar VA, Davies EJ, et al. Exercise-based rehabilitation for heart failure. *Cochrane Syst Rev*. 2014: 4.
20. Lavie, CJ, Menezes, AR, De Schutter, et al. Impact of cardiac rehabilitation and exercise training on psychological risk factors and subsequent prognosis in patients with cardiovascular disease. *Can J Card* 2016; 32: S365eS373
21. Dalal HM, Taylor RS, Jolly K, et al. The effects and costs of home-based rehabilitation for heart failure with reduced ejection fraction: The REACH-HF multicentre randomized controlled trial. *Eur J Prev Cardiol* 2019; 26: 262-72.
22. Carroll S, Tsakirides C, Hobkirk J, et al. Differential Improvements in Lipid Profiles and Framingham Recurrent Risk Score in Patients With and Without Diabetes Mellitus Undergoing Long-Term Cardiac Rehabilitation. *Arch Phys Med Rehab*. 2011; 92: 1382-1387.
23. Gayda M, Brun C, Juneau M, et al. Long-term cardiac rehabilitation and exercise training programs improve metabolic parameters in metabolic syndrome patients with and without coronary heart disease. *Nutr Metab Cardiovasc Dis* 2008;18:142-151.
24. Reibis R, Treszl A, Bestehorn K, et al. Comparable short-term prognosis in diabetic and non-diabetic patients with acute coronary syndrome after cardiac rehabilitation. *Eur J Prev Cardiol* 2012;19: 15-22.
25. Lavie CJ, Morshedi-Meibodi A, et al. Impact of Cardiac Rehabilitation on Coronary Risk Factors, Inflammation, and the Metabolic Syndrome in Obese Coronary Patients. *J Cardiomet Syndr* 2008;3: 36-40.
26. van Tol BAF, Huijsmans RJ, Kroon DW, et al. Effects of exercise training on cardiac performance, exercise capacity and quality of life in patients with heart failure: A meta-analysis. *Eur J Heart Fail* 2006;8: 841-850.
27. Hambrecht R, Adams V, Erbs S, et al. Regular Physical Activity Improves Endothelial Function in Patients With Coronary Artery Disease by Increasing Phosphorylation of Endothelial Nitric Oxide Synthase. *Circulation*. 2003;107:3152-3158.
28. Zheng H, Luo M, Shen Y, et al. Effects of 6 months exercise training on ventricular remodelling and autonomic tone in patients with acute myocardial infarction and percutaneous coronary intervention. *J Rehabil Med* 2008;40: 776-779.

29. Hambrecht R, Walther C, Möbius-Winkler S, et al. Percutaneous Coronary Angioplasty Compared With Exercise Training in Patients With Stable Coronary Artery Disease: A Randomized Trial. *Circulation* 2004;109: 1371-1378.
30. Lewinter C, Doherty P, Gale CP, et al. Exercise-based cardiac rehabilitation in patients with heart failure: a meta-analysis of randomised controlled trials between 1999 and 2013. *Eur J Prev Cardiol.* 2014, 6: 322-335.
31. Sandercock G, Cardoso F, Almodhy M, et al. Cardiorespiratory fitness changes in patients receiving comprehensive outpatient cardiac rehabilitation in the UK: a multicentre study. *Heart.* 2013;99: 785-790.
32. Sandercock G, Hurtado V, Cardoso F. Changes in cardiorespiratory fitness in cardiac rehabilitation patients: A meta-analysis. *Int J Cardiol* 2011 [0]:220-226.
33. Sandercock G, Cardoso F, Almodhy M. Cardiorespiratory fitness changes in patients receiving comprehensive outpatient cardiac rehabilitation in the UK: a multicentre study. *Heart.* 2013;99: 1298-1299.
34. Ingle L, Carroll S. Cardiac rehabilitation and exercise training. *Heart* 2013;56.
35. Khushhal A, Nichols S, Carroll S, et al. Insufficient exercise intensity for clinical benefit? Monitoring and quantification of a community-based Phase III cardiac rehabilitation programme: A United Kingdom perspective. *Plos One* 2019;14: e0217654.
36. Kotseva K, De Backer G, De Bacquer D, et al. Lifestyle and impact on cardiovascular risk factor control in coronary patients across 27 countries: Results from the European Society of Cardiology ESC-EORP EUROASPIRE V registry. *Eur J Prev Cardiol* 2019;26: 824-835.
37. NACR. National Audit of Cardiac Rehabilitation [NACR] Quality and Outcomes Report 2018. Available from: <https://www.bhf.org.uk/information-support/publications/statistics/national-audit-of-cardiac-rehabilitation-quality-and-outcomes-report-2018>. [accessed Jan 2020].
38. West RR, Jones DA, Henderson AH. Rehabilitation after myocardial infarction trial [RAMIT]: multi-centre randomised controlled trial of comprehensive cardiac rehabilitation in patients following acute myocardial infarction. *Heart* 2012;98: 637-644.
39. Naar-King, S., Earnshaw, P, Breckon, J. Towards a Universal Maintenance Intervention: Integrating Motivational Interviewing with Cognitive-Behavioural Strategies for Maintenance of Behaviour Change. *Cog Behav Practice* 2013: 27, 126-137.
40. Foster, C., Hillsdon, M., Thorogood, M., et al Interventions for promoting physical activity. *Cochrane Database of Systematic Reviews* 2005: Issue 1. Art. No.: CD003180. DOI: 10.1002/14651858.CD003180.pub2.
41. Marcus, B., Forsyth, L., Stone, E., et al. Physical activity behavior change: Issues in adoption and maintenance. *Health Psych* 2000: 19: 32–41.
42. Scott, S., Breckon, J.D, Copeland, R. An integrated motivational interviewing and cognitive behavioural intervention promoting physical activity maintenance for adults with chronic health conditions: A feasibility study. *Chronic Illness.* 2018: doi:10.1177/1742395318769370.
43. Hutchison, A., Johnston, L. & Breckon, J. Grounded Theory Based Research within Exercise Psychology: A Critical Review. *Qual Res Psych* 2011: 8: 247-272.
44. Breckon, J.D. Motivational Interviewing to promote physical activity and nutrition behaviour change. In, *Doing Exercise Psychology* [M. Anderson & S. Hanrahan, Eds.]. 2015: Human Kinetics, Champaign, Ill.
45. Ajzen, I. The theory of planned behavior. *Organ Behav Human Decision Processes* 1991; 50, 179–211.

46. Fishbein, M., & Ajzen, I. Predicting and changing behavior: The reasoned action approach. 2011: New York, NY: Taylor & Francis.
47. Schwarzer, R., Lippke, S., Luszczynska, A. Mechanisms of health behavior change in persons with chronic illness or disability: The health action process approach [HAPA]. *Rehabil Psych* 2011; 56, 161–170.
48. Knittle, K., Nurmi, J., Crutzen, R., et al. How can interventions increase motivation for physical activity? A systematic review and meta-analysis. *Health Psych Rev* 2018; 12: 211-230.
49. Hilton, C., Johnston, L. Health psychology: It's not what you do, it's the way that you do it. *Health Psych Open* 2017: 1-10.
50. Lundby C, Montero D, Joyner M. Biology of VO2 max: looking under the physiology lamp. *Acta Physiologica* 2017;220: 218-28.
51. Kavanagh T, Mertens DJ, Hamm LF, et al. Prediction of Long-Term Prognosis in 12 169 Men Referred for Cardiac Rehabilitation. *Circulation* 2002;106: 666-671.
52. Kavanagh T, Mertens DJ, Hamm LF, et al. Peak oxygen intake and cardiac mortality in women referred for cardiac rehabilitation. *J Am Coll Cardiol* 2003;42: 2139-2143.
53. Mancini DM, Eisen H, Kussmaul W, et al. Value of peak exercise oxygen consumption for optimal timing of cardiac transplantation in ambulatory patients with heart failure. *Circulation*. 1991;83: 778-786.
54. Brubaker PH, Kitzman DW. Chronotropic Incompetence: Causes, Consequences, and Management. *Circulation*. 2011;123: 1010-1020.
55. Belardinelli R, Lacalaprice F, Carle F, et al. Exercise-induced myocardial ischaemia detected by cardiopulmonary exercise testing. *Eur Heart J* 2003;24: 1304-1313.
56. Poole DC, Hirai DM, Copp SW, et al. Muscle oxygen transport and utilization in heart failure: implications for exercise [in] tolerance. *Am J Physiol Heart Circ Phys* 2012;302: H1050-H1063.
57. Clark AL, Poole-Wilson PA, Coats AJS. Exercise limitation in chronic heart failure: Central role of the periphery. *J Am Coll Cardiol* 1996;28: 1092-1102.
58. Nichols S, O'Doherty AF, Taylor C, et al. Low skeletal muscle mass is associated with low aerobic capacity and increased mortality risk in patients with coronary heart disease - a CARE CR study. *Clin Physiol Funct Imaging*. 2018, 22:234-240.
59. Fülster S, Tacke M, Sandek A, et al. Muscle wasting in patients with chronic heart failure: results from the studies investigating co-morbidities aggravating heart failure [SICA-HF]. *Eur Heart J*; 2013;34:512-519.
60. Aspenes ST, Nilsen TI, Skaug EA, et al. Peak oxygen uptake and cardiovascular risk factors in 4631 healthy women and men. *Med Sci Sports Exerc* 2011;43:1465-1473.
61. Keteyian SJ, Brawner CA, Savage PD, et al. Peak aerobic capacity predicts prognosis in patients with coronary heart disease. *Am Heart J* 2008;156:292-300.
62. Shephard RJ. Maximal oxygen intake and independence in old age. *Brit J Sports Med* 2009;43: 342-346.
63. Nichols S, Taylor C, Page R, et al. Is Cardiorespiratory Fitness Related to Cardiometabolic Health and All-Cause Mortality Risk in Patients with Coronary Heart Disease? A CARE CR Study. *Sports Med Open* 2018;4:22-32.
64. Vanhees L, Fagard R, Thijs L, et al. Prognostic value of training-induced change in peak exercise capacity in patients with myocardial infarcts and patients with coronary bypass surgery. *Am J Cardiol* 1995; 76: 1014-1019.

65. Martin BJ, Arena R, Haykowsky M, et al. Cardiovascular fitness and mortality after contemporary cardiac rehabilitation. *Mayo Clin Proc* 2013;88:455-463.
66. Nichols S, Taylor C, Goodman T, et al. Routine exercise-based cardiac rehabilitation does not increase aerobic fitness: A CARE CR Study, *Int J Cardiol* 2020, in press.
67. Price KJ, Gordon BA, Bird SR, et al. A review of guidelines for cardiac rehabilitation exercise programmes: Is there an international consensus? *Eur J Prev Cardiol.* 2016;23: 1715-1733.
68. Turk-Adawi K, Supervia M, Pesah E, et al. Availability and delivery of cardiac rehabilitation in the Eastern Mediterranean Region: How does it compare globally? *Int J Cardiol* 2019;285:147-53.
69. British Heart Foundation. National Audit of Cardiac Rehabilitation [NACR] Quality and Outcomes Report 2019. Available at: <https://www.bhf.org.uk>. [accessed Jan 2020].
70. Smith KM, Arthur HM, McKelvie RS, et al Differences in sustainability of exercise and health-related quality of life outcomes following home or hospital-based cardiac rehabilitation. *Eur J Prev Cardiol* 2004; 11: 313-319.
71. Kodama S, Saito K, Tanaka S, et al. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA.* 2009;301: 2024-2035.
72. Martin B-J, Arena R, Haykowsky M, et al. Cardiovascular Fitness and Mortality After Contemporary Cardiac Rehabilitation. *Mayo Clinic Proceedings* 2013;88: 455-463.
73. ACPICR. Standards for physical activity and exercise in the cardiac population 2015. Available at: <http://acpicr.com> [accessed Nov 2019].
74. Mezzani A, Hamm LF, Jones AM, et al. Aerobic exercise intensity assessment and prescription in cardiac rehabilitation: a joint position statement of the European Association for Cardiovascular Prevention and Rehabilitation, the American Association of Cardiovascular and Pulmonary Rehabilitation and the Canadian Association of Cardiac Rehabilitation. *Eur J Prev Cardiol.* 2013;20: 442-467.
75. Garber CE, Blissmer B, Deschenes MR, et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc* 2011;43: 1334-1359.
76. Franklin BA, Thompson PD, Al-Zaiti SS, et al. Exercise-Related Acute Cardiovascular Events and Potential Deleterious Adaptations Following Long-Term Exercise Training: Placing the Risks Into Perspective-An Update: A Scientific Statement From the American Heart Association. *Circulation.* 2020:CIR0000000000000749-CIR.
77. Liou K, Ho S, Fildes J, et al. High Intensity Interval versus Moderate Intensity Continuous Training in Patients with Coronary Artery Disease: A Meta-analysis of Physiological and Clinical Parameters. *Heart Lung Circ* 2016;25:166-174.
78. Gomes Neto M, Duraes AR, Conceicao LSR, et al. High intensity interval training versus moderate intensity continuous training on exercise capacity and quality of life in patients with heart failure with reduced ejection fraction: A systematic review and meta-analysis. *Int J Cardiol* 2018;261:134-141.
79. Gomes-Neto M, Duraes AR, Reis H, et al. High-intensity interval training versus moderate-intensity continuous training on exercise capacity and quality of life in patients

- with coronary artery disease: A systematic review and meta-analysis. *Eur J Prev Cardiol* 2017;24:1696-1707.
80. Elliott AD, Rajopadhyaya K, Bentley DJ, et al. Interval training versus continuous exercise in patients with coronary artery disease: a meta-analysis. *Heart Lung Circ* 2015;24:149-157.
  81. Quindry JC, Franklin BA, Chapman M, et al. Benefits and Risks of High-Intensity Interval Training in Patients With Coronary Artery Disease. *Am J Cardiol* 2019;123:1370-1377.
  82. Wewege MA, Ahn D, Yu J, et al. High-Intensity Interval Training for Patients With Cardiovascular Disease-Is It Safe? A Systematic Review. *J Am Heart Assoc.* 2018;7:e009305.
  83. Rognmo O, Moholdt T, Fau - Bakken H, et al. Cardiovascular risk of high- versus moderate-intensity aerobic exercise in coronary heart disease patients. *Circulation* 2012; 136-140.
  84. Conraads VM, Pattyn N, De Maeyer C, et al. Aerobic interval training and continuous training equally improve aerobic exercise capacity in patients with coronary artery disease: the SAINTEX-CAD study. *Int J Cardiol* 2015;179:203-210.
  85. Ellingsen O, Halle M, Conraads V, et al. High-Intensity Interval Training in Patients With Heart Failure With Reduced Ejection Fraction. *Circulation* 2017;135: 839-849.
  86. Taylor JL, Holland DJ, Spathis JG, et al. Guidelines for the delivery and monitoring of high intensity interval training in clinical populations. *Prog Cardiovasc Dis* 2019;62:140-146.
  87. Ribeiro PAB, Boidin M, Juneau M, et al. High-intensity interval training in patients with coronary heart disease: Prescription models and perspectives. *Ann Phys Rehabil Med* 2017;60: 50-57.
  88. Pymer S, Nichols S, Prosser J, et al. Does exercise prescription based on estimated heart rate training zones exceed the ventilatory anaerobic threshold in patients with coronary heart disease undergoing usual-care cardiovascular rehabilitation? A United Kingdom perspective. *Eur J Prev Cardiol.* 2019:2047487319852711.
  89. Hansen D, Bonne K, Alders T, et al. Exercise training intensity determination in cardiovascular rehabilitation: Should the guidelines be reconsidered? *Eur J Prev Cardiol.* 2019;26: 1921-1928.
  90. Nichols S, Gleadall-Siddall DO, Antony R, et al. Estimated peak functional capacity: an accurate method for assessing change in peak oxygen consumption after cardiac rehabilitation? *Clin Phys Funct Imaging* 2018; 38: 681-688.
  91. De Schutter A, Kachur S, Lavie CJ, et al. Cardiac rehabilitation fitness changes and subsequent survival. *Eur Heart J Qual Care Clin Outcomes.* 2018;4: 173-179.
  92. Gevaert AB, Adams V, Bahls M, et al. Towards a personalised approach in exercise-based cardiovascular rehabilitation: How can translational research help? A 'call to action' from the Section on Secondary Prevention and Cardiac Rehabilitation of the European Association of Preventive Cardiology. *Eur J Prev Cardiol.* 2019:2047487319877716.
  93. Sharman JE, La Gerche A, Coombes JS. Exercise and cardiovascular risk in patients with hypertension. *Am J Hypertension* 2015; 28: 147–158

