

**Disorganized attachment predicts body mass index via
uncontrolled eating**

WILKINSON, L.L., ROWE, A.C. and MILLINGS, Abigail

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/27339/>

This document is the Accepted Version [AM]

Citation:

WILKINSON, L.L., ROWE, A.C. and MILLINGS, Abigail (2020). Disorganized attachment predicts body mass index via uncontrolled eating. *International Journal of Obesity*, 44 (2), 438-446. [Article]

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

1
2 Disorganized attachment predicts body mass index via uncontrolled eating
3

4 Laura L. Wilkinson^a, Angela C. Rowe^b, Abigail Millings^c
5

6 ^aDepartment of Psychology, Swansea University, Swansea, UK
7

8 ^bSchool of Psychological Science, University of Bristol, Bristol, UK
9

10 ^cDepartment of Psychology, University of Sheffield, Sheffield, UK
11

12 *Corresponding author: L.L.Wilkinson@Swansea.ac.uk
13

14 Running header: Disorganized attachment, eating behaviour and BMI
15

16 Competing interests statement: The authors have no competing interests to declare.
17

18
19
20
21
22
23
24
25

26 **Abstract**

27 **Background/Objectives:** Previous research has demonstrated relationships
28 between attachment orientations (expectations of ourselves and others in
29 interpersonal relationships), eating behaviours and obesity. However, such research
30 has been limited to investigations of 'organised' forms of attachment orientations
31 (reflecting coherent and predictable patterns of behaviour). Theoretically, aberrant
32 eating behaviours and body mass index, should also be related to 'disorganized
33 attachment.'

34 **Subjects:** Here we test these relationships for the first time in a general population.
35 Secondary data analyses of a pre-existing dataset were conducted ($N = 537$).

36 **Methods used:** Questionnaire measures of organised (avoidant and anxious) and
37 disorganized attachment were included alongside eating behaviour measures
38 (emotional eating, uncontrolled eating and cognitive restraint) and body mass index
39 (BMI).

40 **Results:** Parallel multiple mediation analysis (PROCESS) showed that uncontrolled
41 eating (but not emotional eating or cognitive restraint) significantly mediated a
42 relationship between disorganized attachment and body mass index (significant
43 indirect relationship; LLCI = .02 ULCI = .16) when both attachment anxiety and
44 avoidance were included as covariates.

45 **Conclusions:** We suggest that the mechanism underpinning this indirect
46 relationship is a form of maladaptive affect regulation, but that the behavioural
47 motivators differ from those observed in anxiously attached individuals. Rather than
48 eating being a premeditated strategy used by individuals high in disorganized
49 attachment to manage emotion, opportunities to eat are simply taken as they present
50 themselves. Professionals engaged in addressing eating problems and weight

51 management should consider attachment orientations in their patient assessments
52 and be mindful that attachment disorganized individuals are especially likely to
53 engage in uncontrolled eating behaviours that are associated with a higher BMI.

54

55 **Key words:** disorganized attachment; attachment anxiety; uncontrolled eating;
56 emotional eating; cognitive restraint; body mass index

57

58

59 **Introduction**

60 Recent estimates have suggested that overweight and obesity will affect 2.7 billion
61 people worldwide by 2025 ¹. Overweight and obesity are associated with an
62 increased risk of cardiovascular disease, type 2 diabetes and cancer amongst other
63 health issues ². As the upward trend in obesity continues ³, the need for effective
64 interventions is of high priority ⁴.

65 ‘Attachment orientation’ is one psychological factor that has been shown to
66 relate to obesity and overweight in adults ⁵. Attachment orientation is a term used to
67 describe the set of expectations that we all possess about how we and others
68 behave in inter-personal relationships. These mental models are thought to be
69 established early in life and reflect interactions with caregivers ⁶. Adult attachment
70 orientations reflects the dynamics and feelings associated with our most important
71 long-term relationships in life, including those from early life ⁷.

72 Currently, adult attachment orientations tends to be conceptualised in terms of
73 two dimensions ⁷. These are attachment *anxiety* which is characterised by a fear of
74 abandonment and attachment *avoidance* which is characterised by a fear of
75 intimacy. A low score on both of these dimensions indicates ‘attachment security’.

76 Whilst a high score on either or both of these dimensions indicates 'attachment
77 insecurity'.

78 A recent meta-analysis showed that, in a general population, higher
79 attachment insecurity was associated with more unhealthy eating behaviours (e.g.,
80 disinhibited eating, uncontrolled eating, emotional eating amongst others)⁸. The
81 strongest relationship was between attachment anxiety and unhealthy eating
82 behaviours, with effects tending to be of medium effect size. Importantly, disinhibited
83 eating (a general propensity to engage in periodic overeating⁹) and emotional eating
84 (episodic overeating in response to negative affect¹⁰) have been found to mediate a
85 relationship between attachment and BMI,^{11, 12} respectively. This meta-analysis⁸
86 also showed that higher attachment avoidance related to more unhealthy eating
87 behaviours, however, the effect size for this relationship was small and the weakest
88 amongst those examined (attachment insecurity, anxiety, avoidance and
89 fearfulness).

90 The mechanism underpinning these associations is thought to be affect
91 regulation; anxiously attached individuals are relatively poor at managing their
92 emotions by comparison to their attachment secure counterparts. When upsetting
93 events take place, they may seek support, but the interaction is characterised by
94 mistrust and fear of rejection rather than reassurance. Moreover, the attachment
95 system is hyperactivated leading to a hyper-vigilance to potentially upsetting stimuli
96^{13, 14}. To 'break the cycle' of hyperactivation, highly anxiously attached individuals
97 rely on external sources of affect regulation such as food¹⁵. Indeed, in recent work, it
98 was shown that attachment anxiety was specifically related to an inability to both
99 disengage from the source of upset and engage in goal-oriented behaviour, which
100 was in turn related to eating in response to stress and body mass index¹². By

101 contrast, attachment avoidance is associated with deactivating strategies, that is, the
102 avoidance of emotions and suppression of stress and help-seeking (Mikulincer and
103 Orbach, 1995). It has been suggested that this maladaptive approach does not
104 actually eliminate stress and therefore remaining physiological stress markers and
105 negative affect precipitate a need to engage in external affect regulation (i.e., eating).
106 It should be noted that support for this theoretical explanation linking attachment
107 avoidance and eating behaviour is scant compared with the better understood
108 relationship between attachment anxiety and eating behaviour.

109 However, these findings focus only on ‘organised’ forms of attachment
110 orientations. That is, where mental models and strategies are ‘coherent’ and allow
111 individuals to select (which may be explicit or implicit) strategies to manage
112 situations that are adaptive and predictable based on their experience of inter-
113 personal relationships ¹⁶. For example, it is logical and adaptive for the child of a
114 neglectful caregiver to deactivate their attachment system as in attachment
115 avoidance and seek to minimise intimacy. Even when individuals are high in both
116 attachment avoidance and anxiety, an orientation called fearful-avoidance, they will
117 use the hyperactivating and deactivating strategies intermittently. By contrast, adult
118 disorganized attachment is characterised by a general fear of romantic attachment
119 figures and refers to a lack of coherent and predictable strategies to manage the self
120 and others ¹⁶. This leads to contradictory and confused behaviour in response to
121 distress; approach behaviours may be initiated but are left incomplete due to fear
122 and a simultaneous desire to distance themselves ¹⁷. A recent development has
123 been the measurement of disorganisation as a construct that is separate and
124 additional to the organised dimensions of adult attachment ¹⁶.

125 While attachment avoidance reflects punitive or neglectful relationship
126 experiences and attachment anxiety reflects inconsistency in relational experiences,
127 disorganized attachment is related to child abuse, loss, trauma and otherwise
128 frightening interpersonal experiences. Research has shown that 80% of maltreated
129 children had a disorganized attachment pattern ¹⁸. Such adverse childhood
130 experiences are also relevant to the understanding of obesity ¹⁹. A systematic review
131 and meta-analysis ²⁰ not only showed a clear relationship between childhood
132 experience of abuse and adult obesity but also severe abuse was significantly more
133 related to obesity than less severe abuse. Suggested mechanisms underpinning this
134 relationship included maladaptive coping responses, stress and emotional
135 perturbations.

136 Considering the shared aetiology of disorganized attachment and obesity,
137 with respect to the experience of abuse in childhood, we investigated whether a
138 relationship between disorganized attachment, eating behaviour and BMI existed.
139 Here, our approach was to focus on a general population and sub-clinical eating
140 behaviours, of the kind investigated previously in relation to attachment anxiety and
141 its relationship with BMI ¹². The ability to investigate this relationship in a large
142 general population has been facilitated by the recent development of the adult
143 disorganized attachment scale ¹⁶, which assesses the construct of disorganisation as
144 a continuous dimension. Importantly, for the assessment of disorganized attachment
145 in a general population, this approach can assess a range of levels, including
146 relatively low scores.

147 Therefore, we hypothesised that higher disorganized attachment scores
148 would relate to higher BMI and that this relationship would be mediated by measures
149 of disinhibited eating. Specifically, we expected this relationship to exist independent

150 of attachment anxiety (i.e., when attachment anxiety is controlled for). Following
151 previous research, we examined the construct of disinhibited eating with measures
152 of uncontrolled eating and emotional eating. Uncontrolled eating is a measure of
153 opportunistic eating in response to both internal (e.g., extreme hunger) and external
154 (e.g., the smell of a tempting food) cues to eat ¹⁰. This is a separate construct to
155 eating as a response to negative emotion (emotional eating).

156 Given the contradictory nature of patterns of behaviour associated with
157 disorganized attachment, we also tentatively hypothesised that a relationship with
158 BMI might also be mediated by dietary restraint. Dietary restraint refers to the
159 cognitive control over intake in order to influence body weight or shape ¹⁰ and has
160 been conceptualised as antagonistic to disinhibited eating ²¹. One possibility is that
161 the hypothesised use of eating to regulate affect by attachment disorganized
162 individuals might precipitate reactionary dietary restraint (which has been shown to
163 ironically lead to further disinhibited eating behaviours ³¹).

164 In addition, using an exploratory approach based on the primary hypotheses
165 being supported, we produced a comprehensive model that incorporated both
166 disorganized attachment *and* attachment anxiety as predictors of body mass index
167 via disinhibited eating behaviours.

168

169

170 **Methods**

171 This study is a secondary data analysis of an unpublished dataset collected for a
172 different purpose. A brief description of the primary study and its results can be
173 found in the supplementary information. The primary hypotheses were pre-registered
174 with the Open Science Framework (osf.io/2dr74) following data collection but before

175 the data analyses associated with this paper. The dataset has been deposited for
176 open access with the Open Science Framework
177 (https://osf.io/2dr74/?view_only=a7bfeea11614401ca9464545ab4f620c).

178

179 *Participants*

180 A total of 752 participants began the study but 163 participants failed to
181 complete all of the measures and were therefore excluded. We excluded a further 52
182 participants from our dataset because either they reported a height and weight that
183 yielded an ambiguous or spurious BMI value (e.g., unlikely to be possible) or a BMI
184 that was valid but considered to be below healthy-range (i.e., below 18) to minimise
185 risk of inadvertently including disordered individuals in our sample. This resulted in a
186 sample size of 537 (Male = 126, female = 404, other = 6, prefer not to say = 1; mean
187 age = 25.5 years old, $SD = 9.9$). This sample size met with our requirements for
188 adequately powering our analyses to detect effects; considering the novelty of our
189 hypotheses a prudent approach was to power for small effect sizes within our model.
190 Following Fritz and MacKinnon's ²² guidance for detecting mediated effects with
191 sample sizes providing .8 power using bias-corrected bootstrap approaches, we
192 required at least a sample size of at least 462. To be eligible for the study,
193 participants had to be over the age of 18 years old and have no current or previous
194 diagnosis of an eating disorder.

195 Participants were recruited in two cohorts. In the first cohort ($n = 130$
196 completers), a consortium-led approach (see Button *et al.* ²³) was taken by
197 university students based at the University of Sheffield ($n = 63$ completers) and
198 Swansea University ($n = 67$ completers). The cohort 1 sample consisted mainly of
199 university students and staff, as the study was advertised to potential participants

200 who could attend a session at the universities via campus posters and student-study
201 participation systems only. The second cohort ($n = 459$) provided only self-report
202 height and weight and therefore were able to complete the study entirely online.
203 They were recruited via social media, posters and student-study participation
204 systems.

205 For cohort 1, ethical approval was obtained from local human research ethics
206 committees at both sites. For cohort 2, an ethical amendment outlining a change to
207 an online only approach was approved by local human research ethics committees
208 at both sites. All participants provided informed consent.

209

210 *Measures*

211 *Disorganized attachment* The 9-item Disorganized Attachment scale¹⁶ was
212 used. This consists of a single subscale and for the current study Cronbach's alpha
213 for this measure was .88. Participants rated their agreement to statements about
214 their general experience of relationships (as opposed to a specific current
215 relationship). For example, "I never know who I am with romantic partners".
216 Responses were provided on a 7-point Likert scale, from 1 (strongly disagree) to 7
217 (strongly agree).

218

219 *Attachment anxiety and avoidance*. The 12-item short-form Experiences in
220 Close Relationships questionnaire²⁴ was used to assess attachment anxiety (6
221 items) and attachment avoidance (6 items). For the current study, Cronbach's alpha
222 for the attachment anxiety subscale was .88 and for the attachment avoidance
223 subscale was .84. On a seven-point scale ranging from strongly disagree to strongly
224 agree, participants stated their level of agreement with statement referring to the

225 experiences of interpersonal relationships. For example, "I worry a fair amount about
226 losing my relationships" (attachment anxiety) and "I don't feel comfortable opening
227 up to others" (attachment avoidance).

228

229 *Eating style.* The 18-item short-form three factor eating questionnaire ¹⁰ was
230 used to assess three constructs, cognitive restraint, emotional eating and
231 uncontrolled eating. Participants responded with the extent to which statements
232 applied to them on a 4-point scale ranging from definitely true to definitely false or a
233 variant of this scale depending on the question. For the cognitive restraint scale (6
234 items; Cronbach's alpha for this study = .85), statements concerned the extent to
235 which they consciously apply restraint to their eating behaviour (e.g., "I consciously
236 hold back at meals in order not to gain weight"). For the uncontrolled eating scale (9
237 items; Cronbach's alpha for this study = .86), statements concerned the extent to
238 which they lost control over their eating behaviour (e.g., "Sometimes when I start
239 eating, I just can't seem to stop"). For the emotional eating scale (3 items;
240 Cronbach's alpha for this study = .87), statements concerned eating in response to
241 emotional states (e.g., "When I feel lonely, I console myself with food").

242

243 *Anthropometric measures:* For cohort 1, participants attended a lab session
244 where their height and weight were measured by a researcher using a portable
245 stadiometer and non-medical grade weighing scales respectively. For cohort 2,
246 current height and weight were self-reported by participants using their preferred
247 units. These were converted to metric measures. Across both cohorts, height and
248 weight measures were used to calculate BMI (kg/m^2).

249

250 *Procedure*

251 The study was hosted on Qualtrics survey software (Qualtrics, Provo, UT, USA) and
252 participants had to click an anonymised link to access the study. They were asked to
253 read an information sheet and informed consent screen and to tick a box if they
254 consented. Following this they were asked to complete all measures (those outlined
255 here for the current study as well as those outlined in the supplementary information
256 for the full version of the study) as well as basic demographic questions. Upon
257 completion participants were either asked about their availability and contact
258 information for a follow-up session for the measurement of height and weight and
259 then debriefed (cohort 1) or asked to self-report their height and weight and then
260 debriefed.

261

262 *Data analysis*

263 Following Price *et al.*²⁵, we merged the BMI data from the two cohorts. This
264 decision was made on the basis that self-reported BMI and researcher-measured
265 BMI have been found to be highly-correlated for both younger²⁶ and older adults²⁷.
266 Nevertheless, height can be overestimated and weight can be underestimated,
267 therefore cohort (i.e., self-report vs. measured) was entered as a covariate into our
268 analyses, however, as it was not a significant covariate and made no difference to
269 the results of our analyses it was trimmed from our final models.

270 In order to examine our primary hypotheses, a parallel mediation model was
271 conducted. This approach was selected because it allows for the simultaneous entry
272 of multiple mediators within a single model, such that the independent contributions
273 of each mediator as part of an indirect pathway can be assessed. It also allowed for
274 covariates to be entered into the model.

275 Our model was conducted with disorganized attachment (predictor), BMI
276 (outcome) and parallel multiple mediators (cognitive restraint, emotional eating and
277 uncontrolled eating). In order to isolate an effect of disorganized attachment we
278 controlled for both attachment avoidance and anxiety. Also, following Wilkinson *et al.*
279 ^{11, 12}, we controlled for gender and age. This model was conducted using PROCESS
280 v3.1 ²⁸. The covariates were controlled for at the level of both the mediator and the
281 outcome.

282 Following this, a comprehensive model was produced that incorporated both
283 findings associated with disorganized attachment (i.e., the results of the first
284 analysis) and a replication of the significant indirect relationship between attachment
285 anxiety, disinhibited eating measures and BMI reported in previous research ^{11, 12}.
286 This approach allows, for the first time, for the indirect effect of attachment anxiety
287 on BMI to be tested whilst also controlling for disorganized attachment.

288 Therefore, path analysis was conducted using IBM SPSS AMOS. v.25.0. We
289 included disorganized attachment, attachment anxiety, attachment avoidance, age
290 and gender as exogenous variables, and uncontrolled eating, emotional eating and
291 BMI as endogenous variables. Cognitive restraint was not included, as our primary
292 analysis (above) failed to support a role for this factor. Covariance relationships were
293 specified between each of the exogenous variables and error terms were included
294 for each endogenous variable. Direct relationships were specified from each
295 exogenous variable to each endogenous variable; therefore both direct and indirect
296 relationships could be examined. Both of our model ran 5000 bootstrap samples and
297 95% confidence intervals are reported.

298 **Results**

299 Means and standard deviations for each measure and correlations between them
300 can be found in Tables 1 and 2 respectively.

301

302 *Confirmatory analyses of primary pre-registered hypotheses: Disorganized*
303 *attachment, eating behaviours (cognitive restraint, emotional and uncontrolled*
304 *eating) and BMI*

305

306 Our parallel multiple mediation model (Fig 1) showed that there was no significant
307 direct relationship between disorganized attachment and BMI when no mediators
308 were included in the model (total effects) and this remained the case when mediators
309 were included (direct effects). However, a significant indirect effect that ran between
310 disorganized attachment and BMI via uncontrolled eating was identified. There were
311 no significant indirect effects associated with cognitive restraint or emotional eating.

312

313 *Path analysis testing a comprehensive model of the relationships between*
314 *attachment orientations, disinhibited eating behaviours and BMI*

315

316 Our path analysis revealed the significant indirect pathway between disorganized
317 attachment and BMI via uncontrolled eating that was observed within our primary
318 analysis. Also, a significant indirect pathway between attachment anxiety and BMI
319 via emotional eating was identified. No direct effects between attachment
320 orientations and BMI (without mediators included) were observed. Figure 2 provides
321 a visualisation of our model, however, for the sake of legibility we have only included
322 key relationships and statistics. Comprehensive information regarding direct

323 relationships relating to our specific hypotheses can be found in Tables 3 and 4 and
324 other direct relationships can be found in the supplementary information file.

325 Overall a good model fit was indicated with a chi square goodness of fit
326 statistic that was not significant ($p = .672$), a comparative fit index (CFI) of 1.00 and a
327 root mean square error of approximation (RMSEA) of .00.

328

329 **Discussion**

330 This study has identified disorganized attachment as a novel predictor of BMI in a
331 non-clinical population via the mechanism of uncontrolled eating. Our hypotheses
332 were partially confirmed. The relationship between disorganized attachment and BMI
333 was mediated by uncontrolled eating only (and not emotional eating or cognitive
334 restraint). This highlights the importance of accounting for uncontrolled eating in
335 individuals who present to health professionals with disorganized attachment and a
336 high BMI.

337 For the first time, we can provide evidence to support a comprehensive
338 theoretical model that incorporates both organised and disorganized attachment
339 orientations and their relationship to eating psychopathology and BMI (Fig 2).
340 Importantly, this speaks to a broader theoretical model that links attachment
341 orientations to health outcomes more generally ¹⁵, suggesting that an extension is
342 required, such that it goes beyond organised forms of attachment orientations to
343 include effects of disorganisation on health outcomes.

344 The most theoretically sound explanation for why disorganized attachment
345 relates to BMI via uncontrolled eating is that these behaviours serve affect regulatory
346 need. Specifically, whilst the motivator for eating might not be emotion, it is some
347 other eating-related cue, the incidental reward associated with such eating

348 behaviours may have the unintended consequence of regulating affect. This
349 suggestion is consistent with Paetzold *et al.*'s¹⁶ view of disorganized attachment in
350 two ways; firstly, disorganized attachment is associated with problematic emotion
351 regulation - anger and hostility. When these are experienced they are associated
352 with impulsivity and general negative emotionality, which in turn may precipitate a
353 need for affect regulation. However, due to their conflict around support seeking (a
354 general fear of relationships "encourages simultaneous and confused approach and
355 avoidance of the attachment figure for support and solace in times of need"¹⁶),
356 highly disorganized individuals are likely to receive less support and perceive
357 support as poorer than less disorganized individuals. External forms of affect
358 regulation, such as disinhibition related to food consumption, may provide one of the
359 few ways for disorganized individuals to manage their emotions.

360 Secondly, the characteristic incoherence of a disorganized attachment
361 strategy is borne out here; specifically, it is likely that the affect regulatory effects of
362 uncontrolled eating behaviours are reinforced but have failed to be translated into a
363 coherent strategy such as 'emotional eating'. Future research is required to test
364 these theoretical pathways; specifically, approaches to emotion regulation that might
365 explain the relationship between disorganized attachment orientation and such
366 eating behaviour. Moreover, the extent to which there is shared (or not) aetiology
367 with attachment anxiety merits scrutiny.

368 For anxiously attached individuals, the indirect relationship between
369 attachment anxiety and BMI via emotional eating is consistent with previous
370 research¹². However, it is notable, that it in our model it was necessary to specify a
371 relationship running from emotional eating to uncontrolled eating (for improved
372 model fit). This is consistent with the broader literature; recent commentary by Van

373 Strien ²⁹ reports on the co-occurrence of such constructs (this paper refers to
374 external eating which is conceptually similar to uncontrolled eating) and explains it in
375 terms of the ‘escape-of-self-awareness’ theory ³⁰, whereby emotional eaters narrow
376 their attention to their immediate environment. Indeed, previous work suggests that
377 attachment anxiety is associated with an inability to disengage with a source of upset
378 ¹² and eating in this context allows for escape/ disengagement.

379 Contrary to one of our hypotheses, there was no significant relationship
380 between disorganized attachment and cognitive restraint. This work is inconsistent
381 with traditional ‘restraint theory’ ³¹, which suggests that disinhibited eating is a
382 response to restraint. Rather, relationships with disinhibited eating behaviours were
383 shown without respective relationships with cognitive restraint. This is more
384 consistent with psychosomatic theory, which suggests that personality traits and
385 psychopathology can cause disinhibited eating irrespective of restraint status ¹⁰. One
386 speculation is that the interplay between restrained eating and disinhibited eating
387 described within traditional restraint theory actually represents a relatively coherent
388 strategy, whereby disinhibition is a predictable response to restraint and restraint is a
389 predictable response to disinhibition. Such a coherent pattern of behaviour is
390 uncharacteristic of attachment disorganized individuals. Given the novelty of this
391 finding and our limited understanding of disorganized attachment relative to
392 organised attachment orientations, future research should attempt to replicate this
393 finding.

394 Some of our findings here are also contrary to a recent meta-analysis ⁸ which
395 showed a significant relationship between attachment avoidance and emotional
396 eating. We failed to find a significant relationship of this kind despite our larger
397 sample size. A greater understanding of the role of attachment avoidance in models

398 of attachment orientations and eating behaviour is needed, especially given recent
399 results suggesting that attachment avoidance actually relates to restricted eating via
400 the mechanism of 'emotional cut-off' ³². This latter finding and the failure to find a
401 significant relationship between attachment avoidance and emotional eating in the
402 current study are generally more consistent with our theoretical understanding of
403 attachment avoidance; attachment avoidant individuals engage in deactivating
404 strategies that actively minimise the experience of negative affect and cognitions ³³,
405 therefore there is no requirement to affect regulate using food in the first place.

406 It is notable that our findings were evident even when the mean disorganized
407 attachment score for the overall sample was relatively low on the scale (2.82
408 arbitrary units; a.u.) and the maximum (6.67 a.u.) was just below the top of the scale
409 (7 a.u.). Stronger relationships may be evident in a more clinically-relevant sample,
410 for example bariatric-metabolic surgery patients. Previous research has shown that
411 attachment insecurity is more prevalent in bariatric-metabolic surgery patients than in
412 lean control populations ^{34, 35} and that there is a relationship between attachment
413 anxiety and binge eating mediated by emotion-regulation difficulties for surgery
414 candidates ³⁶. Future weight loss surgery research should consider a role for
415 disorganized attachment given that eating psychopathologies (including those
416 related to uncontrolled eating) initially decrease following surgery but then
417 significantly and problematically increase in follow ups beyond a year after surgery
418 ^{37, 38}.

419 Previous research has shown that maternal attachment anxiety is associated
420 with the use of feeding practices that promote child emotional eating, amongst other
421 bidirectional effects ³⁹. Future research might consider exploring the effect of
422 parental disorganized attachment on child feeding practices. More generally it has

423 been found that parents with disorganized attachment behave in ways that may
424 engender disorganized attachment in a child ⁴⁰. Whether eating behaviours form
425 some part of this complex relationship remains to be explored.

426 More generally it should be noted that only a relatively small percentage of the
427 variance associated with BMI was explained by our models (16 -18%), which is in
428 line with previous studies ^{11, 12}. Furthermore, only correlational relationships were
429 examined here and therefore causal inferences cannot be made. One possibility is
430 reverse causality which would suggest that a higher BMI leads to disinhibited eating
431 behaviours and this affects attachment orientations. In all likelihood, a more complex
432 non-recursive relationship exists which future longitudinal work might consider. A
433 strength of our study is the inclusion of a sub-set of researcher measured height and
434 weight to derive the body mass index rather than just self-reported height and weight
435 alone.

436 This work represents a theoretical advancement of this area in line with the
437 general attachment literature which is growing with respect to our understanding of
438 adult disorganized attachment. Future work should build on the models presented
439 here to include additional demographic and individual difference level factors, for
440 example socio-economic status, which is associated with both attachment ⁴¹ and
441 BMI ⁴² and/ or neuroticism which is also associated with both eating behaviours ⁴³
442 and attachment orientations ⁴⁴.

443 In addition, given that attachment orientations are fundamentally rooted in
444 experiences with close others, it is vital that future research consider roles for social
445 relationship level variables (e.g., relationship status and social network perceptions).
446 Of particular interest is that perceived social support has been shown to moderate
447 the relationship between attachment anxiety and health-related measures ⁴⁵.

448 Specifically, future work could examine whether, like highly anxiously attached
449 individuals, highly attachment disorganized individuals' health is poorer despite a
450 high level of perceived social support compared to low attachment disorganized
451 individuals.

452 Moreover, this work should inform our broader understanding of the
453 association between attachment orientations and health outcomes ⁴⁶, models *must*
454 consider the role of disorganized attachment over and above organised forms of
455 attachment. Clinicians engaged in addressing eating problems and weight
456 management should consider attachment orientations in their patient assessments
457 and be mindful that attachment disorganized individuals are especially likely to
458 engage in uncontrolled eating behaviours that are associated with a higher BMI.
459 Future work might also consider whether specific interventions are more or less
460 appropriate for use with individuals who have different attachment profiles.

461

462 Supplementary information is available at International Journal of Obesity's
463 website.

464

465 Acknowledgements: We thank Lameise Al-Hashimi and Rebecca Buckley
466 (Swansea University), and Laurie Hutchinson, Imogen Titjen & Ee Jing Loh
467 (University of Sheffield) for their assistance with data collection for the primary study.

468

469 Competing Interests: The authors have no competing interests to declare.

470

471

472 References:

473

- 474 1. World Obesity Federation. Obesity infographics. World Obesity Federation
475 website 2017. <https://www.obesityday.worldobesity.org/world-obesity-day->
476 2017
- 477
- 478 2. World Health Organisation. Obesity and overweight, WHO website 2016.
479 <https://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight>.
- 480
- 481 3. NCD Risk Factor Collaboration. Trends in adult body-mass index in 200
482 countries from 1975 to 2014: a pooled analysis of 1698 population-based
483 measurement studies with 19· 2 million participants. *The Lancet* 2016;
484 **387**(10026): 1377-1396.
- 485
- 486 4. Ells LJ, Demaio A, Farpour-Lambert N. Diet, genes, and obesity. *BMJ: British
487 Medical Journal (Online)* 2018; **360**.
- 488
- 489 5. Diener MJ, Geenen R, Koelen JA, Aarts F, Gerdes VEA, Brandjes DPM et al.
490 The Significance of Attachment Quality for Obesity: A Meta-Analytic Review.
491 *Canadian Journal of Behavioural Science-Revue Canadienne Des Sciences
492 Du Comportement* 2016; **48**(4): 255-265.
- 493
- 494 6. Bowlby J. *Attachment and Loss: Attachment*; John Bowlby, Basic Books,
495 1969.
- 496
- 497 7. Brennan KA, Clark CL, Shaver PR. Self-report measurement of adult
498 attachment: An integrative overview. 1998.

499

500 8. Faber A, Dubé L, Knäuper B. Attachment and eating: A meta-analytic review
501 of the relevance of attachment for unhealthy and healthy eating behaviors in
502 the general population. *Appetite* 2017.

503

504 9. Westenhoefer J, Broeckmann P, Münch A-K, Pudel V. Cognitive control of
505 eating behavior and the disinhibition effect. *Appetite* 1994; **23**(1): 27-41.

506

507 10. Karlsson J, Persson L-O, Sjöström L, Sullivan M. Psychometric properties and
508 factor structure of the Three-Factor Eating Questionnaire (TFEQ) in obese
509 men and women. Results from the Swedish Obese Subjects (SOS) study.
510 *International Journal of Obesity & Related Metabolic Disorders* 2000; **24**(12).

511

512 11. Wilkinson LL, Rowe AC, Bishop RJ, Brunstrom JM. Attachment anxiety,
513 disinhibited eating, and body mass index in adulthood. *International Journal of
514 Obesity* 2010; **34**(9): 1442-1445.

515

516 12. Wilkinson LL, Rowe AC, Robinson E, Hardman CA. Explaining the
517 relationship between attachment anxiety, eating behaviour and BMI. *Appetite*
518 2018; **127**: 214-222.

519

520 13. Mikulincer M. Adult attachment style and affect regulation: strategic variations
521 in self-appraisals. *Journal of personality and social psychology* 1998; **75**(2):
522 420.

523

524 14. Mikulincer M, Florian V. The relationship between adult attachment styles and
525 emotional and cognitive reactions to stressful events. 1998.

526

527 15. Maunder R, Hunter J. Attachment and psychosomatic medicine:
528 developmental contributions to stress and disease. *Psychosomatic medicine*
529 2001; **63**(4): 556-567.

530

531 16. Paetzold RL, Rholes WS, Kohn JL. Disorganized attachment in adulthood:
532 Theory, measurement, and implications for romantic relationships. *Review of*
533 *General Psychology* 2015; **19**(2): 146.

534

535 17. Main M, Solomon J. Procedures for identifying infants as
536 disorganized/disoriented during the Ainsworth Strange Situation. *Attachment*
537 *in the preschool years: Theory, research, and intervention* 1990; **1**: 121-160.

538

539 18. Cicchetti D, Barnett D. Attachment organization in maltreated preschoolers.
540 *Development and psychopathology* 1991; **3**(4): 397-411.

541

542 19. Felitti VJ, Anda RF, Nordenberg D, Williamson DF, Spitz AM, Edwards V et al.
543 Relationship of childhood abuse and household dysfunction to many of the
544 leading causes of death in adults: The Adverse Childhood Experiences (ACE)
545 Study. *American journal of preventive medicine* 1998; **14**(4): 245-258.

546

- 547 20. Hemmingsson E, Johansson K, Reynisdottir S. Effects of childhood abuse on
548 adult obesity: a systematic review and meta-analysis. *Obesity Reviews* 2014;
549 **15**(11): 882-893.
- 550
- 551 21. Van Strien T, Cleven A, Schippers G. Restraint, tendency toward overeating
552 and ice cream consumption. *International Journal of Eating Disorders* 2000;
553 **28**(3): 333-338.
- 554
- 555 22. Fritz MS, MacKinnon DP. Required sample size to detect the mediated effect.
556 *Psychological science* 2007; **18**(3): 233-239.
- 557
- 558 23. Button K, Lawrence N, Chambers C, Munafò M. Instilling
559 scientific rigour at the grassroots. In: *The Psychologist*. British Psychological
560 Society, 2016. pp 158 -167.
- 561
- 562 24. Lafontaine M-F, Brassard A, Lussier Y, Valois P, Shaver PR, Johnson SM.
563 Selecting the best items for a short-form of the Experiences in Close
564 Relationships questionnaire. *European Journal of Psychological Assessment*
565 2016; **32**(2): 140.
- 566
- 567 25. Price M, Higgs S, Lee M. Self-reported eating traits: underlying components of
568 food responsivity and dietary restriction are positively related to BMI. *Appetite*
569 2015; **95**: 203-210.
- 570

- 571 26. Pursey K, Burrows TL, Stanwell P, Collins CE. How accurate is web-based
572 self-reported height, weight, and body mass index in young adults? *Journal of*
573 *medical Internet research* 2014; **16**(1).
- 574
- 575 27. Ng SP, Korda R, Clements M, Latz I, Bauman A, Bambrick H et al. Validity of
576 self-reported height and weight and derived body mass index in middle-aged
577 and elderly individuals in Australia. *Australian and New Zealand journal of*
578 *public health* 2011; **35**(6): 557-563.
- 579
- 580 28. Hayes AF. *Introduction to mediation, moderation, and conditional process*
581 *analysis: A regression-based approach*, Guilford Press, 2013.
- 582
- 583 29. van Strien T. Causes of emotional eating and matched treatment of obesity.
584 *Current diabetes reports* 2018; **18**: 1-8.
- 585
- 586 30. Heatherton TF, Baumeister RF. Binge eating as escape from self-awareness.
587 *Psychological bulletin* 1991; **110**(1): 86.
- 588
- 589 31. Herman CP, Mack D. Restrained and unrestrained eating 1. *Journal of*
590 *personality* 1975; **43**(4): 647-660.
- 591
- 592 32. Han S, Kahn JH. Attachment, Emotion Regulation Difficulties, and Disordered
593 Eating Among College Women and Men. *The Counseling Psychologist* 2017;
594 **45**(8): 1066-1090.
- 595

- 596 33. Mikulincer M, Shaver PR, Pereg D. Attachment theory and affect regulation:
597 The dynamics, development, and cognitive consequences of attachment-
598 related strategies. *Motivation and emotion* 2003; **27**(2): 77-102.
- 599
- 600 34. Nancarrow A, Hollywood A, Ogden J, Hashemi M. The Role of Attachment in
601 Body Weight and Weight Loss in Bariatric Patients. *Obesity surgery* 2018;
602 **28**(2): 410-414.
- 603
- 604 35. Wilkinson L, Rowe A, Sheldon C, Johnson A, Brunstrom J. Disinhibited eating
605 mediates differences in attachment insecurity between bariatric surgery
606 candidates/recipients and lean controls. *International Journal of Obesity* 2017;
607 **41**(12): 1831.
- 608
- 609 36. Shakory S, Van Exan J, Mills JS, Sockalingam S, Keating L, Taube-Schiff M.
610 Binge eating in bariatric surgery candidates: The role of insecure attachment
611 and emotion regulation. *Appetite* 2015; **91**: 69-75.
- 612
- 613 37. Devlin MJ, King WC, Kalarchian MA, Hinerman A, Marcus MD, Yanovski SZ
614 et al. Eating pathology and associations with long-term changes in weight and
615 quality of life in the longitudinal assessment of bariatric surgery study.
616 *International Journal of Eating Disorders* 2018.
- 617
- 618 38. Nasirzadeh Y, Kantarovich K, Wnuk S, Okrainec A, Cassin SE, Hawa R et al.
619 Binge Eating, Loss of Control over Eating, Emotional Eating, and Night Eating

- 620 After Bariatric Surgery: Results from the Toronto Bari-PSYCH Cohort Study.
621 *Obesity surgery* 2018; 1-8.
- 622
- 623 39. Hardman CA, Christiansen P, Wilkinson LL. Using food to soothe: Maternal
624 attachment anxiety is associated with child emotional eating. *Appetite* 2016;
625 **99**: 91-96.
- 626
- 627 40. Madigan S, Bakermans-Kranenburg MJ, Van IJzendoorn MH, Moran G,
628 Pederson DR, Benoit D. Unresolved states of mind, anomalous parental
629 behavior, and disorganized attachment: A review and meta-analysis of a
630 transmission gap. *Attachment & human development* 2006; **8**(2): 89-111.
- 631
- 632 41. Van IJzendoorn MH, Bakermans-Kranenburg MJ. Invariance of adult
633 attachment across gender, age, culture, and socioeconomic status? *Journal*
634 *of social and personal relationships* 2010; **27**(2): 200-208.
- 635
- 636 42. Spinosa J, Christiansen P, Dickson J, Lorenzetti V, Hardman C. From socio-
637 economic disadvantage to obesity: The mediating role of psychological
638 distress and emotional eating. *Obesity* in press.
- 639
- 640 43. Ellickson-Larew S, Naragon-Gainey K, Watson D. Pathological eating
641 behaviors, BMI, and facet-level traits: the roles of conscientiousness,
642 neuroticism, and impulsivity. *Eating behaviors* 2013; **14**(4): 428-431.
- 643

644 44. Crawford TN, Shaver PR, Goldsmith HH. How affect regulation moderates the
645 association between anxious attachment and neuroticism. *Attachment &*
646 *Human Development* 2007; **9**(2): 95-109.

647

648 45. Stanton SC, Campbell L. Psychological and physiological predictors of health
649 in romantic relationships: An attachment perspective. *Journal of personality*
650 2014; **82**(6): 528-538.

651

652 46. Lewczuk K, Kobylińska D, Marchlewska M, Krysztofiak M, Glica A, Moiseeva
653 V. Adult attachment and health symptoms: The mediating role of emotion
654 regulation difficulties. *Current Psychology* 2018: 1-14.

655

656 Figure legends:

657

658 Figure 1. Unstandardised regression coefficients are shown with standard error in
659 brackets $B(SE)$. Values in brackets are direct effects when controlling for indirect
660 effects. Significant indirect relationships between disorganized attachment and BMI
661 are denoted by an asterisk and were found via uncontrolled eating ($B = .08$, $(SE =$
662 $.04)$, $LLCI = .02$, $ULCI = .16$) but not via cognitive restraint ($B = -.0004$, $(SE = .01)$,
663 $LLCI = -.02$, $ULCI = .02$) or emotional eating ($B = .006$, $(SE = .04)$, $LLCI = -.08$, $ULCI$
664 $= .1$). The overall R^2 for the model was .18.

665

666 Figure 2. An updated theoretical model of the relationship between attachment
667 orientations and BMI via eating behaviour is presented that for the first time includes
668 disorganized attachment. Unstandardised regression coefficients are shown with

669 standard error in brackets B (SE). Significant indirect relationships are denoted by an
670 asterisk and were found for disorganized attachment and BMI via uncontrolled eating
671 ($B = .08$, $LLCI = .02$, $ULCI = .17$, $p = .002$) and attachment anxiety and BMI via
672 emotional eating ($B = .19$, $LLCI = .09$, $ULCI = .32$, $p < .001$).

673

674

Figure 1.




Figure 2.




Table 1: For each measure, mean scores, standard deviation (SD) and score range are shown. Units are shown including indication when scores are arbitrary units (a.u.)

	Mean scores	SD	Score range
Disorganised attachment (a.u.)	2.8	1.2	1 - 7
Attachment anxiety (a.u.)	4.3	1.4	1 - 7
Attachment avoidance (a.u.)	3.5	1.2	1 - 7
Uncontrolled eating (a.u.)	20.2	4.9	9 - 36
Emotional eating (a.u.)	7.1	2.4	3 - 12
Cognitive restraint (a.u.)	12	3.8	6 - 24
BMI (kg/m2)	23.7	4.3	18.04 - 47.5

Table 2 Pearson's correlations between each measure are shown
 $(* p < .05 \text{ } ** p < .001)$

	Emotional eating	Cognitive restraint	Attachment anxiety	Attachment avoidance	Disorganised attachment
Uncontrolled eating	.53**	0.06	.24**	0.03	.23**
Emotional eating		.13**	.32**	0.07	.14**
Cognitive restraint			.2**	0.07	0.08
Attachment anxiety				.095*	.36**
Attachment avoidance					.37**
Disorganised attachment					
BMI					
Age					

BMI	Age	Gender
.14**	-.24**	0.05
.204 **	-.17**	.17 **
0.05	0.02	.16**
-0.05	-.31**	.12**
-0.004	0.01	-.11**
-0.05	-.19**	0.003
	.31 **	-0.03
		-.1*

Table 3. Unstandardized path coefficients, standard errors, and *t*-values for path analysis.

Path	Estimate	SE	<i>t</i>	<i>p</i>
Disorganised attachment to UE	0.7	0.18	3.9	<.001
Disorganised attachment to EE	0.02	0.1	0.15	0.88
Disorganised attachment to BMI	-0.15	0.16	-0.92	0.36
Attachment anxiety to UE	-0.01	0.15	-0.09	0.93
Attachment anxiety to EE	0.47	0.08	5.96	<.001
Attachment anxiety to BMI	-0.04	0.14	-0.31	0.76
UE to BMI	0.11	0.04	2.58	0.01
EE to BMI	0.4	0.09	4.5	<.001

*Additional information about other direct pathways can be found in the supplementary information file.

Table 4: Specific indirect effects and their respective confidence intervals and p-values for the path model

	Indirect effect	Bias-corrected 95% CI	p
Disorganised attachment to UE to BMI	0.08	.02, .17	0.002
Disorganised attachment to EE to BMI	0.01	-.08, .1	0.91
Attachment anxiety to UE to BMI	-0.001	-.04, .04	0.87
Attachment anxiety to EE to BMI	0.19	.09, .32	<.001