

**Change in health, wellbeing and physical activity levels during the COVID-19 pandemic: a longitudinal cohort of parkrun participants in the United Kingdom**

QUIRK, Helen <<http://orcid.org/0000-0003-2716-4681>>, HAAKE, Steve <<http://orcid.org/0000-0002-4449-6680>>, GOYDER, Elizabeth, BULLAS, Alice, GRANEY, Mike and WELLINGTON, Chrissie

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/29873/>

---

This document is the Accepted Version [AM]

**Citation:**

QUIRK, Helen, HAAKE, Steve, GOYDER, Elizabeth, BULLAS, Alice, GRANEY, Mike and WELLINGTON, Chrissie (2022). Change in health, wellbeing and physical activity levels during the COVID-19 pandemic: a longitudinal cohort of parkrun participants in the United Kingdom. Health Promotion International. [Article]

---

**Copyright and re-use policy**

See <http://shura.shu.ac.uk/information.html>

1 **Change in health, wellbeing and physical activity levels during the COVID-19**  
2 **pandemic: a longitudinal cohort of *parkrun* participants in the United Kingdom**

3 Quirk, Helen: School of Health and Related Research, The University of Sheffield, 30 Regent St, Sheffield,  
4 S1 4DA, United Kingdom; [h.quirk@sheffield.ac.uk](mailto:h.quirk@sheffield.ac.uk)

5 Haake, Steve\*: Advanced Wellbeing Research Centre, Sheffield Hallam University, Sheffield, United  
6 Kingdom; [s.j.haake@shu.ac.uk](mailto:s.j.haake@shu.ac.uk)

7 Goyder, Elizabeth: School of Health and Related Research, The University of Sheffield, 30 Regent St,  
8 Sheffield, S1 4DA, United Kingdom; [e.goyder@sheffield.ac.uk](mailto:e.goyder@sheffield.ac.uk)

9 Bullas, Alice: Advanced Wellbeing Research Centre, Sheffield Hallam University, Sheffield, United  
10 Kingdom; [a.bullas@shu.ac.uk](mailto:a.bullas@shu.ac.uk)

11 Graney, Mike: *parkrun* UK, Middlesex, United Kingdom; [mike.graney@parkrun.com](mailto:mike.graney@parkrun.com)

12 Wellington, Chrissie: *parkrun* UK, Middlesex, United Kingdom; [chrissie.wellington@parkrun.com](mailto:chrissie.wellington@parkrun.com)

13

14 \*Corresponding author: Steve Haake [s.j.haake@shu.ac.uk](mailto:s.j.haake@shu.ac.uk)

15

# 1    **Summary**

## 2    **Introduction**

3    Lockdown restrictions imposed across the UK in response to the COVID-19 pandemic had a profound  
4    impact on many people’s health and wellbeing. People were encouraged to be active, but population surveys  
5    suggest some groups found this easier than others. We explored the changes in health, wellbeing and  
6    physical activity levels among a sample in the UK who experienced the sudden loss of a weekly community-  
7    based physical activity opportunity, *parkrun*.

## 8    **Methods**

9    A sample of UK *parkrun* participants responded to two surveys: pre-COVID-19 in January/February 2019  
10    and during the COVID-19 pandemic in September 2020. Outcomes were happiness, life satisfaction,  
11    connections with others, physical health, mental health and physical activity. The sample was stratified by  
12    gender, age, deprivation status, physical activity and number of *parkruns* completed. Demographics were  
13    reported using descriptive statistics. Distributions between sub-groups were compared using Chi-square  
14    tests while differences in outcomes were determined using the Mann-Whitney U test. Open text responses  
15    were also analysed.

## 16    **Findings**

17    Happiness, life satisfaction, connections with others, physical health and mental health of 450 *parkrun*  
18    participants were negatively impacted for all sub-groups, although the impact was not experienced equally.  
19    The COVID-19 pandemic negatively impacted the wellbeing of a greater proportion of females, younger  
20    adults, inactive people, those from higher deprivation areas, and those who had completed fewer *parkruns*.

## 21    **Conclusions**

22    There is evidence that the wellbeing of those who were more active, and those more involved in a  
23    community-based physical activity initiative pre-pandemic, was less negatively affected during the COVID-  
24    19 lockdown.

25

26    **Key words:** COVID-19, mental health, physical activity, longitudinal study, *parkrun*, community,  
27    inequalities

28

## 29    **LAY SUMMARY**

30    A sample of 450 UK *parkrun* participants responded to two surveys: one before the COVID-19 pandemic  
31    and one during the pandemic. Outcomes were happiness, life satisfaction, connections with others, physical

health, mental health and physical activity. Physical activity fell by 6% while happiness and life satisfaction fell by 12%. People experienced the worst negative impact on their connections with others. The pandemic was found to affect more women, younger adults, those from more deprived neighbourhoods, those who were least active at *parkrun* registration and those who had completed a lower number of *parkrun* events in the 12 months prior to the close of *parkrun* events. The role that community-based physical activity initiatives will have in bringing people's mental health, connections with others, happiness and life satisfaction back to pre-COVID-19 levels in post-lockdown periods needs further investigation and ongoing monitoring.

## INTRODUCTION

In March 2020, a nationwide 'lockdown' in the United Kingdom (UK) in response to coronavirus disease 2019 (COVID-19), placed stringent restrictions on travel, social interaction, and access to public spaces with the aim of slowing the spread of the virus and protecting healthcare services. People were advised to 'stay at home', only leaving for essential reasons. The closure of 'non-essential' businesses, organisations and spaces included leisure and fitness centres, gyms, swimming pools, physical activity events and sports clubs. This had a profound impact on the quality and quantity of social interactions and individual lifestyles with detrimental consequences to social isolation and loneliness (Bu et al., 2020), mental distress (Banks and Xu, 2020), happiness and life satisfaction (Krekel et al., 2020), especially among women, younger adults, people from black and minority ethnic backgrounds and those with lower household income (Fancourt et al., 2020a).

Despite the closure of sport, exercise and physical activity facilities, physical activity came into the spotlight as governments across the world encouraged people to become and stay active as an 'essential activity' for their health and wellbeing (Payne, 2020; World Health Organisation, 2020). Much interest was given to population level changes in physical activity (Stockwell et al., 2021). Research from the beginning of lockdown in March 2020 suggested that higher proportions of the UK population were self-reporting meeting physical activity guidelines compared to preceding years (Smith et al., 2021), which was supported by Google Trends data from the UK (Ding et al., 2020). Conversely, Sport England data from across the COVID-19 pandemic suggests that the lockdown restrictions had a negative impact on the type and volume of activity people were doing – especially during initial stages of the pandemic (between mid-March and mid-May) (Sport England 2021). The proportion of the population classed as "active" dropped by 7.1% (over 3 million fewer active adults) compared to the 12 months before (Sport England 2021).

Collectively, the available evidence into physical activity change is difficult to compare, generalise and interpret due to methodological differences, seasonal variation in activity levels and the changing COVID-19 lockdown restrictions over place and time. A consistent finding was that physical activity levels differed depending on sociodemographic characteristics such as age, sex, socioeconomic status, disability status, ethnicity and pre-lockdown physical activity level (Smith et al., 2021a; Stockwell et al., 2021; Sport

England, 2021; Falkner et al., 2021). Given the importance of these sociodemographic factors, Marteau et al. (2021) have highlighted the importance of addressing both social and behavioural factors to ensure that interventions are more likely to be successful for improving population health and reducing the gap between the richest and poorest in society.

The COVID-19 pandemic restrictions not only meant changes in the levels and type of physical activity but also a loss of social interaction. Feeling a sense of belonging to a social group is a protective mechanism against social isolation, loneliness and poor mental health (Holmes et al., 2020). The social element of participation is likely to have been lost due to lockdown measures. It is therefore important to explore any changes in health, wellbeing and physical activity levels among those who had their community-based physical activity opportunities abruptly removed during lockdown restrictions.

We examine this issue in the context of *parkrun*, a community-based physical activity opportunity that suspended its 2,200+ worldwide events in March 2020 (over 1,000 of which take place in the UK). *parkruns* are free, weekly, 5 kilometre events where people can participate as a runner, walker or volunteer ([www.parkrun.com](http://www.parkrun.com)). In the UK, before events were closed due to the COVID-19 pandemic, around 170,000 people were taking part each week. *parkrun* has removed many of the barriers to physical activity, encouraging participation by women (Stevinson and Hickson, 2013), older people (Grunseit et al., 2013), people with long-term health conditions [blinded for review], people who were previously inactive [blinded for review] and those living in areas of high deprivation (Smith et al., 2020b). Research suggests that the health and wellbeing gains of participation are derived from the friendly, welcoming and social nature of the events (Grunseit et al., 2020). With the abrupt cancellation of *parkrun* events in March 2020, the *parkrun* population provides a unique opportunity to explore change over time in health and wellbeing among relatively active people.

In this study, we sought to understand how the health, wellbeing and physical activity level of UK *parkrun* participants changed during the COVID-19 pandemic and the extent to which people from different sub-groups differed.

## METHODS

Ethical approval for the original Health and Wellbeing Survey was granted by Sheffield Hallam University Research Ethics Committee on 24/07/ 2018 (reference number ER7034346). Ethical approval for this secondary data analysis study was granted by the same ethics committee on 4/12/2020 (reference number ER29077901).

### Study samples

This study uses a single sample of *parkrun* participants responding to surveys at 2 time points, described below.

*The Health and Wellbeing Survey* (labelled “pre-COVID”)

In 2018, *parkrun* commissioned the Advanced Wellbeing Research Centre (AWRC) at Sheffield Hallam University (UK) to conduct a study into the health and wellbeing of the UK *parkrun* community [blinded for review]. This paper reports data from new *parkrun* registrants who completed the survey during January/February 2019 (i.e., “pre-COVID”).

The Health and Wellbeing Survey measured happiness, life satisfaction, self-reported physical activity level, motives for participation, health status, healthcare usage, mental wellbeing, perceived impact of *parkrun* and the impact of *parkrun* on social opportunities. Participants in the Health and Wellbeing Survey gave permission for their anonymised responses to be used for further research.

#### *The parkrun COVID-19 survey (labelled “COVID”)*

During the COVID-19 pandemic in September 2020, 20 months after the *parkrun* Health and Wellbeing survey was distributed, *parkrun* sent a COVID-19 survey to *parkrun* participants in the UK, including participants in England, Scotland, Wales and Northern Ireland. The online *parkrun* COVID-19 survey was sent via email to a stratified random sample balanced for gender, age and number of *parkrun* walk/runs completed in the 12 months prior to 18<sup>th</sup> March 2020. This represented 57,941 *parkrun* participants and included 2,560 respondents from the pre-COVID Health and Wellbeing Survey. The *parkrun* COVID-19 survey aimed to understand the impact of the COVID-19 pandemic on the health and wellbeing of *parkrun* participants and their thoughts about returning to *parkrun* when events were relaunched in the UK. Participants in the *parkrun* COVID-19 survey gave permission for their responses to be shared with researchers for the purposes of further research.

#### *Combined dataset used in this secondary analysis*

Responses to the Health and Wellbeing Survey and the *parkrun* COVID-19 survey were matched at the person-level using *parkrun* Athlete ID (provided to all *parkrun* registrants to identify them on the *parkrun* database and enable the collation of all their *parkrun* participation data) and date of birth across the two databases. This resulted in a combined (linked) dataset of 450 respondents who had completed both surveys and thus allowed a comparison of responses over time (before and during the pandemic).

#### *Demographic variables*

Additional demographic variables not collected in the surveys were extracted from the *parkrun* database after the matching process. These were:

- Gender (female and male);
- Age derived from date of birth;
- Index of multiple deprivation (IMD) derived from postcode;
- Self-reported physical activity level at *parkrun* registration;
- Number of *parkrun* events completed before *parkrun* events closed in March 2020.

## **Outcomes**

## 1 *Health and wellbeing*

2 Mental wellbeing was captured using questions on happiness, life satisfaction, mental health and connections  
3 with others. The pre-COVID and COVID surveys both used the Office of National Statistics (ONS) personal  
4 wellbeing scales questions for happiness and life satisfaction: i) *Overall, how happy did you feel yesterday?*  
5 and ii) *Overall, how satisfied are you with your life nowadays?* Respondents were asked to respond on a  
6 scale of 0 to 10, where 0 is “not at all” and 10 is “completely”.

7 In the COVID survey, participants were asked: *How has your i) happiness, and ii) satisfaction with life iii)*  
8 *connections with others in your community, iv) physical health, and v) mental health been impacted by the*  
9 *COVID-19 pandemic?* On a 5-point Likert scale, respondents were given the following options: major  
10 positive impact, moderate positive impact, no impact, moderate negative impact, major negative impact.

## 11 *Self-reported physical activity level*

12 The pre-COVID and COVID surveys both used the Milton, Bull & Bauman (2011) single item physical  
13 activity question which asked the following: *In the past week, on how many days have you done a total of 30*  
14 *minutes or more of physical activity, which was enough to raise your breathing rate? This may include sport,*  
15 *exercise, and brisk walking or cycling for recreation or to get to and from places, but should not include*  
16 *housework or physical activity that may be part of your job.* Respondents could answer: 0 days, 1 day, 2 days  
17 etc up to 7 days.

## 18 *Open text responses*

19 The COVID survey gave respondents the option of providing an open-text response to the question: *“Is there*  
20 *anything you want to add about the impact of the pandemic, and the absence of parkrun events on your*  
21 *health and wellbeing?”*

## 22 **Data analysis**

23 Data was visually checked in Microsoft Excel by one researcher (SH) and analysed using frequency counts,  
24 means, standard deviations, medians, minimum and maximum and inter-quartile range. For categorical data  
25 N and % were used.

## 26 *Stratification*

27 The sample was stratified by the following:

- 28 • **Gender:** female and male (Supplementary Data 1a);
- 29 • **Age:** derived from the date of birth and segregated into ‘younger adults’ (less than 55 years of age;  
30 mean age 41.2) and ‘older adults’ (55 years or over; mean age 62.4) (Supplementary Data 1b);
- 31 • **Socioeconomic status (SES):** using the indices of multiple deprivation (IMD), classified into four  
32 quartiles (Q1, Q2, Q3, Q4) and segregated into ‘Low IMD’ (those in the most deprived areas; IMD

Q1 and Q2) and ‘High IMD’ (those in the least deprived areas; IMD Q3 and Q4) (Supplementary Data 1c);

- **Activity level:** derived from a physical activity question asked at *parkrun* registration and segregated into ‘lower activity’ (those reporting 0, 1 or 2 days per week of at least 30 minutes moderate exercise) and ‘higher activity’ (those reporting 3 and 4 or more days per week of at least 30 minutes moderate exercise) (Supplementary Data 1d);
- ***parkrun* engagement level:** derived from *parkrun* participation records and segregated either side of the median into ‘low *parkruns*’ ( $\leq 9$  *parkruns* completed in the previous 12 months; mean number of *parkruns* 3.7) and ‘high *parkruns*’ ( $> 9$  *parkruns* completed in the previous 12 months; mean number of *parkruns* 23.2) (Supplementary Data 1e).

The change in physical activity between the pre-COVID and COVID surveys was determined using the single item activity question with a maximum change of  $\pm 7$  days of activity per week.

Distributions between sub-groups were compared using Chi-square tests with the significance of specific categories analysed using partitioned Chi-square tests. Happiness, life satisfaction and the single item physical activity level were classified as ordinal data with differences determined using the Mann-Whitney U test. Effect sizes were calculated using Cohen’s *d* using pooled standard deviation. All statistical tests were analysed using SPSS (v26).

The open-ended survey responses were analysed in Excel using content analysis and inductive coding (O’Cathain and Thomas, 2004). One researcher (HQ), an experienced qualitative researcher, devised a coding frame inductively from the data and manually assigned codes to the verbatim responses that captured what the respondent was saying (i.e., the thematic content of the response). Content analysis stopped when the researcher had reached a point of having summarised all the responses into themes. Themes were presented as numbers and proportions. Verbatim comments were extracted to illustrate the themes.

## FINDINGS

### Sample characteristics

Table 1 shows the demographics of the full sample; the demographics of all sub-groups are given in Supplementary Data 1. The mean age of the sample was 47.6 years with a slight skew towards younger respondents. The age range was 16 to 80 years and 55.3% were female. The proportion of the sample increased linearly with IMD quartile from 11.2% for quartile 1 (most deprived) to 35.1% for quartile 4 (least deprived). 7.4% were inactive at *parkrun* registration (i.e. reported doing less than one day of least 30 minutes of moderate exercise per week) with the mode at three days of activity per week (31.7% of the cohort).

[insert Table 1]



1 In the year prior to *parkrun* closing due to the COVID-19 pandemic (13 to 14 months after the pre-COVID  
2 survey), participants had done a mean of 13.3 *parkruns*, i.e. just over one per month; the distribution was  
3 highly skewed, with a median of 9 *parkruns* and an inter-quartile range of 3 to 21 *parkruns*.

#### 4 **Happiness, life satisfaction and physical activity**

##### 5 *Full cohort*

6 Table 2 shows happiness, life satisfaction and physical activity at the pre-COVID and at COVID surveys for  
7 the full cohort (all) and the sub-groups. Happiness fell from 7.48 before the COVID-19 pandemic to 6.60  
8 during the COVID-19 pandemic by a mean of -0.88; similarly, life satisfaction fell from 7.48 to 6.56 by a  
9 mean of -0.92. Values of happiness and life satisfaction during the COVID-19 pandemic were significantly  
10 lower for all sub-groups compared to before the COVID-19 pandemic (Table 2:  $p<0.01$  or  $p<0.001$  with  
11 moderate to large effect sizes). The physical activity level for the full cohort fell from 3.47 to 3.22 days per  
12 week by 0.21 days per week (Table 2:  $p<0.05$  with a small effect size).

13 The following sections describe the statistically significant findings for each sub-group.

14 *[insert Table 2]*

##### 15 *Females vs males*

16 Females had higher happiness and life satisfaction before the COVID-19 pandemic than during the COVID-  
17 19 pandemic. Although the differences between genders were not significantly different between time points,  
18 the *change* in life satisfaction from before to during the COVID-19 pandemic was, i.e. for females it dropped  
19 by 1.17 while for men it dropped by 0.62 (Table 2: effect size=0.26,  $p<0.01$ ). There was no statistically  
20 significant difference in physical activity levels between females and males.

##### 21 *Younger vs older*

22 Happiness and life satisfaction were statistically higher for the older sub-group compared to the younger sub-  
23 group both before the COVID-19 pandemic and during the COVID-19 pandemic (Table 2:  $p<0.01$ ). There  
24 was no significant difference in physical activity levels between the two sub-groups at either time point.

##### 25 *Low IMD (most deprived) vs high IMD (least deprived)*

26 Happiness and life satisfaction appeared to be lower at both time points for the low IMD group compared to  
27 the high IMD group, although this was only significant for happiness during the COVID-19 pandemic (Table  
28 2: 6.30 vs 7.45, effect size=0.23,  $p<0.05$ ). There was no significant difference for physical activity levels  
29 between the two sub-groups at either time point, although the *change* in physical activity level from before to  
30 during the COVID-19 pandemic was significantly larger for the low IMD group compared to the high IMD  
31 group, i.e. the activity level of the low IMD group fell by 0.52 days per week while the high IMD group fell  
32 by 0.14 days per week (Table 2: effect size 0.19,  $p<0.05$ ).

##### 33 *Low vs high activity at registration*

1 Happiness, life satisfaction and physical activity were lower for the low activity group compared to the high  
2 activity group before and during the COVID-19 pandemic. The *change* in activity from before to during the  
3 COVID-19 pandemic was greater for the high activity sub-group compared to the low activity group (Table  
4 2: -0.57 vs 0.10, effect size 0.34,  $p<0.05$ ).

#### 5 *Low vs high number of parkruns*

6 Happiness and life satisfaction tended to be higher before the COVID-19 pandemic for the low *parkruns*  
7 sub-group compared to the high *parkruns* sub-group; conversely these variables were lower for the low  
8 *parkruns* sub-group during the COVID-19 pandemic. Although the differences between sub-groups were not  
9 significant, the *change* in happiness was significantly greater for the low *parkruns* sub-group with a drop of -  
10 1.10 compared to -0.70 (Table 2: effect size 0.19,  $p<0.05$ ).

11

### 12 **Perceived impact of the COVID-19 pandemic**

13 Table 3 shows the perceived impact of the COVID-19 pandemic with supplementary data given in Table S1.  
14 The most reported negative impact overall was on connections with others (66% to 77% depending upon  
15 sub-group), while physical health had the lowest negative impact (34% to 50%) and the largest positive  
16 impact (23% to 31% depending upon sub-group). Around a third of respondents reported no impact of the  
17 COVID-19 pandemic on either their physical or mental health. The following sections describe the  
18 statistically significant findings for each sub-group: overall distributions are analysed using the Chi-square  
19 test with the significance of negative impact, no impact and positive impact for each measure determined  
20 using partitioned Chi-square tests.

21 *[insert Table 3]*

#### 22 *Females vs males*

23 There was little statistical difference between females and males although there were indications that a larger  
24 proportion of females improved their connections with others during the COVID-19 pandemic (Table 3: 17%  
25 vs 9%,  $p<0.05$ ) and a larger proportion of females reported worse physical health (Table 3: 47% vs 34%,  
26  $p<0.05$ ).

#### 27 *Younger vs older adults*

28 A larger proportion of younger adults reported a negative impact of the COVID-19 pandemic on their  
29 connections with others (Table 3: 77% vs 66%,  $p<0.05$ ) and on their mental health (Table 3: 65% vs 42%,  
30  $p<0.001$ ). There are also indications that a larger proportion of younger adults reported a major negative  
31 impact to happiness, life satisfaction and mental health (Supplementary Data 1; Tables S1a, S1b and S1e).

32 A larger proportion of older adults reported no impact to their physical health than younger adults (Table 3:  
33 39% vs 30%,  $p<0.05$ ); this was also true for mental health (Table 3: 52% vs 28%,  $p<0.001$ ).

### 1    *Low IMD (most deprived) vs high IMD (least deprived)*

2    A larger proportion of those from the low IMD sub-group reported a negative impact of the COVID-19  
3    pandemic on their physical health when compared to the high IMD sub-group (Table 3: 50% vs 37%,  
4     $p<0.01$ ). This was also true for mental health (Table 3: 66% vs 55%,  $p<0.05$ ). Conversely, a larger  
5    proportion of those from the high IMD sub-group reported no impact to their life satisfaction than those from  
6    the low IMD sub-group (Table 3: 22% vs 14%,  $p<0.05$ ); this was also true for mental health (Table 3: 38%  
7    vs 28%,  $p<0.05$ ).

### 8    *Low vs high activity*

9    A larger proportion of those who had low activity levels at registration reported a negative impact of the  
10    COVID-19 pandemic on their physical health when compared to those with higher levels of physical activity  
11    (Table 3: 48% vs 35%,  $p<0.01$ ).

### 12    *Low vs high number of parkruns*

13    A larger proportion of those who did a low number of *parkruns* reported a negative impact of the COVID-19  
14    pandemic on their happiness when compared to those who did a high number of *parkruns* (Table 3: 74% vs  
15    63%,  $p<0.05$ ).

### 16    **Open-text responses**

17    125 respondents (28% of the COVID survey sample) provided an open-text response. 80% of those  
18    providing an open text response (100 respondents) described aspects of *parkrun* that they missed. Data  
19    coding led to the generation of 11 themes that captured how people had responded to the absence of *parkrun*,  
20    to the COVID-19 pandemic and other comments about *parkrun* in relation to its anticipated return (Table 4).  
21    The top two themes related to missing the *parkrun* community and the lack of incentive for physical activity  
22    that *parkrun* engenders.

### 23    **Discussion**

24    We have been able to analyse changes in health, wellbeing and physical activity among a sample of *parkrun*  
25    participants who had completed surveys before and during the COVID-19 pandemic. Happiness and life  
26    satisfaction dropped by about 12% in the 20-month period between *parkrun* registration (pre-COVID) and  
27    during the COVID-19 pandemic. The happiness and life satisfaction scores fell by almost 1 point below the  
28    pre-COVID-19 national averages for England and Wales 2019-2020 (Office of National Statistics (ONS),  
29    2018) though were higher than those reported in other studies from England during the COVID-19 pandemic  
30    (Carson et al., 2020).

31    Whilst the happiness and life satisfaction among all sub-groups were impacted negatively, this was not  
32    experienced similarly across groups. Happiness levels fell more among younger, female and those from more  
33    deprived areas. Life satisfaction levels fell more among females, more deprived and lower activity level

1 respondents. These findings are consistent with the reports of younger adults and females in the UK  
2 demonstrating worse mental health symptoms and larger deteriorations in mental health compared to older  
3 adults and males during the COVID-19 pandemic (Fancourt et al., 2020; Pierce et al., 2020; Krekel et al.,  
4 2020). The gender differences are consistent with pre-existing health inequalities (Pierce et al., 2020) and  
5 have been attributed in part to informal caring responsibilities and childcare responsibilities held alongside  
6 working commitments by females during the COVID-19 pandemic (Mak et al., 2020).

7 Just over half of our sample reported a negative impact of the pandemic on mental health with 6% reporting  
8 a positive impact of the pandemic on mental health. Again, younger adults were more likely to report a  
9 negative impact of the pandemic on their mental health than older adults, which supports other findings  
10 (O'Connor et al., 2020; Office of National Statistics (ONS) 2020; Pierce et al., 2020). We did not find any  
11 differences in the mental wellbeing impact of the pandemic on people from more deprived neighbourhoods  
12 compared to those in less deprived neighbourhoods which could be attributed to higher physical activity  
13 levels (Johansson et al., 2019), though this needs investigating further.

14 Our data show that the greatest negative impact of the COVID-19 pandemic among our sample was on  
15 people's connections with others. Younger adults were more detrimentally impacted. Our open-text  
16 responses captured how people missed the socialisation and community *parkrun* provides, perhaps more so  
17 than the physical activity itself. This is supported by previous *parkrun* research that has highlighted that the  
18 community and social connections are both major appeal and positive outcome of *parkrun* participation  
19 (Grunseit et al., 2020).

20 Our findings suggest that, given many respondents were able to maintain their level of physical activity  
21 during the COVID-19 lockdown, physical activity on its own was not enough to support mental wellbeing,  
22 showing that the lack of social connections had the most detrimental impact. The importance of maintaining  
23 social connections during the COVID-19 pandemic has been strongly advocated as a potential buffer against  
24 negative physical and mental health outcomes (Nitschke et al., 2020). This suggests that a return to *parkrun*  
25 may mitigate some of the negative mental health effects of lockdown. Further research is needed to find out  
26 if this is the case.

27 Less than half of respondents reported a negative impact of the pandemic on their physical health and around  
28 a quarter reported a positive impact of the pandemic on their physical health. This may be attributed to  
29 physical activity levels and our sample's ability to roughly maintain their activity level during the pandemic  
30 (still around 3 days a week of activity). Physical activity levels fell across the whole sample by about 6%,  
31 though there was evidence that some people increased their activity level whilst others decreased, which is  
32 consistent with the existing, but somewhat mixed evidence base (Bann et al., 2020).

33 The open-text comments suggest that people's physical activity response to the pandemic may have been  
34 influenced by motivation (i.e., having an incentive to be active alone) and opportunity (i.e., time in relation  
35 to other commitments), which varied according to living, working and caring arrangements. *parkrun*

1 provided some people with motivation and incentive to be active and whereas others lacked sufficient  
2 incentive to remain active in the absence of *parkrun* events.

3 Participating in events like *parkrun*, when they return, could contribute to the enhancement of mental  
4 wellbeing, especially among younger female participants during future lockdowns, in the ‘back to normal’  
5 transition and ‘post-lockdown’ periods (Sallis et al., 2020). Further research is needed to find out if this is the  
6 case.

## 7 **Methodological considerations**

8 Findings should be interpreted in the context of the following methodological considerations. Firstly, the  
9 self-reported measures may have been biased by measurement errors and reporting biases. Secondly, the  
10 surveys were conducted at different times of the year (January/February and September) so the findings  
11 should be interpreted with consideration of potential seasonality effects. Thirdly, it is possible that those who  
12 provided a response could be different from other *parkrun* participants, and therefore caution must be taken  
13 when extrapolating these findings to a wider population.

14 In our exploration of potential inequalities, it is important to note the following limitations. The  
15 socioeconomic status of respondents was not inferred from employment, income etc. but was inferred from  
16 IMD which was sourced by the postcode provided at *parkrun* registration. This gives an average for the area  
17 lived in when the respondent first registered with *parkrun*, it does not guarantee that it is specific to the  
18 person. A further limitation of our analysis is that we did not consider the impact of the COVID-19  
19 pandemic on ethnic minority groups which have shown inequalities in physical activity levels during the  
20 COVID-19 pandemic (Bann et al., 2020).

21 We did not control for the potential confounding factors in the analysis and cannot draw any conclusions as  
22 to whether the observed associations between participation and outcomes are causally related. Additional  
23 analysis in Supplementary Data 2 identified the key confounding variables. Further adjusted analysis using  
24 logistic regression could explore the extent to which the observed associations may be explained by the  
25 demographic characteristics associated with participation, rather than participation per se. Finally, our  
26 analysis was unable to distinguish the impact of the pandemic from the impact of the lockdown policy on  
27 health and wellbeing (Foa et al., 2020).

## 28 **CONCLUSIONS**

29 The overall wellbeing of a cohort of 450 *parkrun* participants declined during the COVID-19 pandemic.  
30 Physical activity fell by 6% while happiness and life satisfaction fell by 12%. The *parkrun* participants  
31 perceived that the most notable detrimental impact of the pandemic was on their connections with others.  
32 The pandemic was found to affect more women, younger adults, those from more deprived neighbourhoods,  
33 those who were least active at registration and those who had completed a lower number of *parkrun* events  
34 in the 12 months prior to the close of *parkrun* events. The role that community-based physical activity  
35 initiatives will have in bringing people’s mental health, connections with others, happiness and life

1 satisfaction back to pre-COVID-19 levels in post-lockdown periods needs further investigation and ongoing  
2 monitoring.

## 4 **References**

- 5 BANKS, J. & XU, X. 2020. The mental health effects of the first two months of lockdown and social  
6 distancing during the Covid-19 pandemic in the UK. IFS Working Papers.
- 7 BANN, D., VILLADSEN, A., MADDOCK, J., HUGHES, A., PLOUBIDIS, G., SILVERWOOD, R. & PATALAY, P. 2020.  
8 Changes in the behavioural determinants of health during the coronavirus (COVID-19) pandemic:  
9 gender, socioeconomic and ethnic inequalities in 5 British cohort studies. *medRxiv*.
- 10 BU, F., STEPTOE, A. & FANCOURT, D. 2020. Who is lonely in lockdown? Cross-cohort analyses of predictors  
11 of loneliness before and during the COVID-19 pandemic. *Public Health*, 186, 31-34.
- 12 CARSON, J., PRESCOTT, J., ALLEN, R. & MCHUGH, S. 2020. Winter is coming: age and early psychological  
13 concomitants of the Covid-19 pandemic in England. *Journal of Public Mental Health*.
- 14 DING, D., DEL POZO CRUZ, B., GREEN, M. A. & BAUMAN, A. E. 2020. Is the COVID-19 lockdown nudging  
15 people to be more active: a big data analysis. BMJ Publishing Group Ltd and British Association of  
16 Sport and Exercise Medicine.
- 17 FANCOURT, D., BU, F., MAK, H. W. & STEPTOE, A. 2020a. *Covid-19 Social Study Results Release 27* [Online].  
18 Available:[http://allcatsrgrey.org.uk/wp/download/public\\_health/mental\\_health/3d9db5\\_56829e721](http://allcatsrgrey.org.uk/wp/download/public_health/mental_health/3d9db5_56829e7218df4524b304636d226a6198.pdf)  
19 [8df4524b304636d226a6198.pdf](http://allcatsrgrey.org.uk/wp/download/public_health/mental_health/3d9db5_56829e7218df4524b304636d226a6198.pdf) [Accessed 11/02/2021].
- 20 FANCOURT, D., STEPTOE, A. & BU, F. 2020b. Trajectories of anxiety and depressive symptoms during  
21 enforced isolation due to COVID-19: longitudinal analyses of 36,520 adults in England. *medRxiv*.
- 22 FAULKNER, J., O'BRIEN, W. J., MCGRANE, B., WADSWORTH, D., BATTEN, J., ASKEW, C. D., BADENHORST, C.,  
23 BYRD, E., COULTER, M. & DRAPER, N. 2021. Physical activity, mental health and well-being of adults  
24 during initial COVID-19 containment strategies: A multi-country cross-sectional analysis. *Journal of*  
25 *science and medicine in sport*, 24, 320-326.
- 26 FOA, R., GILBERT, S. & FABIAN, M. O. 2020. COVID-19 and subjective well-being: Separating the effects of  
27 lockdowns from the pandemic. *Available at SSRN 3674080*.
- 28 GRUNSEIT, A., RICHARDS, J. & MEROM, D. 2018. Running on a high: parkrun and personal well-being. *BMC*  
29 *public health*, 18, 59.
- 30 GRUNSEIT, A. C., RICHARDS, J., REECE, L., BAUMAN, A. & MEROM, D. 2020. Evidence on the reach and  
31 impact of the social physical activity phenomenon parkrun: A scoping review. *Preventive Medicine*  
32 *Reports*, 101231.

1 HOLMES, E. A., O'CONNOR, R. C., PERRY, V. H., TRACEY, I., WESSELY, S., ARSENEAULT, L., BALLARD, C.,  
2 CHRISTENSEN, H., SILVER, R. C. & EVERALL, I. 2020. Multidisciplinary research priorities for the  
3 COVID-19 pandemic: a call for action for mental health science. *The Lancet Psychiatry*.  
4 JOHANSSON, L. M., LINGFORS, H., GOLSÄTER, M., KRISTENSON, M. & FRANSSON, E. I. 2019. Can physical  
5 activity compensate for low socioeconomic status with regard to poor self-rated health and low  
6 quality-of-life? *Health and quality of life outcomes*, 17, 33.  
7 KREKEL, C., SWANKE, S., DE NEVE, J. E. & FANCOURT, D. 2020. Are happier people more compliant? Global  
8 evidence from three large-scale surveys during Covid-19 lockdowns.  
9 MAK, H. W., BU, F. & FANCOURT, D. 2021. Mental health and wellbeing amongst people with informal  
10 caring responsibilities across different time points during the COVID-19 pandemic: A population-  
11 based propensity score matching analysis. *medRxiv*.  
12 MARTEAU, T., RUTTER, H. & MARMOT, M. 2021. Changing behaviour: an essential component of tackling  
13 health inequalities. *BMJ*, 372, n332.  
14 MILTON, K., BULL, F. & BAUMAN, A. 2011. Reliability and validity testing of a single-item physical activity  
15 measure. *British journal of sports medicine*, 45, 203-208.  
16 NITSCHKE, J. P., FORBES, P. A., ALI, N., CUTLER, J., APPS, M. A., LOCKWOOD, P. L. & LAMM, C. 2020.  
17 Resilience during uncertainty? Greater social connectedness during COVID-19 lockdown is associated  
18 with reduced distress and fatigue. *British Journal of Health Psychology*.  
19 O'CATHAIN, A. & THOMAS, K. J. 2004. "Any other comments?" Open questions on questionnaires—a bane or  
20 a bonus to research? *BMC medical research methodology*, 4, 1-7.  
21 O'CONNOR, R. C., WETHERALL, K., CLEARE, S., MCCLELLAND, H., MELSON, A. J., NIEDZWIEDZ, C. L.,  
22 O'CARROLL, R. E., O'CONNOR, D. B., PLATT, S. & SCOWCROFT, E. 2020. Mental health and well- being  
23 during the COVID-19 pandemic: longitudinal analyses of adults in the UK COVID-19 Mental Health &  
24 Wellbeing study. *The British Journal of Psychiatry*, 1-8.  
25 OFFICE OF NATIONAL STATISTICS (ONS). 2018. *Personal well-being in the UK: July 2017 to June 2018*  
26 [Online]. Available:  
27 [https://www.ons.gov.uk/peoplepopulationandcommunity/wellbeing/bulletins/measuringnational](https://www.ons.gov.uk/peoplepopulationandcommunity/wellbeing/bulletins/measuringnationalwellbeing/july2017tojune2018)  
28 [wellbeing/july2017tojune2018](https://www.ons.gov.uk/peoplepopulationandcommunity/wellbeing/bulletins/measuringnationalwellbeing/july2017tojune2018) [Accessed 23 Mar 2019].  
29 OFFICE OF NATIONAL STATISTICS (ONS) 2020. Coronavirus and Depression in Adults.  
30 PAYNE, R. 2020. Will the COVID-19 outbreak propel the demand for active spaces or scare the public away?  
31 *Cities & Health*, 1-4.  
32 PIERCE, M., HOPE, H., FORD, T., HATCH, S., HOTOPF, M., JOHN, A., KONTOPANTELLIS, E., WEBB, R., WESSELY,  
33 S. & MCMANUS, S. 2020. Mental health before and during the COVID-19 pandemic: a longitudinal  
34 probability sample survey of the UK population. *The Lancet Psychiatry*, 7, 883-892.  
35 [blinded for review]

1 SALLIS, J. F., ADLAKHA, D., OYEYEMI, A. & SALVO, D. 2020. An international physical activity and public  
2 health research agenda to inform COVID-19 policies and practices. *Journal of Sport and Health*  
3 *Science*.

4 SMITH, L., JACOB, L., BUTLER, L., SCHUCH, F., BARNETT, Y., GRABOVAC, I., VERONESE, N., CAPERCHIONE, C.,  
5 LOPEZ-SANCHEZ, G. F. & MEYER, J. 2020a. Prevalence and correlates of physical activity in a sample  
6 of UK adults observing social distancing during the COVID-19 pandemic. *BMJ Open Sport & Exercise*  
7 *Medicine*, 6, e000850.

8 SMITH, R., SCHNEIDER, P., BULLAS, A., HAAKE, S., QUIRK, H., COSULICH, R. & GOYDER, E. 2020b. Does ethnic  
9 density influence community participation in mass participation physical activity events? The case of  
10 parkrun in England. *Wellcome Open Res*, 5, 9.

11 SPORT ENGLAND. 2021. *The impact of coronavirus on activity levels revealed* [Online]. Available:  
12 <https://www.sportengland.org/news/impact-coronavirus-activity-levels-revealed> [Accessed  
13 20/05/2021].

14 STEVINSON, C. & HICKSON, M. 2013. Exploring the public health potential of a mass community  
15 participation event. *Journal of public health*, 36, 268-274.

16 STOCKWELL, S., TROTT, M., TULLY, M. A., SHIN, J. I., BARNETT, Y. A., BUTLER, L. T., MCDERMOTT, D. T.,  
17 SCHUCH, F. B. & SMITH, L. 2021. Changes in physical activity and sedentary behaviours from before  
18 to during the COVID-19 pandemic lockdown: A Systematic Review. *BMJ Open Sport & Exercise*  
19 *Medicine*.

20 WORLD HEALTH ORGANIZATION. 2020. *Stay physically active during self-quarantine* [Online].  
21 Available: [https://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-](https://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/publications-and-technical-guidance/noncommunicable-diseases/stay-physically-active-during-self-quarantine)  
22 [covid- 19/publications-and-technical-guidance/noncommunicable-diseases/stay-physically-active-](https://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/publications-and-technical-guidance/noncommunicable-diseases/stay-physically-active-during-self-quarantine)  
23 [during- self-quarantine](https://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/publications-and-technical-guidance/noncommunicable-diseases/stay-physically-active-during-self-quarantine) [Accessed 15/01/2020].