

Potential of the triad of fatty acids, polyphenols, and prebiotics from cucurbita against COVID-19 in diabetic patients: a review

ALZAHRANI, Q.E., GILLIS, Richard <<http://orcid.org/0000-0002-7607-6808>>, HARDING, S.E., PINTO, L.H., GULATI, M., KAPOOR, B., RANI, P., SINGH, S.K. and ADAMS, G.G.

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/31133/>

This document is the Published Version [VoR]

Citation:

ALZAHRANI, Q.E., GILLIS, Richard, HARDING, S.E., PINTO, L.H., GULATI, M., KAPOOR, B., RANI, P., SINGH, S.K. and ADAMS, G.G. (2022). Potential of the triad of fatty acids, polyphenols, and prebiotics from cucurbita against COVID-19 in diabetic patients: a review. *Journal of Reports in Pharmaceutical Sciences*, 11 (1), 28-40. [Article]

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

Review Article

Potential of the Triad of Fatty Acids, Polyphenols, and Prebiotics from Cucurbita against COVID-19 in Diabetic Patients: A Review

Abstract

Though the scientific community of the entire world has been struggling to create preventive and therapeutic drugs for coronavirus disease 2019 (COVID-19), the role of nutraceuticals has been hitherto neglected. Established role of fatty acids and polyphenols in combating lifestyle disease can be harnessed to play a significant role in the prevention of this disease. The synergistic effect of these phytonutrients and prebiotics is anticipated to prove beneficial for prevention as well as attenuation of COVID-19 infection. Presence of fatty acids, polyphenols and prebiotics in vegetables from the Cucurbitaceae family makes them an attractive choice for being used as a nutritional supplement during COVID-19. These are known to attenuate the excessive immune response which may prove to be beneficial in preventing and mitigating COVID-19. Use of prebiotics to promote the growth of probiotics has also been recommended for the prevention and cure of COVID-19. However, no such report exists in literature that throws light on such role of cucurbita plants. The present review focuses on the role of the triad of fatty acids, prebiotics and polyphenols present in cucurbita plants in controlling systemic inflammation and endothelial damage, the two main etiopathological factors involved in COVID-19. Cucurbita plants are rich in all these components and their inclusion in diet would be an effective strategy to combat COVID-19. The main focus of the review is to discuss the role of various components of the plants of Cucurbita family, taken as dietary component, in prevention and control of the ongoing pandemic COVID19.

Keywords: Diabetes mellitus, fatty acids, microbiome, polyphenols, prebiotics, SARS-CoV-2

Introduction

Diabetes mellitus

Diabetes mellitus (DM) which is a group of metabolic disease has acquired epidemic proportions in the twenty-first century. More than 460 million people in 2019 are reported to be suffering from some form of DM and the number is anticipated to reach approximately 580 million by the end of this decade.^[1] Diabetic patients are susceptible to a number of co-morbidities, generally resulting in reduced quality of life. It also predisposes patients to a number of opportunistic infections.^[2]

DM is a chronic metabolic disorder of endocrine origin arising because of the failure of the body to respond to the elevated blood glucose levels. This abnormality in carbohydrate metabolism is attributed either to deficiency of insulin secretion or to dysfunction of pancreatic β cells responsible for producing, storing and releasing insulin. Failure of the body to utilize insulin because of insulin resistance can be

another etiopathological factor.^[3,4] However, apart from insulin deficiency or lack of utilization, there are many other causes of DM including various pathophysiological changes.^[5] In fact, the causes of diabetes popularly known as “Dirty Dozen” include pancreatic β -cell failure, insulin resistance, hepatic gluconeogenesis, deranged adipocyte metabolism, hyperglucagonemia, incretin defect, increased renal glucose reabsorption, neurotransmitter dysfunction, central appetite dysregulation, gut microbiota, deregulation of the immune system and abnormal activity of hormones, namely dopamine, testosterone, vitamin D, and renin-angiotensin system.^[6]

Diabetes can be categorized into are four types as classified by The American Diabetic Association. These include two forms of idiopathic diabetes, that is, type 1 (insulin-dependent) and type 2 (non-insulin-dependent) diabetes. Third category is that of gestational diabetes which develops during pregnancy while the fourth one, that is, secondary diabetes is associated with other specific conditions.^[7,8] Fourth type, that is, pre-diabetes is a state in

Qushmua E. Alzahrani¹, Richard B. Gillis², Stephen E. Harding^{3,4}, Luciano Henrique Pinto⁵, Monica Gulati⁶, Bhupinder Kapoor⁶, Pooja Rani⁶, Sachin Kumar Singh⁶, Gary G. Adams²

¹Department of Pharmacy/Nursing/Medicine Health and Environment, University of the Region of Joinville (UNIVILLE) volunteer researcher, Joinville, Santa Catarina, Brazil,

²University of Nottingham, Faculty of Medicine & Health Sciences, Queens Medical Centre, Nottingham, United Kingdom,

³University of Nottingham, National Centre for Macromolecular Hydrodynamics, School of Biosciences, Loughborough, United Kingdom, ⁴Universitetet I Oslo, Oslo, Norway, ⁵Department of Pharmacy/Nursing/Medicine Health and Environment, University of the Region of Joinville (UNIVILLE), Joinville, Santa Catarina, Brazil, ⁶School of Pharmaceutical Sciences, Lovely Professional University, Phagwara, Punjab, India

Received: 06 Oct 2021

Accepted: 09 May 2022

Published: 29 Jun 2022

Address for correspondence:

Dr. Qushmua E. Alzahrani, Department of Pharmacy/Nursing/Medicine Health and Environment, University of the Region of Joinville (UNIVILLE) volunteer researcher, Joinville, SC, 89219-710, Brazil. E-mail: drqushmua.alzahrani@hotmail.com

Access this article online

Website:

www.jrpsjournal.com

DOI:10.4103/jrpts.JRPTS_144_21

Quick Response Code:



How to cite this article: Alzahrani QE, Gillis RB, Harding SE, Pinto LH, Gulati M, Kapoor B, et al. Potential of the triad of fatty acids, polyphenols, and prebiotics from cucurbita against COVID-19 in diabetic patients: A review. J Rep Pharma Sci 2022;11:28-40.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

which blood glucose levels are high but enough to be diagnosed as type 2. It can be prevented for developing into type 2 diabetes by weight loss, life style changes, and medicine.^[9]

Advent of DM leads to certain microvascular and macrovascular complications including diabetic neuropathy, diabetic retinopathy, cardiac myopathy, and diabetic foot ulcers.^[10,11] Coprevalence of these diseases is related to extent of hyperglycemia, its management and age.

Immune system in the human body is the complex host defense system that protects it against diseases. Its main function is to protect the body from any outside invaders such as viruses, bacteria, fungi and toxins as well as from intrinsic ones like tumors.^[12] The primary components of the immune system include thymus, spleen, component system, bone marrow, and lymph nodes.^[13] Immune malfunction is observed in cases of diabetic patients who are generally more prone to infection.^[14] Due to the compromise in the immune function, the whole physiological system is not able to respond appropriately to a physiological insult.^[15]

SARS-CoV-2 and diabetes

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is a highly virulent and fatal human pathogen that emerged in Wuhan, China in 2019 and has led to a global pandemic.^[16-18] Diabetic patients have been reported to be more prone to SARS-CoV-2 with higher rate of mortality and co-morbidity.^[19,20] Mortality in people with diabetes has been reported to be 7.3% vis-à-vis 2.3% overall by Chinese Centre for Disease Control and Prevention in a report on 72,314 cases of Coronavirus disease 2019 (COVID-19).^[21] Though a relationship between diabetes and infection has been clinically well established,^[22] diabetes has definitely emerged as a risk factor which contributes to the severity and mortality of COVID-19.^[23]

Diabetes and cardiovascular diseases are considered amongst the highest risk factors associated with SARS-CoV-2 fatality according to a study that included more than 17 million UK adults.^[20]

Microbiome

A number of studies have reported that commensal gut microbiota forms a complex ecosystem that is vital for the maintenance of human health.^[24-26]

Interestingly, the cells of human gut microbiome outnumber the total number of cells in the human body and have an essential influence on human health stimulating the immune system, managing gastrointestinal infections and protecting neuronal damage.^[27,28] In certain pathogenic conditions, microbial imbalance of gut microbiome which is known as dysbiosis occurs that leads to a number of diseases such as diabetes, neuropsychiatric conditions, auto-immune diseases, tumors as well as allergic disorders.^[29] Moreover, this microbiome imbalance leads to the transmission of pathogen-related molecular that may lead to an abnormal immune response

to the host.^[28] Villanueva-Millán *et al.* reported that the majority of gut bacteria in healthy persons belong to four phyla, that is, *Actinobacteria*, *Bacteroidetes*, *Firmicutes* and *Proteobacteria*.^[30] Interestingly, these four phyla are also present in the lung.^[31,32] Existence of “gut-lung axis” has also been reported which demonstrates that there is communication between the microbiome in the lung and that in the gut. Dysbiosis or inflammation in the gut, therefore, may lead to pulmonary morbidities and *vice versa*.^[33,34] Gastrointestinal symptoms have been extensively reported amongst SARS-CoV-2 patients indicating gut inflammation.^[35] The SARS-CoV-2, an enveloped virus, enters host cells by engaging with the angiotensin-converting enzyme 2 (ACE2) as the entry receptor which can be blocked by the cellular serine protease TMPRSS2 inhibitor.^[36] ACE2 receptors are found in both respiratory and gastrointestinal tract.^[37] Consequently, SARS-CoV-2 is found in the gastrointestinal tract as this virus can bind to ACE2 receptors. In a study reported in fifteen SARS-CoV-2 patients, Zuo *et al.* found the possible relationship between the presence of some gut microbiota and SARS-CoV-2. The study showed a negative correlation between *Bacteroidetes* species which suppress the ACE2 expression and SARS-CoV-2 severity. Other common species *Bacteroides dorei*, *B. thetaiotaomicron*, *B. massiliensis*, and *B. ovatus*, on the other hand, showed negative correlation with fecal viral load of SARS-CoV-2.^[37]

Pro-inflammatory T helper 17 cells (Th17 cells) have a potential pathogenic role in many metabolic syndromes including type 2 DM secreted cytokines Interleukin 17 (IL-17A), IL-17F and IL-22 that are known to reduce insulin signaling, leading to insulin resistance and development of type 2 DM.^[38,39] Many of these biochemical changes are also associated with gut dysbiosis.^[39]

Materials and Methods

Available electronic databases such as Web of Science, ScienceDirect, PubMed, Scopus and Google Scholar were used to search for the *in vitro* as well as *in vivo* effects of fatty acids, polyphenols and prebiotics from the plants of Cucurbitaceae family. Keywords used for literature search were “Cucurbitaceae”, “Cucurbita”, “fatty acids”, “polyphenols”, “prebiotics”, “fibers”, individual plants of Cucurbitaceae family and their combination. The articles were screened for relevance by going through their abstract. All the articles pertaining to physicochemical, physiological, neuroprotective effect of lycopene have been included in the manuscript.

Fatty Acids in Cucurbita and Their Role

Fatty acids, which are the building blocks of the fat in the bodies of humans and animals, are molecules with long straight chains of carbon atoms with a carboxyl group at the its end.^[40] Being an essential component of cell membranes, they generally exist in the form of phospholipids and glycolipids. They play vital structural, functional and physiological roles in the body, they are classified into two types, that is, essential

and non-essential. The essential fatty acids are polyunsaturated fatty acids (PUFA), which cannot be synthesized in the human body and need to be taken through diet source.^[40,41] Saturated fatty acids and unsaturated fatty acids are described below.

Saturated fatty acids

These are vital for a number of biological functions of the human body. SFAs are synthesized by the human body and are also obtained from dietary sources like vegetable oils or animal fats.^[41] SFAs act as energy suppliers and storage material, building blocks of certain hormones and cell membranes. They also contribute towards mechanical functions like shock absorption and thermal insulation.^[42-44] Moreover, SFAs are involved in some signaling pathways involved in maintenance of homeostasis.^[45,46] During their metabolism, SFAs are converted to triacylglycerols by esterification. This takes place in the intestinal mucosa cells. After that, they become a part of chylomicrons, bind to albumin in the circulatory system and enter the cells through Fatty Acid Transport Protein.^[47,48] Palmitic acid, myristic acid and lauric acid, which are C16:0, C14:0 and C12:0, respectively, are important in such signaling pathways.

SFAs are essential for the normal functioning of white blood cells. They are also required for the maintenance of stability of proteins.^[49] Certain SFAs like lauric acid possess antimicrobial properties also.^[50]

Short chain saturated fatty acids (SCFAs) are aliphatic fatty acids of chain length ranging from 1–6 carbons. These include acetate, propionate, butyrate, iso-butyrate, valerate, iso-valerate and hexanoate and are generally produced by anaerobic microflora of the gut from indigestible dietary fibers. SCFAs are reported to provide energy and act as substrates for synthesis of sugars as well as lipids. They are also used by cytokines to modulate certain metabolic activities.^[51] They are widely reported to reduce the serum levels of glucose, insulin resistance as well as inflammation, and enhance the protective Glucagon-like peptide-1 secretion.^[52]

Various characteristics of these fatty acids, have been described in Table 1.

Monounsaturated fatty acids (MUFAs)

These fatty acids contain only a single point of unsaturation, in the form of a double bond connecting two carbon atoms. MUFAs are reported to reduce the risk of heart disease by decreasing the low-density lipoproteins (LDL).^[72] They are also involved in supply of vitamin E to the body as vitamin E levels in plasma have been reported to be increased with inclusion of MUFAs in diet.^[73] They are reported to play a significant role in the absorption and metabolism of fat-soluble vitamins, that is, vitamin A, D, E and K.^[74,75] Further details can be found in Table 1.

MUFAs are known to increase the levels of the high-density lipoproteins (HDL) while lowering the levels of LDL. By this dual effect, they exert vasculoprotective as

well as cardioprotective effects. MUFAs are needed during the development of nervous system in children and are also involved in improving the brain function in adults.^[76] Oleic acid, a MUFA of long chain length, has been widely reported as a healthy fat.^[77,78]

Polyunsaturated fatty acids (PUFAs)

These fatty acids have two or more carbon-carbon double bonds in their chain and are the common constituents of lipid membranes. These are reported to play a significant role in both the development as well as functioning of the nervous system. Similar to MUFAs, they are also reported to decrease LDL levels while increasing the of HDL plasma levels thereby preserving the health of the cardiovascular system.^[72]












PUFAs include linoleic and linolenic acids, which are described as ‘essential’ because of the fact that the human body is not able to synthesize these two fatty acids [see Table 1]. They are vital components of cell membranes and are involved in regulation of blood pressure as well as inflammation.^[79] Both linoleic acid and linolenic acid along with arachidonic acid act as ligands for peroxisome proliferator-activated receptors, that is, PPAR γ and PPAR α . They are thus, involved in modulation of PPAR γ dependent regulation of anti-inflammatory and anti-diabetic effects.^[60,61] They also contribute towards modulation of angiogenesis, apoptosis and immune response.^[80]

Role of cucurbita fatty acids

One of the most dangerous risk factors which can lead to the morbidity and mortality of SARS-CoV-2 in the patients is the Heart Fatty acids binding protein (HFABP).^[81] Linolenic acid has been reported to inhibit adipocyte differentiation which may lead to a decrease in the expression of HFABP.^[82,83] Zhang and Liu (2020) reported that linoleic acid, by RNA export machinery, could decrease influenza virus replication. However, clinical studies are required to extrapolate these findings to SARS-CoV-2.^[84] It has been claimed that PUFAs have an essential role to stop the virus from invading the cell.^[85] However, this claim is based on animal studies and its application to human cells needs scientific validation.

Consuming high-fat and high-sugar diet impacts the gut microbiome and might lead to circadian rhythm disruption.^[86] Backhed *et al.* were the first to demonstrate the gut microbiome in metabolic regulation in mice. Mice lacking a gut microbiome were characterized by a lower fat mass whereas transplanted gut microbiome mice resulted in higher production of leptin and in increased fat mass.^[87] Stimulating the gut microbiome to produce SCFA is an essential role, to manage metabolic disorder as SCFA improve insulin response after oral glucose-tolerance test.^[88] Recently Coutzac *et al.* discovered that SCFAs that are produced by gut microbiota are able to produce an antitumour effect in patients. This study shows the benefit of using fatty acids as immunomodulators.^[89] Moreover, with respect to SARS-CoV-2 as an enveloped virus, fatty acids chain side such as in ginkgolic acids could be used to treat SARS-CoV-2.^[90]

Table 1: Properties and functions of various fatty acids

Fatty acid (Carbon:unsaturation)	Chemical structure	Functions	Reference
Eicosapentaenoic acid (20:5)		Immunomodulator	[53]
Linolenic acid (18:3) ω -3/6		Immunomodulator, glycaemic control, cardioprotective, anti-inflammatory, ↓ risk of AD	[54-57]
Linoleic acid (18:2) ω -6		Regulates blood pressure, blood clotting, blood lipid level, ↓ the risk of CHD	[58,59]
Oleic acid (18:1) ω -9		Fatty acid storage, long-term source of energy, layer of insulation, antihyperglycemic, ↓ metabolic LDL, HDL	[60,61]
Stearic acid (18:0)		Hypocholesterolemic, might prevent Parkinson's disease	[62,63]
Palmitic acid (16:0)		Antioxidant, special structural and functional roles in utero and in infancy, glycemic control	[64]
Palmitoleic acid (16:1)		Antihyperglycemic, anti-inflammatory	[65]
Myristic acid (14:0)		Possible treatment of immunodeficiency diseases, ↓ risk of diabetes	[66]
Lauric acid (12:0)		Antimicrobial activity, anti-inflammatory, antitumor	[67,68]
Butyric acid (4:0)		Anti-inflammatory, influences brain function, improve energy metabolism	[69]
Propionic acid (3:0)		Promotion of satiety and reduction in cholesterol, anti-inflammatory, antihyperglycemic	[70,71]

AD: Alzheimer's disease; CHD: Coronary heart disease; LDL: Low-density lipoprotein; HDL: high-density lipoprotein

It is possible to use nutritional sources to help patients with DM to either prevent, or treat, infection with SARS-CoV-2, and reduce both morbidity and mortality in such CoV-2 patients with co-morbidities.^[91,92] Potential source of these components are the plants belonging to the *Cucurbita* genus. This consists of squashes/gourds such as pumpkin (*C. pepo*) and butternut (*C. moschata*). The superior health promoting effects of phytonutrients in the form of plant products as compared to those in their *per se* form accrue from a complex interplay of cell signaling, biochemistry and physiology. Utilizing the concept of food synergy to increase the bioavailability of a phytonutrients is widely reported in literature.^[93] Presence of beneficial fatty acids along with fiber content is expected to have an effect on glycemic index.^[94]

Butternut squash seed oil contains three types of fatty acids, that is, SFAs, MUFAs and PUFAs. Among the SFAs, palmitic acid constitutes 25% of total fat content (TFC) while stearic acid constitutes 3% TFC. As far as MUFAs are concerned, oleic acid constitutes about 10% of TFC. Among PUFAs, linoleic acid constitutes 23% TFC while linolenic acid makes up 36% TFC.^[95]

Pumpkin seed oil is reported to have a high content of both essential and non-essential fatty acids. While the SFAs include lauric acid, myristic acid, palmitic acid and stearic acid, MUFAs are majorly constituted by palmitoleic acid and oleic acid as MUFAs and linoleic acid and linolenic acid are the major PUFAs.^[96]

The presence of high amounts of oleic and linoleic acids help in significant reduction of the plasma cholesterol levels on regular consumption. It is known to decrease the plasma levels of LDL and increase those of HDL.^[58] The oil obtained from the pumpkin seeds has been reported to have certain therapeutic activities such as antifungal, antidiabetic, and anti-inflammatory activities.^[97] Anti-inflammatory activity of oil obtained from *C. pepo* has also been reported in its wound healing effect. In a study on rats, the skin appendages and well-organized collagen fibers without inflammatory cells were found to reappear with the use of *C. pepo* extract.^[98] *Cucurbita* has been used as traditional medication in most regions of the world.^[99] Despite the fact that the seeds of butternut squash and pumpkin have shown promising therapeutic potential, only a few detailed reports are available on their composition and

the properties of their oil for achieving euglycemia. The oil extracted from pumpkin was reported to exhibit hypoglycemic effects in alloxan-induced diabetic rats.^[100] A number of studies indicate that the compounds found in pumpkins are useful in the management of insulin levels in the plasma.^[101,102]

Ezuruike and Prieto (2014) reported Linoleic acid, Oleic acid, Linolenic acid, Myristic acid, and Palmitoleic acid, in the plant of Cucurbita, as relevant components.^[103] Interestingly, a recent study by Xiao *et al.* discovered the present of SARS-CoV-2 in the faeces of infected patient.^[104] Li *et al.* reported that microbiome in the body have a suppressive role for invading virus through different mechanisms.^[105] However, Ghaffari, *et al.* reported that fatty acids such as oleoylethanolamide inhibit the pathway of SARS-CoV-2 infection.^[106] Recently, the KD Pharma Group and its partner SLA Pharma announced that the new drug candidate which is a long-chain omega-3 marine fatty acids derived from eicosapentaenoic acid will soon enter clinical trials as a potential treatment for the treatment of patients with symptoms of SARS-CoV-2.^[107] For non-severe patients, fatty acids could be included in their diets as a nutritional component.^[108] Hence, including Cucurbita FAs in the diet of diabetic people is recommended.^[109]

Cucurbita flowers have been reported to have the ability to stimulate humoral immune response *in vivo*.^[110] It has also been

stated that pumpkin seeds have immunomodulatory effects which can be used as an immunonutrient and anti-atherogenic hypolipidemic.^[111,112] However, these studies were performed in animal models and the whole pumpkin powder seed was used. Therefore, more clinical studies need to be carried out.

Polyphenols in Cucurbita and Their Role

Polyphenols, the largest class of phyto-bioactives, are generally produced as secondary metabolites in plants. They are reported to exert protective action against ultraviolet radiations, pathogen infection, and oxidative stress.^[113] They generally contain one or more phenolic ring and can be classified into flavonoids (anthoxanthins, flavanones, flavanonols, flavans, antocyanidins, isoflavonoids and neoflavonoids), phenolic acids (caffeic, carnosic, ferulic, gallic, *p*-coumaric, rosmarinic, vanillic), polyphenolic amides (capsaicinoids, avenanthramides) and other polyphenolic compounds like stilbenes and lignan [see Table 2].^[114-116]

In addition to their widely reported antioxidant and anti-inflammatory activities, a number of reports are available on the antiviral potential of polyphenols against several pathogens such as Epstein-Barr virus,^[117,118] enterovirus,^[119] herpes simplex virus,^[120] influenza virus^[121] and other virus causing respiratory tract-related infections.^[122] The antiviral action of polyphenols

Table 2: Quantitative analysis of polyphenols present in some of the plants belonging to Cucurbitaceae family

Polyphenol	Polyphenol content (mg/100g) Benincasa hispida	Lagenaria siceraria	Momordica charantia	Trichosanthes anguina	Cucurbita maxima	Reference
Gallic acid	52.5	9.2	338.6	2.15	2.58	[134-138]
Protocatechuic acid	126	3.2	3.59	10.18	5.09	[135,138-141]
4-Hydroxybenzoic acid	-	7174	0.18	1.473	10.04	[138,139,141-143]
Vanillic acid	-	6337	2.15	0.740	8.81	[135,138,141,142,144]
Chlorogenic acid	-	364	10.73	-	6.85	[135,138,142]
Caffeic acid	79	3	0.42	0.953	17.75	[138-141,143]
<i>p</i> -coumaric acid	-	3120	0.47	0.541	0.03	[135,138,139,141,142]
Ferulic acid	0.66	7670	0.18	0.66	13.60	[136,138,139,143,145]
Syringic acid	-	-	2.10	1.460	2.04	[135,139,141]
Sinapic acid	-	-	-	3.758	32.0	[138,141]
Rutin	1.28	-	2.9	1.95	51.92	[136,138,146]
Kaempferol	-	-	0.27	0.35	0.046	[136,139]
Isoquercetin	-	-	-	-	5.54	[138]
Astragalin	-	-	-	-	28.03	[138]
Myricetin	-	-	-	-	9.04	[138]
Quercetin	0.4	6.1	8.0	0.46	6.97	[136,137,139]
Fatty acids						
Linolenic acid	-	9.0	5870	-	2.24	[147-149]
Linoleic acid	-	61.4	6310	-	49.41	[147-149]
Oleic acid	-	4.0	3920	-	25.41	[147-149]
Stearic acid	-	660	-	-	4.51	[147-149]
Palmitic acid	-	2.0	7360	-	20.78	[147-149]
Palmitoleic acid	-	100	210	-	-	[147-149]
Myristic acid	-	100	90	-	0.009	[147-149]
Lauric acid	-	-	250	-	1.336	[149]

is attributed to inhibition of viral replication, protein synthesis, gene expression, and nucleic acid synthesis.^[117]

Till date, the world is struggling to treat Covid-19 infection which enforces the physicians to use widely known antivirals and corticosteroids. In addition to them, herbal extracts of a number of Chinese medicinal plants such as *Cibotium barometz*, *Gentiana scabra*, *Dioscorea batatas*, *Cassia tora* and *Taxillus chinensis* were found to inhibit SARS-CoV replication.^[123] Only a few reports are available on the *in vitro* studies of the effect of polyphenols (forsythoside A, (-)-catechin gallate, (-)-gallocatechin gallate, resveratrol as well as polyphenols from *Broussonetia papyrifera* and *Sambucus nigra*) against coronavirus.^[124-128]

Pumpkin is rich source of carotenoids (lutein, most abundant), phenolic acids, flavonols, saponins, mineral components and vitamin C.^[129-131] The various phenolic acids found in pumpkin varieties are gallic acid, protocatechuic acid, 4-hydroxybenzoic acid, 4-hydroxybenzaldehyde, vanilic acid, chlorogenic acid, caffeic acid, *p*-coumaric acid, ferulic acid, syringic acid and sinapic acid.^[132] The content of flavonols in pumpkin was reported to be less than phenolic acids. Among the various flavonols, rutin and kaempferol are the most abundant flavonols found in all the pumpkin varieties. The others flavonols are isoquercetin, astragalin, myricetin and quercetin.^[133] The chemical structures of these phenolic acids and flavonols present in pumpkin are depicted in [Figure 1].

In another study, the antioxidant and radical scavenging effect of *Cucurbita* fruits (18 cultivars of the species *C. maxima* Duch., *C. moschata* Duch., *C. pepo* L., and *C. ficifolia* Bouché) were assessed. All the species were found to show remarkable antioxidant and radical scavenging properties mainly attributed to the presence of polyphenols. It is pertinent to note here that both these properties are of immense use in combating the COVID19 infection.^[113,139]

Polyphenolic compounds have been reported to exhibit potency against Covid virus in a number of studies, and can be considered as potential nutraceutical(s) for the treatment of SARS-CoV-2 infection. Polyphenols have been reported to block the production of cytokines by senescent cells (senescence-associated secretory phenotype) and adipocytes, as well as modify the ACE-1/ACE-2 ratio, which can potentially result in beneficial effects in COVID-19^[150] [Figure 2]. Rutin, one of the major flavonols present in pumpkin varieties, has been reported to exhibit good binding characteristics with SARS-CoV-2 main protease and host toll-like receptors in *in silico* studies, indicating it as a novel therapeutic option, having its impact *via* virus-based and host-based anti-CoV strategies.^[151-153] Similarly, the other flavonoids found in pumpkin, that is, kaempferol and quercetin were reported to SARS-CoV-2 main protease in molecular docking studies, and therefore may act as anti-COVID-19 agents.^[154,155] However, further research is necessary to investigate their potential medicinal use.

Prebiotics in Cucurbita and Their Role

Prebiotics are the indigestible plant fibers that act as growth substrates specific for beneficial microorganisms. They

are widely reported to reduce gut inflammation, obesity, and lipid accumulation. These not only improve gut health but also lead to inhibition of pathogens and stimulation of immune system owing to their ability to modulate the type, numbers, composition and even activity of human microbiota.^[156] Administration of prebiotics is known to reduce the inflammation and infection in lungs as well as other respiratory ailments such as asthma and chronic obstructive pulmonary disease.^[157] This interplay is facilitated by the gut lung axis. Prebiotics include fructans, oligosaccharides, arabinooligosaccharides, isomaltooligosaccharides, xylooligosaccharides, resistant starch, lactosucrose, lactobionic acid, galactomannan, polyphenols and polyunsaturated fatty acids. An inverse correlation has been established between prebiotic consumption and serum levels of inflammatory cytokines including C-reactive protein, interleukin (IL)-6, IL-18 and tumor necrosis factor- α (TNF α).^[158] The event of acute rise in these very cytokines called as cytokine storm has been reported to play a crucial role in severity of the disease and mortality therefrom.^[159] SCFAs like acetate, propionate and butyrate are the main components of immunomodulatory signals which are produced by gut microbiome.^[160] A daily intake of just 6g of prebiotics has been reported to increase the levels of SCFAs substantially.^[161] No information, however, is currently available on the direct effect of prebiotics on the COVID-19 infections, although an indirect relationship may be drawn. Further, a number of clinical trials have proven the efficacy of prebiotics in diabetes management by improving the barrier function and translocation of bacterial lipopolysaccharide.^[162]

As mentioned previously, prebiotics are considered as dietary fiber which stimulate health promoting microorganisms.^[163] Role of prebiotics in handling the COVID-19 infection has been attributed to the involvement of gut lung microbiota axis.^[158] A number of studies have reported the effect of prebiotic component of various *Cucurbita* plants.^[164] Prebiotic effect of *C. maxima* was demonstrated by Lokuge *et al.* in pigs showing an increase in beneficial microorganisms and a decrease in coliform bacteria.^[163] Pectin extracted from *C. maxima* was found to exhibit strong bifidogenic effect. An anti-hypertensive effect of a combination of pectin from *C. maxima* along with whey protein hydrolysate was also observed.^[165] In fact, the prebiotic potential of oligosaccharides obtained from *C. moschata* pulp were shown to be better than even inulin (considered as golden standard prebiotic) in terms of their resistance to hydrolysis by gastric juice and equivalent in terms of stimulation of the growth of *Lactobacilli*.^[166] Even waste derived from pumpkin peel and pulp (30:70 w/w) were found to exhibit excellent prebiotic properties for the growth of *Lactobacillus casei*.^[167]

A novel protein-bound polysaccharide obtained from the seeds of *C. maxima* exhibited α -amylase and α -glucosidase inhibitory activity and thus can be used as an antidiabetic agent.^[168] In another study, seed flour from *C. maxima* was reported to




Figure 1: Chemical structures of phenolic acids and flavonols present in pumpkin

interfere in the rat metabolism decreasing significantly the serum glucose and triacylglycerides levels.^[169]

Production of SCFAs is one of the key roles played by prebiotics in association with the colonic bacteria. No information, however, is currently available on the direct effect of prebiotics on the COVID-19 infections, although an indirect relationship may be drawn. Further, a number of clinical trials have proven the efficacy of prebiotics in diabetes management by improving the barrier function and translocation of bacterial lipopolysaccharide.^[170] The SCF producing ability of five cucurbita plants, that is, *Benincasa hispida*, *Lagenaria siceraria*, *Momordica charantia*, *Trichosanthes anguina*, and *Cucurbita maxima* was evaluated and compared with wheat fiber as control. The SCFA production by all four plants was found to be higher than that by wheat fiber using different probiotic strains.^[171]

Conclusions

The triumvirate of fatty acids, polyphenols and prebiotics is an essential component of our diet and a balance in interplay among these leads to anti-oxidant, anti-inflammatory and immunomodulatory effects that are of high significance in both morbidity and mortality associated with COVID-19 infection. Their anti-diabetic nuances further render them even more suitable for consumption by diabetic people to reduce the risk of the complications of COVID-19. The concept of food synergy is widely reported wherein the phytonutrients in the presence of other phytoconstituents are believed to provide higher health benefits as compared to when they are administered alone.^[172] This has been largely attributed to enhancement in the bioavailability of phytonutrients by interplay of biochemical, physiological and signaling factors. Therefore, consumption of various parts of plants belonging to the family Cucurbitaceae




Figure 2: Pathophysiological events of COVID-19 leading to cytokine storm and the potential role of cucurbita phytoconstituents in attenuating the cytokine storm. ACE2: angiotensin I converting enzyme 2; TLR: toll-like receptors; NF-kB: Nuclear factor kappa B; IRF: Interferon-regulatory factors

may prove to be more effective as compared to consumption of the isolated nutrients. We conclude that certain cucurbita plants come across as convincing alternatives by regulating the production and release of proinflammatory cytokines, interfering with the development of the virus and modifying certain molecular pathways related to the pathophysiology. Though the nutraceutical role of the cucurbita plants has been extensively discussed, attributing these effects to specific phytotherapeutics has not been done due to lack of reports in this direction. Based on the fact, that COVID-19 has a plethora of symptoms varying from patient to patient in intensity, it is suggested to work further to elucidate the role of each class of compounds in combating the different symptoms. Such clinical studies will pave a way for making personalized dietary recommendations involving the inclusion of different cucurbita plants in patients manifesting different symptoms.

Looking at the formidable morbidity and mortality of COVID-19, we suggest that these plants should be rigorously tested to protect the COVID-19 patients, especially those belonging to the vulnerable groups.

Acknowledgment

We would like to thank Dr. Frederik H. Verbrugge (Department of Cardiovascular Medicine, Mayo Clinic, Rochester, MN (FHV, BAB) and Biomedical Research Institute, Faculty of Medicine and Life Sciences, Hasselt University, Belgium (FHV) for helpful discussions.

Financial support and sponsorship

No funding was received for this work.

Conflicts of interest

There are no conflicts of interest.

References

- Williams R, Karuranga S, Malanda B, Saeedi P, Basit A, Besançon S, *et al.* Global and regional estimates and projections of diabetes-related health expenditure: results from the international diabetes federation diabetes atlas, 9th edition. *Diabetes Res Clin Pract* 2020;162:108072.
- Casqueiro J, Casqueiro J, Alves C. Infections in patients with diabetes mellitus: A review of pathogenesis. *Indian J Endocrinol Metab* 2012;16:S27-36.
- Adams GG, Imran S, Wang S, Mohammad A, Kok S, Gray DA, *et al.* The hypoglycaemic effect of pumpkins as anti-diabetic and functional medicines. *Food Res Int* 2011;44:862-7.
- Kerrison G, Gillis RB, Jiwani SI, Alzahrani Q, Kok S, Harding SE, *et al.* The effectiveness of lifestyle adaptation for the prevention of prediabetes in adults: A systematic review. *J Diabetes Res* 2017;2017:8493145.
- Kusuhara S, Fukushima Y, Ogura S, Inoue N, Uemura A. Pathophysiology of diabetic retinopathy: the old and the new. *Diabetes Metab J* 2018;42:364-76.
- Kalra S, Chawla R, Madhu SV. The dirty dozen of diabetes. *Indian J Endocrinol Metab* 2013;17:367-9.
- American Diabetes Association. Classification and diagnosis of diabetes: Standards of medical care in Diabetes—2021. *Diabetes Care* 2021;44(Supplement_1):S15-S33.
- Teugwa CM, Boudjeko T, Tchinda BT, Mejiato PC, Zofou D. Anti-hyperglycaemic globulins from selected cucurbitaceae seeds used as antidiabetic medicinal plants in Africa. *Bmc Complement Altern Med* 2013;13:63.

9. Hostalek U. Global epidemiology of prediabetes: Present and future perspectives. *Clin Diabetes Endocrinol* 2019;5:5.
10. Cade WT. Diabetes-related microvascular and macrovascular diseases in the physical therapy setting. *Phys Ther* 2008;88: 1322-35.
11. Chawla A, Chawla R, Jaggi S. Microvascular and macrovascular complications in diabetes mellitus: distinct or continuum? *Indian J Endocrinol Metab* 2016;20:546-51.
12. Sompayrac LM. *How the Immune System Works*. Oxford: John Wiley & Sons; 2019.
13. Pinti M, Appay V, Campisi J, Frasca D, Fülöp T, Sauce D, *et al.* Aging of the immune system: focus on inflammation and vaccination. *Eur J Immunol* 2016;46:2286-301.
14. Zhou X, Chen Y, Mok KY, Zhao Q, Chen K, Chen Y, *et al.*; Alzheimer's Disease Neuroimaging Initiative. Identification of genetic risk factors in the chinese population implicates a role of immune system in Alzheimer's disease pathogenesis. *Proc Natl Acad Sci USA* 2018;115:1697-706.
15. Bossi F, Bernardi S, Zauli G, Secchiero P, Fabris B. Trail modulates the immune system and protects against the development of diabetes. *J Immunol Res* 2015;2015:680749.
16. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 2020;579:270-3.
17. World Health Organization. Coronavirus disease (COVID-19): situation report, 203. 2020.
18. Kapoor B, Kochhar RS, Gulati M, Rani P, Gupta R, Kumar Singh S, *et al.* Triumvirate to treat mucormycosis: Interplay of pH, metal ions and antifungal drugs. *Medical Hypotheses* 2022;159:110748.
19. Xu Z, Shi L, Wang Y, Zhang J, Huang L, Zhang C, *et al.* Pathological findings of Covid-19 associated with acute respiratory distress syndrome. *Lancet Respir Med* 2020;8:420-2.
20. Williamson EJ, Walker AJ, Bhaskaran K, Bacon S, Bates C, Morton CE, *et al.* Factors associated with Covid-19-related death using opensafely. *Nature* 2020;584:430-6.
21. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (Covid-19) outbreak in china: summary of a report of 72 314 cases from the chinese center for disease control and prevention. *Jama* 2020;323:1239-42.
22. Pearson-Stuttard J, Blundell S, Harris T, Cook DG, Critchley J. Diabetes and infection: assessing the association with glycaemic control in population-based studies. *Lancet Diabetes Endocrinol* 2016;4:148-58.
23. Abdi A, Jalilian M, Sarbarzeh PA, Vlaisavljevic Z. Diabetes and Covid-19: A systematic review on the current evidences. *Diabetes Res Clin Pract* 2020;166:108347.
24. Wang CY, Chen YW, Tain YL, Chang SKC, Huang LT, Hsieh CW, *et al.* Fast quantification of short-chain fatty acids in rat plasma by gas chromatography. *J Food Sci* 2020;85:1932-8.
25. Liang D, Leung RK, Guan W, Au WW. Involvement of gut microbiome in human health and disease: brief overview, knowledge gaps and research opportunities. *Gut Pathog* 2018;10:3.
26. Gulati M, Singh SK, Corrie L, Chandwani L, Singh A, Kapoor B, *et al.* Fecal microbiota transplant: latest addition to arsenal against recurrent clostridium difficile infection. *Recent Adv Anti-Infective Drug* 2021;16:2-12.
27. Zhang H, Yeh C, Jin Z, Ding L, Liu BY, Zhang L, *et al.* Prospective study of probiotic supplementation results in immune stimulation and improvement of upper respiratory infection rate. *Synth Syst Biotechnol* 2018;3:113-20.
28. Dhar D, Mohanty A. Gut microbiota and covid-19- possible link and implications. *Virus Res* 2020;285:198018.
29. Cani PD, Jordan BF. Gut microbiota-mediated inflammation in obesity: a link with gastrointestinal cancer. *Nat Rev Gastroenterol Hepatol* 2018;15:671-82.
30. Villanueva-Millán MJ, Pérez-Matute P, Oteo JA. Gut microbiota: a key player in health and disease. A review focused on obesity. *J Physiol Biochem* 2015;71:509-25.
31. Bingula R, Filaire M, Radosevic-Robin N, Bey M, Berthon JY, Bernalier-Donadille A, *et al.* Desired turbulence? Gut-lung axis, immunity, and lung cancer. *J Oncol* 2017;2017:5035371.
32. Zhang D, Li S, Wang N, Tan HY, Zhang Z, Feng Y. The cross-talk between gut microbiota and lungs in common lung diseases. *Front Microbiol* 2020;11:301.
33. Keely S, Talley NJ, Hansbro PM. Pulmonary-intestinal cross-talk in mucosal inflammatory disease. *Mucosal Immunol* 2012;5:7-18.
34. Dumas A, Bernard L, Poquet Y, Lugo-Villarino G, Neyrolles O. The role of the lung microbiota and the gut-lung axis in respiratory infectious diseases. *Cell Microbiol* 2018;20:e12966.
35. Cheung KS, Hung IFN, Chan PPY, Lung KC, Tso E, Liu R, *et al.* Gastrointestinal manifestations of Sars-cov-2 infection and virus load in fecal samples from a hong kong cohort: systematic review and meta-analysis. *Gastroenterology* 2020;159:81-95.
36. Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, *et al.* Sars-cov-2 cell entry depends on Ace2 and Tmprss2 and is blocked by a clinically proven protease inhibitor. *Cell* 2020;181:271-80.e8.
37. Zuo T, Zhan H, Zhang F, Liu Q, Tso EYK, Lui GCY, *et al.* Alterations in fecal fungal microbiome of patients with Covid-19 during time of hospitalization until discharge. *Gastroenterology* 2020;159:1302-10.e5.
38. Abdel-Moneim A, Bakery HH, Allam G. The potential pathogenic role of Il-17/th17 cells in both type 1 and type 2 diabetes mellitus. *Biomed Pharmacother* 2018;101:287-92.
39. Tan TG, Sefik E, Geva-Zatorsky N, Kua L, Naskar D, Teng F, *et al.* Identifying species of symbiotic bacteria from the human gut that, alone, can induce intestinal th17 cells in mice. *Proc Natl Acad Sci USA* 2016;113:E8141-50.
40. Sirimongkolgal N, Phankosol S, Puttala S, Chum-in T, Krisnangkura K. Gibbs energy additivity approaches in estimation of density of fatty. *J Phys: Conf Ser* 2018;1144:012187.
41. Legrand P, Rioux V. The complex and important cellular and metabolic functions of saturated fatty acids. *Lipids* 2010;45: 941-6.
42. Calder PC. Functional roles of fatty acids and their effects on human health. *J Parent Enteral Nutr* 2015;39:18s-32s.
43. Bos DJ, van Montfort SJ, Oranje B, Durston S, Smeets PA. Effects of omega-3 polyunsaturated fatty acids on human brain morphology and function: what is the evidence? *Eur Neuropsychopharmacol* 2016;26:546-61.
44. Ríos-Covián D, Ruas-Madiedo P, Margolles A, Gueimonde M, de Los Reyes-Gavilán CG, Salazar N. Intestinal short chain fatty acids and their link with diet and human health. *Front Microbiol* 2016;7:185.
45. Rocha DM, Caldas AP, Oliveira LL, Bressan J, Hermsdorff HH. Saturated fatty acids trigger Tlr4-mediated inflammatory response. *Atherosclerosis* 2016;244:211-5.
46. Dalile B, Van Oudenhove L, Vervliet B, Verbeke K. The role of short-chain fatty acids in microbiota-gut-brain communication. *Nat Rev Gastroenterol Hepatol* 2019;16:461-78.
47. Tokuyama S, Nakamoto K. Chapter 12 - Pain as modified by polyunsaturated fatty acids. In: Watson RR, De Meester F,

- editors. Omega-3 Fatty Acids in Brain and Neurological Health. Boston, MA: Academic Press; 2014. p. 131-46.
48. Saini RK, Keum YS. Omega-3 and omega-6 polyunsaturated fatty acids: dietary sources, metabolism, and significance - A review. *Life Sci* 2018;203:255-67.
49. Arpón A, Milagro FI, Razquin C, Corella D, Estruch R, Fitó M, *et al.* Impact of consuming extra-virgin olive oil or nuts within a mediterranean diet on DNA methylation in peripheral white blood cells within the PREDIMED-Navarra randomized controlled trial: a role for dietary lipids. *Nutrients* 2017;10:15.
50. Huang CB, Alimova Y, Myers TM, Ebersole JL. Short- and medium-chain fatty acids exhibit antimicrobial activity for oral microorganisms. *Arch Oral Biol* 2011;56:650-4.
51. Feng W, Ao H, Peng C. Gut microbiota, short-chain fatty acids, and herbal medicines. *Front Pharmacol* 2018;9:1354.
52. Puddu A, Sanguineti R, Montecucco F, Viviani GL. Evidence for the gut microbiota short-chain fatty acids as key pathophysiological molecules improving diabetes. *Mediators Inflamm* 2014;2014:162021.
53. Ceccarelli V, Ronchetti S, Marchetti MC, Calvitti M, Riccardi C, Grignani F, *et al.* Molecular mechanisms underlying eicosapentaenoic acid inhibition of Hdac1 and Dnmt expression and activity in carcinoma cells. *Biochim Biophys Acta Gene Regul Mech* 2020;1863:194481.
54. Calder PC, Carr AC, Gombart AF, Eggersdorfer M. Optimal nutritional status for a well-functioning immune system is an important factor to protect against viral infections. *Nutrients* 2020;12:1181.
55. Brown TJ, Brainard J, Song F, Wang X, Abdelhamid A, Hooper L; PUFAH Group. Omega-3, omega-6, and total dietary polyunsaturated fat for prevention and treatment of type 2 diabetes mellitus: systematic review and meta-analysis of randomised controlled trials. *Bmj* 2019;366:l4697.
56. Veselinovic M, Vasiljevic D, Vucic V, Arsic A, Petrovic S, Tomic-Lucic A, *et al.* Clinical benefits of n-3 PUFA and α -Linolenic acid in patients with rheumatoid arthritis. *Nutrients* 2017;9.
57. Gutiérrez S, Svahn SL, Johansson ME. Effects of Omega-3 fatty acids on immune cells. *Int J Mol Sci* 2019;20:5028.
58. Ramsden CE, Zamora D, Leelarthaepin B, Majchrzak-Hong SF, Faurot KR, Suchindran CM, *et al.* Use of dietary linoleic acid for secondary prevention of coronary heart disease and death: evaluation of recovered data from the sydney diet heart study and updated meta-analysis. *BMJ* 2013;346:e8707.
59. Parhizkar S, Latif LA. Supplementary health benefits of linoleic acid by improvement of vaginal cornification of ovariectomized rats. *Adv Pharm Bull* 2013;3:31-6.
60. Yary T, Voutilainen S, Tuomainen TP, Ruusunen A, Nurmi T, Virtanen JK. Serum n-6 polyunsaturated fatty acids, $\Delta 5$ - and $\Delta 6$ -desaturase activities, and risk of incident type 2 diabetes in men: the kuopio ischaemic heart disease risk factor study. *Am J Clin Nutr* 2016;103:1337-43.
61. Wu JHY, Marklund M, Imamura F, Tintle N, Ardisson Korat AV, de Goede J, *et al.*; Cohorts for Heart and Aging Research in Genomic Epidemiology (CHARGE) Fatty Acids and Outcomes Research Consortium (FORCE). Omega-6 fatty acid biomarkers and incident type 2 diabetes: pooled analysis of individual-level data for 39 740 adults from 20 prospective cohort studies. *Lancet Diabet Endocrinol* 2017;5:965-74.
62. Burns JL, Nakamura MT, Ma DWL. Differentiating the biological effects of linoleic acid from arachidonic acid in health and disease. *Prostaglandins Leukot Essent Fatty Acids* 2018;135:1-4.
63. Bajracharya R, Bustamante S, O Ballard JW. Stearic acid supplementation in high protein to carbohydrate (P:C) ratio diet improves physiological and mitochondrial functions of drosophila melanogaster parkin null mutants. *J Gerontol A Biol Sci Med Sci* 2019;74:1564-72.
64. Agostoni C, Moreno L, Shamir R. Palmitic acid and health: introduction. *Crit Rev Food Sci Nutr* 2016;56:1941-2.
65. de Souza CO, Valenzuela CA, Baker EJ, Miles EA, Rosa Neto JC, Calder PC. Palmitoleic acid has stronger anti-inflammatory potential in human endothelial cells compared to oleic and palmitic acids. *Mol Nutr Food Res* 2018;62:e1800322.
66. Iwata K, Sakai H, Takahashi D, Sakane F. Myristic acid specifically stabilizes diacylglycerol kinase δ protein in C2c12 skeletal muscle cells. *Biochim Biophys Acta Mol Cell Biol Lipids* 2019;1864:1031-8.
67. Yamamoto Y, Morikawa T, Kawai T, Nonomura Y. Selective bactericidal activity of divalent metal salts of lauric acid. *Acs Omega* 2017;2:113-21.
68. Lappano R, Sebastiani A, Cirillo F, Rigracciolo DC, Galli GR, Curcio R, *et al.* The lauric acid-activated signaling prompts apoptosis in cancer cells. *Cell Death Discov* 2017;3:17063.
69. Bourassa MW, Alim I, Bultman SJ, Ratan RR. Butyrate, neuroepigenetics and the gut microbiome: can a high fiber diet improve brain health? *Neurosci Lett* 2016;625:56-63.
70. Louis P, Flint HJ. Formation of propionate and butyrate by the human colonic microbiota. *Environ Microbiol* 2017;19:29-41.
71. Tirosh A, Calay ES, Tuncman G, Claiborn KC, Inouye KE, Eguchi K, *et al.* The short-chain fatty acid propionate increases glucagon and FABP4 production, impairing insulin action in mice and humans. *Sci Trans Med* 2019;1.
72. Schwingshackl L, Hoffmann G. Monounsaturated fatty acids and risk of cardiovascular disease: synopsis of the evidence available from systematic reviews and meta-analyses. *Nutrients* 2012;4:1989-2007.
73. Zingg JM. Vitamin E: regulatory role on signal transduction. *Iubmb Life* 2019;71:456-78.
74. Harrison EH, Kopec RE. Chapter 50 - Digestion and Intestinal Absorption of Dietary Carotenoids and Vitamin A☆. In: Said HM, editor. *Physiology of the Gastrointestinal Tract*. 6th ed. Amsterdam: Elsevier; 2018. p. 1133-51.
75. Yamanashi Y, Takada T, Kurauchi R, Tanaka Y, Komine T, Suzuki H. Transporters for the intestinal absorption of cholesterol, vitamin E, and vitamin K. *J Atheroscler Thromb* 2017;24:347-59.
76. Bentsen H. Dietary polyunsaturated fatty acids, brain function and mental health. *Microb Ecol Health Dis* 2017;28:1281916.
77. Ramsden CE, Faurot KR, Carrera-Bastos P, Cordain L, De Lorgeril M, Sperling LS. Dietary fat quality and coronary heart disease prevention: a unified theory based on evolutionary, historical, global, and modern perspectives. *Curr Treat Options Cardiovasc Med* 2009;11:289-301.
78. Martin C. The interface between plant metabolic engineering and human health. *Curr Opin Biotechnol* 2013;24:344-53.
79. Gammone MA, Riccioni G, Parrinello G, D'Orazio N. Omega-3 polyunsaturated fatty acids: benefits and endpoints in sport. *Nutrients* 2018;11:46.
80. Bassaganya-Riera J, Reynolds K, Martino-Catt S, Cui Y, Hennighausen L, Gonzalez F, *et al.* Activation of Ppar gamma and delta by conjugated linoleic acid mediates protection from experimental inflammatory bowel disease. *Gastroenterology* 2004;127:777-91.

81. Yin L, Mou H, Shao J, Zhu Y, Pang X, Yang J, *et al.* Correlation between heart fatty acid binding protein and severe Covid-19: A case-control study. *Plos One* 2020;15:e0231687.
82. Furuhashi M, Hiramitsu S, Mita T, Omori A, Fuseya T, Ishimura S, *et al.* Reduction of circulating Fabb4 level by treatment with omega-3 fatty acid ethyl esters. *Lipids Health Dis* 2016;15:5.
83. Gossell-Williams M, Lyttle K, Clarke T, Gardner M, Simon O. Supplementation with pumpkin seed oil improves plasma lipid profile and cardiovascular outcomes of female non-ovariectomized and ovariectomized sprague-dawley rats. *Phytother Res* 2008;22:873-7.
84. Zhang L, Liu Y. Potential interventions for novel coronavirus in china: A systematic review. *J Med Virol* 2020;92:479-90.
85. Das UN. Can bioactive lipids inactivate coronavirus (Covid-19)? *Arch Med Res* 2020;51:282-6.
86. Voigt RM, Forsyth CB, Green SJ, Mutlu E, Engen P, Vitaterna MH, *et al.* Circadian disorganization alters intestinal microbiota. *PLoS One* 2014;9:e97500.
87. Bäckhed F, Ding H, Wang T, Hooper LV, Koh GY, Nagy A, *et al.* The gut microbiota as an environmental factor that regulates fat storage. *Proc Natl Acad Sci U S A* 2004;101:15718-23.
88. Sanna S, van Zuydam NR, Mahajan A, Kurilshikov A, Vich Vila A, Vösa U, *et al.* Causal relationships among the gut microbiome, short-chain fatty acids and metabolic diseases. *Nat Genet* 2019;51:600-5.
89. Coutzac C, Jouniaux JM, Paci A, Schmidt J, Mallardo D, Seck A, *et al.* Systemic short chain fatty acids limit antitumor effect of Ctlα-4 blockade in hosts with cancer. *Nat Commun* 2020;11:2168.
90. Borenstein R, Hanson BA, Markosyan RM, Gallo ES, Narasipura SD, Bhutta M, *et al.* Ginkgolic acid inhibits fusion of enveloped viruses. *Sci Rep* 2020;10:4746.
91. Mahluji S, Jalili M, Ostadrahimi A, Hallajzadeh J, Ebrahimzadeh-Attari V, Saghaei-Asl M. Nutritional management of diabetes mellitus during the pandemic of COVID-19: a comprehensive narrative review. *J Diabet Metab Disord* 2021;20:1-10.
92. Huq AKO, Bazlur Rahim ANM, Moktadir SMG, Uddin I, Manir MZ, Siddique MAB, *et al.* Integrated nutritional supports for diabetic patients during COVID-19 infection: A comprehensive review. *Curr Diabet Rev* 2022;18:52-60.
93. Jacobs DR Jr, Haddad EH, Lanou AJ, Messina MJ. Food, plant food, and vegetarian diets in the Us dietary guidelines: conclusions of an expert panel. *Am J Clin Nutr* 2009;89:1549-52S.
94. Fujii H, Iwase M, Ohkuma T, Ogata-Kaizu S, Ide H, Kikuchi Y, *et al.* Impact of dietary fiber intake on glycemic control, cardiovascular risk factors and chronic kidney disease in Japanese patients with type 2 diabetes mellitus: the fukuoka diabetes registry. *Nutr J* 2013;12:159.
95. Neelamma G, Swamy BD, Dhamodaran P. Phytochemical and pharmacological overview of Cucurbita maxima and future perspective as potential phytotherapeutic agent. *Eur J Pharm Med Res* 2016;3:277-87.
96. Montesano D, Blasi F, Simonetti MS, Santini A, Cossignani L. Chemical and nutritional characterization of seed oil from Cucurbita maxima L. (var. Berrettina) pumpkin. *Foods* 2018;7:30.
97. Rahim S, Dawar S, Tariq M, Zaki MJ. Mycoflora associated with the seed samples of Cucurbita pepo L. collected from Pakistan. *Pak J Bot* 2013;45:2173-9.
98. Bardaa S, Ben Halima N, Aloui F, Ben Mansour R, Jabeur H, Bouaziz M, *et al.* Oil from pumpkin (cucurbita pepo L.) Seeds: evaluation of its functional properties on wound healing in rats. *Lipids Health Dis* 2016;15:73.
99. Salehi B, Capanoglu E, Adrar N, Catalkaya G, Shaheen S, Jaffer M, *et al.* Cucurbits plants: A key emphasis to its pharmacological potential. *Molecules (Basel, Switzerland)* 2019;24:1854.
100. Gutierrez RP. Review of Cucurbita pepo (pumpkin) its phytochemistry and pharmacology. *Med Chem* 2016;6:12-21.
101. Yoshinari O, Sato H, Igarashi K. Anti-diabetic effects of pumpkin and its components, trigonelline and nicotinic acid, on gotokakizaki rats. *Biosci Biotechnol Biochem* 2009;73:1033-41.
102. Mahmoodpoor A, Medghalchi M, Nazemiyeh H, Asgharian P, Shadvar K, Hamishehkar H. Effect of cucurbita maxima on control of blood glucose in diabetic critically ill patients. *Adv Pharm Bull* 2018;8:347-51.
103. Ezuruike UF, Prieto JM. The use of plants in the traditional management of diabetes in nigeria: pharmacological and toxicological considerations. *J Ethnopharmacol* 2014;155:857-924.
104. Xiao F, Sun J, Xu Y, Li F, Huang X, Li H, *et al.* Infectious Sars-cov-2 in feces of patient with severe Covid-19. *Emerg Infect Dis* 2020;26:1920-2.
105. Li N, Ma WT, Pang M, Fan QL, Hua JL. The commensal microbiota and viral infection: A comprehensive review. *Front Immunol* 2019;10:1551.
106. Ghaffari S, Roshanravan N, Tutunchi H, Ostadrahimi A, Pouraghaei M, Kafil B. Oleoylethanolamide, A bioactive lipid amide, as A promising treatment strategy for coronavirus/ Covid-19. *Arch Med Res* 2020;51:464-7.
107. KD Pharma Group. SLA Pharma initiate clinical trial for COVID-19. Available from: <https://www.kdpharmagroup.com/en/news/kd-pharma-groups-la-pharma-initiate-clinical-trial-covid-19>. [Last accessed 2022 May 16].
108. Caccialanza R, Laviano A, Lobascio F, Montagna E, Bruno R, Ludovisi S, *et al.* Early nutritional supplementation in non-critically ill patients hospitalized for the 2019 novel coronavirus disease (Covid-19): rationale and feasibility of a shared pragmatic protocol. *Nutrition* 2020;74:110835.
109. Jayawardena R, Sooriyaarachchi P, Chourdakis M, Jeewandara C, Ranasinghe P. Enhancing immunity in viral infections, with special emphasis on Covid-19: A review. *Diabet Metab Syndr* 2020;14:367-82.
110. Morittu VM, Musco N, Mastellone V, Bonesi M, Britti D, Infascelli F, *et al.* In vitro and in vivo studies of Cucurbita pepo L. flowers: chemical profile and bioactivity. *Nat Prod Res* 2021;35:2905-9.
111. Iwo MI, Insanu M, Dass CAS. Development of immunonutrient from pumpkin (Cucurbita Moschata Duchense Ex. Lamk.) seed. *Procedia Chem* 2014;13:105-11.
112. Barakat LA, Mahmoud RH. The antiatherogenic, renal protective and immunomodulatory effects of purslane, pumpkin and flax seeds on hypercholesterolemic rats. *N Am J Med Sci* 2011;3:411-7.
113. Annunziata G, Sanduzzi Zamparelli M, Santoro C, Ciampaglia R, Stornaiuolo M, Tenore GC, *et al.* May polyphenols have a role against coronavirus infection? An overview of in vitro evidence. *Front Med (Lausanne)* 2020;7:240.
114. Kiokias S, Proestos C, Oreopoulou V. Phenolic acids of plant origin-a review on their antioxidant activity in vitro (O/W Emulsion Systems) along with their in vivo health biochemical properties. *Foods (Basel, Switzerland)* 2020;9:534.
115. Tsao R. Chemistry and biochemistry of dietary polyphenols. *Nutrients* 2010;2:1231-46.
116. Kapoor B, Gulati M, Gupta R, Singh KS, Gupta M, Nabi A, *et al.* A review on plant flavonoids as potential anticancer agents. *Curr Org Chem* 2021;25:737-47.

117. Abba Y, Hassim H, Hamzah H, Noordin MM. Antiviral activity of resveratrol against human and animal viruses. *Adv Virol* 2015;2015:184241.
118. Zakaryan H, Arabyan E, Oo A, Zandi K. Flavonoids: promising natural compounds against viral infections. *Arch Virol* 2017;162:2539-51.
119. Lalani S, Poh CL. Flavonoids as Antiviral Agents for Enterovirus A71 (EV-A71). *Viruses* 2020;12:184.
120. El-Toumy SA, Salib JY, El-Kashak WA, Marty C, Bedoux G, Bourgougnon N. Antiviral effect of polyphenol rich plant extracts on herpes simplex virus type 1. *Food Sci Human Wellness* 2018;7:91-101.
121. Perez R. Antiviral activity of compounds isolated from plants. *Pharm Biol* 2003;41:107-57.
122. Arora R, Chawla R, Marwah R, Arora P, Sharma RK, Kaushik V, *et al.* Potential of complementary and alternative medicine in preventive management of novel H1N1 flu (swine flu) pandemic: thwarting potential disasters in the bud. *Evid Based Complement Alternat Med* 2011;2011:586506.
123. Chojnacka K, Witek-Krowiak A, Skrzypczak D, Mikula K, Młynarz P. Phytochemicals containing biologically active polyphenols as an effective agent against covid-19-inducing coronavirus. *J Funct Foods* 2020;73:104146.
124. Li H, Wu J, Zhang Z, Ma Y, Liao F, Zhang Y, *et al.* Forsythoside a inhibits the avian infectious bronchitis virus in cell culture. *Phytother Res* 2011;25:338-42.
125. Roh C. A facile inhibitor screening of Sars coronavirus N protein using nanoparticle-based Rna oligonucleotide. *Int J Nanomed* 2012;7:2173-9.
126. Lin SC, Ho CT, Chuo WH, Li S, Wang TT, Lin CC. Effective inhibition of Mers-cov infection by resveratrol. *Bmc Infect Dis* 2017;17:144.
127. Park JY, Yuk HJ, Ryu HW, Lim SH, Kim KS, Park KH, *et al.* Evaluation of polyphenols from broussonetia papyrifera as coronavirus protease inhibitors. *J Enzyme Inhib Med Chem* 2017;32:504-15.
128. Chen C, Zuckerman DM, Brantley S, Sharpe M, Childress K, Hoiczky E, *et al.* Sambucus nigra extracts inhibit infectious bronchitis virus at an early point during replication. *Bmc Vet Res* 2014;10:24.
129. Kulczyński B, Sidor A, Gramza-Michałowska A. Antioxidant potential of phytochemicals in pumpkin varieties belonging to Cucurbita moschata and Cucurbita pepo species. *CyTA – J Food* 2020;18:472-84.
130. Peiretti PG, Meineri G, Gai F, Longato E, Amarowicz R. Antioxidative activities and phenolic compounds of pumpkin (cucurbita pepo) seeds and amaranth (amaranthus caudatus) grain extracts. *Nat Prod Res* 2017;31:2178-82.
131. Patel S, Santani D, Shah M, Patel V. Anti-hyperglycemic and anti-hyperlipidemic effects of bryonia laciniosa seed extract and its saponin fraction in streptozotocin-induced diabetes in rats. *J Young Pharm* 2012;4:171-6.
132. Krimer-Malešević V. Chapter 37 - Pumpkin Seeds: Phenolic Acids in Pumpkin Seed (Cucurbita pepo L.). In: Preedy VR, Watson RR, editors. *Nuts and Seeds in Health and Disease Prevention*. 2nd ed. Amsterdam: Elsevier; 2020. p. 533-42.
133. Kulczyński B, Gramza-Michałowska A. The profile of secondary metabolites and other bioactive compounds in Cucurbita pepo L. and Cucurbita moschata pumpkin cultivars. *Molecules* 2019;24:2945.
134. Fatahiah Z, Zulkhairuazha T, Rosli WW. Quantitative HPLC analysis of gallic acid in Benincasa hispida prepared with different extraction techniques. *Sains Malaysian* 2014;43:1181-7.
135. Sorifa A. Nutritional compositions, health promoting phytochemicals and value added products of bitter gourd: a review. *Int Food Res J* 2018;25.
136. Busuioc AC, Botezatu A-VD, Furdui B, Vinatoru C, Maggi F, Caprioli G, *et al.* Comparative study of the chemical compositions and antioxidant activities of fresh juices from Romanian Cucurbitaceae varieties. *Molecules* 2020;25:5468.
137. Essien EE, Antia BS, Peter N. Lagenaria siceraria (molina) standley. total polyphenols and antioxidant activity of seed oils of bottle gourd cultivars. *World J Pharmaceut Res* 2015;4:274-85.
138. Kulczyński B, Gramza-Michałowska A. The profile of carotenoids and other bioactive molecules in various pumpkin fruits (Cucurbita maxima Duchesne) cultivars. *Molecules* 2019;24:3212.
139. Kostecka-Gugała A, Kruczek M, Ledwożyw-Smoleń I, Kaszycki P. Antioxidants and health-beneficial nutrients in fruits of eighteen cucurbita cultivars: analysis of diversity and dietary implications. *Molecules* 2020;25:1792.
140. Mohan R, Birari R, Karmase A, Jagtap S, Bhutani KK. Antioxidant activity of a new phenolic glycoside from lagenaria siceraria stand. *Fruits. Food Chem* 2012;132:244-51.
141. Okonwu K, Muonekwu J. Potentials of underexploited seed of Trichosanthes cucumerina Linn. *J Appl Sci Environ Manag* 2019;23:791-7.
142. Attar UA, Ghane SG. In vitro antioxidant, antidiabetic, antiacetylcholine esterase, anticancer activities and RP-HPLC analysis of phenolics from the wild bottle gourd (Lagenaria siceraria (Molina) Standl.). *S Afr J Bot* 2019;125:360-70.
143. Nyam K, Tan C, Lai O, Long K, Man YC. Physicochemical properties and bioactive compounds of selected seed oils. *LWT-Food Sci Technol* 2009;42:1396-403.
144. Rezig L, Chouaibi M, Msaada K, Hamdi S. Chemical composition and profile characterisation of pumpkin (Cucurbita maxima) seed oil. *Indus Crop Prod* 2012;37:82-7.
145. Chanda J, Mukherjee PK, Biswas R, Singha S, Kar A, Halder PK. Lagenaria siceraria and its bioactive constituents in carbonic anhydrase inhibition: A bioactivity guided Lc-MS/MS approach. *Phytochem Anal* 2021;32:298-307.
146. Deng Y, Tang Q, Zhang Y, Zhang R, Wei Z, Tang X, *et al.* Protective effect of momordica charantia water extract against liver injury in restraint-stressed mice and the underlying mechanism. *Food Nutr Res* 2017;61:1348864.
147. Lim TK. *Edible Medicinal and Non-Medicinal Plants*. Dordrecht: Springer; 2012.
148. Amin MZ, Islam T, Uddin MR, Uddin MJ, Rahman MM, Satter MA. Comparative study on nutrient contents in the different parts of indigenous and hybrid varieties of pumpkin (cucurbita maxima linn.). *Heliyon* 2019;5:e02462.
149. Yuwai KE, Rao KS, Kaluwin C, Jones GP, Rivett DE. Chemical composition of Momordica charantia L. fruits. *J Agric Food Chem* 1991;39:1762-3.
150. Santos JC, Ribeiro ML, Gambero A. The impact of polyphenols-based diet on the inflammatory profile in Covid-19 elderly and obese patients. *Front Physiol* 2020;11:612268.
151. Hu X, Cai X, Song X, Li C, Zhao J, Luo W, *et al.* Possible SARS-coronavirus 2 inhibitor revealed by simulated molecular docking to viral main protease and host toll-like receptor. *Future Virol* 2020;15:359-68.

152. Al-Zahrani AA. Rutin as a promising inhibitor of main protease and other protein targets of COVID-19: In silico study. *Nat Prod Commun* 2020;15:1934578X20953951.
153. Xu Z, Yang L, Zhang X, Zhang Q, Yang Z, Liu Y, *et al.* Discovery of potential flavonoid inhibitors against Covid-19 3cl proteinase based on virtual screening strategy. *Front Mol Biosci* 2020;7:556481.
154. Verma S, Twilley D, Esmear T, Oosthuizen CB, Reid AM, Nel M, *et al.* Anti-Sars-cov natural products with the potential to inhibit Sars-cov-2 (Covid-19). *Front Pharmacol* 2020;11:561334.
155. Khaerunnisa S, Kurniawan H, Awaluddin R, Suhartati S, Soetjipto S. Potential inhibitor of COVID-19 main protease (Mpro) from several medicinal plant compounds by molecular docking study. *Preprints* 2020;20944:1-14.
156. Lomax AR, Calder PC. Prebiotics, immune function, infection and inflammation: a review of the evidence. *Br J Nutr* 2009;101:633-58.
157. Anand S, Mande SS. Diet, microbiota and gut-lung connection. *Front Microbiol* 2018;9:2147.
158. Conte L, Toraldo DM. Targeting the gut-lung microbiota axis by means of a high-fibre diet and probiotics may have anti-inflammatory effects in Covid-19 infection. *Ther Adv Respir Dis* 2020;14:1753466620937170.
159. Hu B, Huang S, Yin L. The cytokine storm and COVID-19. *J Med Virol* 2021;93:250-6.
160. Rooks MG, Garrett WS. Gut microbiota, metabolites and host immunity. *Nat Rev Immunol* 2016;16:341-52.
161. Sasaki D, Sasaki K, Ikuta N, Yasuda T, Fukuda I, Kondo A, *et al.* Low amounts of dietary fibre increase in vitro production of short-chain fatty acids without changing human colonic microbiota structure. *Sci Rep* 2018;8:435.
162. Robertson MD. Prebiotics and type 2 diabetes: targeting the gut microbiota for improved glycaemic control? *Pract Diabet* 2020;37:133-7.
163. Lokuge GM, Vidanarachchi J, Thavarajah P, Siva N, Thavarajah D, Liyanage R, *et al.* Prebiotic carbohydrate profile and in vivo prebiotic effect of pumpkin (*Cucurbita maxima*) grown in Sri Lanka. *J Nat Sci Found Sri Lanka* 2018;46.
164. Sharma P, Kaur G, Kehinde BA, Chhikara N, Panghal A, Kaur H. Pharmacological and biomedical uses of extracts of pumpkin and its relatives and applications in the food industry: a review. *Int J Veg Sci* 2020;26:79-95.
165. Agarkova EY, Kruchinin AG, Glazunova OA, Fedorova TV. Whey protein hydrolysate and pumpkin pectin as nutraceutical and prebiotic components in a functional mousse with antihypertensive and bifidogenic properties. *Nutrients* 2019;11:2930.
166. Du B, Song Y, Hu X, Liao X, Ni Y, Li Q. Oligosaccharides prepared by acid hydrolysis of polysaccharides from pumpkin (*Cucurbita moschata*) pulp and their prebiotic activities. *Int J Food Sci Technol* 2011;46:982-7.
167. Genevois C, Flores S, de Escalada Pla M. Byproduct from pumpkin (*Cucurbita moschata* Duchesne ex poiret) as a substrate and vegetable matrix to contain *Lactobacillus casei*. *J Function Foods* 2016;23:210-9.
168. Kushawaha DK, Yadav M, Chatterji S, Srivastava A, Watal G. α -Amylase and α -glucosidase inhibitory activity assessment of *Cucurbita maxima* seeds—a LIBS based study. *Int J Phytomed* 2016;8:312-8.
169. Cerqueira PMd, Freitas MCJ, Pumar M, Santangelo SB. The pumpkin (*Cucurbita maxima*, L.) seed flour effect on the rat glucose and lipid metabolism. *Revista de Nutrição* 2008;21:129-36.
170. Tan J, McKenzie C, Potamitis M, Thorburn AN, Mackay CR, Macia L. The role of short-chain fatty acids in health and disease. *Adv Immunol* 2014;121:91-119.
171. Sreenivas K, Lele S. Prebiotic activity of gourd family vegetable fibres using in vitro fermentation. *Food Bioscience* 2013;1:26-30.
172. Jacobs DR Jr, Tapsell LC. Food, not nutrients, is the fundamental unit in nutrition. *Nutr Rev* 2007;65:439-50.