

**The effect of transcranial direct current stimulation (tDCS) on food craving, reward and appetite in a healthy population.**

BEAUMONT, Jordan <<http://orcid.org/0000-0002-9861-3379>>, DAVIS, Danielle, DALTON, Michelle, NOWICKY, Alexander, RUSSELL, Mark <<http://orcid.org/0000-0002-7305-1090>> and BARWOOD, Martin J

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/31270/>

---

This document is the Accepted Version [AM]

**Citation:**

BEAUMONT, Jordan, DAVIS, Danielle, DALTON, Michelle, NOWICKY, Alexander, RUSSELL, Mark and BARWOOD, Martin J (2021). The effect of transcranial direct current stimulation (tDCS) on food craving, reward and appetite in a healthy population. *Appetite*, 157: 105004. [Article]

---

**Copyright and re-use policy**

See <http://shura.shu.ac.uk/information.html>

**The effect of transcranial direct current stimulation (tDCS) on food craving, reward and appetite in a healthy population.**

Jordan D. Beaumont<sup>a,\*</sup>, Danielle Davis<sup>a</sup>, Michelle Dalton<sup>a</sup>, Alexander Nowicky<sup>b</sup>, Mark Russell<sup>a</sup> and Martin J. Barwood<sup>a</sup>

<sup>a</sup>*School of Social and Health Sciences, Leeds Trinity University, Leeds, LS18 5HD, UK*

<sup>b</sup>*Centre for Cognitive Neuroscience, Department of Clinical Sciences, College of Health and Life Sciences, Brunel University London, Uxbridge, UB8 3PH, UK*

*\*Corresponding author, email: [j.beaumont@leedstrinity.ac.uk](mailto:j.beaumont@leedstrinity.ac.uk)*

## ABSTRACT

The ability to control hedonic appetite is associated with executive functioning originating in the prefrontal cortex (PFC). The rewarding components of food can override homeostatic mechanisms, potentiating obesogenic behaviours. Indeed, those susceptible to overconsumption appear to have PFC hypo-activation. Transcranial direct current stimulation (tDCS) over the dorsolateral PFC (DLPFC) has been shown to reduce food craving and consumption, potentially via attenuating this reward response. We examined the effects of stimulation on food reward and craving using a healthy-weight cohort. This study is amongst the first to explore the effects of tDCS on explicit and implicit components of reward for different food categories. Twenty-one healthy-weight participants ( $24 \pm 7$  years,  $22.8 \pm 2.3$  kg·m<sup>-2</sup>) completed two sessions involving double-blind, randomised and counterbalanced anodal or sham tDCS over the right DLPFC, at 2 milliampere for 20 minutes. Food craving (Food Craving Questionnaire-State), reward (Leeds Food Preference Questionnaire), and subjective appetite (100 mm visual analogue scales) were measured pre- and post-tDCS. Eating behaviour trait susceptibility was assessed using the Three Factor Eating Questionnaire-Short Form, Control of Eating Questionnaire, and Food Craving Questionnaire-Trait-reduced. Stimulation did not alter food craving, reward or appetite in healthy-weight participants who displayed low susceptibility to overconsumption, with low trait craving, good craving control, and low uncontrolled eating and emotional eating behaviour. Implicit and explicit reward were reliable measures of hedonic appetite, suggesting these are robust targets for future tDCS research. These findings suggest that applying tDCS over the DLPFC does not change food reward response in individuals not at risk for overconsumption, and future work should focus on those at risk of overconsumption who may be more responsive to the effects of tDCS on hedonic appetite.

37 **KEYWORDS**

38 Appetite control; Dorsolateral prefrontal cortex; Neuromodulation; Food craving; Food  
39 reward

40

41 **HIGHLIGHTS**

- 42 • We consider the effects of tDCS on implicit and explicit reward using a validated task
- 43 • High reliability in reward measures suggest a robust target for future tDCS studies
- 44 • Previously findings are limited by high variation within food-related variables
- 45 • Effects of tDCS may be dependent on participant eating behaviour traits
- 46 • Future work should screen participants using validated psychometric questionnaires

## 1. INTRODUCTION

Obesity is a global health epidemic that affects more than 650 million adults worldwide, and is associated with an increased risk of developing many other health conditions (World Health Organisation, 2020). The aetiology of obesity involves a complex relationship between behavioural, biological and environmental factors, contributing to the dysregulation of energy balance (Hill, 2006). Hedonic appetite can potentiate this dysregulation, with the rewarding components of food overriding homeostatic mechanisms (Boswell & Kober, 2016; Kober & Boswell, 2018). The ability to control hedonic appetite is associated with executive functioning, which originate in the prefrontal cortex (PFC) and inhibit impulsive actions in favour of goal-directed behaviours (Joseph, Alonso-Alonso, Bond, Pascual-Leone, & Blackburn, 2011). Altered PFC activity in response to food stimuli has been identified in individuals with obesity, especially those displaying binge eating symptoms (Boeka & Lokken, 2011; Karhunen, et al., 2000). It is proposed that a reduction of activity in the right dorsolateral PFC (DLPFC) could facilitate obesogenic behaviours through poor appetite control (Alonso-Alonso & Pascual-Leone, 2007). Indeed, dysregulation of the DLPFC has been linked with greater impulsive behaviours, often leading to overconsumption (Gluck, Viswanath, & Stinson, 2017). Increasing DLPFC activity may improve the ability to control hedonic appetite, providing a novel paradigm in the treatment of obesity (Alonso-Alonso, 2013).

The modulation of cortical activity is possible through the use of non-invasive brain stimulation techniques, such as transcranial direct current stimulation (tDCS). This form of stimulation involves the application of a weak electrical current, typically up to 2 milliamperes (mA), to a specific region of the brain via two electrodes that are placed over the scalp (Nitsche & Paulus, 2000). The current is emitted from a battery-powered device, where it is delivered to the brain through an anode electrode and returns to the device through a cathode electrode. The current intensity is not sufficient to cause neuronal firing, but results in the polarity-dependent subthreshold modulation of resting membrane potentials (Filmer,

Dux, & Mattingley, 2014; Jamil & Nitsche, 2017). Although the exact mechanisms are not fully understood, it appears the current inhibits neurotransmitters at the synapse; the anode is associated with the inhibition of gamma-aminobutyric acid (GABA) whereas the cathode is associated with the inhibition of glutamate (Filmer, et al., 2014; Stagg, Antal, & Nitsche, 2018). The inhibition of these neurotransmitters increases or decreases the likelihood of spontaneous neuronal firing, respectively. In addition to these acute effects, tDCS also appears to elicit changes in cortical activity beyond the stimulation duration. For example, in an early study by Nitsche and Paulus (2001), anodal tDCS lasting 13 minutes resulted in greater activity in the motor cortex for up to 90 minutes post-stimulation.

When identifying the effects of tDCS on hedonic appetite, many studies have focussed on measuring state food craving. The first study to identify the impact of tDCS on hedonic appetite compared anodal stimulation to the left and right DLPFC in 21 healthy-weight individuals with frequent food cravings, defined as experiencing 3 or more strong urges to consume high-calorie foods per day (Fregni, et al., 2008). When applying 2 mA stimulation for 20 minutes, a significant reduction in food craving was observed following tDCS over the right DLPFC, but not when applied to the left hemisphere. This reduction in state craving score was replicated in a second study that used the same stimulation parameters and recruited a similar participant cohort ( $n = 19$ ) (Goldman, et al., 2011).

In two recent publications, Burgess and colleagues highlight a potential eating behaviour trait-dependent effect of tDCS (Burgess, et al., 2016; Ray, et al., 2017). Thirty participants who were obese and met the diagnostic criteria for binge eating disorder (BED) underwent 20 minutes of 2 mA tDCS to the right DLPFC, which resulted in a significant decrease in state food craving and in-laboratory food consumption (Burgess, et al., 2016). In contrast, these effects were not significant when this protocol was replicated in 18 participants with frank obesity (i.e. non-binge eating) (Ray, et al., 2017). This suggests that the effects of tDCS may be dependent on individual variation in the level of susceptibility to reward-driven

overconsumption. Consistent with this, previous research has demonstrated that individuals with BED are hyper-responsive to the rewarding aspects of food (Davis, 2013; Davis, et al., 2009). The estimated prevalence of BED in the general population ranges from 0.7 – 3.0%, and is commonly comorbid with overweight and obesity (Kessler, et al., 2013). Recurrent episodes of binge eating behaviour are estimated to occur in 10 – 20% of individuals who are healthy weight, overweight or obese, and constitutes a trait that can be assessed psychometrically and applied to a non-clinical population. Similar to findings in individuals with BED, individuals with eating behaviour trait susceptibility to overconsume (i.e. binge eating and emotional eating) have been found to be hyper-responsive to the rewarding aspects of food (Dalton, Blundell, & Finlayson, 2013a). Therefore, including validated measures of food reward and eating behaviour trait susceptibility may be important when considering the effect of tDCS on food consumption, reward and craving. To date, no study has identified the effects of tDCS on implicit and explicit components of reward across different food categories.

Although there are many promising findings, not all studies have found an effect of tDCS on measures of hedonic appetite. This may be due to the inconsistent application of stimulation parameters (e.g. variation in target electrode placement and current intensity), inadequate experimental blinding, and large variation in experimental measures (Hall, Vincent, & Burhan, 2018; Tremblay, et al., 2014). The most consistently used measure of hedonic appetite in tDCS research is food craving, which is commonly assessed using the Food Craving Questionnaire-State (FCQ-S) (Cepeda-Benito, Gleaves, Williams, & Erath, 2000). Although significant effects of tDCS on food craving have been identified (Fregni, et al., 2008; Goldman, et al., 2011), this has not been consistently shown (Georgii, Goldhofer, Meule, Richard, & Blechert, 2017; Sedgmond, et al., 2019). Across studies there is large variation in state food craving scores, ranging from 0.40% to 41.67% following the active condition (Fregni, et al., 2008; Goldman, et al., 2011; Kekic, et al., 2014; Ljubisavljevic, Maxood, Bjekic, Oommen, & Nagelkerke, 2016), which may be due to the poor reliability of

these measures. Developmental publications of the FCQ-S suggest low-to-moderate reliability ( $r = 0.39 - 0.56$ ) (Cepeda-Benito, et al., 2000; Meule, Teran, et al., 2014). Measures of food consumption have also been utilised, primarily using *ad libitum* buffets of highly palatable foods (Burgess, et al., 2016; Georgii, et al., 2017; Gluck, et al., 2015; Ray, et al., 2017; Sedgmond, et al., 2019). Although greater craving control is associated with improved weight loss outcomes (Dalton, et al., 2017), the effects of tDCS on craving and consumption are not correlated (Burgess, et al., 2016), suggesting other targets are required to validate tDCS as an intervention to alter eating behaviour. Food reward plays a more pivotal role in the dysregulation of energy balance (Boswell & Kober, 2016; Kober & Boswell, 2018). Therefore, it is important to look beyond the measure of food craving and identify the role of tDCS in modulating food reward.

The present study examined how measures of food craving, reward and appetite would change following the inducement of hyper-activity of the right DLPFC through tDCS in a healthy-weight cohort. We hypothesised stimulation would reduce state food craving and subjective appetite, based on previous findings utilising healthy participant groups (Fregni, et al., 2008; Goldman, et al., 2011; Kekic, et al., 2014; Lapenta, Sierve, de Macedo, Fregni, & Boggio, 2014). We also hypothesised that participants' preference for and perceived rewarding value of high-fat and sweet foods would be diminished following anodal tDCS. We also looked to establish the reliability of these measures, including both implicit and explicit components of reward, prior to tDCS with a view to establishing the viability of their future use in our research.

## **2. METHODS**

### **2.1 Participants**

The study was approved by an institutional ethics committee, and all participants provided written informed consent. Sample size was determined using G\*Power 3.0.10 (Faul, Erdfelder, Lang, & Buchner, 2007). An effect size  $f$  of 0.33 was based on mean percentage




difference from baseline in food craving scores following single session tDCS (mean difference between conditions  $-22.22 \pm 33.68\%$ ) (Fregni, et al., 2008; Goldman, et al., 2011; Kekic, et al., 2014; Ljubisavljevic, et al., 2016). Using  $\alpha$  error probability of 0.05, 1 group with 2 measurements, a correlation among repeated measures equal to 0.5, and non-sphericity correlation  $\epsilon$  of 1, sample size calculations determined a minimum sample size of 21, with actual power of 0.82, was required. Twenty-one participants ( $24 \pm 7$  years,  $22.8 \pm 2.3 \text{ kg}\cdot\text{m}^{-2}$ ) were recruited via email and poster advertisements. Interested individuals were initially screened with an online questionnaire. Eligible participants were male or female between 18 and 60 years of age who presented with a body mass index (BMI) between 18.5 and 24.9. All participants were free of neurological, cardiovascular, metabolic and joint disease, and potential participants were excluded if they presented with low mood or depressive symptoms, as indicated using the Centre for Epidemiologic Studies Short Depression Scale (CESD-10) (Andresen, Malmgren, Carter, & Patrick, 1994; Radloff, 1977). Female participants who were pregnant or wishing to conceive were also excluded from the study. Participants were naïve to tDCS protocols, non-smokers and were not recreational drug users or taking any medications at the time of data collection.

## **2.2 Experimental Design**

The study utilised a double-blind, within-participant, repeated measures design. Participants attended the laboratory on three separate occasions (Figure 1). During the first visit, participants completed a series of psychometric questionnaires, and height and body composition were measured. Visits 2 and 3 were experimental trials where all participants received either active or sham tDCS in a randomised and counterbalanced order. Randomisation was determined using a permuted block paradigm and completed by an independent party. The participants and researcher conducting stimulation were blind to the tDCS condition, adhering to a double-blind design, which was maintained using a pin-protected device. All sessions were scheduled at the same time of day within-subject, occurring between 09:00 and 15:00, and with a minimum interval of 48 hours between

sessions to prevent any residual effects of stimulation (Alonzo, Brassil, Taylor, Martin, & Loo, 2012).



**Figure 1** Study Procedure

FCQ-S, Food Craving Questionnaire-State; LFPQ, Leeds Food Preference Questionnaire; tDCS, transcranial direct current stimulation; VAS, visual analogue scales.

### 2.3 Procedure

Participants were required to fast for a minimum of 4 hours prior to each visit, where they were asked to refrain from consuming any food or drink other than water. In addition, they were asked to refrain from consuming products containing caffeine and alcohol in the 12 or 24 hours prior to each visit, respectively. Adherence to this fasting criteria was self-reported at the start of each visit. On arrival at the laboratory, participants were instructed to turn off their mobile phones and remove any metallic objects from their person in adherence with our tDCS protocol.

During visit 1, participants completed the Three Factor Eating Questionnaire-Short Form (TFEQ-r18), Control of Eating Questionnaire (CoEQ), and Food Craving Questionnaire-Trait-reduced (FCQ-T-r); see 2.5.1. Height was measured using a portable stadiometer (SECA

Limited, Birmingham, UK) to the nearest mm. Measurements were taken following inhalation, with the participant standing straight, their back to the rule, and their eyeline level with ear canal. Body composition, including weight and BMI, was assessed using a Tanita BC-418MA analyser (Tanita Europe B.V., Amsterdam). Weight was measured to the nearest 0.1 kg, and body fat percentage to the nearest 0.1%. Participants were then shown the food images used in the Leeds Food Preference Questionnaire (LFPQ; see 2.5.4), and their familiarity and acceptance of each food item was assessed. Any food items that were unfamiliar or had low acceptance (i.e. disliked or not consumed in the normal diet) were substituted with images from a database of additional items with similar nutritional and sensory properties (Oustric, et al., 2020).

During visits 2 and 3, participants completed a series of questionnaires immediately pre-tDCS (Figure 1). These included appetite visual analogue scales (VAS), the FCQ-S and the LFPQ; see 2.5. Each participant then underwent 20 minutes of active or sham tDCS. Questionnaires were then repeated immediately post-stimulation. Visits 2 and 3 were identical, apart from the stimulation condition. At the end of visit 3, participants were debriefed. They were informed of the sham stimulation condition and were asked whether they were able to differentiate between the active and sham conditions, and in which visit they believe active tDCS was delivered.

#### **2.4 Stimulation Protocol**

Stimulation was delivered using the HDCstim direct current stimulator (Newronika s.r.l., Milan, Italy) by a trained researcher. Anodal stimulation was used to target the right DLPFC, in accordance with the International Standards for Electroencephalography 10-20 system (Klem, Lüders, Jasper, & Elger, 1999). A 25 cm<sup>2</sup> anode electrode was placed over the frontal area 4 (F4) and a 51 cm<sup>2</sup> cathode electrode placed over occipital zero point (Oz). Cathode placement over the Oz reduces the impact of associated inhibitory effects on study measures (Bestmann, de Berker, & Bonaiuto, 2015; Galetta, 2017), and decreases the

likelihood of current shunting across the scalp (Rush & Driscoll, 1968). Rubber electrodes were housed in sponge pads, pre-soaked in 0.9% sodium chloride. A constant current of 2 mA was delivered through the anode electrode, culminating in a current density of 0.08 mA·cm<sup>-2</sup>. The current was ramped up over a 30-second period, and active tDCS was then delivered for 20 minutes, with a 30-second ramp down. Stimulation was delivered offline (i.e. no task was performed during tDCS), and participants were instructed to remain seated, relaxed and awake. Sham stimulation involved the same setup as active tDCS, but the current was only delivered for 36 seconds (3% active tDCS duration). This is associated with similar sensations (e.g. itching, tingling) (Brunoni, et al., 2011; Nikolin, Huggins, Martin, Alonzo, & Loo, 2018), but has a limited neuromodulatory effect (Gandiga, Hummel, & Cohen, 2006).

The effectiveness of sham as a blinding technique was assessed during debrief. Impedance was measured at the start of stimulation, and periodically checked thereafter. It is recommended that impedance should remain below 5 kilo-ohm (kΩ) (DaSilva, Volz, Bikson, & Fregni, 2011; Thair, Holloway, Newport, & Smith, 2017). The occurrence of sensations and adverse events were measured using a standardised questionnaire (Brunoni, et al., 2011) immediately following stimulation and at regular intervals for a minimum of 45 minutes, in accordance with our standardised procedure.

## **2.5 Measurements**

### **2.5.1 Psychometric Questionnaires**

A series of psychometric questionnaires were used to assess eating behaviour traits of participants in the screening session (Figure 1). The TFEQ-r18 (Karlsson, Persson, Sjöström, & Sullivan, 2000) measures three subscales of eating behaviour; cognitive restraint, uncontrolled eating, and emotional eating. Scores range from 0 to 100 for each subscale, with higher scores indicating a greater presence of problematic eating behaviour. The TFEQ-r18 has good internal validity, with a Cronbach's alpha (α) of 0.82, and

comparable construct validity to the full TFEQ ( $r = 0.71 - 0.99$ ) (Karlsson, et al., 2000). The CoEQ (Dalton, Finlayson, Hill, & Blundell, 2015) considers the frequency, intensity and severity of food cravings experienced over the previous 7 days. Items are assessed using 100 mm VAS, with scores averaged across items to provide an individual score for craving control, craving for sweet foods, craving for savoury foods, and positive mood. Cronbach's  $\alpha$  highlighted acceptable internal validity; craving control  $\alpha = 0.88$ , craving for sweet foods  $\alpha = 0.67$ , craving for savoury foods  $\alpha = 0.66$ , positive mood  $\alpha = 0.74$ . Finally, general and habitual food cravings were measured using the 15-item FCQ-T-r (Meule, Hermann, & Kübler, 2014). This questionnaire assesses lack of control over eating, emotions experienced before or during food craving and consumption, and guilt from cravings and/or giving in to cravings. A higher score suggests more frequent cravings and a total score greater than 50 highlights clinically relevant trait cravings (Meule, 2018). The FCQ-T-r has high internal validity ( $\alpha = 0.94$ ).

### 2.5.2 Appetite Visual Analogue Scales (VAS)

Four 100 mm VAS were used to assess subjective ratings of appetite (Blundell, et al., 2010), which are sensitive to experimental manipulation and considered reliable and valid measures of subjective appetite (Beechy, Galpern, Petrone, & Das, 2012). Scales measured hunger ("How hungry do you feel right now?"), fullness ("How full do you feel right now?"), prospective consumption ("How much food could you eat right now?"), and the desire to eat ("How strong is your desire to eat right now?"). Scores range from 0 to 100, with higher scores indicating greater prevalence of the appetite measure.

### 2.5.3 Food Craving Questionnaire-State (FCQ-S)

The FCQ-S (Cepeda-Benito, et al., 2000) measures subjective food craving, and is responsive to situational changes (Cepeda-Benito, et al., 2000; Meule, Teran, et al., 2014). This questionnaire assesses the desire to eat, craving for food, and emotional responses to food and consumption over 15 statements. Participants rate each statement on a 5-point

scale, where 1 corresponds with “*Strongly disagree*” and 5 corresponds with “*Strongly agree*”. Corresponding scores are totalled, with a minimum score of 15 and a maximum of 75; higher scores equating to greater momentary craving. Similar to the FCQ-T-r, the state FCQ has good internal validity (Cronbach’s  $\alpha = 0.94$ ) (Cepeda-Benito, et al., 2000).

#### 2.5.4 Leeds Food Preference Questionnaire (LFPQ)

The LFPQ (Dalton & Finlayson, 2014; Finlayson, King, & Blundell, 2007) is a validated computer-based assessment of hedonic preference for food, measuring explicit liking and wanting and implicit wanting as components of reward. “Liking” can be defined as the subjective pleasure elicited by food or related cues, whereas “wanting” is the motivational component of reward that refers to subjective desire or craving for foods (see Finlayson and Dalton (2012) for review). Liking operates at an explicit level (i.e. conscious, introspective), and wanting at both explicit and implicit (i.e. subconscious, automatic) levels (Finlayson & Dalton, 2012). The task uses a standardised set of 16 images depicting ready-to-eat foods that are common in the diet (Table 1), and reward is assessed according to the fat content and taste of these foods. Food images illustrate items that are either high (>40% energy) or low (<20% energy) in fat, and either sweet or savoury. Food items are split into four categories; high-fat savoury (HFSA), high-fat sweet (HFSW), low-fat savoury (LFSA), and low-fat sweet (LFSW). The food items are comparable in protein content, palatability and familiarity (Oustric, et al., 2020).

**Table 1** Standardised food images used in the LFPQ.

HFSA	HFSW	LFSA	LFSW
Garlic bread	Chocolate biscuits	Green salad	Mixed berry salad
Fries	Glazed doughnut	Broccoli	Skittles
Crisps	Blueberry muffin	Vegetable rice	Haribo
Sausage roll	Milk chocolate	Bread roll	Banana

HFSA, high-fat savoury; HFSW, high-fat sweet; LFSA, low-fat savoury; LFSW, low-fat sweet

The LFPQ involves two tasks, where food items are displayed in pairs (forced-choice task) or individually (single-food task). The forced-choice task measures the implicit wanting for foods. Participants are required to choose the food they most want to consume “right now” from two items presented on a computer screen. Scores for implicit wanting are calculated using a frequency-weighted algorithm, by combining reaction time and the frequency of choosing or avoiding a food (Dalton & Finlayson, 2014). In the single-food task, participants are presented with each of the 16 food items individually and asked to rate “*How much do you want some of this food right now?*” and “*How pleasant would it be to taste some of this food right now?*” on 100 mm VAS. This second task measures explicit wanting and liking, respectively, for each food item. In addition, fat appeal bias (FAB) and taste appeal bias (TAB) scores are calculated by subtracting mean scores across food groups (e.g. mean low-fat scores subtracted from mean high-fat scores), and provide scores for explicit liking, explicit wanting and implicit wanting.

## 2.6 Data Analysis

Mean and standard deviations (SD) were calculated at each time point (pre- and post-stimulation) under active and sham tDCS conditions. Normality of data were assessed using Shapiro-Wilks test, and reliability of baseline measures were determined using Pearson’s  $r$

correlations. The effects of tDCS on FCQ-S and LFPQ scores were evaluated using a 2 (condition; active or sham) \* 2 (time point; pre-stimulation vs post-stimulation scores) repeated-measures analysis of variance (ANOVA). Post-hoc significant effects were determined using pair-wise comparisons with Bonferroni correction. Although fasting protocols were standardised, significant differences in baseline scores across all appetite VAS measures were found. To control for this difference, scores were transformed to difference from baseline and analysed using a paired-samples t-test. To determine the impact of difference in baseline hunger scores on other test variables, analysis of covariance (ANCOVA) were performed with baseline hunger as a covariate. ANCOVA were additionally performed to control for behaviour trait scores. Adverse events were compared using further paired-samples t-tests. Statistical analyses were performed using SPSS version 21 and 26 (IBM, New York, USA). Data are presented as mean  $\pm$  SD to an alpha level of 0.05.

To interpret the null findings and assess the strength of evidence, Bayesian statistics were computed using JASP (version 0.13.1; University of Amsterdam). The classification scheme by Lee and Wagenmakers (2013) provides descriptive labels for Bayes factors ( $BF_{10}$ ), and was used to interpret values. In brief, scores greater than 1 provide evidence in favour of the alternative hypothesis, whereas scores below 1 provide evidence in favour of the null hypothesis. Scores are labelled as anecdotal (score between 1 and 3 or 1 and 0.33), moderate (score between 3 and 10 or 0.33 and 0.10), strong (score between 10 and 30 or 0.10 and 0.03), very strong (score between 30 and 100 or 0.03 and 0.01), or extreme (score greater than 100 or lesser than 0.01).

### 3. RESULTS

Participant anthropometric and psychometric characteristics are displayed in Table 2. Participants were weight stable ( $\pm 5\%$ ) for 3 months prior to the study, and displayed “healthy” eating behaviour trait profiles as identified by FCQ-T, CoEQ and TFEQ-r18 scores. Scores for the FCQ-T-r were below the 50-point cut-off for clinically relevant trait craving



361 (Meule, 2018), with CoEQ and TFEQ-r18 scores comparable to healthy-weight individuals in  
362 other studies (Anglé, et al., 2009; De Lauzon-Guillain, et al., 2009; Fleurbaix Laventie Ville  
363 Sante Study, 2004; Wardle, et al., 2018).

364

365 **Table 2** Mean, standard deviation and range for participant anthropometric and eating behaviour trait characteristics

	Female	Male	All
N	11	10	21
Age (years)	25 ± 9 (19 – 52)	23 ± 4 (20 – 29)	24 ± 7 (19 – 52)
Height (cm)	165 ± 6 (155 – 175)	179 ± 6 (170 – 189)	172 ± 9 (155 – 189)
Weight (kg)	60.1 ± 7.4 (49.6 – 71.4)	76.5 ± 7.1 (66.6 – 88.9)	67.9 ± 11.0 (49.6 – 88.9)
BMI (kg·m <sup>-2</sup> )	22.0 ± 2.1 (18.5 – 25.0)	23.8 ± 2.2 (20.1 – 27.7*)	22.8 ± 2.3 (18.5 – 27.7*)
Body fat (kg)	16.3 ± 4.3 (10.9 – 23.3)	12.9 ± 4.9 (6.4 – 20.7)	14.7 ± 4.8 (6.4 – 23.3)
Body fat (%)	26.8 ± 4.3 (20.6 – 33.1)	16.6 ± 5.5 (9.2 – 26.0)	21.9 ± 7.1 (9.2 – 33.1)
CESD-10 (AR)	5 ± 3 (0 – 10)	5 ± 4 (0 – 10)	5 ± 3 (0 – 10)
FCQ-T-r (AR)	36 ± 8 (22 – 49)	34 ± 10 (20 – 47)	35 ± 9 (20 – 49)
TFEQ-r18 Cognitive Restraint (AR)	34 ± 19 (5.6 – 61.1)	33 ± 21 (11.1 – 77.8)	40 ± 20 (5.6 – 77.8)
TFEQ-r18 Uncontrolled Eating (AR)	33 ± 11 (7.4 – 44.4)	34 ± 18 (3.7 – 66.7)	33 ± 14 (3.7 – 66.7)
TFEQ-r18 Emotional Eating (AR)	24 ± 24 (0.0 – 66.7)	20 ± 23 (0.0 – 66.7)	22 ± 22 (0 – 66.7)
CoEQ Craving Control (mm)	66 ± 18 (36.0 – 96.2)	68 ± 18 (36.4 – 94.1)	65 ± 18 (36.0 – 96.2)
CoEQ Craving for Sweet Foods (mm)	30 ± 16 (3.0 – 59.7)	28 ± 21 (2.3 – 67.0)	29 ± 18 (2.3 – 67.0)
CoEQ Craving for Savoury Foods (mm)	54 ± 19 (16 – 78)	46 ± 26 (2.0 – 79.3)	51 ± 23 (2.0 – 79.3)
CoEQ Positive Mood (mm)	51 ± 16 (20 – 84)	54 ± 13 (31.0 – 68.3)	52 ± 14 (20 – 84)

\* n = 1 with BMI >24.9 due to high fat-free mass (weight 88.9 kg, fat-free mass 74.2 kg, fat mass 14.7 kg / 16.5%).


BMI, Body Mass Index; CESD-10, Centre for Epidemiologic Studies Short Depression Scale; FCQ-T-r, Food Craving Questionnaire-Trait reduced form; TFEQ-r18, Three Factor Eating Questionnaire 18-item version; CoEQ, Control of Eating Questionnaire.

367

### 368 **3.1 Appetite Visual Analogue Scales (VAS)**

369 Baseline hunger scores were significantly higher in the active session ( $63.1 \pm 21.4$  mm),  
370 when compared to the sham session ( $51.9 \pm 25.8$  mm) ( $t_{(20)} = 2.567$ ,  $p = 0.018$ ). Similarly,  
371 scores for fullness ( $t_{(20)} = 2.925$ ,  $p = 0.008$ ), prospective consumption ( $t_{(20)} = 3.196$ ,  $p =$   
372  $0.005$ ), and desire to eat ( $t_{(20)} = 2.756$ ,  $p = 0.012$ ) were greater at baseline in the active  
373 versus sham session. Baseline hunger scores did not significantly affect fullness,  
374 prospective consumption or desire to eat ( $p$ 's  $> 0.05$ ). There were no significant changes in  
375 subjective hunger ( $t_{(20)} = 0.572$ ,  $p = 0.574$ ,  $BF_{10} = 0.264$ ), fullness ( $t_{(20)} = 0.146$ ,  $p = 0.886$ ,  
376  $BF_{10} = 0.230$ ), prospective consumption ( $t_{(20)} = 0.969$ ,  $p = 0.344$ ,  $BF_{10} = 0.345$ ), or desire to  
377 eat ( $t_{(20)} = 1.772$ ,  $p = 0.092$ ,  $BF_{10} = 0.858$ ) when comparing pre- and post-stimulation in the  
378 active and sham tDCS conditions (Figure 2). Bayes factors show moderate evidence in  
379 favour of the null hypothesis over the alternative hypothesis for hunger and fullness,  
380 whereas prospective consumption and the desire to eat were supported only by anecdotal  
381 evidence in favour of the null hypothesis. When controlling for behaviour traits scores, the  
382 effects of tDCS on the desire to eat only neared significance ( $p = 0.062 - 0.076$ ), and  
383 remained non-significant for other subjective appetite measures ( $p$ 's  $> 0.32$ ).

384



**Figure 2** Mean  $\pm$  SD appetite visual analogue scale (VAS) scores prior to and following tDCS intervention ( $n = 21$ ).

### 3.2 Food Craving Questionnaire-State (FCQ-S)

There were no significant differences in state food craving from pre- to post-stimulation under active (pre-tDCS  $47.2 \pm 9.9$  AU, post-tDCS  $47.8 \pm 12.2$  AU) or sham conditions (pre-tDCS  $43.8 \pm 10.2$  AU, post-tDCS  $44.9 \pm 9.0$  AU) ( $F_{(1, 19)} = 0.069$ ,  $p = 0.797$ ). Bayes factor highlights moderate evidence in favour of the null hypothesis over the alternative hypothesis ( $BF_{10} = 0.272$ ). In addition, this effect remained non-significant when controlling for baseline hunger ( $F_{(1, 38)} = 0.037$ ,  $p = 0.849$ ) and behaviour trait scores ( $p > 0.74$ ).

### 3.3 Leeds Food Preference Questionnaire (LFPQ)

Stimulation did not alter measures of implicit or explicit food reward, with the condition  $\times$  time point interactions for the liking and wanting of HFSA, LFSA, HFSW and LFSW categories showing no significant effect ( $p > 0.05$ ) (Table 3), which is supported by moderate-to-strong evidence in favour of the null hypothesis over the alternative hypothesis ( $BF_{10} = 0.041$  –

0.168). The interactions remained non-significant when controlling for baseline hunger ( $p$ 's > 0.10) and behaviour trait scores ( $p$ 's > 0.11). In addition, tDCS did not significantly change implicit or explicit TAB, with non-significant condition \* time point interactions for explicit liking ( $F_{(1, 18)} = 0.079$ ,  $p = 0.782$ ,  $BF_{10} = 0.030$ ), explicit wanting ( $F_{(1, 18)} = 0.902$ ,  $p = 0.355$ ,  $BF_{10} = 0.078$ ), and implicit wanting ( $F_{(1, 17)} = 0.786$ ,  $p = 0.388$ ,  $BF_{10} = 0.076$ ). Again, this is supported by strong evidence in favour of the null hypothesis over the alternative hypothesis and the effects remained non-significant when controlling for baseline hunger ( $p$ 's > 0.40) and behaviour trait scores ( $p$ 's > 0.42). Similar non-significant condition \* time point interactions were seen for FAB explicit wanting ( $F_{(1, 18)} = 0.136$ ,  $p = 0.716$ ,  $BF_{10} = 0.183$ ) and implicit wanting ( $F_{(1, 17)} = 0.646$ ,  $p = 0.433$ ,  $BF_{10} = 0.111$ ). These scores remained non-significant when controlling for baseline hunger ( $p = 0.823$  and  $0.236$ , respectively) and behaviour trait scores ( $p$ 's > 0.24). However, there was a significant time point ( $F_{(1, 18)} = 6.785$ ,  $p = 0.018$ ) and condition \* time point interaction for FAB explicit liking ( $F_{(1, 18)} = 7.374$ ,  $p = 0.014$ ,  $BF_{10} = 0.545$ ); scores increased following both active and sham tDCS, and to a greater extent following active stimulation (Table 3). After controlling for baseline hunger scores this effect was no longer significant ( $F_{(1, 36)} = 2.944$ ,  $p = 0.095$ ,  $BF_{10} = 0.313$ ). Similarly, when controlling for baseline behaviour trait scores no significant effects were identified ( $p$ 's > 0.08).

422 **Table 3** Pre-stimulation and post-stimulation LFPQ scores ( $n = 21$ ).

		Explicit Liking (mm)		Explicit Wanting (mm)		Implicit Wanting (AU)	
		Pre-tDCS	Post-tDCS	Pre-tDCS	Post-tDCS	Pre-tDCS	Post-tDCS
Active	HFSA	52.9 ± 23.9	64.7 ± 25.1	49.9 ± 23.8	59.9 ± 25.7	-0.7 ± 31.0	15.7 ± 47.7
	LFSA	54.0 ± 20.9	53.7 ± 22.0	53.3 ± 22.1	53.6 ± 19.3	-1.4 ± 24.9	-15.1 ± 38.7
	HFSW	48.2 ± 24.6	54.6 ± 23.0	48.0 ± 24.2	48.8 ± 24.4	-6.5 ± 27.9	-3.6 ± 26.6
	LFSW	60.2 ± 20.8	60.5 ± 21.8	60.0 ± 19.5	57.1 ± 19.5	10.8 ± 28.7	3.8 ± 26.5
	FAB	<b>-6.6 ± 24.1*</b>	<b>2.5 ± 21.4*</b>	-7.8 ± 25.0	-1.0 ± 23.3	-7.1 ± 45.3	12.1 ± 53.7
	TAB	0.7 ± 18.2	-1.6 ± 20.0	2.4 ± 15.5	-3.9 ± 19.3	3.8 ± 20.2	-9.5 ± 39.1
Sham	HFSA	53.8 ± 26.6	57.2 ± 25.2	51.8 ± 27.7	55.1 ± 25.8	9.6 ± 33.1	14.4 ± 28.5
	LFSA	49.7 ± 18.4	49.8 ± 18.1	49.7 ± 20.1	49.8 ± 18.2	-2.3 ± 25.2	-3.6 ± 23.5
	HFSW	49.0 ± 27.6	47.1 ± 26.4	42.6 ± 28.0	45.9 ± 26.1	-9.5 ± 29.5	-6.9 ± 30.9
	LFSW	57.4 ± 22.3	55.4 ± 19.0	56.8 ± 20.1	53.5 ± 20.5	5.5 ± 30.0	-0.6 ± 29.4
	FAB	<b>-2.1 ± 26.3*</b>	<b>-0.4 ± 23.2*</b>	-6.1 ± 29.6	-1.1 ± 23.8	0.2 ± 45.0	7.6 ± 41.9
	TAB	1.4 ± 18.9	-2.3 ± 12.9	-1.0 ± 16.3	-2.8 ± 11.6	-5.3 ± 29.4	-8.7 ± 25.6

Mean ± SD. HFSA, high-fat savoury; LFSA, low-fat savoury; HFSW, high-fat sweet; LFSW, low-fat sweet; FAB, fat appeal bias; TAB, taste appeal bias. \* Indicates significant difference between active and sham conditions ( $p < 0.05$ ).

424

425 **3.4 Test-Retest Analysis**

426 With the exception of desire to eat ( $r = 0.382$ ,  $p = 0.088$ ), all variables were significantly  
427 correlated between baseline assessment. Twelve of the 23 variables assessed (across  
428 measurement instruments) displayed a strong correlation ( $r = >0.7$ ), with LFPQ implicit  
429 wanting and FAB appearing most consistent. Some baseline measures, particularly FCQ-S  
430 and appetite VAS measures, indicated only moderate reliability ( $r = 0.5$  to  $0.7$ ; Table 4).

431

432 **Table 4** Correlations between baseline (pre-tDCS) measures during visits 2 and 3 ( $n = 21$ ).

			<i>r</i>	<i>p</i>
Food Craving Questionnaire-State			0.549	0.010
Appetite VAS	Hunger		0.654	0.001
	Fullness		0.588	0.005
	Prospective Consumption		0.841	<0.001
	Desire to Eat		0.382	0.088
LFPQ	Implicit Wanting	HFSA	0.837	<0.001
		LFSA	0.795	<0.001
		HFSW	0.882	<0.001
		LFSW	0.718	0.001
	Explicit Liking	HFSA	0.652	0.002
		LFSA	0.664	0.002
		HFSW	0.781	<0.001
		LFSW	0.784	<0.001
	Explicit Wanting	HFSA	0.698	0.001
		LFSA	0.751	<0.001
		HFSW	0.712	0.001
		LFSW	0.668	0.002
	Fat Appeal Bias	Explicit Liking	0.853	<0.001
		Explicit Wanting	0.887	<0.001
		Implicit Wanting	0.677	0.001
	Taste Appeal Bias	Explicit Liking	0.536	0.018
		Explicit Wanting	0.555	0.014
		Implicit Wanting	0.737	<0.001

HFSA, high-fat savoury; HFSW, high-fat sweet; LFPQ, Leeds Food Preference

Questionnaire; LFSA, low-fat savoury; LFSW, low-fat sweet; VAS, visual analogue scale.



433

### 434 **3.5 Responses to tDCS**

435 Successful delivery of the electric current occurred across all 42 stimulation sessions, with  
436 mean impedance levels of  $8 \pm 4 \text{ k}\Omega$  at the start of stimulation which reduced to  $3 \pm 2 \text{ k}\Omega$   
437 within the initial five minutes of current delivery. Stimulation was well-tolerated by  
438 participants with only common adverse events, particularly cutaneous sensations,  
439 experienced during tDCS. The most common sensations reported were tingling, itching,  
440 sleepiness and a burning sensation (Table 5). Tingling ( $p = 0.016$ ), itching ( $p = 0.021$ ) and  
441 sleepiness ( $p = 0.021$ ) were reported by significantly more participants following active  
442 tDCS, when compared with sham stimulation. No other sensations were significantly  
443 different between conditions. Despite experiencing more minor adverse events, participants  
444 were unable to identify the active tDCS session above the level of chance, with only 38%  
445 (8/21) of participants able to successfully distinguish between conditions.

446

**Table 5** Frequency of adverse events experienced immediately post-stimulation.

	Active	Sham	p
Headache	7 (33%)	4 (19%)	0.186
Neck pain	0 (0%)	0 (0%)	-
Scalp pain	3 (14%)	1 (5%)	0.329
Tingling	14 (67%)	7 (33%)	<b>0.016*</b>
Itching	11 (52%)	6 (29%)	<b>0.021*</b>
Burning sensation	9 (43%)	2 (10%)	0.267
Skin redness	5 (24%)	2 (10%)	0.186
Sleepiness	12 (57%)	7 (33%)	<b>0.021*</b>
Trouble concentrating	5 (24%)	3 (14%)	0.329
Acute mood change	2 (10%)	2 (10%)	1.000

\* Indicates significant difference between active and sham conditions.

#### 4. DISCUSSION

The current study examined the effect of induced hyper-activity of the right DLPFC through tDCS on food craving, reward and subjective appetite measures in a healthy-weight cohort. It is important to note that the sample used in the current study appeared to show low susceptibility to hedonic-driven overconsumption, evidenced by their scores on several measures of eating behaviour traits linked to overconsumption. We also sought to examine the reliability of measures prior to tDCS with a view to establishing the viability of their future use in our research. We report strong relationships between key variables, particularly implicit wanting and FAB, when preparatory procedures prior to tDCS were standardised. These variables may prove to be sensitive targets for detecting significant effects in future eating behaviour-focussed tDCS research. Other variables, particularly food craving measures, proved less stable and may require tighter experimental control or larger sample sizes to reveal differences. Collectively our findings are novel to tDCS research.

462

463 Prior work has mainly focussed on measuring food craving and in-laboratory consumption  
464 with equivocal findings (Fregni, et al., 2008; Georgii, et al., 2017; Goldman, et al., 2011;  
465 Sedgmond, et al., 2019). The present study is favourable by comparison in sample size,  
466 study design (i.e. sham-controlled and double-blind) and stimulation parameters (Burgess, et  
467 al., 2016; Fregni, et al., 2008; Goldman, et al., 2011; Ray, et al., 2017). Recently published  
468 meta-analyses have cast doubt in the ability of tDCS to alter measures of food craving (Hall  
469 & Lowe, 2018; Lowe, Vincent, & Hall, 2017), which may be due to the poor test-retest  
470 reliability of food craving measures (Cepeda-Benito, et al., 2000; Meule, Teran, et al., 2014).  
471 This is in agreement with our data which highlighted only moderate reliability of baseline  
472 FCQ-S scores. In comparison, our data show strong relationships between baseline  
473 measures of implicit and explicit reward. In developing the LFPQ, Dalton and Finlayson  
474 (2014) reported a reliability coefficient of 0.6 – 0.7 for implicit wanting and 0.8 – 0.9 for  
475 explicit liking measures, with our data supporting this moderate-to-strong reliability. The  
476 LFPQ has been utilised in several settings, and is considered a sensitive measure for  
477 individual eating behaviour traits (Dalton, et al., 2013a; Dalton, Blundell, & Finlayson, 2013b;  
478 Finlayson, Arlotti, Dalton, King, & Blundell, 2011), and a good predictor of in-laboratory and  
479 real-world food choice and consumption (French, Mitchell, Finlayson, Blundell, & Jeffery,  
480 2014; Griffioen-Roose, Mars, Finlayson, Blundell, & de Graaf, 2011). The present study is  
481 the first to extend the use of the LFPQ to include tDCS procedures, and the reliability of this  
482 questionnaire suggests it is a robust measure and should be explored in future research.

483

484 It is logical that the significant interaction between tDCS condition and time point for FAB  
485 explicit liking was removed when controlling for baseline hunger as the excitatory effects of  
486 anodal tDCS are not associated with increased preference for high-fat foods (Goldman, et  
487 al., 2011; Jauch-Chara, et al., 2014). In addition, healthy individuals are likely to have a  
488 normative response to food stimuli and are able to sufficiently integrate rewarding signals  
489 with other appetitive signals to select appropriate eating behaviours (see Alonso-Alonso and

Pascual-Leone (2007) for review). Healthy individuals are also unlikely to have structural deficits observed in obesity and binge eating, which are associated with alteration in reward response (Balodis, Grilo, & Potenza, 2015; Lowe, Reichelt, & Hall, 2019). It is probable that stimulation would have no additive effects in healthy individuals (Burgess, et al., 2016). The greater baseline hunger score likely heightened the rewarding value of high-calorie foods, particularly those high in fat, that participants were exposed to during the computer-based task (Cameron, Goldfield, Finlayson, Blundell, & Doucet, 2014; Finlayson, King, & Blundell, 2008; Mehta, et al., 2012).

In addition to the equivocal findings for food craving and consumption, previous work has been inconsistent in the recruitment of participants and some of the variation in results may be due to participants' eating behaviour traits. Two previous studies have directly linked tDCS effects as occurring in those with abnormal eating behaviours (Burgess, et al., 2016; Ray, et al., 2017), and when comparing further studies that utilise similar tDCS parameters (i.e. 2 mA for 20 minutes over the right DLPFC), a trait-dependent effect is evident. Studies that recruited participants with frequent food cravings found a consistent reduction in measures of state food craving (Fregni, et al., 2008; Goldman, et al., 2011; Kekic, et al., 2014; Lapenta, et al., 2014). In comparison, studies that did not measure behaviour traits or report comparable traits between healthy and overweight populations, fail to find a significant effect of stimulation on craving (Bravo, et al., 2016; Sedgmond, et al., 2019). Our data supports the robustness of healthy eating behaviours against perturbation in cortical activity within the DLPFC, which is assumed to occur in populations that are obese or with BED (Boeka & Lokken, 2011; Karhunen, et al., 2000). We therefore speculate that there is a diminishing return for attempting to increase neuronal activity within the DLPFC when participants are already able to control their eating behaviours. Hyper-activity in this cortical region may have a ceiling effect beyond which no further improvement is seen. This may account for the null effect we found for food craving, reward and appetite following tDCS,

and can be supported by the moderate-to-strong evidence in favour of the null hypothesis as highlighted by our Bayesian statistical approach.

The present study is not without limitation. It is understood that males and females experience different eating behaviours, and may express differing behavioural traits (Rolls, Fedoroff, & Guthrie, 1991). The present study recruited both male and female participants, which may have influenced the effects of tDCS, and provided an additional source of variation across data. However, this is not without precedent as prior studies that recruited male and female participants have shown an experimental effect (Burgess, et al., 2016; Carvalho, et al., 2019; Goldman, et al., 2011). Given the novelty of using the LFPQ it was important to consider the wider effects of tDCS on this variable before focussing on specific sex. Second, our original hypotheses did not consider the impact of eating behaviour traits and as such these were not controlled for during screening. Our inclusion criteria focussed on weight status, but the participants recruited displayed behaviour traits that do not suggest susceptibility to overconsumption, as discussed above; notably, all participants scored below the 50-point cut-off for trait food craving. Third, prior studies have induced hyperactivity in the DLPFC through bilateral and unilateral stimulation of the cortex (Carvalho, et al., 2019; Fassini, et al., 2019; Lapenta, et al., 2014; Ljubisavljevic, et al., 2016). Although these montages have been shown to improve measures of hedonic appetite (Fregni, et al., 2008; Goldman, et al., 2011), the efficacy of such electrode placement has been debated due to the simultaneous effects of anodal and cathodal stimulation on the same cortical region (Bestmann, et al., 2015). The inhibitory effects associated with cathodal stimulation during traditional montages may also impact hedonic appetite measures, as the left DLPFC is implicated in dietary control and food choice behaviour (Higuera-Hernández, et al., 2018). Similar to the right DLPFC, there is some support for reduced activity in the left DLPFC in response to food, when comparing individuals who are lean with those who are obese (Le, et al., 2006; Le, et al., 2007). In the present study, a prefrontal-occipital montage was used, utilising a similar montage seen in previous work (Marron, et al., 2018; Vitor-Costa, et al.,

2015). The ability of this montage to induce hyperactivity in the DLPFC has been confirmed in several recent computational models (Marron, et al., 2018; Zheng, et al., 2016; Zheng, et al., 2017). Moreover, we verified that the electric current was being delivered in a consistent manner across all 42 stimulation sessions by checking impedance. Finally, the effectiveness of common sham procedures as a blinding technique has been debated due to significantly greater sensations often reported following active tDCS (Horvath, 2015). Indeed, in the present study participants reported significantly greater itching, tingling and sleepiness following active stimulation. However, the inability of participants to identify the active protocol beyond the level of chance, despite these heightened sensations, provides further support for the use of standardised sham protocols as a blinding technique in tDCS research (Ambrus, et al., 2012).

## **5. CONCLUSION**

Our study is the first to report the effects of tDCS on components of food reward in sample of healthy individuals with no susceptibility to overconsume, and we show no significant change in these measures. Prior to examining these effects, we established an indication of data reliability and revealed some plausible targets for future effects through tDCS exposure. In the present sample these effects were transient for the most part and, in line with the work by Burgess and colleagues (Burgess, et al., 2016; Ray, et al., 2017), this highlights a behaviour trait-dependent effect of stimulation. Future work should focus on populations who are at risk of reward-driven overconsumption and weight gain, such as those showing recurrent binge eating behaviours, as these individuals may be responsive to the effects of tDCS on hedonic appetite.

## **ACKNOWLEDGEMENTS**

The authors wish to thank Elizabeth Senior and Lee Newburn for their help with recruitment and data collection, and Dr Jo Corbett of the University of Portsmouth for advising on protocol development.

573

574 **AUTHOR CONTRIBUTIONS**

575 **Jordan Beaumont:** Conceptualisation, Methodology, Validation, Formal analysis,

576 Investigation, Data curation, Writing – original draft, Writing – review and editing,

577 Visualisation, Project administration.

578 **Danielle Davis:** Conceptualisation, Methodology, Resources, Writing – review and editing,

579 Supervision.

580 **Michelle Dalton:** Conceptualisation, Methodology, Software, Formal analysis, Resources,

581 Data curation, Writing – review and editing.

582 **Alexander Nowicky:** Methodology, Writing – review and editing.

583 **Mark Russell:** Methodology, Writing – review and editing.

584 **Martin Barwood:** Conceptualisation, Methodology, Validation, Resources, Writing – original

585 draft, Writing – review and editing, Supervision.

586

587 **DECLARATION OF INTEREST**

588 None.

589

590 **FUNDING**

591 This research did not receive any specific grant from funding agencies in the public,

592 commercial or not-for-profit sectors.

593

## REFERENCES

- Alonso-Alonso, M. (2013). Translating tDCS into the field of obesity: mechanism-driven approaches. *Frontiers in Human Neuroscience*, 7, 512.
- Alonso-Alonso, M., & Pascual-Leone, A. (2007). The Right Brain Hypothesis for Obesity. *JAMA*, 297, 1819-1822.
- Alonzo, A., Brassil, J., Taylor, J. L., Martin, D., & Loo, C. K. (2012). Daily transcranial direct current stimulation (tDCS) leads to greater increases in cortical excitability than second daily transcranial direct current stimulation. *Brain Stimulation*, 5, 208-213.
- Ambrus, G. G., Al-Moyed, H., Chaieb, L., Sarp, L., Antal, A., & Paulus, W. (2012). The fade-in – Short stimulation – Fade out approach to sham tDCS – Reliable at 1 mA for naïve and experienced subjects, but not investigators. *Brain Stimulation*, 5, 499-504.
- Andresen, E. M., Malmgren, J. A., Carter, W. B., & Patrick, D. L. (1994). Screening for depression in well older adults: Evaluation of a short form of the CES-D. *American Journal of Preventive Medicine*, 10, 77-84.
- Anglé, S., Engblom, J., Eriksson, T., Kautiainen, S., Saha, M.-T., Lindfors, P., Lehtinen, M., & Rimpelä, A. (2009). Three factor eating questionnaire-R18 as a measure of cognitive restraint, uncontrolled eating and emotional eating in a sample of young Finnish females. *International Journal of Behavioral Nutrition and Physical Activity*, 6, 41.
- Balodis, I. M., Grilo, C. M., & Potenza, M. N. (2015). Neurobiological features of binge eating disorder. *CNS spectrums*, 20, 557-565.
- Beechy, L., Galpern, J., Petrone, A., & Das, S. K. (2012). Assessment tools in obesity — Psychological measures, diet, activity, and body composition. *Physiology & Behavior*, 107, 154-171.
- Bestmann, S., de Berker, A. O., & Bonaiuto, J. (2015). Understanding the behavioural consequences of noninvasive brain stimulation. *Trends in Cognitive Sciences*, 19, 13-20.
- Blundell, J., De Graaf, C., Hulshof, T., Jebb, S., Livingstone, B., Lluch, A., Mela, D., Salah, S., Schuring, E., Van Der Knaap, H., & Westerterp, M. (2010). Appetite control: methodological aspects of the evaluation of foods. *Obesity Reviews*, 11, 251-270.
- Boeka, A. G., & Lokken, K. L. (2011). Prefrontal systems involvement in binge eating. *Eating and Weight Disorders - Studies on Anorexia, Bulimia and Obesity*, 16, e121-e126.
- Boswell, R. G., & Kober, H. (2016). Food cue reactivity and craving predict eating and weight gain: a meta-analytic review. *Obesity Reviews*, 17, 159-177.
- Bravo, G. L., Poje, A. B., Perissinotti, I., Marcondes, B. F., Villamar, M. F., Manzardo, A. M., Luque, L., LePage, J. F., Stafford, D., Fregni, F., & Butler, M. G. (2016). Transcranial direct current stimulation reduces food-craving and measures of hyperphagia



631 behavior in participants with Prader-Willi syndrome. *American Journal of Medical*  
632 *Genetics Part B: Neuropsychiatric Genetics*, 171, 266-275.

633 Brunoni, A. R., Amadera, J., Berbel, B., Volz, M. S., Rizzerio, B. G., & Fregni, F. (2011). A  
634 systematic review on reporting and assessment of adverse effects associated with  
635 transcranial direct current stimulation. *International Journal of*  
636 *Neuropsychopharmacology*, 14, 1133-1145.

637 Burgess, E. E., Sylvester, M. D., Morse, K. E., Amthor, F. R., Mrug, S., Lokken, K. L.,  
638 Osborn, M. K., Soleymani, T., & Boggiano, M. M. (2016). Effects of transcranial direct  
639 current stimulation (tDCS) on binge-eating disorder. *International Journal of Eating*  
640 *Disorders*, 49, 930-936.

641 Cameron, J. D., Goldfield, G. S., Finlayson, G., Blundell, J. E., & Doucet, É. (2014). Fasting  
642 for 24 Hours Heightens Reward from Food and Food-Related Cues. *PLOS ONE*, 9,  
643 e85970.

644 Carvalho, S., Sampaio, A., Mendes, A. J., Lema, A., Vieira, D., Gonçalves, Ó. F., & Leite, J.  
645 (2019). Polarity Specific Effects of Cross-Hemispheric tDCS Coupled With Approach-  
646 Avoidance Training on Chocolate Craving. *Frontiers in Pharmacology*, 9, 1500.

647 Cepeda-Benito, A., Gleaves, D. H., Williams, T. L., & Erath, S. A. (2000). The development  
648 and validation of the state and trait food-cravings questionnaires. *Behavior Therapy*,  
649 31, 151-173.

650 Dalton, M., Blundell, J., & Finlayson, G. (2013a). Effect of BMI and Binge Eating on Food  
651 Reward and Energy Intake: Further Evidence for a Binge Eating Subtype of Obesity.  
652 *Obesity Facts*, 6, 348-359.

653 Dalton, M., Blundell, J., & Finlayson, G. (2013b). Examination of food reward and energy  
654 intake under laboratory and free-living conditions in a trait binge eating subtype of  
655 obesity. *Frontiers in Psychology*, 4, 757.

656 Dalton, M., & Finlayson, G. (2014). Psychobiological examination of liking and wanting for fat  
657 and sweet taste in trait binge eating females. *Physiology & Behavior*, 136, 128-134.

658 Dalton, M., Finlayson, G., Hill, A., & Blundell, J. (2015). Preliminary validation and principal  
659 components analysis of the Control of Eating Questionnaire (CoEQ) for the  
660 experience of food craving. *European Journal Of Clinical Nutrition*, 69, 1313.

661 Dalton, M., Finlayson, G., Walsh, B., Halseth, A. E., Duarte, C., & Blundell, J. E. (2017).  
662 Early improvement in food cravings are associated with long-term weight loss  
663 success in a large clinical sample. *International Journal Of Obesity*, 41, 1232-1236.

664 DaSilva, A. F., Volz, M. S., Bikson, M., & Fregni, F. (2011). *Electrode Positioning and*  
665 *Montage in Transcranial Direct Current Stimulation*.  
666 <[https://www.jove.com/video/2744/electrode-positioning-montage-transcranial-direct-](https://www.jove.com/video/2744/electrode-positioning-montage-transcranial-direct-current)  
667 [current](https://www.jove.com/video/2744/electrode-positioning-montage-transcranial-direct-current)> Accessed 09 June 2020.

668 Davis, C. (2013). Compulsive Overeating as an Addictive Behavior: Overlap Between Food  
669 Addiction and Binge Eating Disorder. *Current Obesity Reports*, 2, 171-178.

670 Davis, C., Levitan, R. D., Reid, C., Carter, J. C., Kaplan, A. S., Patte, K. A., King, N., Curtis,  
671 C., & Kennedy, J. L. (2009). Dopamine for "Wanting" and Opioids for "Liking": A  
672 Comparison of Obese Adults With and Without Binge Eating. *Obesity*, 17, 1220-  
673 1225.

674 De Lauzon-Guillain, B., Romon, M., Musher-Eizenman, D., Heude, B., Basdevant, A.,  
675 Charles, M. A., & Fleurbaix-Laventie Ville Santé Study, G. (2009). Cognitive restraint,  
676 uncontrolled eating and emotional eating: correlations between parent and  
677 adolescent. *Maternal & Child Nutrition*, 5, 171-178.

678 Fassini, P. G., Das, S. K., Suen, V. M. M., Magerowski, G., Marchini, J. S., da Silva Junior,  
679 W. A., Changyu, S., & Alonso-Alonso, M. (2019). Appetite effects of prefrontal  
680 stimulation depend on COMT Val158Met polymorphism: A randomized clinical trial.  
681 *Appetite*, 140, 142-150.

682 Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G\*Power 3: A flexible statistical  
683 power analysis program for the social, behavioral, and biomedical sciences. *Behavior*  
684 *Research Methods*, 39, 175-191.

685 Filmer, H. L., Dux, P. E., & Mattingley, J. B. (2014). Applications of transcranial direct current  
686 stimulation for understanding brain function. *Trends in Neurosciences*, 37, 742-753.

687 Finlayson, G., Arlotti, A., Dalton, M., King, N., & Blundell, J. E. (2011). Implicit wanting and  
688 explicit liking are markers for trait binge eating. A susceptible phenotype for  
689 overeating. *Appetite*, 57, 722-728.

690 Finlayson, G., & Dalton, M. (2012). Hedonics of Food Consumption: Are Food 'Liking' and  
691 'Wanting' Viable Targets for Appetite Control in the Obese? *Current Obesity Reports*,  
692 1, 42-49.

693 Finlayson, G., King, N., & Blundell, J. (2008). The role of implicit wanting in relation to explicit  
694 liking and wanting for food: Implications for appetite control. *Appetite*, 50, 120-127.

695 Finlayson, G., King, N., & Blundell, J. E. (2007). Is it possible to dissociate 'liking' and  
696 'wanting' for foods in humans? A novel experimental procedure. *Physiology &*  
697 *Behavior*, 90, 36-42.

698 Fleurbaix Laventie Ville Sante Study, G. (2004). The Three-Factor Eating Questionnaire-R18  
699 Is Able to Distinguish among Different Eating Patterns in a General Population. *The*  
700 *Journal of Nutrition*, 134, 2372-2380.

701 Fregni, F., Orsati, F., Pedrosa, W., Fecteau, S., Tome, F. A., Nitsche, M. A., Mecca, T.,  
702 Macedo, E. C., Pascual-Leone, A., & Boggio, P. S. (2008). Transcranial direct current  
703 stimulation of the prefrontal cortex modulates the desire for specific foods. *Appetite*,  
704 51, 34-41.

- 705 French, S. A., Mitchell, N. R., Finlayson, G., Blundell, J. E., & Jeffery, R. W. (2014).  
706 Questionnaire and laboratory measures of eating behavior. Associations with energy  
707 intake and BMI in a community sample of working adults. *Appetite*, 72, 50-58.
- 708 Galetta, S. L. (2017). Occipital Lobe☆. In *Reference Module in Neuroscience and*  
709 *Biobehavioral Psychology*. Elsevier.
- 710 Gandiga, P. C., Hummel, F. C., & Cohen, L. G. (2006). Transcranial DC stimulation (tDCS):  
711 A tool for double-blind sham-controlled clinical studies in brain stimulation. *Clinical*  
712 *Neurophysiology*, 117, 845-850.
- 713 Georgii, C., Goldhofer, P., Meule, A., Richard, A., & Blechert, J. (2017). Food craving, food  
714 choice and consumption: The role of impulsivity and sham-controlled tDCS  
715 stimulation of the right dlPFC. *Physiology & Behavior*, 177, 20-26.
- 716 Gluck, M. E., Alonso-Alonso, M., Piaggi, P., Weise, C. M., Jumpertz-von Schwartzberg,  
717 R., Reinhardt, M., Wassermann, E. M., Venti, C. A., Votruba, S. B., & Krakoff, J.  
718 (2015). Neuromodulation targeted to the prefrontal cortex induces changes in energy  
719 intake and weight loss in obesity. *Obesity (Silver Spring)*, 23, 2149-2156.
- 720 Gluck, M. E., Viswanath, P., & Stinson, E. J. (2017). Obesity, Appetite, and the Prefrontal  
721 Cortex. *Curr Obes Rep*, 6, 380-388.
- 722 Goldman, R. L., Borckardt, J. J., Frohman, H. A., O'Neil, P. M., Madan, A., Campbell, L. K.,  
723 Budak, A., & George, M. S. (2011). Prefrontal cortex transcranial direct current  
724 stimulation (tDCS) temporarily reduces food cravings and increases the self-reported  
725 ability to resist food in adults with frequent food craving. *Appetite*, 56, 741-746.
- 726 Griffioen-Roose, S., Mars, M., Finlayson, G., Blundell, J. E., & de Graaf, C. (2011). The  
727 effect of within-meal protein content and taste on subsequent food choice and  
728 satiety. *British Journal of Nutrition*, 106, 779-788.
- 729 Hall, P. A., & Lowe, C. J. (2018). Cravings, currents and cadavers: What is the magnitude of  
730 tDCS effects on food craving outcomes? *Nutritional Neuroscience*, 1-4.
- 731 Hall, P. A., Vincent, C. M., & Burhan, A. M. (2018). Non-invasive brain stimulation for food  
732 cravings, consumption, and disorders of eating: A review of methods, findings and  
733 controversies. *Appetite*, 124, 78-88.
- 734 Higuera-Hernández, M. F., Reyes-Cuapio, E., Gutiérrez-Mendoza, M., Rocha, N. B., Veras,  
735 A. B., Budde, H., Jesse, J., Zaldívar-Rae, J., Blanco-Centurión, C., Machado, S., &  
736 Murillo-Rodríguez, E. (2018). Fighting obesity: Non-pharmacological interventions.  
737 *Clinical Nutrition ESPEN*, 25, 50-55.
- 738 Hill, J. O. (2006). Understanding and Addressing the Epidemic of Obesity: An Energy  
739 Balance Perspective. *Endocrine Reviews*, 27, 750-761.
- 740 Horvath, J. C. (2015). Are current blinding methods for transcranial direct current stimulation  
741 (tDCS) effective in healthy populations? *Clinical Neurophysiology*, 126, 2045-2046.

- 742 Jamil, A., & Nitsche, M. A. (2017). What Effect Does tDCS Have on the Brain? Basic  
743 Physiology of tDCS. *Current Behavioral Neuroscience Reports*, 4, 331-340.
- 744 Jauch-Chara, K., Kistenmacher, A., Herzog, N., Schwarz, M., Schweiger, U., & Oltmanns, K.  
745 M. (2014). Repetitive electric brain stimulation reduces food intake in humans. *The*  
746 *American Journal of Clinical Nutrition*, 100, 1003-1009.
- 747 Joseph, R. J., Alonso-Alonso, M., Bond, D. S., Pascual-Leone, A., & Blackburn, G. L. (2011).  
748 The neurocognitive connection between physical activity and eating behaviour.  
749 *Obesity Reviews*, 12, 800-812.
- 750 Karhunen, L. J., Vanninen, E. J., Kuikka, J. T., Lappalainen, R. I., Tiihonen, J., & Uusitupa,  
751 M. I. J. (2000). Regional cerebral blood flow during exposure to food in obese binge  
752 eating women. *Psychiatry Research: Neuroimaging*, 99, 29-42.
- 753 Karlsson, J., Persson, L. O., Sjöström, L., & Sullivan, M. (2000). Psychometric properties  
754 and factor structure of the Three-Factor Eating Questionnaire (TFEQ) in obese men  
755 and women. Results from the Swedish Obese Subjects (SOS) study. *International*  
756 *Journal Of Obesity*, 24, 1715-1725.
- 757 Kekic, M., McClelland, J., Campbell, I., Nestler, S., Rubia, K., David, A. S., & Schmidt, U.  
758 (2014). The effects of prefrontal cortex transcranial direct current stimulation (tDCS)  
759 on food craving and temporal discounting in women with frequent food cravings.  
760 *Appetite*, 78, 55-62.
- 761 Kessler, R. C., Berglund, P. A., Chiu, W. T., Deitz, A. C., Hudson, J. I., Shahly, V., Aguilar-  
762 Gaxiola, S., Alonso, J., Angermeyer, M. C., Benjet, C., Bruffaerts, R., de Girolamo,  
763 G., de Graaf, R., Maria Haro, J., Kovess-Masfety, V., O'Neill, S., Posada-Villa, J.,  
764 Sasu, C., Scott, K., Viana, M. C., & Xavier, M. (2013). The Prevalence and  
765 Correlates of Binge Eating Disorder in the World Health Organization World Mental  
766 Health Surveys. *Biological Psychiatry*, 73, 904-914.
- 767 Klem, G. H., Lüders, H. O., Jasper, H. H., & Elger, C. (1999). The ten-twenty electrode  
768 system of the International Federation. The International Federation of Clinical  
769 Neurophysiology. *Electroencephalography and clinical neurophysiology. Supplement*,  
770 52, 3-6.
- 771 Kober, H., & Boswell, R. G. (2018). Potential psychological & neural mechanisms in binge  
772 eating disorder: Implications for treatment. *Clinical Psychology Review*, 60, 32-44.
- 773 Lapenta, O. M., Sierve, K. D., de Macedo, E. C., Fregni, F., & Boggio, P. S. (2014).  
774 Transcranial direct current stimulation modulates ERP-indexed inhibitory control and  
775 reduces food consumption. *Appetite*, 83, 42-48.
- 776 Le, D. S. N. T., Pannacciulli, N., Chen, K., Del Parigi, A., Salbe, A. D., Reiman, E. M., &  
777 Krakoff, J. (2006). Less activation of the left dorsolateral prefrontal cortex in response  
778 to a meal: a feature of obesity. *The American Journal of Clinical Nutrition*, 84, 725-  
779 731.

- 780 Le, D. S. N. T., Pannacciulli, N., Chen, K., Salbe, A. D., Hill, J. O., Wing, R. R., Reiman, E.  
781 M., & Krakoff, J. (2007). Less activation in the left dorsolateral prefrontal cortex in the  
782 reanalysis of the response to a meal in obese than in lean women and its association  
783 with successful weight loss. *The American Journal of Clinical Nutrition*, 86, 573-579.
- 784 Lee, M., & Wagenmakers, E. (2013). *Bayesian Cognitive Modeling: A Practical Course*.  
785 United Kingdom: Cambridge University Press.
- 786 Ljubisavljevic, M., Maxood, K., Bjekic, J., Oommen, J., & Nagelkerke, N. (2016). Long-Term  
787 Effects of Repeated Prefrontal Cortex Transcranial Direct Current Stimulation (tDCS)  
788 on Food Craving in Normal and Overweight Young Adults. *Brain Stimulation*, 9, 826-  
789 833.
- 790 Lowe, C. J., Reichelt, A. C., & Hall, P. A. (2019). The Prefrontal Cortex and Obesity: A  
791 Health Neuroscience Perspective. *Trends in Cognitive Sciences*, 23, 349-361.
- 792 Lowe, C. J., Vincent, C., & Hall, P. A. (2017). Effects of Noninvasive Brain Stimulation on  
793 Food Cravings and Consumption: A Meta-Analytic Review. *Psychosomatic Medicine*,  
794 79.
- 795 Marron, E. M., Viejo-Sobera, R., Cuatrecasas, G., Redolar-Ripoll, D., Lorda, P. G., Datta, A.,  
796 Bikson, M., Magerowski, G., & Alonso-Alonso, M. (2018). Prefronto-cerebellar  
797 neuromodulation affects appetite in obesity. *International Journal Of Obesity*.
- 798 Mehta, S., Melhorn, S. J., Smeraglio, A., Tyagi, V., Grabowski, T., Schwartz, M. W., & Schur,  
799 E. A. (2012). Regional brain response to visual food cues is a marker of satiety that  
800 predicts food choice. *The American Journal of Clinical Nutrition*, 96, 989-999.
- 801 Meule, A. (2018). Food cravings in food addiction: exploring a potential cut-off value of the  
802 Food Cravings Questionnaire-Trait-reduced. *Eating and Weight Disorders - Studies*  
803 *on Anorexia, Bulimia and Obesity*, 23, 39-43.
- 804 Meule, A., Hermann, T., & Kübler, A. (2014). A short version of the Food Cravings  
805 Questionnaire—Trait: the FCQ-T-reduced. *Frontiers in Psychology*, 5, 190.
- 806 Meule, A., Teran, C. B., Berker, J., Gründel, T., Mayerhofer, M., & Platte, P. (2014). On the  
807 differentiation between trait and state food craving: Half-year retest-reliability of the  
808 Food Cravings Questionnaire-Trait-reduced (FCQ-T-r) and the Food Cravings  
809 Questionnaire-State(FCQ-S). *Journal of Eating Disorders*, 2, 25.
- 810 Nikolin, S., Huggins, C., Martin, D., Alonzo, A., & Loo, C. K. (2018). Safety of repeated  
811 sessions of transcranial direct current stimulation: A systematic review. *Brain*  
812 *Stimulation*, 11, 278-288.
- 813 Nitsche, M. A., & Paulus, W. (2000). Excitability changes induced in the human motor cortex  
814 by weak transcranial direct current stimulation. *The Journal of Physiology*, 527, 633-  
815 639.
- 816 Nitsche, M. A., & Paulus, W. (2001). Sustained excitability elevations induced by transcranial  
817 DC motor cortex stimulation in humans. *Neurology*, 57, 1899.

- 818 Oustric, P., Thivel, D., Dalton, M., Beaulieu, K., Gibbons, C., Hopkins, M., Blundell, J., &  
819 Finlayson, G. (2020). Measuring food preference and reward: Application and cross-  
820 cultural adaptation of the Leeds Food Preference Questionnaire in human  
821 experimental research. *Food Quality and Preference*, 80, 103824.
- 822 Radloff, L. S. (1977). The CES-D Scale: A Self-Report Depression Scale for Research in the  
823 General Population. *Applied Psychological Measurement*, 1, 385-401.
- 824 Ray, M. K., Sylvester, M. D., Osborn, L., Helms, J., Turan, B., Burgess, E. E., & Boggiano,  
825 M. M. (2017). The critical role of cognitive-based trait differences in transcranial direct  
826 current stimulation (tDCS) suppression of food craving and eating in frank obesity.  
827 *Appetite*, 116, 568-574.
- 828 Rolls, B. J., Fedoroff, I. C., & Guthrie, J. F. (1991). Gender differences in eating behavior  
829 and body weight regulation. *Health Psychology*, 10, 133-142.
- 830 Rush, S., & Driscoll, D. A. (1968). Current Distribution in the Brain From Surface Electrodes.  
831 *Anesthesia & Analgesia*, 47.
- 832 Sedgmond, J., Lawrence Natalia, S., Verbruggen, F., Morrison, S., Chambers Christopher,  
833 D., & Adams Rachel, C. (2019). Prefrontal brain stimulation during food-related  
834 inhibition training: effects on food craving, food consumption and inhibitory control.  
835 *Royal Society Open Science*, 6, 181186.
- 836 Stagg, C. J., Antal, A., & Nitsche, M. A. (2018). Physiology of Transcranial Direct Current  
837 Stimulation. *The Journal of ECT*, 34.
- 838 Thair, H., Holloway, A. L., Newport, R., & Smith, A. D. (2017). Transcranial Direct Current  
839 Stimulation (tDCS): A Beginner's Guide for Design and Implementation. *Frontiers in*  
840 *Neuroscience*, 11, 641.
- 841 Tremblay, S., Lepage, J.-F., Latulipe-Loiselle, A., Fregni, F., Pascual-Leone, A., & Théoret,  
842 H. (2014). The Uncertain Outcome of Prefrontal tDCS. *Brain Stimulation*, 7, 773-783.
- 843 Vitor-Costa, M., Okuno, N. M., Bortolotti, H., Bertollo, M., Boggio, P. S., Fregni, F., &  
844 Altimari, L. R. (2015). Improving Cycling Performance: Transcranial Direct Current  
845 Stimulation Increases Time to Exhaustion in Cycling. *PLOS ONE*, 10, e0144916.
- 846 Wardle, R. A., Thapaliya, G., Nowak, A., Radford, S., Dalton, M., Finlayson, G., & Moran, G.  
847 W. (2018). An Examination of Appetite and Disordered Eating in Active Crohn's  
848 Disease. *Journal of Crohn's and Colitis*, 12, 819-825.
- 849 World Health Organisation. (2020). *Obesity and Overweight*. <[https://www.who.int/en/news-](https://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight)  
850 [room/fact-sheets/detail/obesity-and-overweight](https://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight)> Accessed 29 July 2020.
- 851 Zheng, H., Huang, D., Chen, S., Wang, S., Guo, W., Luo, J., Ye, H., & Chen, Y. (2016).  
852 Modulating the Activity of Ventromedial Prefrontal Cortex by Anodal tDCS Enhances  
853 the Trustee's Repayment through Altruism. *Frontiers in Psychology*, 7, 1437.

854 Zheng, H., Wang, S., Guo, W., Chen, S., Luo, J., Ye, H., & Huang, D. (2017). Enhancing the  
855 Activity of the DLPFC with tDCS Alters Risk Preference without Changing  
856 Interpersonal Trust. *Frontiers in Neuroscience*, 11, 52.  
857