

The Pentose Phosphate Pathway Dynamics in Cancer and Its Dependency on Intracellular pH

ALFAROUK, Khalid O <<http://orcid.org/0000-0001-8656-6117>>, AHMED, Samrein <<http://orcid.org/0000-0001-9773-645X>>, ELLIOTT, Robert L, BENOIT, Amanda, ALQAHTANI, Saad S, IBRAHIM, Muntaser E, BASHIR, Adil HH, ALHOUFIE, Sari TS, ELHASSAN, Gamal O, WALES, Christian C, SCHWARTZ, Laurent H, ALI, Heyam S <<http://orcid.org/0000-0002-4441-0525>>, AHMED, Ahmed, FORDE, Patrick F, DEVESA, Jesus <<http://orcid.org/0000-0002-4153-2543>>, CARDONE, Rosa A, FAIS, Stefano, HARGUINDEY, Salvador <<http://orcid.org/0000-0002-1725-757X>> and RESHKIN, Stephan J <<http://orcid.org/0000-0001-9757-5908>>

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/33082/>

This document is the Published Version [VoR]

Citation:






ALFAROUK, Khalid O, AHMED, Samrein, ELLIOTT, Robert L, BENOIT, Amanda, ALQAHTANI, Saad S, IBRAHIM, Muntaser E, BASHIR, Adil HH, ALHOUFIE, Sari TS, ELHASSAN, Gamal O, WALES, Christian C, SCHWARTZ, Laurent H, ALI, Heyam S, AHMED, Ahmed, FORDE, Patrick F, DEVESA, Jesus, CARDONE, Rosa A, FAIS, Stefano, HARGUINDEY, Salvador and RESHKIN, Stephan J (2020). The Pentose Phosphate Pathway Dynamics in Cancer and Its Dependency on Intracellular pH. *Metabolites*, 10 (7): 285. [Article]

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

Review

The Pentose Phosphate Pathway Dynamics in Cancer and Its Dependency on Intracellular pH

Khalid O. Alfarouk ^{1,2,3,*} , Samrein B. M. Ahmed ⁴, Robert L. Elliott ^{5,6}, Amanda Benoit ⁶, Saad S. Alqahtani ⁷, Muntaser E. Ibrahim ⁸, Adil H. H. Bashir ⁸, Sari T. S. Alhoufie ⁹, Gamal O. Elhassan ¹⁰, Christian C. Wales ², Laurent H. Schwartz ¹¹, Heyam S. Ali ¹² , Ahmed Ahmed ¹³, Patrick F. Forde ¹⁴, Jesu Devesa ¹⁵ , Rosa A. Cardone ¹⁶, Stefano Fais ¹⁷, Salvador Harguindey ¹⁸  and Stephan J. Reshkin ¹⁶ 

¹ Alfarouk Biomedical Research LLC, Temple Terrace, FL 33617, USA

² American Biosciences Inc., New York, NY 10913, USA; dwales@americanbiosciences.com

³ Al-Ghad International College for Applied Medical Sciences, Al-Madinah Al-Munawarah 42316, Saudi Arabia

⁴ College of Medicine, University of Sharjah, Sharjah P. O. Box 27272, UAE; samahmed@sharjah.ac.ae

⁵ The Elliott-Elliott-Baucom Breast Cancer Research and Treatment Center, Baton Rouge, LA 70806, USA; drrobertelliott@cox.net

⁶ The Sallie A. Burdine Breast Foundation, Baton Rouge, LA 70806, USA; amanda25304@live.com

⁷ Clinical Pharmacy Department, College of Pharmacy, Jazan University, Jazan 45142, Saudi Arabia; Ssalqahtani@jazanu.edu.sa

⁸ Institute of Endemic Diseases, University of Khartoum, Khartoum 11111, Sudan; Mibrahim@iend.org (M.E.I.); Derma55@yahoo.com (A.H.H.B.)

⁹ Department of Clinical Laboratory Sciences, Faculty of Applied Medical Sciences, Taibah University, Al-Madinah Al-Munwarah 42353, Saudi Arabia; Shoufie@taibahu.edu.sa

¹⁰ Unaizah College of Pharmacy, Qassim University, Unaizah 56264, Saudi Arabia; gamaosma963@gmail.com

¹¹ Assistance Publique des Hôpitaux de Paris, 75003 Paris, France; dr.Laurentschwartz@gmail.com

¹² Department of Pharmaceutics, Faculty of Pharmacy, University of Khartoum, Khartoum 11111, Sudan; Heyam57@hotmail.com

¹³ Department of Oesphogastric and General Surgery, University Hospitals of Leicester, Leicester LE5 4PW, UK; Ahmed.ahmed@doctors.org.uk

¹⁴ CancerResearch@UCC, Western Gateway Building, University College Cork, Cork T12 XF62, Ireland; p.forde@ucc.ie

¹⁵ Scientific Direction, Foltra Medical Centre, Travesía de Montouto 24, 15886 Teo, Spain; jesu.devesa@usc.es

¹⁶ Department of Biosciences, Biotechnologies, and Biopharmaceutics, University of Bari, 90126 Bari, Italy; rosaangela.cardone@uniba.it (R.A.C.); stephanjoel.reshkin@uniba.it (S.J.R.)

¹⁷ Department of Oncology and Molecular Medicine, Istituto Superiore di Sanità, Viale Regina Elena 299, 00161 Rome, Italy; stefano.fais@iss.it

¹⁸ Department of Oncology, Institute for Clinical Biology and Metabolism, 01004 Vitoria, Spain; salvaszh@telefonica.net

* Correspondence: Alfarouk@Hala-Alfarouk.org

Received: 21 May 2020; Accepted: 6 July 2020; Published: 11 July 2020



Abstract: The Pentose Phosphate Pathway (PPP) is one of the key metabolic pathways occurring in living cells to produce energy and maintain cellular homeostasis. Cancer cells have higher cytoplasmic utilization of glucose (glycolysis), even in the presence of oxygen; this is known as the “Warburg Effect”. However, cytoplasmic glucose utilization can also occur in cancer through the PPP. This pathway contributes to cancer cells by operating in many different ways: (i) as a defense mechanism via the reduced form of nicotinamide adenine dinucleotide phosphate (NADPH) to prevent apoptosis, (ii) as a provision for the maintenance of energy by intermediate glycolysis, (iii) by increasing genomic material to the cellular pool of nucleic acid bases, (iv) by promoting survival through increasing glycolysis, and so increasing acid production, and (v) by inducing cellular proliferation by the synthesis of nucleic acid, fatty acid, and amino acid. Each step of the PPP can be upregulated in

some types of cancer but not in others. An interesting aspect of this metabolic pathway is the shared regulation of the glycolytic and PPP pathways by intracellular pH (pHi). Indeed, as with glycolysis, the optimum activity of the enzymes driving the PPP occurs at an alkaline pHi, which is compatible with the cytoplasmic pH of cancer cells. Here, we outline each step of the PPP and discuss its possible correlation with cancer.

Keywords: cancer; metabolism; enzyme; pH; redox

1. Introduction

In 1935 and 1936, Otto Warburg showed that pyridine nucleotide diphosphopyridine nucleotide DPN (currently known as NAD⁺) functioned as an electron carrier [1–3]. He also demonstrated the presence of a second enzyme called triphosphopyridine (TPN), currently known as NAP⁺. He then discovered the Zwischenferment enzyme (known as glucose-6-phosphate dehydrogenase (G6PDH)) [1,2]. Many decades later, the Pentose Phosphate Pathway (PPP) was introduced to biochemistry.

The PPP has also been termed the phosphogluconate pathway or the hexose monophosphate shunt. It is a cytoplasmic pathway consisting of two phases: (i) the oxidative phase, followed by (ii) the nonoxidative phase. This pathway finishes as one of the glycolytic intermediates, e.g., Glyceraldehyde 3-phosphate. The oxidative phase produces two molecules of NADPH, while the nonoxidative phase produces a building block of nucleic acid.

When glucose enters the cell, glucokinase phosphorylates the free glucose in order to anchor it inside the cell by forming glucose-6-phosphate. Glucose-6-phosphate has two fates: either (i) to convert to fructose-6-phosphate via glucose-6-phosphate isomerase and complete the classical glycolytic pathway, or (ii) to convert to 6-phosphogluconate via glucose-6-phosphate dehydrogenase (G6PDH) and complete the Pentose Phosphate Pathway. We previously discussed how cancer cells recruit or manipulate the glycolytic pathway [4]. In the current manuscript, we will discuss the latter fate of glucose-6-phosphate conversion and its effects on cancer (see Figure 1).




Figure 1. Scheme showing the interaction between glycolysis and PPP.

2. Oxidative Phase

2.1. First Step

Upon glucose fixation, glucose-6-phosphate is converted to 6-phosphogluconolactone via glucose-6-phosphate dehydrogenase (G6PDH), which requires NADP⁺ to become NADPH (Figure 1). This step is irreversible and is catalyzed by G6PDH, which has two subtypes: A and B. While there is not a strong requirement for Mg⁺² (it has not been demonstrated to be an absolute requirement for the *Leuconostoc* enzyme), it is slightly stimulated by HCO⁻³ [5,6]. Indeed, the optimum activity of G6PDH occurs at an intracellular alkaline pH (pHi) which is compatible with the cytoplasmic pH of cancer cells [7,8].

The activity of G6PDH is enhanced by spironolactone [9], and insulin [10]; however, whether dexamethasone and glucose alone can increase G6PDH activity or not remains controversial [10,11]. It has been shown that high levels of glucose inhibit G6PDH activity [12] and reduce NADPH. However, such data is controversial, since Sodium Hydrogen Exchanger (NHE) stimulates glycolysis and, therefore, may act as a transient inhibitor since NADPH can also increase the Pentose Phosphate Pathway (PPP).

A study by Catanzaro et al. demonstrated G6PDH overexpression in cisplatin-resistant cancer cells, and that the use of G6PDH competitive inhibitor, 6-aminonicotinamide-AN6, sensitized the cells to cisplatin [13]. The blockade of G6PDH induced the lapatinib (TK inhibitor)-mediated cytotoxicity via enhancing autophagy [14]. Moreover, obstructing G6PDH function reversed cisplatin resistance via interfering with the homeostatic balance of reactive oxygen species (ROS), resulting in less available NADPH to neutralize ROS [15].

Possible inhibitors of G6PDH include P53 [16], ATP, which inhibits G6PDH both directly [17] and by increasing the NADPH/NADP⁺ ratio [18], and Palmitoyl-CoA which also inhibits G6PDH [19]. Moreover, NAD⁺ increases G6PDH activity [20]. Several nutraceutical products have been shown to inhibit G6PDH and could be used therapeutically, like Fermented wheat germ extract (FWGE) (Metatrol[®], Avemar[®], Polydatin or Piceid (major Resveratrol derivative) [21–23].

Evolutionary Consequences of Population Selected G6PDH Deficiency

Interestingly, G6PDH deficiency is an X-linked recessive hereditary disease [24,25]; this deficiency is accompanied by several clinical manifestations, as follows:

G6PDH deficiency confers natural resistance against malaria [26]. From an epidemiological standpoint, G6PDH deficiency is correlated with reduced cancer risk [27–29]. Thus, G6PDH deficiency represents a natural prevention strategy against tumor development. Such a deficiency could be explained as follows:

Free radical scavengers such as NADPH are cytoplasmic reducing agents (see below). In other words, NADPH is a cytoplasmic free radical scavenger. Overproduction of free radicals, reactive oxygen species (ROS) and reactive nitrogen species (RNS) by NAD(P)H oxidase isoforms and NO synthase respectively, induces cellular senescence and apoptosis [30], as well as necrosis [31].

Cancer cells induce anti-apoptotic proteins [32] and are devoid of pro-apoptotic protein expression [33]. Hence, the ratio of the anti-apoptotic/apoptotic proteins is indicative of increased malignancy [34]. NADPH increases this ratio and, therefore, decreases apoptosis [35,36]. Also, the pro-apoptotic protein, BCL-2, has anti-oxidant properties [37], which increase the perturbations of the cellular redox status. So, two questions arise: is it preventive or prompting? And does it depend on the stage of the tumor?

2.2. Second Step

As shown in Figure 1, 6-phosphoglucono- δ -lactone is converted into 6-phosphogluconate via the 6-phosphogluconolactonase (6PGL) enzyme.

The activity of 6-phosphogluconolactonase enzyme increases sharply as intracellular pH increases [38], which is compatible with the cytoplasmic pH of cancer cells.

Epstein Barr virus (EBV) upregulates the expression of 6-phosphogluconolactonase [34], which is crucial to tumor growth [39]. Furthermore, it has been reported that 6PGL correlates with pancreatic cancer resistance to Gemcitabine [40].

The inhibitory abilities of an interesting compound, GP269, were tested against 6-phosphogluconolactonase (6PGL). While this inhibition was attempting to target *Trypanosoma Brucei*, this compound could also be a potential therapeutic tool for breast cancer, since 6PGL was found to be overexpressed in human epidermal growth factor receptor 2 (HER2) positive breast cancer cells [41,42].

2.3. Third Step

In the last step of the oxidative phase, 6-phosphogluconate undergoes decarboxylation by the 6-phosphogluconate dehydrogenase (6PGD) enzyme in the presence of NADP⁺ to produce ribulose 5-phosphate and NADPH (Figure 1). NADPH inhibits 6PGD, so it has a negative mechanism. Again, the optimum pHi for this enzyme is alkaline [43,44].

Consistent with its association with poor prognoses, 6PGD supports tumor growth [39,45], metastasis [46], and chemotherapy resistance [47]. Aberrant expression levels of 6PGD were reported in cervical cancer and were linked to more proliferative and migratory abilities of cancer cells [48]. The deactivation of 6PGD resulted in a reduction of RhoA and Rac1 function, which, in turn, negatively affected the cancerous behavior of the cervical cancer cells [48]. Like G6PDH, 6PGD was reported to help in sensitizing cisplatin-resistant cells in both ovarian and lung cancer. In addition, 6PGD expression was shown to be linked with poor prognoses in breast cancer patients; thus, its inhibition was suggested to improve breast cancer outcomes [49]. Therefore, 6PGD overexpression is a successful adaptive (evolutionary) strategy to confer cancer cell survival. Therefore, it will not be surprising if there is extensive research on 6PGD as a potential target in cancer management [49]. Parietin (Phycione) is an anti-infectious agent [50] and a 6PGD inhibitor [45,51,52]. Therefore, it has been suggested that it be administered directly as an anticancer agent, as well as to resensitize resistant tumors to therapy [51–53]. This applies to solid tumors, but also to hematological malignancies, as it has been demonstrated that inhibition of 6PG sensitizes leukemia cells to the antimalarial drug, dihydroartemisinin [52]

3. Nonoxidative Phase

In contrast to the oxidative phase, in this phase, all the reactions are reversible. In accordance with the steps above, the nonoxidative steps are as follows.

3.1. Fourth Step

3.1.1. Ribose-5 Phosphate Formation

Ribulose-5-phosphate produces Ribose 5-phosphate (R5P) via Ribose-5-phosphate isomerase (Rpi) (Figure 1). The formation of R5P depends on (i) cell growth [54], (ii) the redox state of the cell [55,56], and (iii) the metabolic state of the cell [3]. Moreover, Rpi has its optimum activity at an alkaline pHi [57].

Rpi has two forms: RpiA or RpiB [58,59], with RpiA being mainly expressed in humans [60] and RpiB mainly detected on lower microorganisms [61–63]. It has been shown that ribose-5-phosphate isomerase A overexpression promotes and is associated with several types of cancer, including liver, lung, and breast [64,65]. One of the possible explanations of how Rpi promotes carcinogenesis is through its activation of ERK and β -catenin pathways [64], and perhaps through the inhibition of LC3 (e.g., alteration of autophagy) [66].

3.1.2. Xylulose 5-Phosphate Formation

Xylulose 5-phosphate (Xu5P) is formed from Ribulose-5-phosphate via the metalloproteinase enzyme Ribulose 5-Phosphate 3-Epimerase (RPE) (Figure 1), which has optimal activity at a slightly alkaline pHi [58].

Ribulose 5-Phosphate 3-Epimerase (RPE) has been detected in pancreatic cancer, and its effect might be through the activation of fructose 2,6 bisphosphate (F-2, 6BP), which activates phosphofructose kinase 1 (PFK1), and/or K-ras [67–69].

Hydrogen peroxide (H₂O₂) inactivates RPE, while iron and manganese activate the RPE [70]. It is known that a high glucose load induces intracellular alkalinity and the Warburg Effect [71]. Also, in response to higher glucose, Xu5P activates the carbohydrate-responsive, element-binding protein (ChREBP) [72,73], which is also known as MLX-interacting protein-like (MLXIPL) and, in humans,

is encoded by the MLXIPL gene [74]. ChREBP protein is overexpressed in cancer, and its attenuation is accompanied by the activation of P53 and induction of cell cycle arrest [75–77]. Moreover, ChREBP mediates the activation of L-Pyruvate Kinase (L-PK), which is essential for pyruvate formation [78,79].

3.2. Fifth Step

Here, ribose-5-phosphate and xylulose-5-phosphate interact to form sedoheptulose 7-phosphate to produce sedoheptulose 7-phosphate and glyceraldehyde 3-phosphate using transketolase enzyme (TKT) (Figure 1). Glyceraldehyde 3-phosphate is an essential intermediate to induce glycolysis and the Warburg Effect [4].

Transketolase (TKT) is a thiamine-dependent enzyme [80], again having optimal activity at an alkaline pHi [81,82]. TKT plays a crucial role in regulating the cellular redox state [83]. TKT expression is associated with poor prognoses and metastases, while its suppression decreases tumor metastasis [84]. Oxythiamine is an inhibitor of TKT [85]. Moreover, it has been shown that Furosemide (Lasix[®]) have a drastic inhibitory effect on TKT, as it induces thiamine deficiency [86].

3.3. Sixth Step

Sedoheptulose 7-phosphate interacts with glyceraldehyde 3-phosphate to produce erythrose 4-phosphate and fructose 6-phosphate via the transaldolase enzyme (TADOL) (Figure 1), which has optimal activity at alkaline pHi [87,88].

The TADOL enzyme is a rate-limiting enzyme in the nonoxidative PPP [89–91]. *Transaldolase 1 gene (TADOL1)* is ubiquitously expressed, except in erythrocytes (Red Blood Cells (RBCs)), that might be challenging, because RBCs rely on the PPP in their metabolism.

It had been shown that the TADOL activity was higher in liver tumor rather than in normal control rat liver [90]. TADOL is a master regulator of the redox state of the cell, and so its overexpression might affect not only tumor survival, but also response to therapy. It has been shown that elevated TADOL expression correlates with a decrease in response to HER2 inhibition in breast cancer patients [92].

Starvation decreased TADOL activity, which is beneficial to the prevention and/or slowing of cancer development [90,93,94]. Arabinose 5-phosphate is the aldopentose (a metabolic intermediate in the biosynthesis of lipopolysaccharide) inhibits TADOL [95].

The formation of fructose 1,6-bisphosphate inhibits TADOL activity, which might explain why the PPP and glycolysis do not occur simultaneously.

Some tissues, such as heart and skeletal muscle, have lower transaldolase protein expression [90]; as such, those tissues tend to have lower susceptibility for tumor growth. While we cannot conclude that the TADOL could be cancer's Achilles' Heel, it might open the window regarding PPP activity and tissue susceptibility; this might, in turn, reveal why PPP favors carcinogenesis. Also, these tissues have a higher mitochondrial number compared to other tissues, and so, we propose that PPP activity might be inversely proportional to mitochondrial metabolism.

3.4. Seventh Step

This step could be represented as reinventing the wheel to closely pool glycolytic pathways, like the fifth and sixth steps. In this step, xylulose 5-phosphate interacts with erythrose 4-phosphate to produce glyceraldehyde 3-phosphate and fructose 6-phosphate via transketolase (TKT), which has optimal activity at pHi of 7.5–7.6 [90,96] (Figure 1). For TKT kinetics, see above.

It seems that the net result of the second phase of the PPP (i.e., the nonoxidative phase) is to support the glycolytic pathway via the provision of two molecules of fructose-6-phosphate and one of glyceraldehyde-3-phosphate. These molecules are critical intermediates in the glycolytic pathway. Therefore, in conclusion, the PPP provides the nucleic acid building blocks, maintains the redox state of the cell and supports cellular energy.

3.4.1. NADPH and GSH

How does the PPP maintain oxidation-reduction reactions in the cell (i.e., the Redox state of the cell)? During the first and the third steps (see above), nicotinamide adenine dinucleotide phosphate (NADP⁺) is a cofactor in the formation of NADPH; that is, each round of the PPP forms two molecules of NADPH. NADPH acts as a hydrogen donor to the disulfide bond of the oxidized glutathione (G-S-S-G) bond to form reduced glutathione GSH, while the NADPH becomes NADP⁺ via the glutathione reductase enzyme. Then, GSH donates its H⁺ to the oxidant, e.g., H₂O₂, to produce H₂O, while the GSH converts back into the oxidized form, G-S-S-G, via the activity of glutathione peroxidase enzyme. Therefore, this system acts like a series of gears in that the NADPH gear activates the glutathione gear and is a scavenger of free radicals “Glutathione Redox Cycle” (Figure 2) [97–100]. Therefore, the NADP⁺/NADPH ratio could be expressed indirectly as the GSH/GSSG ratio. If the PPP decreases, NADPH decreases, resulting in an abundant glutathione ratio in the oxidized form.




Figure 2. Glutathione Reductase Cycle.

3.4.2. The Overview of PPP and Cancer

Reactive oxygen species (ROS) and reactive nitrogen species (RNS) are double-edged swords. On the one hand, they lead to aberrant gene expression and genetic instability that are fundamental processes in the development and progression of cancer [101–104]. Furthermore, ROS and NOS, at a certain point, are supportive of evolutionary tumor trajectory via supporting angiogenesis and metastasis, especially during cellular senescence, as they are hallmarks of inflammatory and tumor microenvironments [105–108]. However, the presence of ROS and RNS is a key determinant in the apoptosis of cancer cells [109–113]. Therefore, cancer cells robustly adjust to a very fine ratio of the cellular redox state in order to confer their survival. As a result, we conclude that highly dividing cells are well equipped for tumor development, while senescent cells are more prone to induce metastasis, as these cells lack an efficient PPP [39,114]. This inefficiency during senescence is not accompanied by a decrease in the level of metabolites since metabolites can derive from two sources: (i) the metabolites of the nonoxidative phase, e.g., ribose 5-phosphate, ribulose, xylulose 5-phosphate and sedoheptulose 7-phosphate [115], and (ii) those coming from glycolysis rather than the oxidative phase of the PPP, which reflects the activity of the PPP on the next stage of metastasis and settlement at the distal site (Mesenchymal Epithelial transition (MET)) [116,117]. In other words, cancer cells manipulate and escalate the metabolic pathways, and the PPP is one of those pathways to support their evolutionary fitness.

The Pentose Phosphate Pathway maintains cancer’s redox state, but it also affects cell signaling during the cell cycle, as well as managing the metabolic pathways, e.g., xylulose 5-phosphate (see above) [72,117–120].

Necrosis is another cellular death pathway that has a drastic effect on tumorigenesis. That is why chemotherapy acts via apoptosis and not necrosis [121–125]. However, several mechanisms of drug resistance are associated with the upregulation of the PPP to prevent the formation of free radicals [126]. The role of the PPP is not confined only to antagonizing the effect of chemotherapeutic agents; it also

increases the expression and/or activity of the proteins (e.g., Multidrug-resistance-associated protein (MRP), ATP-binding-cassette (ABC) transporters) that either promote the efflux of anticancer drugs (e.g., vincristine, doxorubicin, daunorubicin) or block the influx of these agents [127–129].

3.4.3. The Possible Crosstalk between the Glycolysis and the Pentose Phosphate Pathway (NADPH Is a Double-edged Sword)

Generally, the presence of NADPH inhibits G6PDH, thereby blocking the PPP negative feedback control) and might block the possibility of nucleic acid pooling. Again, this raises an important question: does NADPH suppress tumor growth? NADPH has a critical role in the completion of the first phase of glycolysis (preparatory phase). Therefore, NADPH shifts the metabolism from PPP to glycolysis. The formation of NADPH will increase the formation of NAD⁺, and therefore, support the formation of pyruvic acid and lactic acid, which represents an evolutionary advantage [130]. Therefore, the role of NADPH in cancer is a conundrum.

In aggressive cancer cells, where the metabolic need for nucleotides exceeds that of NADPH, TKT and TALDO catalyze the reverse reactions and divert glyceraldehyde 3-phosphate and fructose 6-phosphate from glycolysis to the nonoxidative PPP to produce additional ribonucleotides. In this regard, glycolysis itself acts as a backup for the DNA building blocks. Therefore, the nonoxidative phase might represent the bridge between phase 1 of the PPP and phase 1 of glycolysis, that could enhance the metabolic plasticity of cancer cells [131]. An important overlap between these two systems that has, to our knowledge, been overlooked is that both pathways share an exquisite sensitivity to cytoplasmic, intracellular pH (pHi), in that all the enzymes that drive both pathways have their optimum activities at alkaline values (Table 1). This probable parallel regulation by processes that determines pHi underlines the importance of both metabolic pathways in cancer and of the aforementioned crosstalk of phase 1 of the two systems in determining the relative usage of one or the other pathways in a specific cellular context. Indeed, this shared intracellular alkaline optimum of the enzymes that drives glycolysis and the PPP in cancer cells opens new avenues for possible therapeutic opportunities for manipulating this pathway via its sensitivity to changes in pHi. Additionally, the formation of fructose 1,6-bisphosphate inhibits TADOL activity, which might further explain why the PPP and glycolysis do not occur simultaneously.

Table 1. Optimum pHi for each PPP driving enzyme.

Enzyme	Optimum pH
Glucose-6-phosphate dehydrogenase (G6PDH)	7.8 [8]
6-phosphogluconolactonase (6PGL)	7.4 [38]
6-phosphogluconate dehydrogenase (6PGD)	Range from (7–10) depending on several factors including the buffer used in the experiment [43]
Ribose-5-phosphate isomerase (RPI)	8.4 [132]
Ribulose 5-Phosphate 3-Epimerase (RPE)	7.25–7.5 [69]
Transketolase (TKT)	7.5–7.6 [82,90,96]
Transaldolase (TADOL)	8 [133]

4. Concluding Remarks

Cancer as a biological system and/or tissue has unusual physiological, biochemical, and biophysical parameters compared to normal tissues [134]. To maintain viability, cancer has modified several existing metabolic pathways to fulfill its energy demands with as low a cost as possible [135]. One of these metabolic pathways that utilize glucose in the cytoplasm is the Pentose Phosphate Pathway (PPP). In the 1920s, Otto Warburg observed that cancer cells had a higher fermentation rate, and later concluded that cancer relied on the cytoplasmic utilization of glucose rather than mitochondrial

utilization. However, cytoplasmic utilization can occur through glycolysis and/or the PPP. In our previous work, we showed the detailed biochemistry of glycolysis in cancer and its possible therapeutic opportunities and limitations [4]. Here, we present a study of the PPP pathway that suggests that it has a higher impact on carcinogenesis than glycolysis, and that cancer might invest in the PPP rather than glycolysis based on the cellular state and the cell cycle phase, whether it undergoes proliferation or growth. Both glycolysis and the PPP have mutual intermediates, but the PPP has more ramifications through its contribution to the synthesis of nucleic acids and maintains the redox state at the optimum that assures cancer cell survival. Also, the PPP interacts with the mitochondrion and so manipulates the programmed cell death pathway (apoptosis). Therefore, the PPP is paramount in cancer biology. Much work is now focusing on targeted diagnostics and therapeutics by selecting definite steps; however, tumor colonies consist of a heterogeneous population having different modified metabolic strategies that can confound the diagnosis and therapy of cancer. We have tried here to contribute to drawing a complete picture of the cancer puzzle to be in a relatively better position for its management rather than its eradication. Lastly, the described intracellular alkaline optimum of all the enzymes that drive the PPP in cancer cells (Table 1) opens new avenues for understanding the roles, regulations, and possible therapeutic opportunities for manipulating this pathway via its sensitivity to the changes in pH_i that is common to cancer cells.

Author Contributions: K.O.A. involved in conceptualization, investigation, project administration, methodology, Writing-original draft, Writing-review & editing, visualization, validation. S.J.R. involved in supervision, methodology, Writing-review & editing, validation, visualization, conceptualization. S.B.M.A., R.L.E., A.B., S.S.A., M.E.I., A.H.H.B., S.T.S.A. and G.O.E. involved in resources, validation, formal analysis. P.F.F., J.D., C.C.W., L.H.S., H.S.A., A.A., R.A.C., S.F. and S.H. involved in the investigation, formal analysis, writing and funding acquisition. All authors have read and agreed to the published version of the manuscript.

Funding: This work is partially supported by Orchid Life, LLC. The authors declare that this study received funding from Orchid Life, LLC. The funder was not involved in the Writing of this article or the decision to submit it for publication.

Acknowledgments: We thank Linda Aurdal Wales for the excellent discussion on the topic. In memory of Charles Ed Bailey, for his valuable discussions on the current topic.

Conflicts of Interest: Author K.O.A. and C.W. are employed by the American Biosciences Company. The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Abbreviations

G6PDH	Glucose-6-phosphate dehydrogenase
6PGL	6-phosphogluconolactonase
6PGD	6-phosphogluconate dehydrogenase
R5P	Ribose-5-phosphate isomerase
RPE	Ribulose 5-Phosphate 3-Epimerase
TKT	Transketolase
TADOL	Transaldolase
PGI	Phosphoglucose isomerase
PFK1	Phosphofructokinase-1
PFK2	Phosphofructokinase-2
ALDO	Fructose-bisphosphate aldolase
TPI	Triosephosphate isomerase
GAPDH	Glyceraldehyde phosphate dehydrogenase
PGK	Phosphoglycerate kinase
PGM	Phosphoglycerate mutase
ENO	Enolase
PK	Pyruvate Kinase
LDH	Lactate Dehydrogenase

References

1. Warburg, O.; Christian, W.; Griese, A. Wasserstoff(ü)bertragendes Co-Ferment, seine Zusammensetzung und Wirkungsweise. *Biochem. Z.* **1935**, *282*, 157–205.
2. Warburg, O.; Christian, W. Optischer Nachweis der Hydrierung und Dehydrierung des Pyridins im Gärungs-Co-Ferment. *Biochem. Z.* **1936**, *286*, 81.
3. Stincone, A.; Prigione, A.; Cramer, T.; Wamelink, M.M.C.; Campbell, K.; Cheung, E.; Olin-Sandoval, V.; Grüning, N.-M.; Krüger, A.; Alam, M.T.; et al. The return of metabolism: Biochemistry and physiology of the pentose phosphate pathway. *Biol. Rev.* **2015**, *90*, 927–963. [[CrossRef](#)] [[PubMed](#)]
4. Alfarouk, K.O.; Verduzco, D.; Rauch, C.; Muddathir, A.K.; Bashir, A.H.H.; Elhassan, G.O.; Ibrahim, M.E.; Orozco, J.D.P.; Cardone, R.A.; Reshkin, S.J.; et al. Glycolysis, tumor metabolism, cancer growth and dissemination. A new pH-based etiopathogenic perspective and therapeutic approach to an old cancer question. *Oncoscience* **2014**, *1*, 777. [[CrossRef](#)] [[PubMed](#)]
5. Olive, C.; Levy, H.R. The preparation and some properties of crystalline glucose 6-phosphate dehydrogenase from *Leuconostoc mesenteroides*. *Biochemistry* **1967**, *6*, 730–736. [[CrossRef](#)] [[PubMed](#)]
6. Anderson, W.B.; Nordlie, R.C. Glucose dehydrogenase activity of yeast glucose 6-phosphate dehydrogenase. I. Selective stimulation by bicarbonate, phosphate, and sulfate. *Biochemistry* **1968**, *7*, 1479–1485. [[CrossRef](#)]
7. Türkoğlu, V.; Aldemir, S.; Çiftçi, M. Purification and Characterization of Glucose 6-Phosphate Dehydrogenase from Sheep Liver. *Turk. J. Chem.* **2003**, *27*, 395–402.
8. Mahmoud, A.I.; Abdel-Hady, M.G.; Ahmed, M.H.S.; Mohamed, A.G.; Mohamed, M.A.-M. Purification and Characterization of glucose-6-phosphate Dehydrogenase From Camel Liver. *Enzym. Res.* **2014**, *2014*, 714054.
9. Pessôa, B.S.; Peixoto, E.B.M.I.; Papadimitriou, A.; Lopes de Faria, J.M.; Lopes de Faria, J.B. Spironolactone improves nephropathy by enhancing glucose-6-phosphate dehydrogenase activity and reducing oxidative stress in diabetic hypertensive rat. *J. Renin-Angiotensin-Aldosterone Syst.* **2012**, *13*, 56–66. [[CrossRef](#)] [[PubMed](#)]
10. Manos, P.; Nakayama, R.; Holten, D. Regulation of glucose-6-phosphate dehydrogenase synthesis and mRNA abundance in cultured rat hepatocytes. *Biochem. J.* **1991**, *276 Pt 1*, 245–250. [[CrossRef](#)]
11. Salati, L.M.; Adkins-Finke, B.; Clarke, S.D. Free fatty acid inhibition of the insulin induction of glucose-6-phosphate dehydrogenase in rat hepatocyte monolayers. *Lipids* **1988**, *23*, 36–41. [[CrossRef](#)] [[PubMed](#)]
12. Zhang, Z.; Apse, K.; Pang, J.; Stanton, R.C. High glucose inhibits glucose-6-phosphate dehydrogenase via cAMP in aortic endothelial cells. *J. Biol. Chem.* **2000**, *275*, 40042–40047. [[CrossRef](#)] [[PubMed](#)]
13. Catanzaro, D.; Gaude, E.; Orso, G.; Giordano, C.; Guzzo, G.; Rasola, A.; Ragazzi, E.; Caparrotta, L.; Frezza, C.; Montopoli, M. Inhibition of glucose-6-phosphate dehydrogenase sensitizes cisplatin-resistant cells to death. *Oncotarget* **2015**, *6*, 30102–30114. [[CrossRef](#)] [[PubMed](#)]
14. Mele, L.; La Noce, M.; Paino, F.; Regad, T.; Wagner, S.; Liccardo, D.; Papaccio, G.; Lombardi, A.; Caraglia, M.; Tirino, V.; et al. Glucose-6-phosphate dehydrogenase blockade potentiates tyrosine kinase inhibitor effect on breast cancer cells through autophagy perturbation. *J. Exp. Clin. Cancer Res.* **2019**, *38*, 1–13. [[CrossRef](#)] [[PubMed](#)]
15. Hong, W.; Cai, P.; Xu, C.; Cao, D.; Yu, W.; Zhao, Z.; Huang, M.; Jin, J. Inhibition of glucose-6-phosphate dehydrogenase reverses cisplatin resistance in lung cancer cells via the redox system. *Front. Pharmacol.* **2018**, *9*, 43. [[CrossRef](#)] [[PubMed](#)]
16. Jiang, P.; Du, W.; Wang, X.; Mancuso, A.; Gao, X.; Wu, M.; Yang, X. p53 regulates biosynthesis through direct inactivation of glucose-6-phosphate dehydrogenase. *Nat. Cell Biol.* **2011**, *13*, 310–316. [[CrossRef](#)]
17. Horne, R.N.; Anderson, W.B.; Nordlie, R.C. Glucose dehydrogenase activity of yeast glucose 6-phosphate dehydrogenase. Inhibition by adenosine 5'-triphosphate and other nucleoside 5'-triphosphates and diphosphates. *Biochemistry* **1970**, *9*, 610–616. [[CrossRef](#)]
18. Levy, H.R.; Daouk, G.H.; Katopes, M.A. Regulation of coenzyme utilization by the dual nucleotide-specific glucose 6-phosphate dehydrogenase from *Leuconostoc mesenteroides*. *Arch. Biochem. Biophys.* **1979**, *198*, 406–413. [[CrossRef](#)]
19. Kawaguchi, A.; Bloch, K. Inhibition of glucose 6-phosphate dehydrogenase by palmitoyl coenzyme A. *J. Biol. Chem.* **1974**, *249*, 5793–5800.

20. Levy, H.R.; Christoff, M.; Ingulli, J.; Ho, E.M. Glucose-6-phosphate dehydrogenase from *Leuconostoc mesenteroides*: Revised kinetic mechanism and kinetics of ATP inhibition. *Arch. Biochem. Biophys.* **1983**, *222*, 473–488. [[CrossRef](#)]
21. Comín-Anduix, B.; Boros, L.G.G.; Marin, S.; Boren, J.; Callol-Massot, C.; Centelles, J.J.J.; Torres, J.L.L.; Agell, N.; Bassilian, S.; Cascante, M.; et al. Fermented Wheat Germ Extract Inhibits Glycolysis/Pentose Cycle Enzymes and Induces Apoptosis through Poly(ADP-ribose) Polymerase Activation in Jurkat T-cell Leukemia Tumor Cells. *J. Biol. Chem.* **2002**, *277*, 46408–46414. [[CrossRef](#)] [[PubMed](#)]
22. Zhurakivska, K.; Troiano, G.; Caponio, V.; Dioguardi, M.; Arena, C.; Lo Muzio, L.; Zhurakivska, K.; Troiano, G.; Caponio, V.C.A.; Dioguardi, M.; et al. The Effects of Adjuvant Fermented Wheat Germ Extract on Cancer Cell Lines: A Systematic Review. *Nutrients* **2018**, *10*, 1546. [[CrossRef](#)] [[PubMed](#)]
23. Mele, L.; Paino, F.; Papaccio, F.; Regad, T.; Boocock, D.; Stiuso, P.; Lombardi, A.; Liccardo, D.; Aquino, G.; Barbieri, A.; et al. A new inhibitor of glucose-6-phosphate dehydrogenase blocks pentose phosphate pathway and suppresses malignant proliferation and metastasis in vivo. *Cell Death Dis.* **2018**, *9*, 572. [[CrossRef](#)]
24. Guindo, A.; Fairhurst, R.M.; Doumbo, O.K.; Wellems, T.E.; Diallo, D.A. X-linked G6PD deficiency protects hemizygous males but not heterozygous females against severe malaria. *PLoS Med.* **2007**, *4*, e66. [[CrossRef](#)] [[PubMed](#)]
25. Farhud, D.; Yazdanpanah, L. Glucose-6-phosphate dehydrogenase (G6PD) Deficiency. *Iran. J. Public Health* **2008**, *37*, 1–18.
26. Greene, L.S. G6PD deficiency as protection against falciparum malaria: An epidemiologic critique of population and experimental studies. *Am. J. Phys. Anthropol.* **1993**, *36*, 153–178. [[CrossRef](#)]
27. Cocco, P. Does G6PD deficiency protect against cancer? A critical review. *J. Epidemiol. Community Health* **1987**, *41*, 89–93. [[CrossRef](#)]
28. Cocco, P.; Dessì, S.; Avataneo, G.; Picchiri, G.; Heinemann, E. Glucose-6-phosphate dehydrogenase deficiency and cancer in a Sardinian male population: A case-control study. *Carcinogenesis* **1989**, *10*, 813–816. [[CrossRef](#)]
29. Pisano, M.; Cocco, P.; Cherchi, R.; Onnis, R.; Cherchi, P. Glucose-6-phosphate dehydrogenase deficiency and lung cancer: A hospital based case-control study. *Tumori J.* **1991**, *77*, 12–15. [[CrossRef](#)]
30. Kern, J.C.; Kehrer, J.P. Free radicals and apoptosis: Relationships with glutathione, thioredoxin, and the BCL family of proteins. *Front. Biosci.* **2005**, *10*, 1727–1738. [[CrossRef](#)]
31. Valko, M.; Leibfritz, D.; Moncol, J.; Cronin, M.T.D.; Mazur, M.; Telser, J. Free radicals and antioxidants in normal physiological functions and human disease. *Int. J. Biochem. Cell Biol.* **2007**, *39*, 44–84. [[CrossRef](#)] [[PubMed](#)]
32. Certo, M.; Moore, V.D.G.; Nishino, M.; Wei, G.; Korsmeyer, S.; Armstrong, S.A.; Letai, A. Mitochondria primed by death signals determine cellular addiction to antiapoptotic BCL-2 family members. *Cancer Cell* **2006**, *9*, 351–365. [[CrossRef](#)] [[PubMed](#)]
33. Lowe, S.W.; Lin, A.W. Apoptosis in cancer. *Carcinogenesis* **2000**, *21*, 485–495. [[CrossRef](#)] [[PubMed](#)]
34. Beerheide, W.; Tan, Y.J.; Teng, E.; Ting, A.E.; Jedpiyawongse, A.; Srivatanakul, P. Downregulation of proapoptotic proteins Bax and Bcl-X(S) in p53 overexpressing hepatocellular carcinomas. *Biochem. Biophys. Res. Commun.* **2000**, *273*, 54–61. [[CrossRef](#)] [[PubMed](#)]
35. Lee, J.K.; Edderkaoui, M.; Truong, P.; Ohno, I.; Jang, K.; Berti, A.; Pandol, S.J.; Gukovskaya, A.S. NADPH Oxidase Promotes Pancreatic Cancer Cell Survival via Inhibiting JAK2 Dephosphorylation by Tyrosine Phosphatases. *Gastroenterology* **2007**, *133*, 1637–1648. [[CrossRef](#)] [[PubMed](#)]
36. Zou, X.; Feng, Z.; Li, Y.; Wang, Y.; Wertz, K.; Weber, P.; Fu, Y.; Liu, J. Stimulation of GSH synthesis to prevent oxidative stress-induced apoptosis by hydroxytyrosol in human retinal pigment epithelial cells: Activation of Nrf2 and JNK-p62/SQSTM1 pathways. *J. Nutr. Biochem.* **2012**, *23*, 994–1006. [[CrossRef](#)] [[PubMed](#)]
37. Cai, J.; Jones, D.P. Superoxide in apoptosis. Mitochondrial generation triggered by cytochrome c loss. *J. Biol. Chem.* **1998**, *273*, 11401–11404. [[CrossRef](#)]
38. Bauer, H.P.; Srihari, T.; Jochims, J.C.; Hofer, H.W. 6-Phosphogluconolactonase. Purification, Properties and Activities in Various Tissues. *Eur. J. Biochem.* **1983**, *133*, 163–168. [[CrossRef](#)]
39. Sukhatme, V.P.; Chan, B. Glycolytic cancer cells lacking 6-phosphogluconate dehydrogenase metabolize glucose to induce senescence. *FEBS Lett.* **2012**, *586*, 2389–2395. [[CrossRef](#)]
40. Mori-Iwamoto, S.; Kuramitsu, Y.; Ryozaawa, S.; Mikuria, K.; Fujimoto, M.; Maehara, S.-I.; Maehara, Y.; Okita, K.; Nakamura, K.; Sakaida, I. Proteomics finding heat shock protein 27 as a biomarker for resistance of pancreatic cancer cells to gemcitabine. *Int. J. Oncol.* **2007**, *31*, 1345–1350. [[CrossRef](#)]

41. Tran, A.T.; Sadet, A.; Calligari, P.; Lopes, P.; Ouazzani, J.; Sollogoub, M.; Miclet, E.; Abergel, D. Targeting the Pentose Phosphate Pathway: Characterization of a New 6PGL Inhibitor. *Biophys. J.* **2018**, *115*, 2114–2126. [[CrossRef](#)] [[PubMed](#)]
42. Choi, J.; Kim, E.S.; Koo, J.S. Expression of pentose phosphate pathway-related proteins in breast cancer. *Dis. Markers* **2018**, *2018*, 9369358. [[CrossRef](#)] [[PubMed](#)]
43. Ceyhan, D.; Danişan, A.; Ögüş, I.H.; Özer, N. Purification and Kinetic Properties of 6-Phosphogluconate Dehydrogenase from Rat Small Intestine. *Protein J.* **2005**, *24*, 293–301. [[CrossRef](#)] [[PubMed](#)]
44. Adem, S.; Ciftci, M. Purification and biochemical characterization of glucose 6-phosphate dehydrogenase, 6-phosphogluconate dehydrogenase and glutathione reductase from rat lung and inhibition effects of some antibiotics. *J. Enzym. Inhib. Med. Chem.* **2016**, *31*, 1342–1348. [[CrossRef](#)] [[PubMed](#)]
45. Lin, R.; Elf, S.; Shan, C.; Kang, H.-B.; Ji, Q.; Zhou, L.; Hitosugi, T.; Zhang, L.; Zhang, S.; Seo, J.H.; et al. 6-Phosphogluconate dehydrogenase links oxidative PPP, lipogenesis and tumour growth by inhibiting LKB1–AMPK signalling. *Nat. Cell Biol.* **2015**, *17*, 1484–1496. [[CrossRef](#)]
46. Chan, B.; VanderLaan, P.A.; Sukhatme, V.P. 6-Phosphogluconate dehydrogenase regulates tumor cell migration in vitro by regulating receptor tyrosine kinase c-Met. *Biochem. Biophys. Res. Commun.* **2013**, *439*, 247–251. [[CrossRef](#)]
47. Wang, X.; Li, X.; Zhang, X.; Fan, R.; Gu, H.; Shi, Y.; Liu, H. Glucose-6-phosphate dehydrogenase expression is correlated with poor clinical prognosis in esophageal squamous cell carcinoma. *Eur. J. Surg. Oncol.* **2015**, *41*, 1293–1299. [[CrossRef](#)]
48. Zheng, W.; Feng, Q.; Liu, J.; Guo, Y.; Gao, L.; Li, R.; Xu, M.; Yan, G.; Yin, Z.; Zhang, S.; et al. Inhibition of 6-phosphogluconate Dehydrogenase Reverses Cisplatin Resistance in Ovarian and Lung Cancer. *Front. Pharmacol.* **2017**, *8*, 421. [[CrossRef](#)]
49. Guo, H.; Xiang, Z.; Zhang, Y.; Sun, D. Inhibiting 6-phosphogluconate dehydrogenase enhances chemotherapy efficacy in cervical cancer via AMPK-independent inhibition of RhoA and Rac1. *Clin. Transl. Oncol.* **2019**, *21*, 404–411. [[CrossRef](#)]
50. Yang, X.; Peng, X.; Huang, J. Inhibiting 6-phosphogluconate dehydrogenase selectively targets breast cancer through AMPK activation. *Clin. Transl. Oncol.* **2018**, *20*, 1145–1152. [[CrossRef](#)]
51. Basile, A.; Rigano, D.; Loppi, S.; Di Santi, A.; Nebbioso, A.; Sorbo, S.; Conte, B.; Paoli, L.; De Ruberto, F.; Molinari, A.M.; et al. Antiproliferative, Antibacterial and Antifungal Activity of the Lichen *Xanthoria parietina* and Its Secondary Metabolite Parietin. *Int. J. Mol. Med.* **2015**, *16*, 7861–7875. [[CrossRef](#)] [[PubMed](#)]
52. Elf, S.; Lin, R.; Xia, S.; Pan, Y.; Shan, C.; Wu, S.; Lonial, S.; Gaddh, M.; Arellano, M.L.; Khoury, H.J.; et al. Targeting 6-phosphogluconate dehydrogenase in the oxidative PPP sensitizes leukemia cells to antimalarial agent dihydroartemisinin. *Oncogene* **2017**, *36*, 254–262. [[CrossRef](#)] [[PubMed](#)]
53. Zugic, A.; Jeremic, I.; Isakovic, A.; Arsic, I.; Savic, S.; Tadic, V. Evaluation of Anticancer and Antioxidant Activity of a Commercially Available CO₂ Supercritical Extract of Old Man’s Beard (*Usnea barbata*). *PLoS ONE* **2016**, *11*, e0146342. [[CrossRef](#)] [[PubMed](#)]
54. Pilz, R.B.; Willis, R.C.; Boss, G.R. The influence of ribose 5-phosphate availability on purine synthesis of cultured human lymphoblasts and mitogen-stimulated lymphocytes. *J. Biol. Chem.* **1984**, *259*, 2927–2935.
55. Garcia, J.; Han, D.; Sancheti, H.; Yap, L.-P.; Kaplowitz, N.; Cadenas, E. Regulation of mitochondrial glutathione redox status and protein glutathionylation by respiratory substrates. *J. Biol. Chem.* **2010**, *285*, 39646–39654. [[CrossRef](#)] [[PubMed](#)]
56. Foyer, C.H.; Halliwell, B. The presence of glutathione and glutathione reductase in chloroplasts: A proposed role in ascorbic acid metabolism. *Planta* **1976**, *133*, 21–25. [[CrossRef](#)] [[PubMed](#)]
57. Noltmann, E.A. Aldose-ketose isomerases. In *The Enzymes*; Boyer, P.D., Ed.; Academic Press: New York, NY, USA; London, UK, 1972; pp. 272–354.
58. Rangarajan, E.S.; Sivaraman, J.; Matte, A.; Cygler, M. Crystal structure of D-ribose-5-phosphate isomerase (RpiA) from *Escherichia coli*. *Proteins Struct. Funct. Genet.* **2002**, *48*, 737–740. [[CrossRef](#)] [[PubMed](#)]
59. Stern, A.L.; Burgos, E.; Salmon, L.; Cazzulo, J.J. Ribose 5-phosphate isomerase type B from *Trypanosoma cruzi*: Kinetic properties and site-directed mutagenesis reveal information about the reaction mechanism. *Biochem. J.* **2007**, *401*, 279–285. [[CrossRef](#)]
60. Riganti, C.; Gazzano, E.; Polimeni, M.; Aldieri, E.; Ghigo, D. The pentose phosphate pathway: An antioxidant defense and a crossroad in tumor cell fate. *Free Radic. Biol. Med.* **2012**, *53*, 421–436. [[CrossRef](#)]

61. Sørensen, K.I.; Hove-Jensen, B. Ribose catabolism of *Escherichia coli*: Characterization of the *rpiB* gene encoding ribose phosphate isomerase B and of the *rpiR* gene, which is involved in regulation of *rpiB* expression. *J. Bacteriol.* **1996**, *178*, 1003–1011. [[CrossRef](#)]
62. Zhang, R.-G.; Andersson, C.E.; Skarina, T.; Evdokimova, E.; Edwards, A.M.; Joachimiak, A.; Savchenko, A.; Mowbray, S.L. The 2.2 Å resolution structure of RpiB/AlsB from *Escherichia coli* illustrates a new approach to the ribose-5-phosphate isomerase reaction. *J. Mol. Biol.* **2003**, *332*, 1083–1094. [[CrossRef](#)] [[PubMed](#)]
63. Kaur, P.K.; Dinesh, N.; Soumya, N.; Babu, N.K.; Singh, S. Identification and characterization of a novel Ribose 5-phosphate isomerase B from *Leishmania donovani*. *Biochem. Biophys. Res. Commun.* **2012**, *421*, 51–56. [[CrossRef](#)] [[PubMed](#)]
64. Chou, Y.-T.; Chen, L.-Y.; Tsai, S.-L.; Tu, H.-C.; Lu, J.-W.; Ciou, S.-C.; Wang, H.-D.; Yuh, C.-H. Ribose-5-phosphate isomerase A overexpression promotes liver cancer development in transgenic zebrafish via activation of ERK and β -catenin pathways. *Carcinogenesis* **2018**, *40*, 461–473. [[CrossRef](#)] [[PubMed](#)]
65. Chou, Y.-T.; Jiang, J.-K.; Yang, M.-H.; Lu, J.-W.; Lin, H.-K.; Wang, H.-D.; Yuh, C.-H. Identification of a noncanonical function for ribose-5-phosphate isomerase A promotes colorectal cancer formation by stabilizing and activating β -catenin via a novel C-terminal domain. *PLoS Biol.* **2018**, *16*, e2003714. [[CrossRef](#)] [[PubMed](#)]
66. Heintze, J.; Costa, J.R.; Weber, M.; Ketteler, R. Ribose 5-phosphate isomerase inhibits LC3 processing and basal autophagy. *Cell. Signal.* **2016**, *28*, 1380–1388. [[CrossRef](#)] [[PubMed](#)]
67. Ying, H.; Kimmelman, A.C.; Lyssiotis, C.A.; Hua, S.; Chu, G.C.; Fletcher-Sananikone, E.; Locasale, J.W.; Son, J.; Zhang, H.; Coloff, J.L.; et al. Oncogenic Kras maintains pancreatic tumors through regulation of anabolic glucose metabolism. *Cell* **2012**, *149*, 656–670. [[CrossRef](#)]
68. Patra, K.C.; Hay, N. The pentose phosphate pathway and cancer. *Trends Biochem. Sci.* **2014**, *39*, 347–354. [[CrossRef](#)]
69. Gonzalez, S.N.; Valsecchi, W.M.; José, D.M.; Delfino, M.; Cazzulo, J.J.; Maugeri, D.; Delfino, J.M.; Cazzulo, J.J. Structure, kinetic characterization and subcellular localization of the two ribulose 5-phosphate epimerase isoenzymes from *Trypanosoma cruzi*. *PLoS ONE* **2017**, *12*, e0172405. [[CrossRef](#)]
70. Sobota, J.M.; Imlay, J.A. Iron enzyme ribulose-5-phosphate 3-epimerase in *Escherichia coli* is rapidly damaged by hydrogen peroxide but can be protected by manganese. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 5402–5407. [[CrossRef](#)]
71. Alfarouk, K.O.; Shayoub, M.E.A.; Muddathir, A.K.; Elhassan, G.O.; Bashir, A.H.H. Evolution of tumor metabolism might reflect carcinogenesis as a reverse evolution process (dismantling of multicellularity). *Cancers* **2011**, *3*, 3002–3017. [[CrossRef](#)]
72. Kabashima, T.; Kawaguchi, T.; Wadzinski, B.E.; Uyeda, K. Xylulose 5-phosphate mediates glucose-induced lipogenesis by xylulose 5-phosphate-activated protein phosphatase in rat liver. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 5107–5112. [[CrossRef](#)] [[PubMed](#)]
73. Liangpunsakul, S.; Ross, R.A.; Crabb, D.W. Activation of carbohydrate response element binding protein (ChREBP) by ethanol. *J. Investig. Med.* **2013**, *61*, 270. [[CrossRef](#)] [[PubMed](#)]
74. Meng, X.; Lu, X.; Li, Z.; Green, E.D.; Massa, H.; Trask, B.J.; Morris, C.A.; Keating, M.T. Complete physical map of the common deletion region in Williams syndrome and identification and characterization of three novel genes. *Hum. Genet.* **1998**, *103*, 590–599. [[CrossRef](#)] [[PubMed](#)]
75. Tong, X.; Zhao, F.; Mancuso, A.; Gruber, J.J.; Thompson, C.B. The glucose-responsive transcription factor ChREBP contributes to glucose-dependent anabolic synthesis and cell proliferation. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 21660–21665. [[CrossRef](#)] [[PubMed](#)]
76. Diolaiti, D.; McFerrin, L.; Carroll, P.A.; Eisenman, R.N. Functional interactions among members of the MAX and MLX transcriptional network during oncogenesis. *Biochim. Biophys. Acta* **2015**, *1849*, 484–500. [[CrossRef](#)] [[PubMed](#)]
77. Iizuka, K. The transcription factor carbohydrate-response element-binding protein (ChREBP): A possible link between metabolic disease and cancer. *Biochim. Biophys. Acta Mol. Basis Dis.* **2017**, *1863*, 474–485. [[CrossRef](#)]
78. Kawaguchi, T.; Takenoshita, M.; Kabashima, T.; Uyeda, K. Glucose and cAMP regulate the L-type pyruvate kinase gene by phosphorylation/dephosphorylation of the carbohydrate response element binding protein. *Proc. Natl. Acad. Sci. USA* **2001**, *98*, 13710–13715. [[CrossRef](#)] [[PubMed](#)]

79. Burke, S.J.; Collier, J.J.; Scott, D.K. cAMP opposes the glucose-mediated induction of the L-PK gene by preventing the recruitment of a complex containing ChREBP, HNF4 α , and CBP. *J. Am. Fed. Biol. Exp.* **2009**, *23*, 2855–2865. [[CrossRef](#)]
80. Miyamoto, K.; Ohta, H. Enzymatic decarboxylation of synthetic compounds. In *Future Directions in Biocatalysis*; Elsevier: Amsterdam, The Netherlands, 2007; pp. 305–343.
81. Jahromi, R.R.F.; Morris, P.; Martinez-Torres, R.J.; Dalby, P.A. Structural stability of *E. coli* transketolase to temperature and pH denaturation. *J. Biotechnol.* **2011**, *155*, 209–216. [[CrossRef](#)]
82. Datta, A.G.; Racker, E. Mechanism of action of transketolase. I. Properties of the crystalline yeast enzyme. *J. Biol. Chem.* **1961**, *236*, 617–623.
83. Combs, G.F. Thiamin. In *The Vitamins*; Elsevier: Amsterdam, The Netherlands, 2012; pp. 261–276.
84. Tseng, C.-W.; Kuo, W.-H.; Chan, S.-H.; Chan, H.-L.; Chang, K.-J.; Wang, L.-H. Transketolase Regulates the Metabolic Switch to Control Breast Cancer Cell Metastasis via the α -Ketoglutarate Signaling Pathway. *Cancer Res.* **2018**, *78*, 2799–2812. [[CrossRef](#)] [[PubMed](#)]
85. Raš, B.; Comin, B.; Puigjaner, J.; Brandes, J.L.; Creppy, E.; Saboureau, D.; Ennamany, R.; Paul Lee, W.-N.; Boros, L.G.; Cascante, M. Oxythiamine and dehydroepiandrosterone induce a G1 phase cycle arrest in Ehrlich's tumor cells through inhibition of the pentose cycle. *FEBS Lett.* **1999**, *456*, 113–118. [[CrossRef](#)]
86. Sica, D.A. Diuretics. In *Side Effects of Drugs Annual*; Elsevier: Amsterdam, The Netherlands, 2005; pp. 233–243.
87. Venkataraman, R.; Racker, E. Mechanism of action of transaldolase. I. Crystalization and properties of yeast enzyme. *J. Biol. Chem.* **1961**, *236*, 1876–1882. [[PubMed](#)]
88. Huang, S.-Y.; Zhang, Y.-H.P.; Zhong, J.-J. A thermostable recombinant transaldolase with high activity over a broad pH range. *Appl. Microbiol. Biotechnol.* **2012**, *93*, 2403–2410. [[CrossRef](#)] [[PubMed](#)]
89. Banki, K.; Hutter, E.; Colombo, E.; Gonchoroff, N.J.; Perl, A. Glutathione levels and sensitivity to apoptosis are regulated by changes in transaldolase expression. *J. Biol. Chem.* **1996**, *271*, 32994–33001. [[CrossRef](#)] [[PubMed](#)]
90. Heinrich, P.C.; Morris, H.P.; Weber, G. Behavior of transaldolase (EC 2.2.1.2) and transketolase (EC 2.2.1.1) Activities in normal, neoplastic, differentiating, and regenerating liver. *Cancer Res.* **1976**, *36*, 3189–3197. [[PubMed](#)]
91. Perl, A. The pathogenesis of transaldolase deficiency. *IUBMB Life* **2007**, *59*, 365–373. [[CrossRef](#)]
92. Ding, Y.; Gong, C.; Huang, D.; Chen, R.; Sui, P.; Lin, K.H.; Liang, G.; Yuan, L.; Xiang, H.; Chen, J.; et al. Synthetic lethality between HER2 and transaldolase in intrinsically resistant HER2-positive breast cancers. *Nat. Commun.* **2018**, *9*, 4274. [[CrossRef](#)]
93. Bennett, C.F.; Kwon, J.J.; Chen, C.; Russell, J.; Acosta, K.; Burnaevskiy, N.; Crane, M.M.; Bitto, A.; Vander Wende, H.; Simko, M.; et al. Transaldolase inhibition impairs mitochondrial respiration and induces a starvation-like longevity response in *Caenorhabditis elegans*. *PLoS Genet.* **2017**, *13*, e1006695. [[CrossRef](#)]
94. Ogawa, T.; Murakami, K.; Yoshino, M. Inhibition by fructose 1,6-bisphosphate of transaldolase from *Escherichia coli*. *FEMS Microbiol. Lett.* **2016**, *363*, fnw183. [[CrossRef](#)]
95. Light, S.H.; Anderson, W.F. Arabinose 5-phosphate covalently inhibits transaldolase. *J. Struct. Funct. Genom.* **2014**, *15*, 41–44. [[CrossRef](#)] [[PubMed](#)]
96. Egan, R.M.; Sable, H.Z. Transketolase Kinetics. The Slow Reconstitution of the Holoenzyme Is Due to Rate-Limiting Dimerization of the Subunits. *J. Biol. Chem.* **1981**, *256*, 4877–4883. [[PubMed](#)]
97. Ballatori, N.; Krance, S.M.; Notenboom, S.; Shi, S.; Tieu, K.; Hammond, C.L. Glutathione dysregulation and the etiology and progression of human diseases. *Biol. Chem.* **2009**, *390*, 191–214. [[CrossRef](#)] [[PubMed](#)]
98. Zhao, Y.; Seefeldt, T.; Chen, W.; Wang, X.; Matthees, D.; Hu, Y.; Guan, X. Effects of glutathione reductase inhibition on cellular thiol redox state and related systems. *Arch. Biochem. Biophys.* **2009**, *485*, 56–62. [[CrossRef](#)] [[PubMed](#)]
99. Jones, D.P.; Sies, H. The Redox Code. *Antioxid. Redox Signal.* **2015**, *23*, 734–746. [[CrossRef](#)] [[PubMed](#)]
100. Salbitani, G.; Bottone, C.; Carfagna, S. Determination of Reduced and Total Glutathione Content in Extremophilic Microalga *Galdieria phlegrea*. *Bio-Protocol* **2017**, *7*, e2372. [[CrossRef](#)]
101. Baird, A.-M.; O'Byrne, K.J.; Gray, S.G. Reactive Oxygen Species and Reactive Nitrogen Species in Epigenetic Modifications. In *Systems Biology of Free Radicals and Antioxidants*; Springer: Berlin/Heidelberg, Germany, 2014; pp. 437–455.
102. Valko, M.; Izakovic, M.; Mazur, M.; Rhodes, C.J.; Telser, J. Role of oxygen radicals in DNA damage and cancer incidence. *Mol. Cell. Biochem.* **2004**, *266*, 37–56. [[CrossRef](#)]

103. Ríos-Arrabal, S.; Artacho-Cordón, F.; León, J.; Román-Marinetto, E.; Del Mar Salinas-Asensio, M.; Calvente, I.; Núñez, M.I. Involvement of free radicals in breast cancer. *Springerplus* **2013**, *2*, 404. [[CrossRef](#)]
104. Dreher, D.; Junod, A.F. Role of oxygen free radicals in cancer development. *Eur. J. Cancer* **1996**, *32A*, 30–38. [[CrossRef](#)]
105. Liu-Smith, F.; Dellinger, R.; Meyskens, F.L., Jr. Updates of reactive oxygen species in melanoma etiology and progression. *Arch. Biochem. Biophys.* **2014**, *563*, 51–55. [[CrossRef](#)]
106. Augsten, M.; Sjöberg, E.; Frings, O.; Vorrink, S.U.; Frijhoff, J.; Olsson, E.; Borg, A.; Ostman, A. Cancer-Associated Fibroblasts Expressing CXCL14 Rely upon NOS1-Derived Nitric Oxide Signaling for Their Tumor-Supporting Properties. *Cancer Res.* **2014**, *74*, 2999–3010. [[CrossRef](#)] [[PubMed](#)]
107. Yang, W.; Zou, L.; Huang, C.; Lei, Y. Redox Regulation of Cancer Metastasis: Molecular Signaling and Therapeutic Opportunities. *Drug Dev. Res.* **2014**, *75*, 331–341. [[CrossRef](#)] [[PubMed](#)]
108. Kumari, S.; Badana, A.K.; Malla, R. Reactive Oxygen Species: A Key Constituent in Cancer Survival. *Biomark. Insights* **2018**, *13*. [[CrossRef](#)] [[PubMed](#)]
109. Simon, H.-U.; Haj-Yehia, A.; Levi-Schaffer, F. Role of reactive oxygen species (ROS) in apoptosis induction. *Apoptosis* **2000**, *5*, 415–418. [[CrossRef](#)] [[PubMed](#)]
110. Fubini, B.; Hubbard, A. Reactive oxygen species (ROS) and reactive nitrogen species (RNS) generation by silica in inflammation and fibrosis. *Free Radic. Biol. Med.* **2003**, *34*, 1507–1516. [[CrossRef](#)]
111. Ahn, H.J.; Kim, K.I.; Hoan, N.N.; Kim, C.H.; Moon, E.; Choi, K.S.; Yang, S.S.; Lee, J.-S. Targeting Cancer Cells with Reactive Oxygen and Nitrogen Species Generated by Atmospheric-Pressure Air Plasma. *PLoS ONE* **2014**, *9*, e86173. [[CrossRef](#)] [[PubMed](#)]
112. Di Meo, S.; Reed, T.T.; Venditti, P.; Victor, V.M. Role of ROS and RNS Sources in Physiological and Pathological Conditions. *Oxid. Med. Cell. Longev.* **2016**, *2016*, 1–44. [[CrossRef](#)] [[PubMed](#)]
113. He, L.; He, T.; Farrar, S.; Ji, L.; Liu, T.; Ma, X. Antioxidants Maintain Cellular Redox Homeostasis by Elimination of Reactive Oxygen Species. *Cell. Physiol. Biochem.* **2017**, *44*, 532–553. [[CrossRef](#)]
114. Kowalik, M.A.; Columbano, A.; Perra, A. Emerging Role of the Pentose Phosphate Pathway in Hepatocellular Carcinoma. *Front. Oncol.* **2017**, *7*, 87. [[CrossRef](#)]
115. James, E.L.; Michalek, R.D.; Pitiyage, G.N.; de Castro, A.M.; Vignola, K.S.; Jones, J.; Mohney, R.P.; Karoly, E.D.; Prime, S.S.; Parkinson, E.K. Senescent Human Fibroblasts Show Increased Glycolysis and Redox Homeostasis with Extracellular Metabolomes That Overlap with Those of Irreparable DNA Damage, Aging, and Disease. *J. Proteome Res.* **2015**, *14*, 1854–1871. [[CrossRef](#)]
116. Sousa, B.; Pereira, J.; Paredes, J.; Sousa, B.; Pereira, J.; Paredes, J. The Crosstalk Between Cell Adhesion and Cancer Metabolism. *Int. J. Mol. Sci.* **2019**, *20*, 1933. [[CrossRef](#)] [[PubMed](#)]
117. Vizán, P.; Alcarraz-Vizán, G.; Díaz-Moralli, S.; Solovjeva, O.N.; Frederiks, W.M.; Cascante, M. Modulation of pentose phosphate pathway during cell cycle progression in human colon adenocarcinoma cell line HT29. *Int. J. Cancer* **2009**, *124*, 2789–2796. [[CrossRef](#)]
118. Uyeda, K.; Yamashita, H.; Kawaguchi, T. Carbohydrate responsive element-binding protein (ChREBP): A key regulator of glucose metabolism and fat storage. *Biochem. Pharmacol.* **2002**, *63*, 2075–2080. [[CrossRef](#)]
119. Iizuka, K.; Wu, W.; Horikawa, Y.; Takeda, J. Role of glucose-6-phosphate and xylulose-5-phosphate in the regulation of glucose-stimulated gene expression in the pancreatic β cell line, INS-1E. *Endocr. J.* **2013**, *60*, 473–482. [[CrossRef](#)] [[PubMed](#)]
120. De Santis, M.C.; Porporato, P.E.; Martini, M.; Morandi, A. Signaling Pathways Regulating Redox Balance in Cancer Metabolism. *Front. Oncol.* **2018**, *8*, 126. [[CrossRef](#)] [[PubMed](#)]
121. Koceva-Chyła, A.; Jędrzejczak, M.; Skierski, J.; Kania, K.; Jóźwiak, Z. Mechanisms of induction of apoptosis by anthraquinone anticancer drugs aclarubicin and mitoxantrone in comparison with doxorubicin: Relation to drug cytotoxicity and caspase-3 activation. *Apoptosis* **2005**, *10*, 1497–1514. [[CrossRef](#)] [[PubMed](#)]
122. Ricci, M.S.; Zong, W.-X. Chemotherapeutic approaches for targeting cell death pathways. *Oncologist* **2006**, *11*, 342–357. [[CrossRef](#)]
123. Galluzzi, L.; Vitale, I.; Vacchelli, E.; Kroemer, G. Cell Death Signaling and Anticancer Therapy. *Front. Oncol.* **2011**, *1*, 5. [[CrossRef](#)]
124. Mansilla, S.; Llovera, L.; Portugal, J. Chemotherapeutic targeting of cell death pathways. *Anti-Cancer Agents Med. Chem.* **2012**, *12*, 226–238. [[CrossRef](#)]
125. Ocker, M.; Höpfner, M. Apoptosis-Modulating Drugs for Improved Cancer Therapy. *Eur. Surg. Res.* **2012**, *48*, 111–120. [[CrossRef](#)]

126. Alfarouk, K.O.; Stock, C.-M.; Taylor, S.; Walsh, M.; Muddathir, A.K.; Verduzco, D.; Bashir, A.H.H.; Mohammed, O.Y.; Elhassan, G.O.; Harguindey, S.; et al. Resistance to cancer chemotherapy: Failure in drug response from ADME to P-gp. *Cancer Cell Int.* **2015**, *15*, 71. [[CrossRef](#)] [[PubMed](#)]
127. Zaman, G.J.; Lankelma, J.; van Tellingen, O.; Beijnen, J.; Dekker, H.; Paulusma, C.; Oude Elferink, R.P.; Baas, F.; Borst, P. Role of glutathione in the export of compounds from cells by the multidrug-resistance-associated protein. *Proc. Natl. Acad. Sci. USA* **1995**, *92*, 7690–7694. [[CrossRef](#)] [[PubMed](#)]
128. Bagrij, T.; Klokouzas, A.; Hladky, S.B.; Barrand, M.A. Influences of glutathione on anionic substrate efflux in tumour cells expressing the multidrug resistance-associated protein, MRP1. *Biochem. Pharmacol.* **2001**, *62*, 199–206. [[CrossRef](#)]
129. Lai, L.; Tan, T.M.C. Role of glutathione in the multidrug resistance protein 4 (MRP4/ABCC4)-mediated efflux of cAMP and resistance to purine analogues. *Biochem. J.* **2002**, *361*, 497–503. [[CrossRef](#)]
130. Alfarouk, K.O.; Muddathir, A.K.; Shayoub, M.E.A. Tumor acidity as evolutionary spite. *Cancers* **2011**, *3*, 408–414. [[CrossRef](#)]
131. Boros, L.G.; Puigjaner, J.; Cascante, M.; Lee, W.N.P.; Brandes, J.L.; Bassilian, S.; Yusuf, F.I.; Williams, R.D.; Muscarella, P.; Melvin, W.S.; et al. Oxythiamine and dehydroepiandrosterone inhibit the nonoxidative synthesis of ribose and tumor cell proliferation. *Cancer Res.* **1997**, *57*, 4242–4248. [[PubMed](#)]
132. Alfarouk, K.O.; Ahmed, S.B.M.; Ahmed, A.; Elliott, R.L.; Ibrahim, M.E.; Ali, H.S.; Wales, C.C.; Nourwali, I.; Aljarbou, A.N.; Bashir, A.H.H.; et al. The interplay of dysregulated pH and electrolyte imbalance in cancer. *Cancers* **2020**, *12*, 898. [[CrossRef](#)]
133. Alfarouk, K.O.; Bashir, A.H.H. Diabetes mellitus type 2 through oncology lens. *Med. Hypotheses* **2011**, *76*, 761–762. [[CrossRef](#)]
134. Domagk, G.F.; Doering, K.M.; Chilla, R. Purification and Properties of Ribose-Phosphate Isomerase from *Candida utilis*. *Eur. J. Biochem.* **1973**, *38*, 259–264. [[CrossRef](#)]
135. Tsolas, O.; Horecker, B.L. Transaldolase. *Enzymes* **1972**, *7*, 259–280.



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).