

Implementation and Scalability of Physical Activity Interventions Delivered Within Primary Care: A Narrative Review

OSINAIKE, Jimi <<http://orcid.org/0000-0001-8505-243X>>, MYERS, Anna <<http://orcid.org/0000-0001-6432-8628>>, LOWE, Anna <<http://orcid.org/0000-0001-5297-8957>>, COPELAND, Robert <<http://orcid.org/0000-0002-4147-5876>> and HARDCASTLE, Sarah <<http://orcid.org/0000-0001-8378-3781>>

Available from Sheffield Hallam University Research Archive (SHURA) at:
<https://shura.shu.ac.uk/34058/>

This document is the Published Version [VoR]

Citation:

OSINAIKE, Jimi, MYERS, Anna, LOWE, Anna, COPELAND, Robert and HARDCASTLE, Sarah (2024). Implementation and Scalability of Physical Activity Interventions Delivered Within Primary Care: A Narrative Review. *Lifestyle Medicine*, 5 (4). [Article]

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

REVIEW OPEN ACCESS

Implementation and Scalability of Physical Activity Interventions Delivered Within Primary Care: A Narrative Review

Jimi Osinaike¹  | Anna Myers¹  | Anna Lowe²  | Robert J Copeland^{1,2}  | Sarah J Hardcastle¹ 

¹Academy of Sports and Physical Activity, Sheffield Hallam University, Collegiate Crescent, Sheffield, UK | ²Advanced Wellbeing Research Centre, Sheffield Hallam University, Sheffield, UK

Correspondence: Jimi Osinaike (J.Osinaike@shu.ac.uk)

Received: 4 October 2023 | **Revised:** 28 June 2024 | **Accepted:** 12 July 2024

Funding: The authors received no specific funding for this work.

Keywords: implementation | physical activity | primary care | scalability

ABSTRACT

Primary Aim: To describe the features of implementation in the setting of primary care (PC) for physical activity (PA) interventions that improved total and moderate to vigorous PA (MVPA).

Secondary Aim: To assess the scalability potential of effective PC PA interventions.

Method: A comprehensive search was conducted across multiple electronic databases to identify relevant studies published between 2012 and 2023. Implementation-related features were extracted, and the scalability potential of effective PC PA interventions was assessed using the Intervention Scalability Tool (ISAT) as a framework.

Results: Twenty-six studies met the inclusion criteria after screening. Analysis of the implementation-related features revealed that effective PC PA interventions showed promising reach with a mean participation rate of 43%. Effective PC PA interventions that increased PA were mostly delivered by nurses and were underpinned by the behavioural change techniques (BCTs) of goal setting, feedback and self-monitoring. The scalability assessment revealed that remote-based interventions and those delivered by nurses had moderate to high scores in the scalability domains of fidelity, reach and acceptability, delivery setting and workforce, implementation infrastructure and sustainability.

Conclusion: PC PA interventions, whether delivered remotely or face-to-face, show promise for increasing PA, especially when incorporating BCTs like goal setting and feedback. Using the ISAT, most interventions displayed promise for implementation at scale, but further examination is needed concerning the delivery setting, implementation infrastructure and sustainability of these interventions. Remote-based (automated PA advice and mailed instructions) PA interventions and those delivered in contact with a nurse showed high scalability potential. Further work that examines the acceptability and effectiveness of other healthcare professionals in intervention delivery would be worthwhile, and more work is needed to assess the utility and effectiveness of remote-based PA interventions in PC.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Author(s). *Lifestyle Medicine* published by John Wiley & Sons Ltd.

1 | Introduction

The evidence supporting physical activity (PA) for the management and prevention of non-communicable diseases, such as diabetes, hypertension and depression, is compelling [1]. Despite this, approximately 25% of adults globally are insufficiently physically active [2]. Consequently, the World Health Organization (WHO) has set a target to reduce physical inactivity by 15% by 2030 [3]. This target includes a recommendation to integrate PA promotion into primary care (PC) settings [3]. In the United Kingdom, adults visit their general practitioner (GP) approximately five times per year, with half of these contacts related to the management of long-term health conditions [4]. The PC setting is an opportunity to reach a large segment of the population with health promotion efforts.

PA delivered within PC has been identified as a cost-effective strategy to reduce the incidence of non-communicable diseases [5]. According to the National Institute for Health and Care Excellence (NICE), PA promotion in PC can range from verbal advice, discussion, negotiation or encouragement, with or without written or other support [6]. Several reviews have explored the effectiveness of brief PA interventions in the PC setting [7–10]. These reviews found that brief PA advice delivered within PC (either in-person visits or telephone consultations) with or without other supplements (e.g., written prescription, information pamphlet, use of pedometers) have shown were effective in improving PA [7–10]. Despite this, the implementation of interventions into routine practice in PC has been a challenge. In the United Kingdom, only 15% of GPs report providing PA advice and about 35% of patients report receiving PA advice [11–13]. Barriers are reported at individual and organisational levels [14]. At the individual level, health professionals report insufficient confidence and knowledge about PA promotion [15, 16]. Organisational-related barriers relate to a lack of time, lack of training and technical support (e.g., provision of high-quality print materials to reinforce verbal messages) [17–19]. There is a need for practical guidance on how to effectively scale up PC PA interventions. The PRACTIS (PRACTical planning for Implementation and Scale-up) guide addresses this gap by detailing how to anticipate and address potential barriers and facilitators to effective implementation and scale-up [20]. It begins by examining the features of the implementation setting, which include the intervention population (individual level), implementers (provider level), delivery setting (organisational level) and intervention factors (intervention design/underlying principles, delivery format, duration, resources required, fidelity/adaptation). Together these factors determine outcomes relating to intervention reach, acceptability, adoption, delivery and sustainability in practice. A clear understanding of these features for PC will help to promote planning and accountability for future implementation of PA interventions and could enhance scale-up efforts.

Whilst some of the related features of implementation, such as the number of contacts needed to deliver effective PA interventions in PC, have been previously described [10], the intervention reach, acceptability, intervention content, fidelity, adaptation and scale-up of PC PA interventions have not been systematically explored. This narrative review therefore describes the features of implementation in the setting of PC for PA interventions that improved total and moderate to vigorous PA (MVPA). The review also

provides a structured, systematic and comprehensive assessment of the scalability potential of effective PC PA interventions.

2 | Methods

2.1. Eligibility Criteria

Studies were selected based on the following criteria: (i) Participants: Adults aged ≥ 18 years. (ii) Recruitment: through PC (i.e., recruited from patients attending or registered with a general practice, family practice or PC clinic). The PC settings were broadly defined as the first point of contact in the healthcare system providing accessible and coordinated care. (iii) Intervention: PA interventions delivered in person or remotely (e.g., leaflets, email or phone calls). Trials were included if they were embedded within PC (i.e., delivered within or out of PC (e.g., interventions involving written prescriptions sent to patients via email) rather than referred or outsourced to a community or local PA service). We excluded studies examining multifactorial interventions, such as promoting dietary modification in addition to PA. Trials evaluating exercise referral schemes were also excluded because PC health professionals would only be acting as referral mechanisms rather than being directly involved in delivering interventions. (iv) Design: Randomised controlled trials. (v) Outcome: PA intervention designed to increase time spent in light intensity physical activity, MVPA and total physical activity (e.g., metabolic equivalent (MET), steps/day, minutes in total PA). (vi) No restrictions were made on the method used to assess PA with data from self-report (subjective) and device measures (objective) included.

Articles not available in the English Language, unpublished studies, protocols and dissertations were excluded.

2.1 | Search Strategies

The search approach combined intervention and setting keywords with pre-existing randomised controlled trial filters, and this was developed and evaluated in Medline and across the following databases: PsycINFO, CINAHL and The Cochrane Library (CENTRAL) (see Supplementary file 1 for search strategies). The search was conducted between 2012 and 2023 to access recent evidence on PA promotion within PC. In addition, this timeframe allows us to ascertain the most recent developments of interventions since the publication of the NICE guidelines on PA promotion in PC in 2012 [6]. We used RefWorks [21] (a reference management tool) and Microsoft Excel (version 2403) to collect and organise references. As this review was part of a Ph.D. project, a single-handed screening was chosen because it allows for the completion of screening within a limited time frame. The first author (JO) screened the titles and abstracts using the specified inclusion criteria. Two other authors (AM and AL) then reviewed 10% of the titles and abstracts to ensure consistency of screening.

2.2 | Data Extraction

Implementation characteristics were extracted from studies based on three implementation characteristics: (i) individual level: The

size of the target population, eligibility criteria, participation and attrition rates and interventionists; (ii) intervention factors: theories/models, delivery format, duration, outcomes, number of interventions contact resources required (Fitbit, diaries, etc.) and intervention fidelity and or adaptations; (iii) Effectiveness: This was defined in terms of the measure of PA (MVPA or total PA) and how it was assessed (device measured or self-reported).

2.3 | Risk of Bias and Quality of Evidence Assessment

The risk of bias (RoB) was assessed using the Cochrane tool (RoB 2) [22]. The RoB was assessed single-handedly by the first author with 10% of studies independently assessed by a second reviewer and checked for consistency. Any disagreement between the reviewers was discussed with a third reviewer and afterwards resolved through consensus by referring to the full text.

2.4 | Scalability Assessment

For the purposes of the review, scalability is defined as the ability of a health intervention shown to be efficacious on a small scale and or under controlled conditions to be expanded under real-world conditions to reach a greater proportion of the eligible population, while retaining effectiveness [23]. While there are several scalability assessment frameworks/tools, most do not provide a score to compare interventions for scalability [24]. The Intervention Scalability Assessment Tool (ISAT) [25] was chosen because it contains scalability domains consistent with what has been reported in the scalability literature [24–26]. In addition, the ISAT was selected because it offers a practical guide to conduct a structured, systematic and comprehensive scalability assessment.

Table 1 outlines the ISAT domains and the criteria used to assess questions within each scalability domain. The ISAT is designed to evaluate the scalability of interventions by considering various factors across 10 domains. These domains are divided into two parts: Part A focuses on the context of the intervention, while Part B addresses implementation planning and feasibility. After completing Parts A and B, Part C provides a summative assessment based on scoring from the previous sections. This assessment generates a radar plot to visualise the intervention's readiness for scale-up and helps stakeholders make informed decisions about scalability. Final recommendations can be merits scale up, promising but further information/planning is warranted or does not merit scale up.

All sections of the included studies, such as the introduction, methods, results, discussion, conclusions and appendices/supplementary material, were thoroughly reviewed for relevant scalability data. Each domain includes optional questions at the end to assess readiness based on the provided information. These questions help identify strengths and weaknesses across domains. Additionally, the ISAT offers preparatory questions to consider before answering the readiness questions. Questions in the ISAT are scored from 0 to 3, with 0 representing 'not at all scalable' and 3 indicating 'scalable to a large extent'. If a question is not applicable, it can be marked as 'N/A' and not scored. To

calculate the final score for each domain, the average score across the questions (if multiple) is taken.

In the assessment for scalability, the authors initially determined the appropriateness of ISAT domain questions needed to evaluate the scalability of the included studies. Following this, authors (JO, AM and SH) deliberated on each item to achieve consensus regarding the interpretation of the domain questions. While the ISAT offers preparatory questions, they are not directly relevant to PC PA interventions. As a result, suitable criteria were developed (see Table 1) for the context (i.e., PA interventions in PC) when addressing the scalability domain questions. If information was absent to answer applicable questions, these items were deemed indeterminate (IN). After reaching a consensus on question suitability, interpretations and scoring criteria, the first author (JO) extracted relevant information from effective intervention studies and assessed scalability based on the agreed interpretation and criteria. Subsequently, the authors (JO, AM and SH) reviewed the assessment conducted by the first author (JO) and reached a consensus on the scalability assessment.

3 | Results

A total of 3224 studies were identified from searches and 1580 titles, and abstracts were screened after removing duplicates. Twenty-six studies met the inclusion criteria (see Figure 1 for the PRISMA flow diagram of studies included in the review).

3.1 | Quality Assessment

Thirteen (50%) studies in this review showed some concerns in the domain-assessing deviations from intended interventions. Overall, most studies in this review ($n = 21$; 81%) were judged to have some concerns based on the Cochrane RoB tool [22] (see Supplementary file 2).

3.2 | Recruitment Strategies and Reach of the Included Studies

Table 2 describes the study characteristics. The recruitment strategies of participants were either opportunistic (during routine visits to PC) or systematic (mail invites, flyers or phone calls). Twenty-four studies reported on the recruitment processes. Eighteen (69%) studies employed a systematic approach to participant recruitment [27–44]. Systematic recruitment was the dominant method in 11 (85%) [28, 29, 32–34, 36, 38–40, 43, 44] out of the 13 studies that demonstrated effectiveness in improving PA outcomes. The participation rate (PR), a measure of 'reach' was higher with systematic recruitment (15–80%) when compared to opportunistic (12–53%). Amongst the 13 studies that showed effectiveness in improving PA outcomes, eight (62%) [32, 34, 36, 38, 40, 43–45] reported on PR and this was in the range of 18–80% (mean PR = 43%). These interventions [32, 34, 36, 38, 40, 43–45] were all characterised by systematic recruitment, with only one intervention having both opportunistic and systematic recruitment [40]. The largest participation rate (80%) [43] involved participants with diabetes who were recruited systematically.

TABLE 1 | Description of ISAT domains and criteria [25].

Domain	Domain questions	ISAT scoring consideration	Authors scoring considerations ^a
The problem (A1)	Is the problem of sufficient concern to warrant the scale-up of an (the) intervention/programme to address it?	What is the problem and who does it affect? Describe the nature and scope of the problem using epidemiological data. How does the problem impact on the health of the population? What is the current practice to address the issue?	This domain will be scored 3 (largely scalable) considering physical inactivity is a global public health issue and that PA has been shown to improve health outcomes.
The intervention (A2)	Will the outcomes delivered by this intervention address the needs of the target group (and/or) problem?	Describe the aims/objectives and intended outcomes of the intervention proposed for scale-up. Describe the key elements of the primary care PA intervention proposed for scale-up (including the process of delivery). What are the key intervention components? (e.g., frequency and intensity of the intervention, etc.).	This domain will be scored 3 (largely scalable) considering that the health outcomes of the PA intervention (such as prevention and management of non-communicable diseases) will address the health needs of participants
Strategic and political context (A3)	Is addressing the problem consistent with policy/strategic directions or priorities? Will scaling up this programme/intervention up be strategically useful to funders/funding agencies?	Is addressing the problem consistent with national, state or regional policy directions or priorities? Is addressing the problem an identified need of funding agencies? How well will the intervention proposed for scale-up align with the broader strategic and/or political context?	This domain will be scored 3 (largely scalable) across all interventions. This is because PA promotion within the PC care setting is a strategic priority of health organisations such as the WHO [3]

(Continues)

TABLE 1 | (Continued)

Domain	Domain questions	ISAT scoring consideration	Authors scoring considerations ^a
Evidence of effectiveness (A4)	Is there compelling evidence from the literature to indicate that the intervention is effective in addressing the problem in the target population?	A statistically significant difference, though a good start, is not necessarily a difference of policy/clinical significance. What was the size of the intervention effect (if known)? (Mean difference, relative risk, odds ratio, hazard ratio, sensitivity, specificity and statistical significance). Is the effect size of the intervention meaningful from a population health policy perspective? What is the strength of evidence of effectiveness for the intervention in addressing the problem described in domains A1 and A2, based on the literature?	This was scored by comparing with existing evidence in the primary care PA intervention literature. Recent systematic reviews and meta-analysis [10] found that PA interventions delivered or prompted by health professionals in PC increased MVPA by 14 min/week. Thus, our scoring was: Less than 30 min/week increase in MVPA (scored 1) 30–60 min/week increase in MVPA (scored 2) Greater than 60 min/week increase in MVPA (scored 3)
Intervention costs (A5)	Is there evidence that the benefits of the intervention exceeded the costs?	What were the intervention costs reported (if available)? Consider costs associated with start-up (e.g., building infrastructure, conducting training), costs associated with ongoing delivery as well as cost per participant or cost per unit of outcome. Was there any evidence of benefit outweighing the costs?	

(Continues)

TABLE 1 | (Continued)

Domain	Domain questions	ISAT scoring consideration	Authors scoring considerations ^a
Fidelity and adaptation (B1)	<p>Will the core components of the scaled-up intervention be consistent with what was previously shown to be effective?</p> <p>If the core components of the programme/intervention are to be modified from its original form during scale-up, will the impact of the modification likely be favourable? (NA).</p> <p>Can programme fidelity be monitored and/or maintained when implemented at scale?</p>	<p>Will there be any changes and/or adaptations made to the intervention from what was described in Domain A2 if the intervention is scaled up?</p> <p>Are those changes and/or adaptations likely to have any impact on the intended outcomes of the intervention as described in Domain A2?</p> <p>How will intervention fidelity be monitored and maintained?</p>	<p>It is likely that PA interventions delivered face-to-face and underpinned by BCTs (e.g., goal setting, self-efficacy, addressing barriers, etc.) or delivered using MI techniques will be time consuming and may not be consistently delivered per protocol when implemented at scale.</p> <p>Maintaining fidelity monitoring: Consider fidelity monitoring via direct assessment (i.e., observer report using video or audio tape) and indirect assessment (i.e., self-report using pencil and paper surveys, or technology based. It will be more feasible to monitor fidelity in remote-based PA interventions such as automated PA advice.</p> <p>Intervention involving face-to-face contact with the healthcare provider is likely not to be delivered per protocol because of the high chance of human error. On the other hand, interventions delivered through automated PA advice reduce the chance of human error.</p> <p>Based on the number of intervention components, it will be easier to monitor the intervention fidelity of single components when compared to multi-component. For instance, it will be much easier to monitor fidelity of an intervention involving only face-to-face PA advice compared to interventions involving PA advice and follow-up supports such as telephone calls, use of pedometers and referral to a walking group.</p> <p>Interventions without direct human involvement, like automated PA advice, are more likely to be delivered per protocol due to the absence of human interference that could affect fidelity.</p>
Reach and acceptability (B2)	<p>Does the selected intervention have the potential to reach the intended target population at scale?</p> <p>Is the selected intervention likely to be acceptable to the target population?</p>	<p>How will the intended target group be identified and recruited at scale?</p>	<p>Consider the attrition rate of the intervention group. The attrition rate (AR) has been reported as an objective measure of intervention acceptability. Thus, an intervention with an AR less than 10% (3 = largely scalable); 10–20% (2 = somewhat scalable); greater than 20% (1 = scalable to some extent)</p> <p>Identifying target population: Consider the recruitment strategies used to recruit participants for the PC PA intervention. The use of mixed recruitment strategies (opportunistic and systematic) is more likely to recruit a more diverse population. The use of opportunistic recruitment alone will be scored 2. Systematic recruitment alone: scored 1</p>

(Continues)

TABLE 1 | (Continued)

Domain	Domain questions	ISAT scoring consideration	Authors scoring considerations ^a
Delivery setting and workforce (B3)	<p>Is the delivery setting(s) selected to deliver the programme at a scale consistent with that used in previous studies?</p> <p>Is the delivery workforce selected to deliver the programme at a scale consistent with that used in previous studies?</p> <p>Is the intervention likely to be acceptable to the delivery workforce involved in its delivery at scale?</p> <p>If the intervention requires integration into existing organisational or community structures, how likely is it to be acceptable?</p>	<p>Who will deliver the intervention at scale?</p> <p>Is the intervention likely to be acceptable to the delivery workforce involved if implemented at scale?</p> <p>Is the intervention likely to be acceptable to the delivery workforce involved if implemented at scale?</p> <p>Who will deliver the intervention at scale?</p> <p>Are there any foreseeable facilitators and/or barriers for the delivery process? Will the intervention burden health workers by adding to their workload?</p>	<p>Consider the delivery workforce required for the implementation of the PC PA intervention. Are they front-line PC health practitioner (GPs, practice nurses) or allied professionals working in primary care (Health counsellors, exercise professionals and health educators)?</p> <p>Consider the duration and number of contacts with the interventionists involved in the delivery of the primary care PA intervention. Duration of PA intervention has been reported as a major determinant of the promotion of PA within primary care. A very brief intervention (less than 5 min) will be scored (3) because it is likely to be more acceptable to front-line primary care practitioners such as GPs. Contact: Remote-based PA interventions involving no contact are more likely to be acceptable to the delivery workforce. This will be scored 3.</p>
Implementation infrastructure (B4)	<p>Are the implementation infrastructure requirements of the intervention/programme feasible for scale-up?</p>	<p>Describe the infrastructure requirements for the delivery of the programme/ intervention, that is, classrooms, clinic facilities, sporting fields, community centres, IT equipment, etc.</p> <p>Describe the operational requirements for delivery of the intervention, that is, training, education, monitoring and feedback systems, accreditation processes, etc.</p> <p>Are there any foreseeable facilitators and/or barriers to building implementation infrastructure as part of the scale-up process?</p> <p>Facilitators or barriers in this case can be in terms of acceptability to the workforce, changes to practice, workload, etc.</p>	<p>It is likely that interventions involving face-to-face contact with an interventionist or the use of telephone calls are likely to involve training of health personnel and other operational and infrastructural demands.</p> <p>The use of allied health professionals could also mean the need for more funds to employ them.</p> <p>Use of digital support tools such as Fitbit and signposting to PA websites requires less infrastructural support compared to follow-up telephone calls.</p> <p>PA interventions delivered solely remotely (no face-to-face contact with an interventionist) such as use of sending automated PA advice require little infrastructural support.</p> <p>Group sessions are likely to require more operational requirements compared to individual sessions.</p>

(Continues)

TABLE 1 | (Continued)

Domain	Domain questions	ISAT scoring consideration	Authors scoring considerations ^a
Sustainability (B5)	Is the level of integration of the intervention into delivery settings required for implementation at scale sustainable?	What level of integration into existing service delivery settings or organisations will the intervention require if scaled up? Also, consider whether the level of integration is feasible or sustainable.	It will be more likely to sustain intervention involving less infrastructure resources. For example, an intervention involving use of pamphlets is more sustainable than those involving telephone calls or face-to-face interactions with a healthcare provider.
	Is the level of resourcing required to implement the intervention at scale sustainable?	If the intervention is implemented at scale, will it require a large commitment of funds (initial or ongoing)?	It is likely that interventions involving health professionals (such as health counsellors and fitness coaches) will require additional resources in terms of employment and integration into existing PC practice.
	Is the delivery workforce selected for implementation at scale sustainable?	Is the proposed delivery workforce required for implementation at scale sustainable (e.g., financially and/or in terms of supply)?	

Note: 3, largely scalable; 2, somewhat scalable; 1, scalable to some extent; N/A, not applicable.

^aNot part of the ISAT.




FIGURE 1 | PRISMA flow diagram.

3.3 | PA Intervention Contents, Contact, Mode of Delivery and BCTs

Twenty-one (81%) interventions in this review were delivered face-to-face with additional intervention supports (use of activity monitors for self-monitoring, written prescription, walking plan, online and community PA supports). Only five interventions (19%) [27, 28, 35, 36, 46] were delivered remotely. These studies involved the use of an activity device/pedometer for PA self-monitoring, written prescription, automated PA advice, telephone calls, email communication, individualised text message feedback and the use of PA logs or action plans as additional intervention supports. Twelve (46%) studies, [33, 35, 36, 40–44, 47–50] reported on the duration of face-to-face contact with interventionists. The mean time per contact was 26 min for interventions involving face-to-face contact with the interventionist, whilst group-based face-to-face contacts averaged 2 h. The content (see Table 2) of the face-to-face PA counselling included information on recommended PA guidelines, the benefits of PA, health implications of physical inactivity, online and community PA support, goal setting, action planning and feedback from step count or PA questionnaire.

Different types of PC PA interventions were effective including remotely delivered [28, 36] and face-to-face [29, 32–34, 36, 38–

40, 43–45, 51] interventions with varying components such as the use of pedometers for self-monitoring, written PA prescription, follow up telephone calls, use of PA diaries and step charts. Goal setting, feedback and self-monitoring were the most common BCTs reported, and this was evident in 15 (94%) [29, 30, 32, 33, 36, 38, 40–46, 50, 52] out of the 16 studies that reported on BCTs. Among the 13 studies [28, 29, 32–34, 36, 38–40, 43–45, 51] that increased PA, eight (62%) [28, 29, 33, 36, 40, 43, 45, 53] reported the BCTs utilised, while five (38%) [32, 34, 39, 40, 45] outlined the intervention delivery techniques employed. The most prevalent BCTs in these effective interventions were self-monitoring, feedback and goal setting. Effective interventions without a specified BCT made use of the 5As (assess, advise, agree, assist and arrange) of the health promotion delivery model and motivational interviewing (MI).

3.4 | Interventionists

Most interventions were delivered by nurses ($n = 9$) [27, 29, 33, 34, 36, 42, 44, 49, 54] or researchers ($n = 8$) [35, 37, 39–41, 45, 51, 55]. Other interventionists included primary care physicians (PCPs) ($n = 3$) [32, 45, 47], counsellors ($n = 2$), [32, 56], exercise specialists ($n = 2$) [30, 39] and other healthcare providers ($n = 2$) [43, 48]. Effective interventions were characterised by either contact with

TABLE 2 | Characteristics of included studies.

Author, Year/ country	Eligibility criteria	Intervention component and content	Contact duration and number of contacts	Interventionists and PA		PA outcome measure	Participation rate	Attrition rate/ follow-up.	Recruitment procedure	Theory/ Behavioural change technique (BCT)/model	Fidelity
				intervention components delivered							
Reed et al., 2019/USA	19 and 65 years, not meeting recommended PA levels/week	Fitbit and electronic PA action PA and weekly text message reminder plan.	Action plan sent monthly.	Practice nurse set up the Fitbit, sent. Electronic PA action plans and feedback.	Step Count using Fitbit.	15%	10%/12 weeks	Telephone, text message invite following PA screening In routine consultation	5As health promotion model	NR	
Agarwal et al., 2020/Canada	18–79, attended a periodic health assessment.	Face-to-face PA consultation and support (online, community PA prescription	One physical contact at baseline length of 2–5 min	PCP reviewed automated PA prescription sent a customised toolkit with tailored messaging	Total PA measured subjectively	NR	18.9%/24 weeks	During a routine periodic assessment	Health Action Process	Only 25% of the patients received a tool kit	
Riera-Sampol et al., 2021/Spain	age 35–75, with at least two cardiovas- cular risk factors were included Patients with dementia or cognitive impairment, with major active illness, were excluded	MI individualised prescription of PA using health assets	At baseline and four follow-up visits (2, 6, 9 and 12 months). Duration: NR	Primary care nurses assessed and carried out MI	Adherence to at least 150 min of total PA weekly and self-reported PA levels using the IPAQ	NR	Intervention group (44%) control (8%)		Trans- theoretical model of change, MI	NR	

(Continues)

TABLE 2 | (Continued)

Author, Year/ country	Eligibility criteria	Interventionists				PA outcome measure	Participation rate	Attrition rate/ follow-up.	Recruitment procedure	Theory/ Behavioural change technique (BCT)/model	Fidelity
		Intervention component and content	Contact duration and number of contacts	Intervention components delivered	PA outcome measure						
Gao et al., 2015/USA	one primary care visit in the 12 months prior to screening, at least 18 years overweight or obese	Face-to-face PA counselling and electronic feedback reports.	Nine electronic PA feedback	Mailed expert feedback PA counselling, PA self-help booklet and newsletter sent following filled postal PA readiness questionnaire	Spent in moderate intensity PA using accelerometer	43%	13% and 16% at 6 and 12 months, respec- tively/6 and 12 months	Phone calls following medical record screening	NR	NR	NR
Hardeman et al., 2020/UK	40 and 74 years, no diagnosis of vascular disease and not on a care pathway for known risk factors	Face-to-face discussion, provision of a pedometer and accelerometer, a Step Chart for self-monitoring and a Step it Up Booklet.	One face-to-face meeting at baseline Lasting 3 min and 9 s	Healthcare practitioner (HCP) (Specific HCP NR) Provided PA counselling	Total PA and step count (number of minutes per day spent in MVPA measured objectively	17%	16%/12 weeks.	Mail Invite	Goal setting, action planning, feedback and self- monitoring	Goal setting, action planning, feedback and self- monitoring	60%, (9 out of 15 components of the BCT were delivered at 3 months
Harris et al., 2015/UK	60–74 years and had no contra- indications to increasing PA	PA consultations incorporating pedometer step-count and accelerometer PA intensity feedback, and an individual PA diary and walking plan	Four face-to-face contact Duration NR	Nurses Provided PA counselling and supported with the use of pedometers	Time spent in MVPA measured objectively	NR	3% and 6% lost to follow-up at 3 and 12 months, respectively	Mailed Invitation following GP screening	Goal setting, self- monitoring, building self-efficacy and social support, overcoming barriers, preventing relapses and building lasting habits	Goal setting, self- monitoring, building self-efficacy and social support, overcoming barriers, preventing relapses and building lasting habits	NR

(Continues)

TABLE 2 | (Continued)

Author, Year/ country	Eligibility criteria	Intervention component and content	Contact duration and number of contacts	Interventionists and PA		PA outcome measure	Participation rate	Attrition rate/ follow-up.	Recruitment procedure	Theory/ Behavioural change technique (BCT)/model	Fidelity
				Intervention components delivered	GP trainee (who was the researcher) carried out the MI calls						
Heron et al., 2014/UK	35–75 years, inactive based on a questionnaire No contra- indications to PA	Group 1: information on their baseline step count, leaflet on PA, follow-up MI telephone Group 2: Group 1 intervention but with a specific, uniform daily step-count goal (2500 steps/day above baseline)	At baseline, with four follow-up telephone calls	GP trainee (who was the researcher) carried out the MI calls	Total PA pedometer step count	NR	5%/12 weeks	Routine consultation	Theory of planned behaviour’ 5As’ model BCT (establishing rapport, agenda setting, assessing importance and confidence, scaling questions, pros and cons, brainstorming solutions, negotiating attainable goals and follow-up)	NR	
Hinrichs et al., 2016/Switzer- land	70 and older, diagnosed with at least one chronic disease (DM, osteoarthritis, etc.), report mobility consultations	Exercise therapist PA consultation with GP motivational support	Eight consultations with the exercise specialist (five at the GP practice and three over- the-phone consultations) Duration of Consultation NR.	Exercise therapist	Pedometer PA step count/day	65%.	22% and 39% at 6 and 12 months, respectively	Mailed invitation following medical record screening	NR	NR	

(Continues)

TABLE 2 | (Continued)

Author, Year/ country	Eligibility criteria	Intervention component and content	Contact duration and number of contacts	Interventionists and PA components delivered			PA outcome measure	Participation rate	Attrition rate/ follow-up.	Recruitment procedure	Theory/ Behavioural change technique (BCT)/model	Fidelity
Kolt et al., 2012/New Zealand	65 or older, ability to walk and freedom from health conditions that contraindicate participation in PA	Initial face-to-face then followed up telephone counselling sessions by trained physical activity counsellors and Pedometers	One initial counselling At baseline in PC and three telephone counselling Duration of physician initial counselling NR Telephone counselling by trained counsellors lasted 10-30 min	PCP and PA counsellors delivered PA counselling	Moderate PA (MPA) inten- sity/subjective.	57%	13% and 15% at 3/4 and 12 months, respectively	Mail to patients	Trans theoretical model of behaviour change and MI	NR		
Mutrie et al., 2012/UK	Living independently and not meeting PA recommenda- tions	PA consultations and supports (PA booklet and pedometer)	Two face-to-face contacts lasting an average of 30 min	Nurses delivered the consultations and fitted the pedometer	Pedometer and ActivPal (monitor) recorded the average daily step count	NR.	10% of all randomised parti- pants/24 weeks	Letter sent following screening of medical records	Cognitive model of behaviour change	NR		
Yaacob et al., 2018/Malaysia	Aged 35–45 years determined to lead a sedentary lifestyle on the 7-day physical activity questionnaire and who have no chronic illness	Pedometer and support (log diaries and support groups, follow-up monthly meeting)	Four face-to-face contacts	Researcher provided PA counselling and -follow-up meetings	Total PA level (step/day) measured objectively	NR	29% in both intervention and control groups/12 weeks	NR	NR	NR		

(Continues)

TABLE 2 | (Continued)

Author, Year/ country	Eligibility criteria	Intervention component and content	Contact duration and number of contacts	Interventionists and PA		PA outcome measure	Participation rate	Attrition rate/ follow-up.	Recruitment procedure	Theory/ Behavioural change technique (BCT)/model	Fidelity
				Intervention components delivered	PA intervention components						
van der Weegen et al., 2015/The Netherlands	40 and 70 years old, type 2DM or COPD, BMI >25 and not having at least 30 min of MVPA patient with coexisting medical conditions were excluded	Group 1: participants tool (Personal Activity Monitor) and self-management support programme (SSP) (four individual consultations) Group 2: SSP Group 3: Usual care	Four face-to-face contacts Duration: NR	Nurses delivered the PA counselling		Average minutes per day in MVPA with the PA monitor	37%.	Group 1: 14% Group 2: 12% Group 3 (usual care): 9%/12 weeks.	Mail invite	5As model	Group 1: 12 Participants did not receive the minimal intervention (at least three consultations as intended Group 2: Seven participants did not receive the minimal intervention as intended
Vetrovsky et al., 2018/Czech Republic	Over 18 years of age, were physically inactive	Pedometer group (PED): given a pedometer and email (PEMAIL): Pedometer and email communication based on self-monitoring, action planning, goal setting and personalised feedback.	E-mail counselling was sent on average six times	Researcher sent email motivational counselling		Change in mean total daily step count	49%.	No attrition/12 weeks	During routine practice, screening	Based on MI principles BCT self-monitoring, action planning, goal setting and personalised feedback	

(Continues)

TABLE 2 | (Continued)

Author, Year/ country	Eligibility criteria	Intervention component and content	Contact duration and number of contacts	Interventionists			PA outcome measure	Participation rate	Attrition rate/ follow-up.	Recruitment procedure	Theory/ Behavioural change technique (BCT)/model	Fidelity
				and PA intervention components delivered								
Young et al., 2019/USA	Inactive, having either prediabetes or diabetes without insulin	Telephone-based MI and accelerometer and individualised support (tips for motivation and reducing barriers, a 24-week PA log and a PA personal action plan)	Seven MI calls were scheduled. MI calls varied from 5 to 25 min.	Researcher made the calls and delivered MI-based email counselling.	Accelerometry	NR.	21% /24 weeks	Telephone call following electronic medical screening.	Self-determination theory	NR		
Harris et al., 2017/UK	Eligible patients were 45–75 years old without contraindications to increasing MVPA	Postal group: Pedometer and written instruction for 12 weeks posted out to the patient Nurse support: Pedometer written instruction and tailored PA consultation Usual care: instructed to continue with their usual PA	Three face-to-face intervention lasting 10-20 min	Practice nurse provided written PA instruction and counselling	Average daily step-count, assessed by accelerometry and changes in step-counts spent weekly in MVPA in 10-min	42%	8% and 7% for postal and nurse support intervention groups, respectively/12 months	Mail invite following a medical screening	BCT: goals and planning, self-monitoring and feedback and encouraging social support, addressing barriers and relapse anticipation	NR		

(Continues)

TABLE 2 | (Continued)

Author, Year/ country	Eligibility criteria	Intervention component and content	Contact duration and number of contacts	Interventionists and PA			PA outcome measure	Participation rate	Attrition rate/ follow-up.	Recruitment procedure	Theory/ Behavioural change technique (BCT)/model	Fidelity
					intervention components delivered							
Knight et al., 2014/Canada	Adults aged 55–75 years who were generally healthy, with no diagnosis (e.g., hypertensive, diabetic, obese)	Exercise group: (EX),	One face-to-face contact at baseline	Researcher provided a written prescription	Total PA Step count	NR	No attrition/12 weeks	During a scheduled clinic visit	NR	NR		
		participants received PA	Duration of contact NR		measured by a device							
		prescription (high-intensity activity)	Number of exercise prescription sent NR									
		Sedentary behaviour (SB group:										
		participants received PA										
		prescription targeting reductions in sedentary behaviour)										
		Comprehensive counselling group:										
		participants received an activity										
		prescription targeting both increases in high-intensity activity and reductions in sedentary behaviour.										
		All groups had a PA										
		self-monitoring device										

(Continues)

TABLE 2 | (Continued)

Author, Year/ country	Eligibility criteria	Intervention component and content	Contact duration and number of contacts	Interventionists and PA		PA outcome measure	Participation rate	Attrition rate/ follow-up.	Recruitment procedure	Theory/ Behavioural change technique (BCT)/model	Fidelity
				Intervention components delivered	and PA intervention components delivered						
Gallegos-Carrillo et al., 2016/Mexico	Aged 35–70 years, mild hypertension, not meeting up to 150 min of moderate to vigorous PA (MVPA)/week	Brief counselling (BC) group: provision of written and verbal information to the patient, regarding PA benefits and advice on how to increase PA levels safely and progressively Referral group (ER): referral to local PA support	Three counselling session contact lasted 15 min	Brief counselling was provided by a primary care nurse		Accelerometry time spent in MVPA	73%	10% and 9% for BC and ER groups, respectively/24 weeks	NR	NR	NR
Miriam et al., 2012/USA	One visit in the previous 12 months, impaired glucose tolerance be free from a diagnosis of diabetes, have a haemoglobin A1c (HbA1c) below 7% and not be on diabetes medications. A body mass index (BMI) between 25 and 45 kg/m ²	One in-person baseline counselling session and regular telephone counselling, primary care provider endorsement in clinic and monthly automated telephone system for encouragement and tailored mailed PA support feedback	One contact at baseline Three telephone counselling every 2 weeks for 6 weeks followed by monthly calls over the entire 1-year intervention period. Duration of phone call NR	Baseline counselling consultation and telephone counselling with a trained health counsellor (baseline counselling telephone counselling provider not mentioned) Primary care provider (specific provider not mentioned) provided endorsement physically and via telephone		Moderate intensity PA/subjective	21%	9%/12 months	Recruited by phone following screening of medical records	Elements of social cognitive theory (modelling, self-monitoring, goal setting, reinforcement and cognitive reframing) Trans-theoretical model of change	NR

(Continues)

TABLE 2 | (Continued)

Author, Year/ country	Eligibility criteria	Intervention component and content	Contact duration and number of contacts	Interventionists and PA		PA outcome measure	Participation rate	Attrition rate/ follow-up.	Recruitment procedure	Theory/ Behavioural change technique (BCT)/model	Fidelity
				Intervention components delivered	intervention components delivered						
Khunti et al., 2021/UK	40–74 years for White Europeans or 25–74 years for those from an ethnic minority, previously recorded plasma glucose	Walking away (WA); annual group-based support Walking away plus (WAP) walking away and support (group-based support and mobile prompt pedometer use, feedback and biannual telephone calls) Control: information leaflet.	Four sessions: Group educational session at baseline. Four follow-up sessions Text messages were sent at least weekly over the first 6 months and then monthly. Annual follow-on sessions lasted 2.5 h	Trained educators delivered the group PA sessions	Time spent in MVPA/ objective	12.6%	12 months: 25.3% and 25.7% for WA and WAP, respectively, compared. 24 months: 33.8% and 31.1% for WA and WAP, respectively, compared to 18.9% for control/12 and 48 months	Routine visit	Social cognitive theory, common sense model, dual process theory BCT: Consequences of behaviour, identification/ problem- solving barriers, goal setting, action plan, set graded tasks	NR	
Lewis et al., 2020/USA	55–74 years with a body mass index of 25–35 kg/m ² , fewer than 60 min of planned exercise a week	Pedometer group (PED): A's PA counselling and exercise prescription and pedometer and a logbook to record PA steps Electronic activity monitoring (EAM) group: 5A's PA counselling and exercise prescription and activity monitor	One face-to-face contact Duration of PA counselling NR	A researcher with a background in exercise physiology and training in motivational interviewing delivered the 5A's PA counselling	Total PA was measured across using a sense wear armband	NR	10% and 5% at 12 weeks for PED and EAM groups respec- tively/12 weeks	Routine visits and flyers	5A's health promotion model	NR	

(Continues)

TABLE 2 | (Continued)

Author, Year/ country	Eligibility criteria	Interventionists					PA outcome measure	Participation rate	Attrition rate/ follow-up.	Recruitment procedure	Theory/ Behavioural change technique (BCT)/model	Fidelity
		Intervention component and content	Contact duration and number of contacts	Intervention components delivered	and PA intervention							
Rossen et al., 2021/Sweden	40–80 years, prediabetes or diagnosed with type 2 diabetes with a duration of ≥1 year	Multicom- ponent group: Pedometer and directed to a website for self-monitoring and PA consultations Self-monitoring of steps without counselling support	Twelve group consultations and nine face-to-face consultations Multi- component group had 45 min with their diabetes nurse consisting of counselling and study assessments	Researcher and diabetic nurse delivered group PA counselling sessions The diabetic nurse delivered the individualised PA counselling session			Objectively measured MVPA	53%	9% and 11% at 12 and 24 months respectively for all randomised groups/12 and 24 months	During regular visits and mailed invitation	Health belief model, the stages of change model and the social cognitive theory MI techniques	The intervention was delivered as intended as measured by MI logs
Takahashi et al., 2016/USA	18 years old, overweight or obese and had multiple medical comorbid conditions	Immediate pedometer group: Pedometer and setting PA goals based on step count at each visit and a digital versatile disc on exercise and journal for activity self-entry Delayed pedometer group: pedometer, educational materials and behavioural goal setting	Three intervention visits lasting 30–90 min	Researcher delivered PA counselling (goal setting)			Total PA step counts measured by a pedometer	15%	17% and 10% for the immediate and delayed group respectively / 16 weeks	Mailed invite	Social cognitive theory	NR

(Continues)

TABLE 2 | (Continued)

Author, Year/ country	Eligibility criteria	Interventionists			PA outcome measure	Participation rate	Attrition rate/ follow-up.	Recruitment procedure	Theory/ Behavioural change technique (BCT)/model	Fidelity
		Intervention component and content	Contact duration and number of contacts	Intervention components delivered						
Pears et al., 2016/UK	40–74 years and not previously diagnosed with heart disease, stroke, diabetes or kidney disease	All groups received face-to-face consultations and written materials (feedback on current PA) Motivational very brief intervention (MVBI); PA counselling (importance of PA, action plan using a diary) Pedometer VBI (PVBI); PA counselling (the 10,000 steps per/day recom- mendation, use a pedometer to monitor Combined VBI; motivational and pedometer VBI (CVBI) Control: Health check consultation only	Two contacts with interven- tionists (baseline and at 4 weeks) Mean delivery time was 6 min and 48 s for the MVBI, 5 min and for the PVBI and 9 and 35 min for the CVBI	Practice nurses and healthcare assistants delivered PA counselling and written prescription	Step counts/day and average number of minutes per day spent in light moderate and vigorous activities	18%	MVBI: 26.5% PVBI: 31% CVBI: 20% Control: 21.1%/4 weeks	Mailed invite and during a visit	Goal setting (behaviour), action planning, feedback on behaviour and self- monitoring of behaviour	Mean Overall fidelity ranged from 62% for the MVBI, 72 % for the P VBI and 74% for the CVBI

(Continues)

TABLE 2 | (Continued)

Author, Year/ country	Eligibility criteria	Intervention component and content	Contact duration and number of contacts	Interventionists and PA components delivered			PA outcome measure	Participation rate	Attrition rate/ follow-up.	Recruitment procedure	Theory/ Behavioural change technique (BCT)/model	Fidelity
Westland et al., 2020/ The Netherlands	Aged 40–75 years, at risk of CVD, did less than 30 min of MVPA on 5 or more days a week	PA counselling Consultations included (use of an activity log, forms for action planning, information useful websites, apps for PA, awareness) and an accelerometer device to -monitor their PA level	Four consultations with a duration of 20–30 min	Primary care nurses provided PA consultation	Change in minutes of MVPA/ objective	NR	NR	31.1% and 6.8% for intervention and control groups. respec- tively/24 weeks	Mail invite	BCTs (goals setting, action planning, feedback on behaviour, self- monitoring and problem solving)	NR	
Yates et al., 2016/UK	18 and 74 years of age inclusive and above the 90th percentile of the calculated diabetes (Leicester Practice Risk Score) risk score	Structured group educational programme and pedometer and short telephone contact to provide further support	Two Face-to-face group education programme were delivered lasting 2 h. Three telephone contacts (around 15 min)	Educator	Time MVPA intensity (count/min)/ objective	80%	Intervention group: 15.6%, 22.9%, 27.1% at 12, 24 and 36 months respec- tively/12, 24 and 36 months	Mailed invite	BCT (self-efficacy, identifying barriers and promoting self-regulatory skills through pedometer use	NR		

(Continues)

TABLE 2 | (Continued)

Author, Year/ country	Eligibility criteria	Interventionists					Theory/ Behavioural change technique (BCT)/model		
		Intervention component and content	Contact duration and number of contacts	Intervention components delivered	PA outcome measure	Participation rate	Attrition rate/ follow-up.	Recruitment procedure	Fidelity
Huebschmann et al., 2022/USA	Type 2 DM, Not meeting PA Guideline	PA tracking/ accountability through Fitbit and PA logs PA coaching phone calls (identifying PA motives, weighing pros and cons for PA, engaging social support for PA reminders/ rewards, relapse management) PA face-to-face visit (follow-up to ensure safety of exercises) Each patient received a Fitbit	Six phone calls and three in-person monthly primary care visits with a clinician	Clinic staff (PC staff involved in health and well-being) delivered the PA coaching calls Primary care clinician (one physician assistant and one doctor of medicine (MD) supervised the intervention safety	Total PA was measured by an accelerometer	20.6%	6% of all participants (attrition of intervention and control group NR)/12 weeks	NR	BCTs (goal setting, self- monitoring, accountability from others' monitoring with awareness, problem- solving, and social support)

Abbreviations: BCT, Behavioural change technique; MI, motivational interview; NR, Not reported; PA, physical activity; PCP, Primary care physician; UK, United Kingdom; USA, United States of America.




FIGURE 2 | Overview of the scalability domain assessment of effective PC PA interventions. Red denotes that further work is needed to ascertain scalability. Amber denotes mixed findings but promising for scale-up. Green denotes adequate evidence of scalability.

a nurse ($n = 5$) [29, 34, 36, 40, 44], a PCP ($n = 2$) [32, 45], an educator ($n = 1$) [43], a counsellor ($n = 2$), [32, 38], exercise specialist ($n = 1$) [39] or a researcher ($n = 3$) [39, 40, 45].

3.5 | Intervention Fidelity

Intervention fidelity was the least reported implementation characteristic. Out of the 26 studies, only five [34, 40, 44, 47, 48] reported on intervention fidelity. These studies reported fidelity to the intervention in terms of the intervention dose delivered and received by participants. Specifically, only three (23%) [34, 40, 44] out of the 13 effective interventions reported on intervention fidelity. One of the studies [40] reported that the intervention was delivered as intended, while the other [34] reported that the intervention was not received by participants as intended.

3.6 | PA Intervention Outcomes and Measures

Table 3 describes PA outcomes of effective interventions. Out of the 13 effective interventions [28, 29, 32–34, 36, 38–40, 43–45, 51], eight (62%) [28, 29, 32, 34, 36, 38–40] and five (39%) [33, 43–45, 51] included MVPA and total PA steps as PA outcome measures respectively. Eleven (85%) [28, 29, 33, 34, 36, 39, 40, 43, 45, 51] studies that demonstrated effectiveness used device-based measures of PA, while two (15%) [32] used self-reported measures.

3.6.1 | Objective MVPA

Six (46%) [28, 29, 34, 36, 39, 40] out of the 13 studies in this review that demonstrated effectiveness measured MVPA using a device. The majority ($n = 4$) (67%) of these interventions [29, 34, 36, 40] were delivered by a nurse, and they had an increase in the

range of 33–84 min/week of MVPA compared to controls after 3–24 months follow-up. Furthermore, these interventions had an average of three to nine face-to-face contacts (mean = 5), and all involved an activity monitor for self-monitoring as a supporting component. Only two interventions were remotely delivered: one involved pedometer and written PA instructions sent by mail for 12 weeks [36] and the other involved mailed tailored PA feedback according to motivational readiness [28].

3.6.2 | Objective Total PA

Five (39%) [33, 43–45, 51] out of the 13 studies that increased PA assessed PA used device-based measures (i.e., pedometers measuring steps/day). Generally, these interventions were underpinned by BCTs and were characterised by a lesser number of contacts and a greater magnitude of increase in minutes/week spent in total PA compared to effective interventions with objectively measured MVPA. Specifically, the intervention [45] with the largest increase in total PA had an increase of 182 steps/day (~130 min/week) when compared to controls. This intervention was delivered by a GP trainee (contact duration not reported) and delivered using the 5As model of health promotion and MI techniques.

4 | Scalability Assessment and Recommendations

Figure 2 provides an overview of the scalability domain assessment of effective interventions (domains A1–A3 were not included in this figure because all interventions scored the same). Overall, domains A1 (the problem), A2 (intervention), A3 (political/strategic context), and B2 (reach and acceptability) scored highly, demonstrating good evidence for scalability. Fidelity and cost-effectiveness were often not reported, making it difficult to assess scalability in these domains. Mixed scores were seen in the

domains of effectiveness, delivery setting and workforce, implementation infrastructure and sustainability, indicating potential for scale-up in these areas.

Table 3 provides a summary of the implementation characteristics of 15 effective interventions (representing 13 individual studies), their scalability assessment scores and recommendations. Four (27%) [28, 36, 44] (representing three studies) interventions were deemed to be highly scalable (merit scale-up). Two of these interventions [28, 36] were delivered remotely and scored moderate to high in the reach and acceptability domain (B2), and highly (largely scalable) in domains B1 (fidelity and adaptation), B3 (delivery setting and workforce), B4 (implementation infrastructure) and B5 (sustainability). The other interventions (two-arm treatment) [44] were delivered by a nurse (two face-to-face contacts with an average duration of 6 min with a pedometer and PA diary as follow-up support) and scored highly in domains B2 (reach and acceptability), B3 (delivery setting and workforce), B5 (sustainability), moderately in the domain B4 (implementation infrastructure) and low in domain B1 (fidelity and adaptation) (see Table 3 for other domain scoring).

Ten (67%) interventions [29, 32–34, 36, 39, 43, 45, 51, 56] were deemed as promising (requiring further information/planning). These interventions had moderate to high scores in domain B2 (reach and acceptability domain), low to moderate scores in domains B3 (delivery setting and workforce), B4 (implementation infrastructure) and B5 (sustainability). One intervention [40] was deemed not to merit scaling up. This intervention was characterised by an initial 12 group sessions delivered by a researcher followed by nine sessions of face-to-face MI sessions lasting 45 min. A radar plot diagram of scalability domain scoring can be seen in Supplementary file 3.

5 | Discussion

Most of the effective interventions were found to be highly scalable in the reach and acceptability domain, with a 15% average attrition rate, indicating that PC PA interventions are well-received by patients. Despite this, PA interventions delivered within PC yield modest results in improving MVPA [10], which are likely to reduce further when interventions are implemented at scale (scale-up penalty) [57]. While the need to identify PC PA interventions with a significant effect size is evident, exploring the reasons for the scale-up penalty is necessary. A possible reason for this penalty is likely due to the adaptations made during the scale-up process to better fit evidence-based interventions to user needs and delivery contexts [58]. For example, in a 2019 review of 10 obesity trials, adaptations during scale-up led to a 25% scale-up penalty, reducing intervention effects observed in pre-scale trials [59]. This highlights the need to find PC PA interventions with large effect sizes that require minimal adaptation for scale-up. Thus, careful consideration of scalability, delivery infrastructure and compatibility with the local context becomes apparent when designing interventions [60]. Furthermore, future work will also need to examine and report on intervention fidelity considering the lack of reporting on fidelity in studies included in the review.

The most scalable interventions, as assessed by ISAT, were either delivered remotely (automated PA advice and mailed

instructions) or briefly in person by nurses. Digital interventions are cost-effective, reaching many people with fewer resources [61]. Additionally, the automated and algorithmic features of digital interventions ensure protocol-based fidelity, which presents less challenges than when relying on human interventionists [60]. In-person interventions, led by nurses, lasted under 10 min, making them highly feasible for scaling up. Research [62] indicates that nurses, compared to GPs, face fewer time constraints in promoting PA, highlighting their potential in supporting PA promotion within PC. Furthermore, nurse-led interventions were underpinned by BCTs including goal setting, self-monitoring and feedback, which have previously been shown to encourage behavioural change [63]. This further underscores nurses' capacity to deliver or support PA interventions within PC, highlighting the need to train more nurses in behaviour change and the implementation of evidence-based BCTs.

Most of the effective PA interventions in this review display promise to be scaled up but require further planning and consideration in terms of intervention duration, infrastructural demands and sustainability. Time constraints have been reported as a major barrier to the implementation of PA promotion within PC [15, 16]. It has been suggested that interventions that will be feasible to implement within routine PC should be very brief interventions (VBIs) which can be delivered within 5 min [64]. Most of the interventions deemed to be promising did not report on the duration of interventions and thus made it difficult to make a judgement on their feasibility for scale-up. The interventions that reported on intervention duration were in the range of 10–30 min and are likely to be less feasible within routine PC consultations. Lower scores were associated with intensive interventions (i.e., multiple contacts/lengthy contact and follow-up supports such as phone calls and or use of activity devices) due to the time commitments needed for delivery. Additionally, lower scores were linked to the involvement of additional healthcare professionals (such as exercise specialists and counsellors) due to the likely need for extra investment in training and employing these professionals to deliver PA services.

Generally, this review highlights that effective PC PA interventions showed promising reach with an average participation rate of 43%. Similar participation rates have been observed in PA interventions targeting type 2 diabetes patients in healthcare settings [61]. This review suggests that systematic recruitment methods may yield higher recruitment compared to opportunistic approaches. To enhance the reach of PC PA interventions, systematic recruitment of patients with comorbidities through methods such as postal invites, SMS or email for routine health checks is recommended. Periodic health checks could be supported by opportunistic PA counselling during routine consultations which have been shown to facilitate the recruitment of disadvantaged groups when compared to systematic recruitment [65]. Therefore, using multiple recruitment methods has the potential to increase overall reach and engagement, while also maximising recruitment from socially disadvantaged populations.

Furthermore, this review found that fewer interactions with interventionists are needed to improve participation in light intensity PA compared to MVPA, which supports the wider WHO message [66] of 'some movement is good, more is better'. This might also have positive implications for implementation, given

TABLE 3 | Implementation features of effective PA interventions, scalability assessment scores and recommendations.

Study/ follow-up (attrition rate)	Outcome/ measure (objec- tive/subjective)	Eligibility crite- ria/recruitment type/participation rate	Interventionist/ intervention component	Number of contact and duration	BCT/theory/ model and technique of delivery	Fidelity	Effectiveness and magnitude of change/ additional remarks (e.g., qualitative findings)	Scalability domain score/scalability recommendations
Gao et al., 2015/6 (13%) and 12 months (16%)	MVPA/objective	One primary care visit in the 12 months prior to screening, at least 18 years, overweight or obese /Systematic/43%	Mailed expert feedback PA counselling and PA self-help booklet and newsletter sent following filled postal PA readiness questionnaire Use of pedometer for PA self-monitoring	Twelve months postal mail PA counselling feedback	Goal setting, self-efficacy, self-monitoring, decisional balance	NR	Significantly increased MVPA at 6 months ($p = 0.03$) and 12 months ($p = 0.053$) (Actual minutes spent in MVPA NR)	Domain A1 = 3 Domain A2 = 3 Domain A3 = 3 Domain A4 = 1 Domain A5 = IN Domain B1 = 3 Domain B2 = 2 Domain B3 = 2.8 Domain B4 = 3 Domain B5 = 2.7 Recommendation: Merit
Harris et al., 2015/3(3%) and 12 months (6%)	MVPA/objective	60–74 years had no contraindications to increasing PA/systematic/30%	Nurse/PA consultations and PA supports (pedometer for self-monitoring, PA handbook, PA diary and walking plan)	Four face-to-face contacts. Duration: NR	Goal setting, self-monitoring, building self-efficacy and social support, overcoming barriers, preventing relapses and building lasting habits	NR	Increase in time spent in MVPA when compared to control was 66 min/week at 3 months At 12 months increase was 40 min/week MVPA	Domain A1 = 3 Domain A2 = 3 Domain A3 = 3 Domain A4 = 2 Domain A5 = IN Domain B1 = 0.7 Domain B2 = 2.5 Domain B3 = 2 Domain B4 = 2 Domain B5 = 2 Recommendation: Promising but requiring further informa- tion/planning

(Continues)

TABLE 3 | (Continued)

Study/ follow-up (attrition rate)	Outcome/ measure (objec- tive/subjective)	Eligibility crite- ria/recruitment type/participation rate	Interventionist/ intervention component	Number of contact and duration	BCT/theory/ model and technique of delivery	Fidelity	Effectiveness and magnitude of change/ additional remarks (e.g., qualitative findings)	Scalability domain score/scalability recommendations
Heron et al., 2014/ 12 weeks (5%)	Total step ^a Count/objective	35–75 years, inactive based on a questionnaire. No contraindications to PA/opportunistic/ 49.3	GP trainee/ Group 1: MI PA counselling and leaflet on PA, pedometer and follow-up MI telephone No formal training in MI techniques was undertaken beyond that experienced within undergraduate medical training	One contact and four follow-up telephone calls Duration NR	Theory of planned behaviour, '5 As' model, BCT (establishing rapport, agenda Setting, assessing importance and confidence, pros and cons, brainstorming solutions), MI	NR	Group 1 achieved an average increase of over 2600 steps/day (182 min/week) from baseline, which was statistically significant compared with the increase in group 2 (748 steps/day (52 min/week)	Domain A1 = 3 Domain A2 = 3 Domain A3 = 3 Domain A4 = 3 Domain A5 = 1N Domain B1 = 0.7 Domain B2 = 2.5 Domain B3 = 2 Domain B4 = 2 Domain B5 = 2 Recommendation: Promising but requiring further informa- tion/planning
Kolt et al., 2012/4 months (13%) and 12 months (15%)	Moderate PA (MPA) intensity /subjective	65 or older; ability to walk and freedom from health conditions that contraindicate participation in PA/systematic/57%	Primary care physician (PCP) and counsel- lor/pedometer green prescription (PGP) group: initial face-to-face counselling by PCP and telephone counselling by counsellor and pedometers	One face-to-face counselling by PCP (duration NR) and three telephone Counselling sessions over 3–4 months by trained counselling (lasting between 10 and 30 min)	Trans theoretical model, goal setting, preventing relapse, assessing PA progress	NR	The PGP group had a significant increase in time spent in moderate intensity (58 min/week) from baseline to 3/4 month when compared to the standard group increase of 30 min/week This initial increase (at 4 months) was followed by a decrease of 7 and 12 min/week MPA in the PGP and standard group respectively at 12 months	Domain A1 = 3 Domain A2 = 3 Domain A3 = 3 Domain A4 = 1 Domain A5 = 1N Domain B1 = 0.7 Domain B2 = 2.5 Domain B3 = 2 Domain B4 = 2 Domain B5 = 1.3 Recommendation: Promising but requiring further informa- tion/planning

(Continues)

TABLE 3 | (Continued)

Study/ follow-up (attrition rate)	Outcome/ measure (objec- tive/subjective)	Eligibility crite- ria/recruitment type/participation rate	Interventionist/ intervention component	Number of contact and duration	BCT/theory/ model and technique of delivery	Fidelity	Effectiveness and magnitude of change/ additional remarks (e.g., qualitative findings)	Scalability domain score/scalability recommendations
Mutrie et al., 2012/12 weeks (10%) and 24 weeks (10%)	Daily step ^a Count/objective	Living independently and not meeting current PA recommenda- tion/Systematic/32%	Intervention group: PA consultations at baseline and support at 12 weeks (second PA consultation, walking group and PA booklet and pedometer) weeks (second PA consultation, walking group and PA booklet and pedometer)	Two face-to-face contacts with a trained nurse lasting an average of 30 min	Cognitive model of behaviour change.	NR	Significant increase when compared to control during the first 12 weeks (2119 steps/day (148 min/week) 24 weeks (6 months): 190 steps /day (13 min/week PA) increase in steps in the intervention group when compared to control, but not significant Qualitative findings: The findings suggest that the pedometer was easy to use and was an important motivational tool for many 'I think when you've got your meter on, you try to get a wee bit better ... I feel it's like a challenge' The nurses had concerns about time	Domain A1 = 3 Domain A2 = 3 Domain A3 = 3 Domain A4 = 1 Domain A5 = IN Domain B1 = 0.7 Domain B2 = 2.5 Domain B3 = 2 Domain B4 = 2 Domain B5 = 2 Recommendation: Promising but requiring further informa- tion/planning

(Continues)

TABLE 3 | (Continued)

Study/ follow-up (attrition rate)	Outcome/ measure (objec- tive/subjective)	Eligibility crite- ria/recruitment type/participation rate	Interventionist/ intervention component	Number of contact and duration	BCT/theory/ model and technique of delivery	Fidelity	Effectiveness and magnitude of change/ additional remarks (e-g., qualitative findings)	Scalability domain score/scalability recommendations
Yaacob et al., 2018/ 12 weeks (29%)	Step count ^a /objective	aged 35 – 45 years determined to lead a sedentary lifestyle on the 7-day physical activity questionnaire and who have no chronic illness/NR	Researcher/ pedometer and support (log diaries and support groups, follow-up monthly meeting	Four physical contacts Duration: NR	NR	NR	At 12 weeks there was a significant increase in the step counts/day from baseline by 631 steps/day (44 min/week PA) of the intervention group compared to the control which had a decrease of 95 steps/day (7 min/week) from baseline	Domain A1 = 3 Domain A2 = 3 Domain A3 = 3 Domain A4 = 2 Domain A5 = IN Domain B1 = 0.7 Domain B2 = 0.5 Domain B3 = 1.3 Domain B4 = 2 Domain B5 = 1.3 Recommendation: Promising but requiring further informa- tion/planning

(Continues)

TABLE 3 | (Continued)

Study/ follow-up (attrition rate)	Outcome/ measure (objec- tive/subjective)	Eligibility crite- ria/recruitment type/participation rate	Interventionist/ intervention component	Number of contact and duration	BCT/theory/ model and technique of delivery	Fidelity	Effectiveness and magnitude of change/ additional remarks (e.g., qualitative findings)	Scalability domain score/scalability recommendations
Van der Weegen et al., 2015/6 (14%) and 9 months (12%)	Minutes of MVPA/objective	40 and 70 years old, type 2DM or COPD, BMI > 25 and not having at least 30 min of MVPA/Systematic/ 37%	Practice nurses/group 1: activity monitor and self-management support programme (SSP) (individual consultations, automated PA feedback and use of booklet about PA) Practices nurses were trained on the execution of intervention and how to use the 5A model of PA promotion Group 2: SSP Group 3: Usual care	Four face-to-face contacts Duration: NR	5A model involving goal setting, awareness of the risk of physical inactivity	Group 1: 12 participants did not receive the minimal intervention	Group 1 showed a significant increase of 8 min/day (72 min/week) more MVPA than participants in group 2 and 12 min/day (84 min/week) more PA than- group 3 (usual care group) at 4–6 months. Group 2 increased nonsignificantly by 7 min/day (49 min/week) compared to control At 9 months: Group 1 showed a significant increase of 9 min/day (63 min/week) more than group 2 and 11 min/day (77 min/week), more than group 3	Domain A1 = 3 Domain A2 = 3 Domain A3 = 3 Domain A4 = 3 Domain A5 = IN Domain B1 = 0.7 Domain B2 = 2 Domain B3 = 2 Domain B4 = 2 Domain B5 = 2 Recommendation: Promising but requiring further informa- tion/planning

(Continues)

TABLE 3 | (Continued)

Study/ follow-up (attrition rate)	Outcome/ measure (objec- tive/subjective)	Eligibility crite- ria/recruitment type/participation rate	Interventionist/ intervention component	Number of contact and duration	BCT/theory/ model and technique of delivery	Fidelity	Effectiveness and magnitude of change/ additional remarks (e.g., qualitative findings)	Scalability domain score/scalability recommendations
Harris et al., 2017/ 12 Months 8% (postal group), 7% (nurse group)	MVPA/ objective.	Eligible patients were 45–75 years old without contraindications to increasing MVPA /systematic/10%	Postal group: Pedometer, handbook, written instruction for 12 weeks posted out to a patient Nurse support group: Pedometer, PA handbook, PA diary, written instruction, and tailored PA consultation	Nurse group received three physical contacts lasting 10–20 min	BCT: Goals and planning, self-monitoring, encouraging social support, addressing barriers, relapse anticipation	NR	Both nurse and postal group had a significant increase of MVPA (35 and 33 min/week, respectively) at 12 months from baseline compared to control (usual care) (5 min/week)	Postal group Domain A1 = 3 Domain A2 = 3 Domain A3 = 3 Domain A4 = 2 Domain A5 = IN Domain B1 = 3 Domain B2 = 2.5 Domain B3 = 3 Domain B4 = 3 Domain B5 = 2.7 Recommendation: Merit Nurse Group Domain A1 = 3 Domain A2 = 3 Domain A3 = 3 Domain A4 = 2 Domain A5 = IN Domain B1 = 1 Domain B2 = 2.5 Domain B3 = 2 Domain B4 = 2 Domain B5 = 2.3 Recommendation: Promising but requiring further informa- tion/Planning

(Continues)

TABLE 3 | (Continued)

Study/ follow-up (attrition rate)	Outcome/ measure (objec- tive/subjective)	Eligibility crite- ria/recruitment type/participation rate	Interventionist/ intervention component	Number of contact and duration	BCT/theory/ model and technique of delivery	Fidelity	Effectiveness and magnitude of change/ additional remarks (e.g., qualitative findings)	Scalability domain score/scalability recommendations
Miriam et al., 2012/12 months (9%)	Moderate intensity PA/subjective	Impaired glucose tolerance is free from a diagnosis of diabetes /Systematic/99%	Trained health counsellor and primary care provider (specific provider not mentioned) One in-person baseline counselling session, telephone counselling, primary care provider endorsement in clinic, monthly automated telephone system for encouragement and tailored mailed PA support feedback	One face-to-face contact (duration NR) and three telephone calls delivered by a counsellor (duration NR) Duration of PA endorsement by primary care provider) NR	Social cognitive theory (modelling, self-monitoring, goal setting, reinforcement and cognitive reframing) Trans theoretical model of change	NR	Moderate PA increased significantly by 60 min/week when compared to control time (no increase in MVPA)	Domain A1 = 3 Domain A2 = 3 Domain A3 = 3 Domain A4 = 1 Domain A5 = IN Domain B1 = 0.7 Domain B2 = 2 Domain B3 = 3 Domain B4 = 2 Domain B5 = 1.3 Recommendation: Promising but requiring further informa- tion/planning

(Continues)

TABLE 3 | (Continued)

Study/ follow-up (attrition rate)	Outcome/ measure (objec- tive/subjective)	Eligibility crite- ria/recruitment type/participation rate	Interventionist/ intervention component	Number of contact and duration	BCT/theory/ model and technique of delivery	Fidelity	Effectiveness and magnitude of change/ additional remarks (e.g., qualitative findings)	Scalability domain score/scalability recommendations
Lewis et al., 2020/12 weeks 5%	MVPA/objective	55–74 years with a body mass index of 25–35 kg/m ² , fewer than 60 min of planned exercise a week, /Opportunistic and systematic 68%	Researcher (with a background in exercise physiology and training in MI) delivered the 5A's counselling Electronic activity monitoring (EAM) group: 5A's PA counselling and exercise prescription and activity monitor	One face-to-face contact Duration of PA counselling NR	5 A model/MI	NR	The EAM intervention produced a significant increase of 11 min/day (79 min/week) from baseline to 12 weeks compared to the PED intervention which had an increase of 0.2 min/day (1.4 min/week) from baseline to 12 weeks.	Domain A1 = 3 Domain A2 = 3 Domain A3 = 3 Domain A4 = 3 Domain A5 = IN Domain B1 = 0.7 Domain B2 = 3 Domain B3 = 1.3 Domain B4 = 2 Domain B5 = 1 Recommendation: Promising but requiring further informa- tion/Planning
Rossen et al., 2021/12 (9%) and 24 Months (11%)	MVPA/objective	40–80 years, prediabetes or diagnosed with type 2 diabetes with a duration of ≥1 year/Opportunistic and systematic/53%	Researcher and diabetic nurse/ group session led by research staff. The diabetic nurse delivered the individualised PA counselling session based on MI Multicomponent group: Pedometer and directed to a website for self-monitoring and PA consultations	Twelve group consultations (duration NR) and nine face-to-face consultations lasting 45 min	Health belief model, the stages of change model and the social cognitive theory. MI technique	The intervention was delivered as intended as measured for MI logs	Multi-component vs. control group: significant increase of 56 min/week at 12 weeks when compared to control. There was an increase of 77 min/week at 24 months when compared to control	Domain A1 = 3 Domain A2 = 3 Domain A3 = 3 Domain A4 = 1 Domain A5 = IN Domain B1 = 0.7 Domain B2 = 2.5 Domain B3 = 1.3 Domain B4 = 1 Domain B5 = 1 Recommendation: Does not merit scale up

(Continues)

TABLE 3 | (Continued)

Study/ follow-up (attrition rate)	Outcome/ measure (objec- tive/subjective)	Eligibility crite- ria/recruitment type/participation rate	Interventionist/ intervention component	Number of contact and duration	BCT/theory/ model and technique of delivery	Fidelity	Effectiveness and magnitude of change/ additional remarks (e.g., qualitative findings)	Scalability domain score/scalability recommendations
Pears et al., 2016/4 weeks 26.5% (MVBI); 31% (PVBI)	Total PA ^a /objective	40–74 years and not previously diagnosed with heart disease, stroke, diabetes or kidney dis- ease/opportunistic and systematic/18% patients recruited during health check	Nurse/ motivational very brief intervention (MVBI); PA counselling and action plan using a diary) Pedometer VBI (PVBI) PA counselling, pedometer and Step Chart Combined VBI (CVBI) Motivational and pedometer VBI (CVBI)	MVBI: Two face-to-face contact Mean delivery time was 6 min and 48 s Pedometer VBI was delivered within 5 min CVBI: delivered within 9 min and 35 s	Goal setting (behaviour), action planning, feedback on behaviour and self-monitoring of behaviour	Mean overall fidelity ranged from 62% for the MVBI, 72% for the PVBI	MVBI had an increase of 20.3 counts/minute when compared to control at 4 weeks. PVBI had an increase of 23.5 counts/minute when compared to control at 4 weeks CVBI had no increase in count/minute when compared to control at 4 weeks. Cost-effectiveness; £6.83 (US\$10.06) per participant for the MVBI £17.09 (US\$25.17) for the PVBI Qualitative enquiry: Practitioners felt that the PVBI was the easiest and quickest to deliver Participants confirmed that the health check was a good time to discuss PA and that the VBI was a good reminder of the importance of PA	PVBI Group Domain A1 = 3 Domain A2 = 3 Domain A3 = 3 Domain A4 = 1 Domain A5 = 2 Domain B1 = 1.3 Domain B2 = 1.5 Domain B3 = 2.8 Domain B4 = 2 Domain B5 = 2.7 Recommendation: Merit MBVI Domain A1 = 3 Domain A2 = 3 Domain A3 = 3 Domain A4 = 1 Domain A5 = 3 Domain B1 = 1.3 Domain B2 = 2.5 Domain B3 = 2.5 Domain B4 = 2 Domain B5 = 2.7 Recommendation: Merit

(Continues)

TABLE 3 | (Continued)

Study/ follow-up (attrition rate)	Outcome/ measure (objec- tive/subjective)	Eligibility crite- ria/recruitment type/participation rate	Interventionist/ intervention component	Number of contact and duration	BCT/theory/ model and technique of delivery	Fidelity	Effectiveness and magnitude of change/ additional remarks (e.g., qualitative findings)	Scalability domain score/scalability recommendations
Yates et al., 2016/12 (15%), 24(22.9%) and 36 months (27%)	Total PA ^a /objective	18 and 74 years of age inclusive and above the 90th percentile of the calculated diabetes (Leicester Practice Risk/Systematic/ 97%	Educator/group educational programme and support (pedometer and telephone follow-up) Educators were recruited and trained to deliver the counselling sessions	Two group sessions lasting 2 h. Three telephone contacts (around 15 min)	BCT (self-efficacy, identifying barriers and promoting self-regulatory skills)	NR	Significant increase in ambulatory activity of 411 steps/day) (29 min/week) in the intervention group compared with control at 12 months. There was a decline in both groups with no difference between groups by 36 months	Domain A1 = 3 Domain A2 = 3 Domain A3 = 3 Domain A4 = 1 Domain A5 = IN Domain B1 = 0.7 Domain B2 = 2 Domain B3 = 1 Domain B4 = 2 Domain B5 = 1.7 Recommendation: Promising but requiring further informa- tion/planning

Abbreviations: MI, motivational interview; MVPA, moderate to vigorous physical activity; NR, not reported; PA, physical activity; PCP, primary care physician.

^aSteps/day converted to minutes/week: (steps ÷ 100) × 7 (conversion of steps/day to moderate intensity PA).

Scoring: 0 (not at all), 1 (to a small extent), 2 (somewhat scalable), 3 (largely scalable), IN (indeterminate).

The problem (A1), the intervention (A2), strategic /political context (A3), evidence of effectiveness (A4), and intervention cost (A5) and BI (fidelity and adaptation), B2 (reach and acceptability), B3 (delivery workforce and setting), B4 (implementation infrastructure) and B5 (sustainability).

health professionals might be more willing to promote a general message of increased movement, rather than a focused one on exercise intensity (i.e., the importance of MVPA).

5.1 | Strength and Limitations

To the best of the authors' knowledge, this study is the first to evaluate the scalability potential of effective PC PA interventions using a scalability assessment tool. Utilising the ISAT, we demonstrated that effective PC PA interventions exhibit scalability potential. The authors acknowledge the subjective nature of scalability assessment with the ISAT. Further refinement of scalability tools to develop a scoring system for evaluating the scalability of complex interventions such as PA promotion is needed. Despite attempts to locate scalability information from published companion papers, crucial details such as intervention duration, fidelity assessment and cost-effectiveness were often missing, hindering a comprehensive assessment.

Nurses were the main providers of effective PC PA interventions, indicating their potential to deliver or assist in PA interventions within PC. Further research on the utility and effectiveness of other healthcare professionals such as link workers, counsellors and exercise specialists in delivering PA interventions would be worthwhile. While a single-reviewer extraction system and a 10% independent screening check were pragmatically used, there may have been relevant articles that were missed. This narrative review aimed to provide a general overview of PC PA intervention implementation characteristics rather than conduct a systematic review. Despite rigorous methods, some relevant research may have been overlooked.

5.2 | Implication for Practice and Research

The inclusion of BCTs including self-monitoring, goal setting and feedback are likely to be important for increasing PA in PC, despite modest evidence concerning the effectiveness of PC-based PA interventions. PC health professionals are advised to use Moving Medicine resources by the UK Faculty of Sports and Exercise Medicine to deliver evidence-based PA interventions supported by BCTs [67].

Mixed findings exist regarding the optimal number of contacts and duration required for effective PC PA interventions to enhance total PA or MVPA. Considering known time constraints in PC, remote-based PA interventions are worth considering for broader implementation. However, due to limited literature on remote-based PA interventions in PC, further research is needed to assess their effectiveness. Additionally, with the introduction of new roles in PC such as link workers, community pharmacists, exercise specialists and first contact physiotherapists, additional research is necessary to evaluate the feasibility, sustainability and effectiveness of training these other PC staff to deliver PA interventions.

This study suggests that PC PA interventions have satisfactory reach. Promoting light-intensity PA may require less contact time than MVPA. Future research should explore PC healthcare providers' attitudes and practices regarding the promotion of light

versus moderate intensity PA. Finally, future trial designs could be improved by using multiple recruitment strategies to broaden reach, engagement and inclusion of socially disadvantaged populations.

5.3 | Conclusion

PC PA interventions, whether delivered remotely or in person and utilising BCTs such as goal setting, self-monitoring and feedback, show promise for increasing PA. While most interventions demonstrate scalability potential, further examination is needed regarding the delivery settings, implementation infrastructure and sustainability. Remotely delivered PA interventions and those delivered by a nurse showed high scalability potential. Further work that examines the acceptability and effectiveness of other healthcare professionals in intervention delivery would be worthwhile, and more research is needed to assess the utility and effectiveness of remote-based PA interventions in PC.

Acknowledgements

Jimi Osinaike (JO) is a Ph.D. student, and this study is part of his Ph.D. study. RC, AM, AL and SH are part of JO's supervisory team.

Conflicts of Interest

The authors declares no conflicts of interest.

Data Availability Statement

The data described in this article is currently openly available. Please contact the author for further information if required.

References

1. B. K. Pedersen and B. Saltin, "Exercise as Medicine—Evidence for Prescribing Exercise as Therapy in 26 Different Chronic Diseases," *Scandinavian Journal of Medicine and Science in Sports* 25, no. S3 (2015): 1–72.
2. R. Guthold, G. A. Stevens, L. M. Riley, and F. C. Bull, "Worldwide Trends in Insufficient Physical Activity From 2001 to 2016: A Pooled Analysis of 358 Population-Based Surveys With 1.9 Million Participants," *Lancet Global Health* 6, no. 10 (2018): e1077–e1086.
3. World Health Organisation: Global Action Plan for Physical Activity. (2018), assessed March 20, 2024, <https://iris.who.int/bitstream/handle/10665/272722/9789241514187-eng.pdf?sequence=1>.
4. N. Goodwin, N. Curry, C. Naylor, S. Ross, and W. Duldig, *Managing People With Long-Term Conditions* (London: The Kings Fund, 2010), 50–51.
5. World Health Organisation, WHO: WHO Global NCD Action Plan 2013–2030. 2022, assessed March 20, 2024, https://cdn.who.int/media/docs/default-source/ncds/mnd/2022_discussion_paper_01_aug.pdf?sfvrsn=6aa03d21_3.
6. NICE: Physical Activity: Brief Advice for Adults in Primary Care, 2013, assessed March 20, 2024, <https://www.nice.org.uk/guidance/ph44/resources/physical-activity-brief-advice-for-adults-in-primary-care-pdf-1996357939909>.
7. G. Orrow, A. Kinmonth, S. Sanderson, and S. Sutton, "Effectiveness of Physical Activity Promotion Based in Primary Care: Systematic Review and Meta-Analysis of Randomised Controlled Trials," *BMJ* 344, no. 7850 (2012): 808.

8. A. Sanchez, P. Bully, C. Martinez, and G. Grandes, "Effectiveness of Physical Activity Promotion Interventions in Primary Care: A Review of Reviews," *Preventive Medicine* 76 (2015): S56–S67.
9. V. Van der Wardt, C. Di Lorito, and A. Viniol, "Promoting Physical Activity in Primary Care: A Systematic Review and Meta-Analysis," *British Journal of General Practice* 71, no. 706 (2021): e399–e405.
10. V. E. Kettle, C. D. Madigan, A. Coombe, et al., "Effectiveness of Physical Activity Interventions Delivered or Prompted by Health Professionals in Primary Care Settings: Systematic Review and Meta-Analysis of Randomised Controlled Trials," *BMJ* 376 (2022): e068465.
11. R. Chatterjee, T. Chapman, M. G. T. Brannan, and J. Varney, "GPs' Knowledge, Use, and Confidence in National Physical Activity and Health Guidelines and Tools: A Questionnaire-Based Survey of General Practice in England," *British Journal of General Practice* 67, no. 663 (2017): e668–e675.
12. J. Gaglani and P. Oakeshott, "Conducting a Survey of Patient Reported Advice on Diet and Exercise in Primary Care. Education for Primary Care," *Education for Primary Care* 29, no. 3 (2018): 178–179.
13. S. Morton, D. Thompson, P. Wheeler, G. Easton, and A. Majeed, "What Do Patients Really Know? An Evaluation of Patients' Physical Activity Guideline Knowledge Within General practice1," *London Journal of Primary Care* 8, no. 4 (2016): 48–55.
14. L. H. Hall, R. Thorneoloe, R. Rodriguez-Lopez, et al., "Delivering Brief Physical Activity Interventions in Primary Care: A Systematic Review," *British Journal of General Practice* 72, no. 716 (2022): e209–e216.
15. P. C. Wheeler, R. Mitchell, M. Ghaly, and K. Buxton, "Primary Care Knowledge and Beliefs About Physical Activity and Health: A Survey of Primary Healthcare Team Members," *BJGP Open* 1, no. 2 (2017): bjgopen17X100809.
16. A. Lowe, A. Myers, H. Quirk, J. Blackshaw, S. Palanee, and R. Copeland, "Physical Activity Promotion by GPs: A Cross-Sectional Survey in England," *BJGP Open* 6, no. 3 (2022): BJGPO.2021.0227.
17. M. O'Brien, C. Shields, S. Crowell, O. Theou, P. McGrath, and J. Fowles, "The Effects of Previous Educational Training on Physical Activity Counselling and Exercise Prescription Practices Among Physicians Across Nova Scotia: A Cross-Sectional Study," *Canadian Medical Education Journal* 9, no. 4 (2018): e35–e45.
18. J. Windt, A. Windt, J. Davis, R. Petrella, and K. Khan, "Can A 3-Hour Educational Workshop and the Provision of Practical Tools Encourage Family Physicians to Prescribe Physical Activity as Medicine? A Pre–Post Study," *BMJ Open* 5, no. 7 (2015): e007920.
19. E. T. Hébert, M. O. Caughy, and K. Shuval, "Primary Care Providers' Perceptions of Physical Activity Counselling in a Clinical Setting: A Systematic Review," *British Journal of Sports Medicine* 46, no. 9 (2012): 625.
20. H. Koorts, E. Eakin, P. Estabrooks, A. Timperio, J. Salmon, and A. Bauman, "Implementation and Scale Up of Population Physical Activity Interventions for Clinical and Community Settings: The PRAC-TIS Guide," *International Journal of Behavioural Nutrition and Physical Activity* 15, no. 1 (2018): 1–11.
21. ProQuest RefWorks, 2024, assessed June 20, 2024, <https://refworks.proquest.com/learn-more/>.
22. J. A. Sterne, J. Savović, M. J. Page, et al., "RoB 2: A Revised Tool for Assessing Risk of Bias in Randomised Trials," *BMJ* 366 (2019): 14898.
23. A. J. Milat, L. King, A. E. Bauman, and S. Redman, "The Concept of Scalability: Increasing the Scale and Potential Adoption of Health Promotion Interventions Into Policy and Practice," *Health Promotion International* 28, no. 3 (2013): 285–298.
24. A. Ben Charif, H. T. V. Zomahoun, A. Gogovor, et al., "Tools for Assessing the Scalability of Innovations in Health: A Systematic Review," *Health Research Policy and Systems* 20, no. 1 (2022): 34.
25. A. Milat, K. Lee, K. Conte, et al., "Intervention Scalability Assessment Tool: A Decision Support Tool for Health Policy Makers and Implementers," *Health Research Policy and Systems* 18 (2020): 1–17.
26. T. Greenhalgh and C. Papoutsi, "Spreading and Scaling Up Innovation and Improvement," *BMJ* 365 (2019): 12068.
27. J. R. Reed, P. Estabrooks, B. Pozehl, K. Heelan, and C. Wichman, "Effectiveness of the 5A's Model for Changing Physical Activity Behaviours in Rural Adults Recruited From Primary Care Clinics," *Journal of Physical Activity and Health* 16, no. 12 (2019): 1138–1146.
28. S. Gao, R. A. Stone, L. J. Hough, et al., "Physical Activity Counselling in Overweight and Obese Primary Care Patients: Outcomes of the VA-STRIDE Randomized Controlled Trial," *Preventive Medicine Reports* 3 (2016): 113–120.
29. T. Harris, S. M. Kerry, C. R. Victor, et al., "A Primary Care Nurse-Delivered Walking Intervention in Older Adults: PACE (Pedometer Accelerometer Consultation Evaluation)-Lift Cluster Randomised Controlled Trial," *PLoS Medicine* 12, no. 2 (2015): e1001783.
30. T. Hinrichs, B. Bücker, R. Klaaßen-Mielke, et al., "Home-Based Exercise Supported by General Practitioner Practices: Ineffective in a Sample of Chronically Ill, Mobility-Limited Older Adults (the HOME fit Randomized Controlled Trial)," *Journal of the American Geriatrics Society* 64, no. 11 (2016): 2270–2279.
31. E. Knight, M. I. Stuckey, and R. J. Petrella, "Prescribing Physical Activity Through Primary Care: Does Activity Intensity Matter?" *The Physician and Sports Medicine* 42, no. 3 (2014): 78–79.
32. G. S. Kolt, G. M. Schofield, N. Kerse, N. Garrett, T. Ashton, and A. Patel, "Healthy Steps Trial: Pedometer-Based Advice and Physical Activity for Low-Active Older Adults," *The Annals of Family Medicine* 10, no. 3 (2012): 206–212.
33. N. Mutrie, O. Doolin, C. F. Fitzsimons, et al., "Increasing Older Adults' Walking Through Primary Care: Results of a Pilot Randomized Controlled Trial," *Family Practice* 29, no. 6 (2012): 633–642.
34. S. van der Weegen, R. Verwey, M. Spreuwenberg, H. Tange, T. van der Weijden, and L. de Witte, "It's LiFe! Mobile and Web-Based Monitoring and Feedback Tool Embedded in Primary Care Increases Physical Activity: A Cluster Randomized Controlled Trial," *Journal of Medical Internet Research* 17, no. 7 (2015): e4579.
35. D. R. Young, M. K. Nguyen, A. Yamamoto, et al., "Telephone-Based Motivational Interviewing Versus Usual Care in Primary Care to Increase Physical Activity: A Randomized Pilot Study," *Pilot and Feasibility Studies* 5, no. 1 (2019): 1–12.
36. T. Harris, S. M. Kerry, E. S. Limb, et al., "Effect of a Primary Care Walking Intervention With and Without Nurse Support on Physical Activity Levels in 45-to 75-Year-Olds: The Pedometer and Consultation Evaluation (PACE-UP) Cluster Randomised Clinical Trial," *PLoS Medicine* 14, no. 1 (2017): e1002210.
37. E. Knight, M. I. Stuckey, and R. J. Petrella, "Health Promotion Through Primary Care: Enhancing Self-Management With Activity Prescription and mHealth," *The Physician and Sports Medicine* 42, no. 3 (2014): 90–99.
38. M. C. Morey, C. F. Pieper, D. E. Edelman, et al., "Enhanced Fitness: A Randomized Controlled Trial of the Effects of Home-Based Physical Activity Counselling on Glycaemic Control in Older Adults With Prediabetes Mellitus," *Journal of the American Geriatrics Society* 60, no. 9 (2012): 1655–1662.
39. Z. H. Lewis, K. J. Ottenbacher, S. R. Fisher, et al., "Effect of Electronic Activity Monitors and Pedometers on Health: Results From the TAME Health Pilot Randomized Pragmatic Trial," *International Journal of Environmental Research and Public Health* 17, no. 18 (2020): 6800.
40. J. Rossen, K. Larsson, M. Hagströmer, et al., "Effects of a Three-Armed Randomised Controlled Trial Using Self-Monitoring of Daily Steps With and Without Counselling in Prediabetes and Type 2 Diabetes—The Sophia Step Study," *International Journal of Behavioural Nutrition and Physical Activity* 18, no. 1 (2021): 1–11.

41. P. Y. Takahashi, S. M. Quigg, I. T. Croghan, D. R. Schroeder, and J. O. Ebbert, "Effect of Pedometer Use and Goal Setting on Walking and Functional Status in Overweight Adults With Multimorbidity: A Crossover Clinical Trial," *Clinical Interventions in Aging* 11 (2016): 1099.
42. H. Westland, M. J. Schuurmans, I. D. Bos-Touwen, et al., "Effectiveness of the Nurse-Led Activate Intervention in Patients at Risk of Cardiovascular Disease in Primary Care: A Cluster-Randomised Controlled Trial," *European Journal of Cardiovascular Nursing* 19, no. 8 (2020): 721–731.
43. T. Yates, C. L. Edwardson, J. Henson, et al., "Walking Away From Type 2 Diabetes: A Cluster Randomized Controlled Trial," *Diabetic Medicine* 34, no. 5 (2017): 698–707.
44. S. Pears, M. Bijker, K. Morton, et al., "A Randomised Controlled Trial of Three Very Brief Interventions for Physical Activity in Primary Care," *BMC Public Health [Electronic Resource]* 16, no. 1 (2016): 1–13.
45. N. Heron, M. A. Tully, M. C. McKinley, and M. E. Cupples, "Steps to a Better Belfast: Physical Activity Assessment and Promotion in Primary Care," *British Journal of Sports Medicine* 48, no. 21 (2014): 1558–1563.
46. T. Vetrovsky, J. Cupka, M. Dudek, et al., "A Pedometer-Based Walking Intervention With and Without Email Counselling in General Practice: A Pilot Randomized Controlled Trial," *BMC Public Health [Electronic Resource]* 18, no. 1 (2018): 1–13.
47. P. Agarwal, N. Kithulegoda, Z. Bouck, et al., "Feasibility of an Electronic Health Tool to Promote Physical Activity in Primary Care: Pilot Cluster Randomized Controlled Trial," *Journal of Medical Internet Research* 22, no. 2 (2020): e15424.
48. W. Hardeman, J. Mitchell, S. Pears, et al., "Evaluation of a Very Brief Pedometer-Based Physical Activity Intervention Delivered in NHS Health Checks in England: The VBI Randomised Controlled Trial," *PLoS Medicine* 17, no. 3 (2020): e1003046.
49. K. Gallegos-Carrillo, C. García-Peña, J. Salmerón, N. Salgado-de-Snyder, and F. Lobelo, "Brief Counselling and Exercise Referral Scheme: A Pragmatic Trial in Mexico," *American Journal of Preventive Medicine* 52, no. 2 (2017): 249–259.
50. K. Khunti, S. Griffin, A. Brennan, et al., "Promoting Physical Activity in a Multi-Ethnic Population at High Risk of Diabetes: The 48-Month PROPELS Randomised Controlled Trial," *BMC Medicine* 19, no. 1 (2021): 1–11.
51. L. H. Yaacob and A. K. Azidah, "Pedometer-Based Walking Intervention With and Without Group Support Among Sedentary Adults in Primary Care Patients in North-East Malaysia: A Randomized Controlled Trial," *Bangladesh Journal of Medical Science* 17, no. 1 (2018): 52–57.
52. A. G. Huebschmann, R. E. Glasgow, I. M. Leavitt, et al., "Integrating a Physical Activity Coaching Intervention Into Diabetes Care: A Mixed-Methods Evaluation of a Pilot Pragmatic Trial," *Translational Behavioral Medicine* 12, no. 4 (2022): 601–610.
53. S. Pears, M. Bijker, K. Morton, et al., "A Randomised Controlled Trial of Three Very Brief Interventions for Physical Activity in Primary Care," *BMC Public Health [Electronic Resource]* 16, no. 1 (2016): 1–13.
54. A. Riera-Sampol, M. Bennasar-Veny, P. Tauler, and A. Aguilo, "Effectiveness of Physical Activity Prescription by Primary Care Nurses Using Health Assets: A Randomized Controlled Trial," *Journal of Advanced Nursing* 77, no. 3 (2021): 1518–1532.
55. T. Vetrovsky, J. Cupka, M. Dudek, et al., "Mental Health and Quality of Life Benefits of a Pedometer-Based Walking Intervention Delivered in a Primary Care Setting," *Acta Gymnica* 47, no. 3 (2017): 138–143.
56. M. C. Morey, C. F. Pieper, D. E. Edelman, et al., "Enhanced Fitness: A Randomized Controlled Trial of the Effects of Home-Based Physical Activity Counselling on Glycaemic Control in Older Adults With Prediabetes Mellitus," *Journal of the American Geriatrics Society* 60, no. 9 (2012): 1655–1662.
57. D. A. Chambers, R. E. Glasgow, and K. C. Stange, "The Dynamic Sustainability Framework: Addressing the Paradox of Sustainment Amid Ongoing Change," *Implementation Science* 8 (2013): 1–11.
58. A. J. Milat, L. King, R. Newson, et al., "Increasing the Scale and Adoption of Population Health Interventions: Experiences and Perspectives of Policy Makers, Practitioners, and Researchers," *Health Research Policy and Systems* 12 (2014): 1–1.
59. S. McCrabb, C. Lane, A. Hall, et al., "Scaling-Up Evidence-Based Obesity Interventions: A Systematic Review Assessing Intervention Adaptations and Effectiveness and Quantifying the Scale-Up Penalty," *Obesity Reviews* 20, no. 7 (2019): 964–982.
60. G. Marcu, S. J. Ondersma, A. N. Spiller, B. M. Broderick, R. Kadri, and L. R. Buis, "The Perceived Benefits of Digital Interventions for Behavioural Health: Qualitative Interview Study," *Journal of Medical Internet Research* 24, no. 3 (2022): e34300.
61. H. Christensen and K. M. Griffiths, "The Prevention of Depression Using the Internet," *The Medical Journal of Australia* 177, no. 7 (2002): S122.
62. F. Douglas, N. Torrance, E. van Teijlingen, S. Meloni, and A. Kerr, "Primary Care Staff's Views and Experiences Related to Routinely Advising Patients About Physical Activity: A Questionnaire Survey," *BMC Public Health [Electronic Resource]* 6 (2006): 138.
63. G. B. Samdal, G. E. Eide, T. Barth, G. Williams, and E. Meland, "Effective Behaviour Change Techniques for Physical Activity and Healthy Eating in Overweight and Obese Adults: Systematic Review and Meta-Regression Analyses," *International Journal of Behavioral Nutrition and Physical Activity* 14, no. 1 (2017): 42.
64. L. Lammings, S. Pears, D. Mason, et al., "What Do We Know About Brief Interventions for Physical Activity That Could Be Delivered in Primary Care Consultations? A Systematic Review of Reviews," *Preventive Medicine* 99 (2017): 152–163.
65. F. C. Warren, K. Stych, M. Thorogood, et al., "Evaluation of Different Recruitment and Randomisation Methods in a Trial of General Practitioner-Led Interventions to Increase Physical Activity: A Randomised Controlled Feasibility Study With Factorial Design," *Trials* 15, no. 1 (2014): 1–11.
66. F. C. Bull, S. S. Al-Ansari, S. Biddle, et al., "World Health Organization 2020 Guidelines on Physical Activity and Sedentary Behaviour," *British Journal of Sports Medicine* 54, no. 24 (2020): 1451–1462.
67. Faculty of Sports and Exercise Medicine. Moving Medicine, 2024, assessed March 31, 2024, <https://movingmedicine.ac.uk/>.

Supporting Information

Additional supporting information can be found online in the Supporting Information section.