

Editorial: Sedentary behaviour and cardiometabolic health.

AZEVEDO, Liane <<http://orcid.org/0000-0001-9966-9414>>, CUCATO, Gabriel Grizzo and MORSETH, Bente

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/34359/>

This document is the Published Version [VoR]

Citation:

AZEVEDO, Liane, CUCATO, Gabriel Grizzo and MORSETH, Bente (2024). Editorial: Sedentary behaviour and cardiometabolic health. *Frontiers in Cardiovascular Medicine*, 11: 1498410. [Article]

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>



OPEN ACCESS

EDITED AND REVIEWED BY
Elsayed Z. Soliman,
Wake Forest University, United States

*CORRESPONDENCE

Liane Beretta De Azevedo
✉ Lazevedo@shu.ac.uk

RECEIVED 18 September 2024

ACCEPTED 04 October 2024

PUBLISHED 16 October 2024

CITATION

Azevedo LBD, Cucato GG and Morseth B
(2024) Editorial: Sedentary behaviour and
cardiometabolic health.

Front. Cardiovasc. Med. 11:1498410.
doi: 10.3389/fcvm.2024.1498410

COPYRIGHT

© 2024 Azevedo, Cucato and Morseth. This is
an open-access article distributed under the
terms of the [Creative Commons Attribution
License \(CC BY\)](#). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that the
original publication in this journal is cited, in
accordance with accepted academic practice.
No use, distribution or reproduction is
permitted which does not comply with these
terms.

Editorial: Sedentary behaviour and cardiometabolic health

Liane Beretta De Azevedo^{1*}, Gabriel Grizzo Cucato² and
Bente Morseth³

¹College of Health, Wellbeing and Life Sciences, Sheffield Hallam University, Sheffield, United Kingdom, ²School of Design, Northumbria University, Newcastle upon Tyne, United Kingdom, ³School of Sport Sciences, UiT The Arctic University of Norway, Tromsø, Norway

KEYWORDS

sedentary behavior, cardiovascular, cardiometabolic, public health, behaviour change

Editorial on the Research Topic

Sedentary behaviour and cardiometabolic health

Sedentary behaviour is a growing public health issue, with the latest World Health Organisation (WHO) physical activity guidelines now including recommendations regarding sedentary behaviour due to the increasing body of evidence showing the harmful effects of sedentary behaviour on health outcomes (1). In an overview of 18 systematic reviews, Saunders et al., (2) concluded that sedentary behaviour was associated with depression, poor cognitive function and low physical-related quality of life. The review of reviews also highlighted that breaking up sedentary behaviour may improve body composition and markers of cardiometabolic risks. It has been reported that Americans spend 55% of their waking time (7.7 h/day) on sedentary behaviour (3). A more recent epidemiological study in 28 European countries showed an increase in the prevalence of sedentary behaviour (defined by sitting for longer than 4 h 40 min a day) for both men and women between 2002 and 2017, with a higher prevalence in men (4).

Although sedentary behaviour is on the rise, cardiovascular disease and mortality incidence have been declining since 1990, especially in developed countries compared to developing counterparts. This decline is associated with better access to medication and a healthy lifestyle (5). However, cardiovascular disease remains the most prevalent non-communicable disease worldwide. It is estimated that CVD causes 17.9 million deaths per year, representing 32% of global deaths (6).

This Research Topic contains studies that discuss the prevalence of sedentary behaviour and its potential mechanisms. It also explores the impact of breaking up sedentary time on cardiometabolic markers. Lastly, it examines men's perception of the threat of cardiovascular diseases and the effectiveness of behaviour change.

The study by Liang et al., analysed a sample of 5,610 respondents with cardiovascular disease from the US National Health and Nutrition Examination Survey (NHANES). In contrast to the study by Lopes-Valenciano (4), which found an increase in prevalence in European countries, Liang et al. found that the prevalence of sedentary behaviour, physical activity levels, and smoking remained stable between 1990 and 2000 and 2017–2018 in the USA. However, the prevalence of poor diet decreased. Surprisingly, there was a significant increase in obesity, which is difficult to explain considering the reduction in poor diet and maintenance of physical activity and sedentary behaviour. The authors, however, performed a sub-analysis which showed that the group with low education and income were at a higher risk of being exposed to unhealthy lifestyle

factors such as sedentary behaviour compared to those at the other end of the spectrum (i.e., high education and income levels), justifying the needs for targeted interventions for this population.

Two studies in this Research Topic explored the mechanisms of the association between sedentary behaviour and cardiometabolic markers. It is known that high levels of leptin are related to obesity and cardiovascular disease (7). In a cross-sectional study, **Shih et al.** found a positive association between the high leptin group and sedentary behaviour, after adjusting to other behaviours and metabolic markers. The association between sedentary time and leptin levels was observed in individuals with both high BMI and normal BMI, suggesting that BMI was not the sole reason for the increase in leptin levels. The authors suggested that other factors, such as elevated sympathetic tone, may also play a role (8). The authors concluded that sedentary behaviour independently contributes to increased leptin levels and highlighted the importance of reducing sedentary behaviour to lower leptin levels and prevent cardiovascular diseases.

On the other hand, **Vandercappellen et al.** investigated the association between sedentary behaviour and physical activity levels with biomarkers of cardiac injury in a large population (The Maastricht study). After controlling for demographic and other lifestyle risk factors, they did not find an association between physical activity, sedentary time, and cardiac troponins, contradicting previous findings from the literature (9). However, they found that vigorous and moderate to vigorous physical activity were associated with lower levels of NT-pro BNP (a biomarker often associated with cardiac stress and heart failure), suggesting that high-intensity physical activity may benefit cardiovascular function.

The literature has highlighted the adverse effects of continuous and prolonged sedentary behaviour, which is still a prevalent issue (10). **Silva et al.** assessed sedentary behaviour using accelerometers in a sample of 184 adults aged 18–59 years. The authors observed that younger adults spent more time in sedentary bouts (uninterrupted sedentary behaviour) than middle-aged adults. However, time on sedentary breaks (interruption of sedentary bouts with physical activity) was similar. The authors attribute the increase in the use of interactive electronic devices by the younger population as a potential reason for the prolonged time in sedentary bouts. However, while other studies have found an association between cardiovascular markers and sedentary behaviour patterns (11), only age showed a significant correlate in this study.

Domosławska-Zylińska et al. conducted a survey of 1,000 Polish men to assess their perceived susceptibility to

cardiovascular disease and the efficacy of preventive behaviours such as physical activity. Despite 54.2% recognising the severity of cardiovascular disease, only 15.1% perceived themselves as being at risk. Most individuals expressed confidence in the effectiveness of preventive measures, including regular physical activity, a healthy diet, and medication adherence. However, the responsive (high threat, high efficacy) and proactive (low threat, high efficacy) groups reported higher engagement in preventive behaviours compared to the indifferent group (low threat, low efficacy). The study concluded that increasing self-efficacy is essential to improving preventive behaviour uptake in men.

The articles in this Research Topic offer additional evidence regarding the connection between sedentary behaviour and cardiovascular health. This includes insights into the mechanisms of the association, prevalence and patterns and perceived risks of cardiovascular disease, and the efficacy of modifying preventative behaviours. While more research is needed to guide public health guidelines on the relationship between sedentary behaviour and cardiometabolic outcomes, we hope that this issue has contributed to advancing our knowledge in this area.

Author contributions

LA: Conceptualization, Writing – original draft, Writing – review & editing. GC: Conceptualization, Writing – review & editing. BM: Conceptualization, Writing – review & editing.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

References

1. Bull FC, Al-Ansari SS, Biddle S, Borodulin K, Buman MP, Cardon G, et al. World health organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med.* (2020) 54(24):1451–62. doi: 10.1136/bjsports-2020-102955
2. Saunders TJ, McIsaac T, Douillette K, Gaulton N, Hunter S, Rhodes RE, et al. Sedentary behaviour and health in adults: an overview of systematic reviews. *Appl Physiol Nutr Metab.* (2020) 45(10(Suppl. 2)):S197–S217. doi: 10.1139/apnm-2020-0272
3. Matthews CE, Chen KY, Freedson PS, Buchowski MS, Beech BM, Pate RR, et al. Amount of time spent in sedentary behaviors in the United States, 2003–2004. *Am J Epidemiol.* (2008) 167(7):875–81. doi: 10.1093/aje/kwm390
4. López-Valenciano A, Mayo X, Liguori G, Copeland RJ, Lamb M, Jimenez A. Changes in sedentary behaviour in European union adults between 2002 and 2017. *BMC Public Health.* (2020) 20(1):1206–1. doi: 10.1186/s12889-020-09293-1
5. Amini M, Zayeri F, Salehi M. Trend analysis of cardiovascular disease mortality, incidence, and mortality-to-incidence ratio: results from global burden of disease study 2017. *BMC Public Health.* (2021) 21(1):401–0. doi: 10.1186/s12889-021-10429-0
6. World Health Organization. Cardiovascular diseases. (2021). Available online at: <https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-cvds> (accessed September 18, 2024)

7. Katsiki N, Mikhailidis DP, Banach M. Leptin, cardiovascular diseases and type 2 diabetes mellitus. *Acta Pharmacol Sin.* (2018) 39(7):1176–88. doi: 10.1038/aps.2018.40
8. Chantry AJ, Bishop NC, Hamer M, Paine NJ. Sedentary behaviour, but not moderate-to-vigorous physical activity, is associated with respiratory responses to acute psychological stress. *Biol Psychol.* (2023) 177:108510. doi: 10.1016/j.biopsycho.2023.108510
9. Aengevaeren VL, Hopman MTE, Thompson PD, Bakker EA, George KP, Thijssen DHJ, et al. Exercise-induced cardiac troponin I increase and incident mortality and cardiovascular events. *Circulation.* (2019) 140(10):804–14. doi: 10.1161/CIRCULATIONAHA.119.041627
10. Chandrasekaran B, Pesola AJ, Rao CR, Arumugam A. Does breaking up prolonged sitting improve cognitive functions in sedentary adults? A mapping review and hypothesis formulation on the potential physiological mechanisms. *BMC Musculoskelet Disord.* (2021) 22(1):274–5. doi: 10.1186/s12891-021-04136-5
11. English C, Janssen H, Crowfoot G, Callister R, Dunn A, Mackie P, et al. Breaking up sitting time after stroke (BUST-stroke). *Int J Stroke.* (2018) 13(9):921–31. doi: 10.1177/1747493018801222