

Risk factors for the development of knee osteoarthritis across the lifespan: a systematic review and meta-analysis.

DUONG, Vicky, SHAHEED, Christina Abdel, FERREIRA, Manuela L, NARAYAN, Sujita W, VENKATESHA, Venkatesha, HUNTER, David J, ZHU, Jimmy, ATUKORALA, Inoshi, KOBAYASHI, Sarah, GOH, Siew Li, BRIGGS, Andrew M, CROSS, Marita, ESPINOSA-MORALES, Rolando, FU, Kai, GUILLEMIN, Francis, KEEFE, Francis, STEFAN LOHMANDER, L, MARCH, Lyn, MILNE, George J, MEI, Yifang, MOBASHERI, Ali, NAMANE, Mosedi, PEAT, George <<http://orcid.org/0000-0002-9008-0184>>, RISBERG, May Arna, SHARMA, Saurab, SIT, Regina, TELLES, Rosa Weiss, ZHANG, Yuqing and COOPER, Cyrus

Available from Sheffield Hallam University Research Archive (SHURA) at:

<https://shura.shu.ac.uk/35657/>

This document is the Published Version [VoR]

Citation:

DUONG, Vicky, SHAHEED, Christina Abdel, FERREIRA, Manuela L, NARAYAN, Sujita W, VENKATESHA, Venkatesha, HUNTER, David J, ZHU, Jimmy, ATUKORALA, Inoshi, KOBAYASHI, Sarah, GOH, Siew Li, BRIGGS, Andrew M, CROSS, Marita, ESPINOSA-MORALES, Rolando, FU, Kai, GUILLEMIN, Francis, KEEFE, Francis, STEFAN LOHMANDER, L, MARCH, Lyn, MILNE, George J, MEI, Yifang, MOBASHERI, Ali, NAMANE, Mosedi, PEAT, George, RISBERG, May Arna, SHARMA, Saurab, SIT, Regina, TELLES, Rosa Weiss, ZHANG, Yuqing and COOPER, Cyrus (2025). Risk factors for the development of knee osteoarthritis across the lifespan: a systematic review and meta-analysis. *Osteoarthritis and cartilage*, 33 (10), 1162-1179. [Article]

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

Osteoarthritis and Cartilage



Risk factors for the development of knee osteoarthritis across the lifespan: A systematic review and meta-analysis



Vicky Duong ^{# * 1 2}, Christina Abdel Shaheed ^{# † ** 1 3}, Manuela L. Ferreira ^{‡ 4},
 Sujita W. Narayan ^{# † 5}, Venkatesha Venkatesha [§], David J. Hunter ^{# ¶ 6}, Jimmy Zhu ^{¶ || 7},
 Inoshi Atukorala ^{## 8}, Sarah Kobayashi ^{# 9}, Siew Li Goh ^{†† ‡‡ 10}, Andrew M. Briggs ^{§§ 11},
 Marita Cross ^{# ¶ 12}, Rolando Espinosa-Morales ^{¶¶ 13}, Kai Fu ^{||| 14}, Francis Guillemin ^{### 15},
 Francis Keefe ^{††† 16}, L. Stefan Lohmander ^{‡‡‡ 17}, Lyn March ^{# ¶ 18}, George J. Milne ^{§§§ 19},
 Yifang Mei ^{¶¶¶ 20}, Ali Mobasher ^{||||| ##### †††† ‡‡‡‡ 21}, Mosedi Namane ^{§§§§}, George Peat ^{¶¶¶¶ 22},
 May Arna Risberg ^{|||||| ##### 23}, Saurab Sharma ^{††††† ‡‡‡‡‡ §§§§§ 24}, Regina Sit ^{¶¶¶¶¶ 25},
 Rosa Weiss Telles ^{||||||| ##### 26}, Yuqing Zhang ^{†††††† 27}, Cyrus Cooper ^{‡‡‡‡‡ §§§§§ 28}

[#] Sydney Musculoskeletal Health, Kolling Institute, Faculty of Medicine and Health, University of Sydney, Level 10, St Leonards, Sydney 2065, New South Wales, Australia
[†] Faculty of Medicine and Health, School of Public Health, University of Sydney, Level 10N, King George V Building, Royal Prince Alfred Hospital, Sydney 2050, New South Wales, Australia

[‡] The George Institute for Global Health, Faculty of Medicine and Health, The University of New South Wales, Level 18, International Towers 3, 300 Barangaroo Ave, Barangaroo, Sydney 2000, New South Wales, Australia

[§] Northern Sydney Local Health District Executive, Royal North Shore Hospital, The Kolling Institute, Level 10, St Leonards, Sydney 2065, New South Wales, Australia

[¶] Department of Rheumatology, Royal North Shore Hospital, Clinical Administration 7C, St Leonards, Sydney 2065, New South Wales, Australia

^{||} Liverpool Hospital, Sydney, New South Wales, Australia

^{##} Department of Clinical Medicine, University of Colombo, Colombo, Sri Lanka

^{††} Centre for Epidemiology and Evidence-Based Practice, Universiti Malaya, Kuala Lumpur, Malaysia

^{‡‡} Sports and Exercise Medicine Research and Education Group, Faculty of Medicine, Universiti Malaya, Kuala Lumpur, Malaysia

^{§§} Curtin School of Allied Health, Curtin University, Perth 6845, Western Australia, Australia

* Correspondence to: The Kolling Institute, Level 10, St Leonards, Sydney 2065, NSW, Australia.

** Correspondence to: Royal Prince Alfred Hospital, Level 10N, King George V Building, PO Box M179, Missenden Road, Sydney 2050, NSW, Australia.

E-mail addresses: vicky.duong@sydney.edu.au (V. Duong), christina.abdelshaheed@sydney.edu.au (C. Abdel Shaheed).

¹ Co-first authors.

² 0000-0002-4450-8194

³ 0000-0003-1258-5125

⁴ 0000-0002-3479-0683

⁵ 0000-0001-8706-2113

⁶ 0000-0003-3197-752X

⁷ 0009-0006-2628-0225

⁸ 0000-0002-1223-6454

⁹ 0000-0001-7816-2198

¹⁰ 0000-0001-5898-1196

¹¹ 0000-0002-6736-3098

¹² 0000-0003-2717-8960

¹³ 0000-0002-3923-6956

¹⁴ 0000-0002-8721-0734

¹⁵ 0000-0002-9860-7024

¹⁶ 0000-0003-0583-9326

¹⁷ 0000-0002-5424-9448

¹⁸ 0000-0003-1736-8111

¹⁹ 0000-0002-9796-0600

²⁰ 0000-0002-0119-5562

²¹ 0000-0001-6261-1286

²² 0000-0002-9008-0184

²³ 0000-0002-9511-1314

²⁴ 0000-0002-9817-5372

²⁵ 0000-0002-4717-7773

²⁶ 0000-0003-4027-2943

²⁷ 0000-0001-7638-0888

²⁸ 0000-0003-3510-0709

<https://doi.org/10.1016/j.joca.2025.03.003>

1063–4584/© 2025 The Author(s). Published by Elsevier Ltd on behalf of Osteoarthritis Research Society International. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

¶¶ National Institute of Rehabilitation, National University Autonomous of Mexico, Mexico City, Mexico
 ||| Department of Orthopedic Surgery, Shanghai Sixth People's Hospital Affiliated to Shanghai Jiao Tong University School of Medicine, No. 600 Yishan Road, Xuhui District, Shanghai 200233, China
 ### Université de Lorraine, Inserm, INSPIRE, Nancy 54500, France
 ††† Duke Pain Prevention and Treatment Research Program, Department of Psychiatry and Behavioral Sciences, Duke University School of Medicine, Durham, NC, USA
 ‡‡‡ Department of Clinical Sciences Lund, Orthopedics, Lund University, PO Box 117, Lund 22100, Sweden
 §§§ Marshall Centre for Infectious Disease Research and Department of Computer Science and Software Engineering, University of Western Australia, 35 Stirling Highway, Crawley 6009, Western Australia, Australia
 ¶¶¶ Department of Rheumatology and Immunology, The Second Affiliated Hospital of Southern University of Science and Technology/The Third People's Hospital of Shenzhen, 29 Bulan Road, Longgang District, Shenzhen 518000, China
 |||| Research Unit of Health Sciences and Technology, Faculty of Medicine, University of Oulu, Oulu, Finland
 #### State Research Institute Centre for Innovative Medicine, Vilnius, Lithuania
 †††† Department of Joint Surgery, First Affiliated Hospital of Sun Yat-sen University, Guangzhou, China
 ‡‡‡‡ World Health Organization Collaborating Center for Public Health Aspects of Musculoskeletal Health and Aging, Université de Liège, Liège, Belgium
 §§§§ Division of Family Medicine, Department of Family, Community and Emergency Medicine, University of Cape Town, Cape Town, South Africa
 ¶¶¶¶ Centre for Applied Health & Social Care Research (CARE), Sheffield Hallam University, Robert Winston Building, Collegiate Campus, Sheffield, United Kingdom
 ||||| Department of Sport Medicine, Norwegian School Sport Sciences, P. Box 4014 Ullevaal Stadion, 0806 Oslo, Norway
 ##### Division of Orthopaedic Surgery, Oslo University Hospital, Oslo, Norway
 ††††† Pain Management and Research Centre, Royal North Shore Hospital, Northern Sydney Local Health District, Douglas Building, St Leonards, Sydney 2065, New South Wales, Australia
 ‡‡‡‡‡ School of Health Sciences, Faculty of Medicine and Health, University of New South Wales, Sydney, New South Wales, Australia
 §§§§§ Pain Management Research Institute, Kolling Institute, Faculty of Medicine and Health, the University of Sydney and Northern Sydney Local Health District, St Leonard's, NSW, Australia
 ¶¶¶¶¶ The Jockey Club School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong
 ||||| Department of Internal Medicine, Faculty of Medicine, Universidade Federal de Minas Gerais, Belo Horizonte, Brazil
 ##### Rheumatology Service, Hospital das Clínicas da UFMG/Ebserh, Belo Horizonte, Brazil
 †††††† Division of Rheumatology, Allergy and Immunology, Massachusetts General Hospital, Harvard Medical School, 55 Fruit Street, Bulfinch 165, Boston, MA 02114, USA
 ‡‡‡‡‡ University of Southampton, Southampton, United Kingdom
 §§§§§ Institute of Musculoskeletal Science, University of Oxford, Oxfordshire, United Kingdom

ARTICLE INFO

Article history:
 Received 20 September 2024
 Accepted 19 March 2025

Keywords:
 Knee osteoarthritis
 Risk factors
 Cohort studies

SUMMARY

Objective: To identify and quantify risk factors for incident knee osteoarthritis (KOA) across the lifespan.
Methods: This systematic review and meta-analysis identified eligible studies from seven electronic databases and three registries. Longitudinal cohort studies or randomised controlled trials evaluating participants who developed incident symptomatic and/or radiographic KOA were included. Two independent reviewers completed data screening and extraction. Estimates were pooled using a random effects model and reported as odds ratio (OR), hazard ratio, or risk ratio and corresponding 95% confidence intervals (95% CI). Grading of Recommendations, Assessment, Development and Evaluation was used to determine the certainty of evidence. Population attributable fractions were calculated, including risk factors significantly associated with radiographic KOA based on the pooled meta-analysis and where we could determine communality scores using existing clinical datasets.

Results: We identified 131 studies evaluating > 150 risk factors. Previous knee injury, older age and high bone mineral density were associated with an increased risk of incident radiographic KOA based on the pooled analysis [OR (95% CI): 2.67 (1.41, 5.05), 1.15 (1.00, 1.33) and 1.82 (1.12, 2.94), respectively], with moderate-to-high certainty. Two risk factors (overweight/obesity and previous knee injury) accounted for 14% of incident radiographic KOA. Other modifiable risk factors, including occupational physical activity, also contribute to radiographic or symptomatic KOA.

Conclusion: Novel strategies addressing known modifiable risk factors including overweight/obesity, knee injuries and occupational physical activity are needed to reduce overall burden of KOA.

Systematic review registration: PROSPERO ID: CRD42023391187

© 2025 The Author(s). Published by Elsevier Ltd on behalf of Osteoarthritis Research Society International. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Knee osteoarthritis (KOA) is a leading cause of disability.^{1,2} The worldwide age-standardised prevalence of KOA was estimated to be 4307 cases (95% Uncertainty Interval: (3845–4913) per 100,000 population).³ Global estimates point to an increasing prevalence of symptomatic KOA with age, however estimates and certainty of point prevalence vary by age group and country.^{4–7} A review estimated global prevalence of approximately 16.0% (95% confidence interval (CI) 14.3–17.8%) in people aged ≥15 years and approximately 22.9% (95% CI 19.8% to 26.1%) in people aged ≥40 years.^{4–6} Lifetime risk for symptomatic KOA has been estimated to be as high as 45%.⁸ This risk increases to 60.5% in obese individuals compared to normal weight or overweight individuals (30.2% or 46.9%, respectively).⁸ Osteoarthritis (OA) is a global public health problem

due to increasing rates of obesity and an ageing population, urging the implementation of wide-scale population-level preventive strategies.⁹

A 2015 review summarised the literature on risk factors for the onset of KOA in adults.¹⁰ That hitherto largest review evaluated just 16 risk factors for incident KOA across 46 studies, was published in 2015, excluded studies not published in English and focussed on a cohort aged ≥50 years.¹⁰ It did not provide Grading of Recommendations Assessment, Development and Evaluation (GRADE) ratings for effect estimates and pooled across outcomes and measures of effect, resulting in pooled estimates with high heterogeneity and therefore, unclear clinical applicability. Other published reviews are either narrative, focus on a specific risk factor, such as knee extensor strength¹¹ or malalignment¹² or do not provide pooled estimates.¹³

A timely update of the evidence is needed to inform KOA prevention and guideline recommendations¹⁴ as several reports have been published since these reviews. Given that prevention of symptomatic and radiographic KOA remains a priority for population health and individual patient care, the aims of this systematic review were to identify and quantify potential risk factors associated with increased or lowered risk of incident KOA and to determine the relative contribution of identified risk factors for incident symptomatic and radiographic KOA across the lifespan.

Methods

This systematic review was registered in the International Prospective Register of Systematic Reviews PROSPERO (CRD42023391187) and the results are reported according to the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) 2020 guidelines.¹⁵

Search strategy and selection criteria

Searches were conducted in MEDLINE, EMBASE, CINAHL, Cochrane Central Register of Controlled Trials (CENTRAL), PsycINFO, the Chinese National Knowledge Infrastructure and Pan African Clinical Trials Registry, ClinicalTrials.gov, the World Health Organization International Clinical Trials Registry Platform and the European (EU) Clinical Trials Register (www.clinicaltrialsregister.eu) from their inception to June 3, 2024. The MEDLINE search strategy was developed in consultation with a librarian at The University of Sydney. The search strategy syntax was then adjusted for the other databases ([Supplemental File 1](#)). The search strategy was supplemented by consultation with content experts and hand searching reference lists of previous reviews.^{10–13}

Articles in any language whereby individuals of any age without baseline KOA were considered. Studies needed to define incident KOA and provide quantitative data regarding the risk of incident KOA within the sample(s). Both radiographic and symptomatic/clinical diagnoses of incident KOA (as reported by the studies) were considered.¹⁶ Cohort studies, and randomised controlled trials (RCTs) in which prevention of OA was stipulated as the primary aim and which followed participants to determine incident KOA, were considered. All types of exposures/interventions were eligible. We excluded conference abstracts, case-control or cross-sectional studies, studies which analysed data from <200 participants,¹⁷ had follow-up of <1 year, solely included participants with KOA at baseline. Studies were excluded if they assessed knee pain exclusively as a risk factor or assessed KOA using magnetic resonance imaging, as we considered these to be evidence of early-stage OA features.

Citations were exported into EndNote and duplicates were removed. Citations were then imported into Covidence (www.covidence.org) for title, abstract and full text screening¹⁸ by two authors independently (VD, CAS, JZ, IA, SLG, SK). Conflicts were resolved by discussion or by an adjudicator (MLF, DJH).

Data extraction

A custom data extraction form was used which included bibliometric data, study/cohort characteristics, incident KOA assessment and classification, treatment or exposure details and outcome data [odds/risk/hazard ratio (HR) estimates and 95% CIs]. Two authors independently extracted data (VD, CAS, SWN, JZ, IA, SLG, SK) and datasheets were checked for completeness by one author (SWN). Discrepancies were resolved through discussion and a third reviewer (MLF, DJH) was consulted if necessary.

Risk of bias

Risk of bias was assessed independently by two authors (VD, CAS, SWN, JZ, SK, SLG, IA) using the Cochrane Risk of Bias tool (RoB-1) for included RCTs and the Modified Quality In Prognosis Studies (QUIPS)¹⁹ for cohort studies.

Data synthesis and analysis

Following data checking, risk estimates, and 95% CIs were transferred into a separate data extraction sheet for analysis. Meta-analyses were conducted to obtain a pooled risk estimate and corresponding 95% CIs when at least two studies reported on the same risk estimate outcome (odds ratio (OR), risk ratio (RR), hazard ratio (HR)) for a given risk factor, met the following criteria:

- (1) Examined the same risk factor and utilised the same measurement approach,
- (2) Assessed incident KOA using the same operational outcome (symptomatic or radiographic KOA etc.),
- (3) Employed a similar analysis approach (univariate or multivariate),
- (4) Had the same unit of analysis (i.e., at the knee or person level).

For pooled analyses, we used adjusted estimates that had adjusted for similar covariates (e.g., age and sex). Random effects meta-analyses were performed using the DerSimonian-Laird method.²⁰ To account for a small number of studies (<5), the truncated Knapp-Hartung standard error adjustment was applied to the pooled effect.

Assessment of heterogeneity and bias

Tests of heterogeneity were conducted using the Q statistic. Variability due to between-study heterogeneity was quantified using the I^2 statistic. Where the I^2 statistic indicated high between-study heterogeneity ($I^2 > 50\%$), a sensitivity analysis was performed using the leave-one-out meta-analysis approach to help identify potential sources of heterogeneity. All meta-analyses were conducted using STATA, version 18.0.

The overall certainty of evidence was assessed using a modified GRADE framework.²¹ Pooled estimates were assigned a certainty of evidence rating (high, moderate, low, very low). Overall certainty of evidence started at high and was downgraded by one level for: (1) risk of bias; (2) inconsistency; (3) imprecision; (4) publication bias ([Supplemental File 2](#)). We did not downgrade for indirectness, as our review encompassed a specific research question and condition. For single cohort studies, overall certainty of evidence started at moderate and was downgraded one level for risk of bias (modified QUIPS rating of high or moderate risk) or imprecision. We used a modified version of the QUIPS tool to assess risk of bias of cohort studies. The six domains considered were: study participation, study attrition, risk factor measurement, outcome measurement, study confounding, and statistical analysis and reporting.

Publication bias

Publication bias was assessed when ten or more studies contributed to an effect estimate using the Begg's test.²²

Population attributable fraction (PAF) calculation

Our PAF calculations included risk factors that were found to be associated with radiographic KOA based on the pooled effect (computed from the adjusted analysis reported in individual studies) and where we

could determine communality scores using existing clinical datasets. There was insufficient data to allow PAF calculations for symptomatic KOA. The RR estimates from multiple studies were pooled using random effects based meta-analysis. The weighted PAF measure was computed using communality-based weights, based on person-level data.

The communality was computed as part of a principal component analysis. A description of the methodology is described in [Supplemental File 3](#). The Osteoarthritis Initiative (OAI) dataset (incidence cohort) was used to compute the communalities of available risk factors, as this dataset is similar to our study context and scope.²³

Protocol deviations

We present data on KOA only and not hand or hip OA, to avoid potential clinical heterogeneity and support a clear clinical message and recommendations.

Decisions to exclude studies with < 200 participants or follow-up < 1 year were to avoid imprecise estimates due to the heterogeneity of the condition and allow sufficient time to observe our outcome of interest, respectively. We used a modified version of QUIPS instead of the Newcastle-Ottawa scale to assess risk of bias of cohort studies.

Results

Of 14,363 records, 6129 duplicates were removed. A total of 8234 titles and abstracts were screened, full texts of 401 records were inspected and 131 papers were included ([Fig. 1](#)). Excluded studies and reasons for exclusion are found in [Supplemental File 4](#).

General characteristics of the included studies

We included 131 studies.^{24–154} Two papers reported on the same RCT and population group but over different time points (2.5 and 6–7 years, respectively)^{24,25} and 129 were cohort studies, published between 1988 and 2024.^{26–154} The most common cohorts included the OAI (n=26),^{32,35,40,47,51,55,57,69,71,73,80,87,88,90,99,103,106,121–123,127,134,136,137,142,146} the Multicentre Osteoarthritis Study (MOST) (n=17),^{36,38,64,66,72,75,82,90,106,109,114,125,128,129,132,135,146} and the Framingham/Framingham Offspring Cohorts (n=13), all of which were conducted in the USA.^{28,49,59,65,68,74,91,94,100,110,131,141,148}

The sample size in the cohort studies ranged from 215⁶⁰ to ≥1.7 million.²⁹ Age of participants ranged from median interquartile range (IQR) 22.8 (20–26)⁸⁴ to a mean standard deviation (SD) of 80.7 (5.0)²⁸ years, with percentages of females ranging from 20%¹⁰² to 100%.^{125,141} Ethnicity was uncommonly reported. The average length of follow-up ranged from 1 year¹⁴⁵ to 40 years⁷⁴ ([Supplemental Files 5 and 6](#)).

Most studies used ORs to report the risk of incident KOA (n=77).^{26–28,30–32,35,36,38–41,45,47–50,53–55,57,59,61,62,65–68,70–76,78–81,85–87,90,95,97,98,100,103,106–108,110–112,115,117,122–124,127–129,132–136,138,139,141,143,146,148–150,153,154} Twenty-six cohort studies reported risk of incident KOA using HR (20.8%).^{29,33,37,42–44,51,56,58,69,77,83,89,92,93,99,101,105,113,118–120,126,137,142,151} Twenty-one cohort studies reported risk of incident KOA using RR (14.6%).^{34,46,52,63,64,82,88,91,94,96,104,109,114,116,121,125,130,131,140,144,147} Five studies used other measures, such as probability estimates or percentages (3.1%).^{60,84,102,145,152}

Over half of the studies (n=73, 56.2%) defined incident KOA using radiographic criteria,^{26,27,30,32,34,35,39,40,44,45,48–51,54,59–62,67–76,78,80–82,}

PRISMA 2020 flow diagram for new systematic reviews which included searches of databases, registers and other sources




Fig. 1

PRISMA flow diagram.

84,89–94,96,97,100–104,107–112,114,118,120,123–126,129,130,133,135–140,142–144,153

most commonly as Kellgren and Lawrence (KL grade) ≥ 2 . Nine studies reported on symptomatic KOA without evidence of radiographic KOA,^{31,37,63,79,85,86,95,116,141} 26 studies reported symptomatic-radiographic KOA^{28,36,41,47,55,57,64–66,87,88,99,106,117,119,121,122,127,128,131,132,134,145,146,148,150} and nine used a combination of approaches (self-report, physician diagnosed).^{38,43,46,98,105,149,151,152,154} Twelve studies used the International Classification of Disease 9th Revision (ICD-9) or 10th Revision (ICD-10) codes^{29,33,42,52,53,56,58,77,83,113,115,147} (Supplemental Files 5 and 6).

Risk of bias of included studies

Of the 129 cohort studies, 90 were appraised as having low risk of bias, 24 had moderate risk of bias and 15 had high risk of bias. Of the two studies which originated from the same RCT, both were low risk of bias (Supplemental Files 7 and 8).

Pooled analysis

Potentially modifiable risk factors (e.g., overweight/obesity, previous knee injury) were considered separately from non-modifiable risk factors (e.g., age, sex). Data extractions for risk estimates are presented in Supplemental Files 9 to 11. Pooling of effect estimates was possible for ten potentially modifiable and non-modifiable risk factors and are presented in Fig. 2, with significant associations demonstrated for five (detailed below). Supplemental Figs. 1–13 present the forest plots, including individual studies.

Over 150 risk factors were reported. Numerous risk factors were associated with a lower incidence of KOA (moderate to low-quality evidence (Fig. 3)). However, most risk factors identified were associated with a higher incidence of KOA (Figs. 4–6).

We present the results from a select number of risk factors where there were sufficiently large sample sizes. Further results are described in the Supplemental File 12. The GRADE ratings are presented in Figs. 3–6.

Potentially modifiable risk factors

Overweight/obesity

Thirty-two studies evaluated body mass index (BMI) as a risk factor for developing incident KOA. Of these, 20 evaluated radiographic KOA and the remaining evaluated symptomatic KOA.

Radiographic KOA

Pooling was possible for five studies,^{26,32,81,112,149} which evaluated BMI as a continuous exposure. For every unit increase in BMI, individuals were 17% more likely to develop radiographic KOA (adjusted OR 1.17, 95% CI 1.10, 1.24). The between-study heterogeneity was high ($I^2 = 52.9\%$). Sensitivity analysis excluding one study¹⁴⁹ reduced I^2 to 47.6%, yet the estimate remained similar (aOR 1.19, 95% CI 1.12–1.27) (Fig. 2). For studies unable to be pooled,^{45,97,98,111,133} the effect estimates ranged from adjusted odds ratio (aOR) 1.6 (1.0, 2.8)⁹⁸ to 18.3 (5.1, 65.1)¹¹¹ for individuals who were at least overweight (BMI ≥ 25 kg/m²).

Four studies investigated the effect of BMI on RR of incident radiographic KOA.^{82,104,109,114} There was a statistically significant association between BMI and medial tibiofemoral OA but not lateral tibiofemoral OA in people who are overweight or obese.¹⁰⁹

Symptomatic KOA

When repeated measures of BMI were considered across the lifespan in a United Kingdom birth cohort study, the greatest odds of




Fig. 2 Meta-analysis results for all pooled studies. Abbreviations: ROA, radiographic osteoarthritis; SxOA, symptomatic osteoarthritis; ICD OA, osteoarthritis defined using International Classification of Disease codes; Q2/Q3/Q4, quartiles of physical activity.




Fig. 3

Risk factors associated with lower incidence of knee OA. *For referent categories, refer to the [Supplemental Files 9–11](#). Abbreviations: ROA, radiographic osteoarthritis; SxOA, symptomatic osteoarthritis; ICD OA, osteoarthritis defined using International Classification of Disease codes; aMED, alternate Mediterranean diet score; HDL, high-density lipoprotein; HOMA-IR, homeostatic model assessment for insulin; Q2/Q3/Q4, quartiles of fibre consumption; MMSE, Mini-Mental State Examination; mtDNA, mitochondrial DNA.

incident symptomatic KOA was found for high BMI at 43 years of age (aOR 2.64; 95% CI 1.8, 3.87).⁸⁵ However, two studies evaluating symptomatic KOA found conflicting results for individuals with BMI ≥ 25 kg/m² in Korean and Chinese adults, with statistically significant results found in the Korean, but not the Chinese cohort.^{79,133} One study reporting on HRs found Japanese females and males with BMI ≥ 25 kg/m² were at higher risk of developing incident symptomatic KOA, 4.13 (95% CI 2.15, 7.94) and 7.74 (95% CI 1.07, 55.91) respectively.⁹³

Previous knee injury

Eleven studies^{28,56,60,76,98,109–111,113,115,144} investigated the impact of previous knee injuries (self-reported injury resulting in impaired function (e.g., inability to walk for ≥ 2 days))¹⁰⁹ on incident KOA. Most studies investigated anterior cruciate ligament (ACL) injuries with/without meniscus injuries.

Radiographic KOA

Six studies reporting ORs for the association between past knee injury and incident radiographic KOA were pooled and found an almost three-fold increased odds (aOR 2.67 95% CI 1.41, 5.05)^{26,76,78,90,110,111} (Fig. 2). Previous knee injury was associated with incident radiographic KOA in both medial and lateral compartments in the MOST and OAI cohorts, the adjusted risk ratio (aRR) ranged from 1.57 (95% CI 1.13, 2.19) to 2.08 (95% CI 1.3, 3.31).

Habitual/Leisure-time physical activity level

Fourteen studies^{40,58,65,73,74,89,94,98,99,106,110,127,148,153} reported on leisure-time physical activity using various measures. Two studies reporting ORs were pooled.^{74,110} These studies measured physical activity using the Framingham Physical Activity Index. The pooled estimate showed no association between physical activity quartiles and incident radiographic KOA in elderly patients (ranging from




Fig. 4 Risk factors associated with higher incidence of knee OA (relative risk). *For referent categories, refer to Supplemental File 9. Abbreviations: ROA, radiographic osteoarthritis; SxOA, symptomatic osteoarthritis; ICD OA, osteoarthritis defined using International Classification of Disease codes; Q2/Q3/Q4, quartiles of physical activity.

Quartile 1 (sedentary) to Quartile 4 (high levels of leisure-time physical activity) (Fig. 2)). A further seven studies^{40,58,65,89,98,99,106} showed no association between leisure-time physical activity and incident radiographic or symptomatic KOA. Two studies^{94,127} found that high levels of leisure-time physical activity in overweight or obese individuals was associated with incident KOA, with one showing that the weight-associated risk for KOA was increased in women with high levels of physical activity; RR 3.6 (1.7 to 7.7) for the heaviest quintile.⁹⁴ Similar trends were observed for men. One study showed that football participation, including in the teen years was associated with greater risk of incident radiographic or symptomatic KOA.⁷³ Total recreational physical activity and weight bearing physical activity in individuals with low levels of lower-limb muscle mass were found to be associated with incident radiographic KOA; OR 1.20 (1.07, 1.34) and 1.21 (1.09, 1.35) respectively.¹⁵³ Heavy levels of daily habitual physical activity (≥ 1 h per day) were found to be associated with incident radiographic KOA in one study (OR), whereas no associations were found between light or moderate levels of daily

habitual physical activity (≥ 1 h per day) and incident radiographic KOA.¹⁴⁸ **Occupational physical activity** Nine studies investigated occupational risk factors, including occupational physical activity and their association with KOA.^{42,58,80,93,98,100,119,133,146} Eight showed a clear association between occupational physical activity and incident radiographic and/or symptomatic KOA. This included work that required walking while handling materials (compared to jobs mostly involving sitting),⁸⁰ and occupations which combine knee bending and physical demands.¹⁰⁰ One study reporting on HRs, did not find an association between type of work (office work, professional, manual, jobless or housewives) and incident symptomatic KOA in a Japanese cohort.⁹³ Three studies reported on occupational job category and/or workplace physical activity.^{80,133,146} They found that manual work (ranging from light, light manual and heavy), were associated with incident radiographic KOA, symptomatic-radiographic KOA and




Fig. 5

Risk factors associated with higher incidence of knee OA (odds ratio). *For referent categories, refer to [Supplemental File 10](#). Abbreviations: ROA, radiographic osteoarthritis; SxOA, symptomatic osteoarthritis; ICD OA, osteoarthritis defined using International Classification of Disease codes; TF, tibiofemoral; ATP III, National Cholesterol Education Program Adult Treatment Panel III; BP, blood pressure; HDL, high-density lipoprotein.

symptomatic KOA. Men had a two-fold greater risk of incident radiographic KOA if they engaged in heavy, manual occupations (aRR 2.7 95% CI 1.17, 6.26).¹⁴⁶ However, an association was not seen for women.

Manual handling of loads > 20 kg for ≥13 years, was associated with incident KOA (ICD criteria) (adjusted hazard ratio (aHR) 1.71 95% CI 1.03, 2.85).⁵⁸ Another study reported associations between shift work exceeding 20 years and incident radiographic KOA (aHR 1.28 95% CI 1.04, 1.57).¹¹⁹

Bone mineral density (BMD)

Three studies reporting on ORs evaluated the association between areal BMD and radiographic KOA.^{27,49,50} Pooled results from two studies found an association between higher femoral neck BMD (mean femoral neck BMD ≥1.01 g/cm² at baseline) and incident radiographic KOA (aOR 1.82 95% CI 1.12, 2.94). A higher risk of incident radiographic KOA was also reported with the highest quartile

of lumbar spine BMD at baseline in one study (mean BMD: 1.32 g/cm², aOR 4.7 95% CI 2.1, 10.7).⁵⁰

Non-modifiable risk factors

Sex and gender

Nineteen studies assessed female sex or gender as a potential risk factor for KOA.

Radiographic KOA

Pooled results from seven studies found that females had almost double the odds of developing incident radiographic KOA compared to males (aOR 1.93 95% CI 1.25, 2.98).^{26,32,76,90,110,112,133} The between-study heterogeneity was high ($I^2 = 63.5\%$). Sensitivity analysis excluding a study³² reduced I^2 to 30.4% and the effect remained significant (aOR 2.22, 95% CI 1.62, 3.05) (Fig. 2). Another study reporting on HRs found that males were ~40% less likely to develop incident




Fig. 6 Risk factors associated with higher incidence of knee OA (hazard ratio). *For referent categories, refer to Supplemental File 11. Abbreviations: ROA, radiographic osteoarthritis; SxOA, symptomatic osteoarthritis; ICD OA, osteoarthritis defined using International Classification of Disease codes; Q2/Q3/Q4, quartiles of physical activity.

radiographic KOA (aHR 0.59 95% CI 0.49, 0.71).⁹² Two studies unable to be pooled found female sex was associated with radiographic KOA in the OAI cohort,^{92,109} but not in the MOST and the Cohort Hip and Knee (CHECK) cohorts.^{92,109}

Symptomatic KOA

The pooled estimate from two studies evaluating symptomatic KOA was not statistically significant.^{79,133} Males with uncontrolled diabetes (HbA1C ≥ 7%) were more likely to develop incident symptomatic KOA (aHR 1.62 95% CI 1.08, 2.43), compared to their female counterparts (aHR 1.36 95% CI 0.97, 1.93).³⁷ One study evaluating RRs found that females had almost double the risk of developing symptomatic KOA (aRR 1.96 95% CI 1.01, 3.82).¹³¹

Age

Fourteen studies evaluated the association between age and incident KOA.

Radiographic KOA

Three studies reporting on ORs found an association between age and radiographic KOA, with increasing risk as age increased (pooled aOR 1.15 95% CI 1.00, 1.33).^{76,90,112} The between-study heterogeneity was high ($I^2 = 89.3\%$). Sensitivity analysis excluding one study¹¹² reduced the I^2 to 21.8% and the pooled effect remained significant per unit increase in age (aOR 1.06, 95% CI 1.02, 1.10) (Fig. 2). One study reporting on HRs did not find an association between age and incident radiographic KOA.⁹²

Three studies reporting RRs found associations between age and radiographic KOA.^{94,104,109} In one study, associations between age and radiographic KOA were found for females but not males across all age categories.⁹⁴

Symptomatic KOA

One study reporting on ORs, did not find any association between age and symptomatic KOA across any age category.⁷⁹ Two studies reporting on HRs, found conflicting results regarding the association between age and symptomatic KOA. One study found significant

Group	Risk factor for radiographic KOA	Communality	Unweighted PAF	Weighted PAF	Percent weighted PAF
Non-modifiable	Age	–	0.935	–	
	Sex	–	0.413	–	
	Hand OA	–	0.344	–	
	Overall weighted PAF	–	–	–	
Modifiable	Being overweight or obese	0.061	0.119	0.102	77.20
	Previous knee injury	0.045	0.035	0.030	22.80
	Overall weighted PAF	–	–	0.141	100

Risk factor definition for computation of PAF: Age - < 50 years vs ≥ 50 years, Sex- female vs male, Hand OA - yes/no, Obesity/overweight - yes vs. no, Previous knee injury - yes vs no.

Table 1

Principal component analysis/Communality Weighted Population Attributable Fractions for risk factors by modifiable and non-modifiable group.

associations across all comparisons and that females ≥70 years had a higher aHR for developing symptomatic KOA (13.51 95% CI 7.19, 25.4), compared to females aged 60–69 (5.89 95% CI 3.38, 10.25) or females aged 50–59 (3.13 95%CI 1.77, 5.56).⁹³ Males ≥70 years had a higher aHR for developing symptomatic KOA (7.36 95% CI 2.94, 18.43), compared to males aged 60–69 (5.73 95% CI 2.57, 12.77) or males aged 50–59 (4.07 95% CI 1.80, 9.20).⁹³ Another study found no associations across all comparisons except in individuals aged < 65 with uncontrolled diabetes (HbA1c ≥ 7%) where an association was seen (1.51 95% CI 1.12, 2.03).³⁷

Additional findings

Based on moderate certainty evidence, parity (5–12 births), poor sleep quality, high intake of ultra-processed foods and depressive symptoms were associated with higher risk of incident KOA (symptomatic or radiographic) (Figs. 4–6). Conversely, there was moderate certainty evidence that a mediterranean diet, high dietary fibre and consumption of cereal, dark bread and green tea were associated with lower risk of incident KOA (Fig. 3).

PAF

Table 1 presents the communality, unweighted and weighted PAFs adjusted for communality.

Fig. 7 shows the life course model of potentially modifiable risk factors for incident radiographic knee OA. Together, two modifiable risk factors (overweight/obesity and previous knee injury) account for 14% of incident radiographic KOA.

Discussion

This review identified > 150 risk factors related to incident radiographic and/or symptomatic KOA. There is high to moderate certainty evidence that overweight/obesity, previous knee injury, higher femoral neck spine BMD and occupational physical activity (e.g., shift work) are associated with an increased risk of radiographic KOA. There is moderate to low certainty evidence that female sex and older age are associated with incident KOA. Pooled estimates show a lack of association between the following risk factors and incident radiographic KOA: smoking, alcohol intake, leisure-time physical activity and osteoporosis (based on low or very low certainty of evidence). Most of the studies evaluated three large cohorts from the USA (OAI, MOST and Framingham). Two modifiable risk factors (overweight/obesity and previous knee injury) accounted for 14% of overall risk of incident radiographic KOA,

therefore there are opportunities to quantify the contribution of other risk factors towards incident KOA.

Strengths and limitations of this study

Our evaluation of > 150 risk factors associated with increasing or lowered risk of incident KOA, alongside strength of evidence ratings, provides important guidance for future research. We synthesised the evidence for all reported risk factors for incident KOA without date or language restrictions, making this the most comprehensive review in the field. Our methods were rigorous, in line with best practice standards for conducting systematic reviews. Few studies were conducted outside the USA, and in low- and middle-income countries, limiting the generalisability to global populations. We considered different outcomes (e.g., radiographic, symptomatic KOA) separately, which resulted in greater homogeneity across our pooled analyses and enabled more tailored clinical guidance. This is advantageous as there is heterogeneity in defining both symptomatic and/or radiographic OA¹⁵⁵ and the overlap between the two is limited.

Due to the large heterogeneity in exposures, outcomes, and measures of effect, it was not possible to conduct pooled analyses for all risk factors. Some studies reported large OR estimates with unusually wide CIs, suggesting a sparse data bias. The possibility of confounding cannot be eliminated, given that most studies were observational in nature. Furthermore, we could not assess publication bias for any risk factor as too few studies were included in the pooled analysis.

We could not determine a correlation matrix between all the risk factors found to be associated with incident radiographic KOA based on our pooled analysis, as the available data did not allow. This means the contribution of important risk factors, such as occupational physical activity, could not be accounted for in the PAF analysis. Communality values were calculated based on information about risk factors from a single cohort (OAI), which limits the generalisability of the findings. We were unable to calculate PAF for symptomatic KOA as the available data did not permit. Nevertheless, this is an important clinical question, as radiographic KOA does not consistently relate to symptoms. We excluded studies where there was evidence of MRI structural changes, which meant that earlier studies in children with longer term follow-up were not captured.^{156–158}

Finally, it was not possible to determine if the same participants were being reported on in different studies across the OAI, MOST and Framingham cohorts, therefore we cannot exclude potential duplication of participants across studies.




Fig. 7

Life-course model of potentially modifiable risk factors for osteoarthritis across the lifespan.

Nevertheless, we provide more rigorous estimates compared to previous research. For example, the 2015 review¹⁰ did not account for communality (clustering) of risk factors within an individual and reported a larger contribution of obesity and knee injury in the PAF calculations compared to our review.

Meaning of the study and implications

Our review identified moderate quality evidence that modifiable risk factors, including overweight/obesity, previous knee injury and occupational physical activity, are associated with incident KOA and

should be core targets for future preventive strategies. Previous studies have shown no effect of sex on the risk of developing incident KOA,¹⁰ whereas we found that females had a two-fold greater risk of developing incident radiographic KOA, compared to males. However, in individuals with diabetes, males were at greater risk of developing incident KOA than females. Older age was not found to be a strong risk factor for incident KOA, as most incident cases of KOA are likely to be diagnosed at middle age.¹⁵⁹

Societal interventions are urgently needed to address modifiable risk factors associated with incident KOA, including high intake of ultra-processed foods, poor sleep quality, occupational risks, knee injury and overweight/obesity (particularly in females). This includes targeted interventions to prevent weight gain and obesity and funded physical activity, education and nutrition programs during critical developmental years.^{160–162} Given that mid-life obesity is a strong predictor of developing incident KOA, and that obese children and adolescents are five times more likely to be obese in adulthood, it is imperative that these interventions are implemented at an early age.¹⁶³

Our findings reflect the physical activity paradox documented in previous research.¹⁶⁴ There is inconclusive evidence regarding the association between leisure-time physical activity and incident KOA. However, it was measured heterogeneously making it difficult to ascertain the type, dose and frequency of leisure-time physical activity which may be beneficial or adversely associated with incident KOA. Leisure-time physical activity was associated with incident radiographic or symptomatic KOA in individuals with low levels of lower limb muscle mass who engaged in weight-bearing physical activity and in overweight or obese individuals who engaged in heavy physical activity. This suggests that the type of physical activity, and/or the intrinsic capacity of the individual may be important considerations.

Conversely, occupational physical activity was significantly associated with incident radiographic or symptomatic KOA, calling for urgent efforts from workplaces and employers to reduce sedentary work, reduce physical load-bearing, improve strength and help reduce weight in the workforce.¹⁶⁵ Such efforts may have benefits beyond KOA prevention, there may be social, mental health and economic benefits.¹⁶⁴

There is also a need for multi-component injury prevention interventions and solutions for individuals involved in sporting activities from an early age. ACL injuries are the most prevalent form of knee injury and there is evidence to support injury prevention programs which involve focused strengthening, landing and pivoting techniques.¹⁶⁶

Future directions

Our data provides a platform to inform risk prediction models for the development of incident KOA. Given the emphasis on leisure-time physical activity in international clinical guidelines for OA prevention,¹⁴ there is a need to resolve the present KOA risk uncertainty in future high quality studies, both for radiographic and symptomatic KOA. However, it is clear that future mitigation strategies need to target occupational physical activity. Government and workplace interventions aimed at reducing sedentary work, heavy manual handling, particularly over prolonged periods, duration of shift work and consumption of ultra-processed foods may prove effective in curbing the risk of incident KOA across the lifespan.

Furthermore, given the focus on non-drug interventions for the prevention and management of KOA, it is worthwhile establishing the longitudinal association of modifiable risk factors such as smoking, alcohol consumption and dietary risk factors (e.g., intake of cereal, dark bread, and green tea) in large, prospective cohort studies.

Our review identified heterogeneity in the definition and classification of KOA, prompting broader discussion for a consensus-based approach on KOA criteria.

There are limited studies on the incidence of KOA in Hispanics, Africans, or Southeast Asians, warranting research in these populations. The role of epigenetic and potentially modifiable genetic factors that may contribute meaningfully to prevention strategies should be explored. A focus on, and funding of, discovery research in addition to implementing community and lifestyle programs can help to reduce the global burden of KOA.

Conclusion

Obesity and previous knee injury together accounted for 14% of the total risk of KOA, urging novel societal strategies and future trials targeting these risk factors to reduce overall disease burden. Efforts addressing occupational risk factors and future research evaluating the association between potentially important risk factors such as leisure-time physical activity and incident KOA are also needed.

Author contributions

All authors contributed to the conception and design of the study, provided critical input on the draft and approved the final version. VD and CAS carried out the search strategy. VD, CAS, JZ, IA, SLG, SK, SWN carried out data extraction and performed risk of bias assessments. Conflicts were resolved by an adjudicator MLF, DJH. VD, CAS and VV carried out data analysis. VD and CAS led the write up of the draft manuscript.

Declarations of interest

VD and CAS received funding from Sydney Musculoskeletal Health, The University of Sydney to undertake this review. CAS is supported by a NHMRC Emerging Leadership 2 Fellowship. MLF is supported by an NHMRC Investigator Grant Leadership 1 and provided consulting advice on the scientific advisory board for Novartis. DJH is supported by an NHMRC Investigator Grant Leadership 2 (#1194737). DJH provides consulting advice on scientific advisory boards for Pfizer, Lilly, TLCBio, Novartis. AMB has received grants from AO Alliance, Asia Pacific League of Associations for Rheumatology, Australian Rheumatology Association, Curtin University, Pan American League of Associations for Rheumatology, World Federation of Chiropractic, Australian Government, Department of Health Grant, Medical Research Future Fund Grant (Australian National Health and Medical Research Council), Western Australian Government Department of Health Grant, Bone and Joint Decade Foundation (Sweden), Institute for Bone and Joint Research (Australia), Canadian Memorial Chiropractic College and Arthritis and Osteoporosis Western Australia. AMB provides consulting to the World Health Organization. MC has received royalties for joint authorship of a chapter on Up To Date. FG has received a grant from Novartis. LSL provides consulting to Anthro Therapeutics AB. LM has received grants from MRFF Better Use Existing Medicines, MRFF Better Outcomes for Childhood Arthritis and NHMRC CRE Better Outcomes for Inflammatory Arthritis. AM has received grants from the Research Council of Finland, Research Council of Lithuania and the European Commission. AM provides consultation to Grünenthal, Sanofi, Viatrix, HALEON, Kolon TissueGene, Pacira, Consumer Healthcare at Sanofi, Aptissen. GP has an Academic Consultant Contract with Office for Health Improvement & Disparities (formerly Public Health England). MAR has received a National Institutes of Health (NIH) grant: R37-HD037985. SS received Postdoctoral funding from the International Association for the Study of Pain (2021–2023). FK has received support from the following NIH

grants: AG058702-04; AR077360-03; NR019196-01; AG064947. SWN, VV, JZ, IA, SK, SLG, RE, KF, GJM, YM, MN, RS, YZ and CC have nothing to declare.

Funding

Funded by Sydney Musculoskeletal Health, Faculty of Medicine and Health, The University of Sydney. The funders of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at doi:10.1016/j.joca.2025.03.003.

References

- Katz JN, Arant KR, Loeser RF. Diagnosis and treatment of hip and knee osteoarthritis: a review. *JAMA* 2021;325:568–78.
- Bitton R. The economic burden of osteoarthritis. *Am J Manag Care* 2009;15:S230–5.
- Steinmetz JD, Culbreth GT, Haile LM, Rafferty Q, Lo J, Fukutaki KG, et al. Global, regional, and national burden of osteoarthritis, 1990–2020 and projections to 2050: a systematic analysis for the Global Burden of Disease Study 2021. *Lancet Rheumatol* 2023;5:e508–22.
- Cui A, Li H, Wang D, Zhong J, Chen Y, Lu H. Global, regional prevalence, incidence and risk factors of knee osteoarthritis in population-based studies. *EClinicalMedicine* 2020;29:29–30.
- Blanco FJ, Silva-Díaz M, Vila VQ, Seoane-Mato D, Ruiz FP, Juan-Mas A, et al. Prevalence of symptomatic osteoarthritis in Spain: EPISER2016 study. *Reumatol Clín (Engl Ed)* 2021;17:461–70.
- Peat G, Rathod-Mistry T, Paskins Z, Marshall M, Thomas MJ, Menz HB, et al. Relative prevalence and distribution of knee, hand and foot symptomatic osteoarthritis subtypes in an English population. *Musculoskel Care* 2020;18:219–24.
- Ho-Pham LT, Lai TQ, Mai LD, Doan MC, Pham HN, Nguyen TV. Prevalence of radiographic osteoarthritis of the knee and its relationship to self-reported pain. *PLoS One* 2014;9, e94563.
- Murphy L, Schwartz TA, Helmick CG, Renner JB, Tudor G, Koch G, et al. Lifetime risk of symptomatic knee osteoarthritis. *Arthritis Care Res (Hoboken)* 2008;59:1207–13.
- Cross M, Smith E, Hoy D, Nolte S, Ackerman I, Fransen M, et al. The global burden of hip and knee osteoarthritis: estimates from the global burden of disease 2010 study. *Ann Rheum Dis* 2014;73:1323–30.
- Silverwood V, Blagojevic-Bucknall M, Jinks C, Jordan J, Protheroe J, Jordan K. Current evidence on risk factors for knee osteoarthritis in older adults: a systematic review and meta-analysis. *Osteoarthr Cartil* 2015;23:507–15.
- Øiestad BE, Juhl CB, Culvenor AG, Berg B, Thorlund JB. Knee extensor muscle weakness is a risk factor for the development of knee osteoarthritis: an updated systematic review and meta-analysis including 46 819 men and women. *Br J Sports Med* 2022;56:349–55.
- Tanamas S, Hanna FS, Cicuttini FM, Wluka AE, Berry P, Urquhart DM. Does knee malalignment increase the risk of development and progression of knee osteoarthritis? A systematic review. *Arthritis Care Res: Off J Am Coll Rheumatol* 2009;61:459–67.
- Chaganti RK, Lane NE. Risk factors for incident osteoarthritis of the hip and knee. *Curr Rev Musculoskel Med* 2011;4:99–104.
- McAlindon TE, Bannuru RR, Sullivan M, Arden N, Berenbaum F, Bierma-Zeinstra S, et al. OARSI guidelines for the non-surgical management of knee osteoarthritis. *Osteoarthr Cartil* 2014;22:363–88.
- Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Int J Surg* 2021;88, 105906.
- Kellgren JH, Lawrence J. Radiological assessment of osteoarthritis. *Ann Rheum Dis* 1957;16:494–502.
- Nüesch E, Trelle S, Reichenbach S, Rutjes AW, Tschannen B, Altman DG, et al. Small study effects in meta-analyses of osteoarthritis trials: meta-epidemiological study. *BMJ* 2010;341, c3515.
- Covidence systematic review software, Veritas Health Innovation, Melbourne, Australia. Available at: www.covidence.org.
- Hayden JA, van der Windt DA, Cartwright JL, Côté P, Bombardier C. Assessing bias in studies of prognostic factors. *Ann Intern Med* 2013;158:280–6.
- DerSimonian R, Laird N. Meta-analysis in clinical trials revisited. *Contemp Clin Trials* 2015;45:139–45.
- Huguet A, Hayden JA, Stinson J, McGrath PJ, Chambers CT, Tougas ME, et al. Judging the quality of evidence in reviews of prognostic factor research: adapting the GRADE framework. *Syst Rev* 2013;2:71.
- Begg CB, Berlin JA. Publication bias: a problem in interpreting medical data. *J R Stat Soc Ser A: Stat Soc* 1988;151:419–45.
- Eckstein F, Wirth W, Nevitt MC. Recent advances in osteoarthritis imaging—the osteoarthritis initiative. *Nat Rev Rheumatol* 2012;8:622–30.
- Runhaar J, van Middelkoop M, Reijman M, Willemssen S, Oei EH, Vroegindewij D, et al. Prevention of knee osteoarthritis in overweight females: the first preventive randomized controlled trial in osteoarthritis. *Am J Med* 2015;128:888–895.e884.
- de Vos BC, Landsmeer MLA, van Middelkoop M, Oei EHG, Krul M, Bierma-Zeinstra SMA, et al. Long-term effects of a lifestyle intervention and oral glucosamine sulphate in primary care on incident knee OA in overweight women. *Rheumatology* 2017;56:1326–34.
- Yoshimura N, Muraki S, Oka H, Kawaguchi H, Nakamura K, Akune T. Accumulation of metabolic risk factors such as overweight, hypertension, dyslipidemia, and impaired glucose tolerance raises the risk of occurrence and progression of knee osteoarthritis: A 3-year follow-up of the road study. *Osteoporos Int* 2012;22:S121.
- Bergink AP, Rivadeneira F, Bierma-Zeinstra SM, Zillikens MC, Ikram MA, Uitterlinden AG, et al. Are bone mineral density and fractures related to the incidence and progression of radiographic osteoarthritis of the knee, hip, and hand in elderly men and women? The Rotterdam Study. *Arthritis Rheumatol* 2019;71(3):361–9.
- McAlindon T, Zhang Y, Hannan M, Naimark A, Weissman B, Castelli W, et al. Are risk factors for patellofemoral and tibiofemoral knee osteoarthritis different? *J Rheumatol* 1996;23:332–7.
- Reyes C, Leyland KM, Peat G, Cooper C, Arden NK, Prieto-Alhambra D. Association between overweight and obesity and risk of clinically diagnosed knee, hip, and hand osteoarthritis: A population-based cohort study. *Arthritis Rheumatol* 2016;68:1869–75.
- Brouwer GM, van Tol AW, Bergink AP, Belo JN, Bernsen RM, Reijman M, et al. Association between valgus and varus alignment and the development and progression of radiographic osteoarthritis of the knee. *Arthritis Rheum* 2007;56:1204–11.

31. Konstari S, Paananen M, Heliovaara M, Knekt P, Marniemi J, Impivaara O, *et al.* Association of 25-hydroxyvitamin D with the incidence of knee and hip osteoarthritis: a 22-year follow-up study. *Scand J Rheumatol* 2012;41:124–31.
32. Guillen R, Otero F, Mosquera A, Vazquez-Mosquera M, Rego-Perez I, Blanco FJ, *et al.* Association of accelerated dynamics of telomere sequence loss in peripheral blood leukocytes with incident knee osteoarthritis in Osteoarthritis Initiative cohort. *Sci Rep* 2021;11, 15914.
33. Lin SH, Wang TC, Lai CF, Tsai RY, Yang CP, Wong CS. Association of anterior cruciate ligament injury with knee osteoarthritis and total knee replacement: A retrospective cohort study from the Taiwan National Health Insurance Database. *PLoS ONE* 2017;12, e0178292.
34. Koo BS, Song Y, Lee S, Sung YK, Shin KJ, Cho NH, *et al.* Association of Asian mitochondrial DNA haplogroup B with new development of knee osteoarthritis in Koreans. *Int J Rheum Dis* 2019;22:411–6.
35. Salis Z, Gallego B, Nguyen TV, Sainsbury A. Association of decrease in body mass index with reduced incidence and progression of the structural defects of knee osteoarthritis: A prospective multi-cohort study. *Arthritis Rheumatol* 2022;16:16.
36. Rogers-Soeder TS, Lane NE, Walimbe M, Schwartz AV, Tolstykh I, Felson DT, *et al.* Association of diabetes mellitus and biomarkers of abnormal glucose metabolism with incident radiographic knee osteoarthritis. *Arthritis Care Res* 2020;72:98–106.
37. Zheng J, Huang X, Huang J, Meng B, Li F, Liu H, *et al.* Association of diabetes mellitus status and hyperglycemia with symptomatic knee osteoarthritis. *Arthritis Care Res* 2023;75:509–18.
38. Harvey WF, Yang M, Cooke TDV, Segal NA, Lane N, Lewis CE, *et al.* Association of leg-length inequality with knee osteoarthritis: a cohort study. *Ann Intern Med* 2010;152(5):287–95.
39. Karvonen-Gutierrez CA, Harlow SD, Mancuso P, Jacobson J, De Leon CFM, Nan B. Association of leptin levels with radiographic knee osteoarthritis among a cohort of midlife women. *Arthritis Care Res* 2013;65(6):936–44.
40. Chang AH, Lee J, Chmiel JS, Almagor O, Song J, Sharma L. Association of Long-term Strenuous Physical Activity and Extensive Sitting With Incident Radiographic Knee Osteoarthritis. *JAMA Netw Open* 2020;3, e204049.
41. Purser JL, Golightly YM, Feng Q, Helmick CG, Renner JB, Jordan JM. Association of slower walking speed with incident knee osteoarthritis-related outcomes. *Arthritis Care Res* 2012;64:1028–35.
42. Konstari S, Saaksjarvi K, Heliovaara M, Rissanen H, Knekt P, Arokoski JPA, *et al.* Associations of metabolic syndrome and its components with the risk of incident knee osteoarthritis leading to hospitalization: a 32-year follow-up study. *Cartilage* 2021;13:1445S–56S.
43. Lu H, Wang L, Zhou W, Jin S, Chen H, Su Y, *et al.* Bidirectional association between knee osteoarthritis and depressive symptoms: evidence from a nationwide population-based cohort. *BMC Musculoskelet Disord* 2022;23:213.
44. Golightly YM, Marshall SW, Kraus VB, Renner JB, Villaveces A, Casteel C, *et al.* Biomarkers of incident radiographic knee osteoarthritis: do they vary by chronic knee symptoms? *Arthritis Rheum* 2011;63:2276–83.
45. Reijman M, Pols HA, Bergink AP, Hazes JM, Belo JN, Lievense AM, *et al.* Body mass index associated with onset and progression of osteoarthritis of the knee but not of the hip: the Rotterdam Study. *Ann Rheum Dis* 2007;66:158–62.
46. Gelber AC, Hochberg MC, Mead LA, Wang NY, Wigley FM, Klag MJ. Body mass index in young men and the risk of subsequent knee and hip osteoarthritis. *Am J Med* 1999;107:542–8.
47. Dai Z, Reza Jafarzadeh S, Niu J, Felson DT, Jacques PF, Li S, *et al.* Body mass index mediates the association between dietary fiber and symptomatic knee osteoarthritis in the osteoarthritis initiative and the Framingham Osteoarthritis Study. *J Nutr* 2018;148(12):1961–7.
48. Hochberg MC, Lethbridge-Cejku M, Tobin JD. Bone mineral density and osteoarthritis: data from the Baltimore Longitudinal Study of Aging. *Osteoarthritis Cartil* 2004;12 A:S45–8.
49. Zhang Y, Hannan MT, Chaisson CE, McAlindon TE, Evans SR, Aliabadi P, *et al.* Bone mineral density and risk of incident and progressive radiographic knee osteoarthritis in women: the Framingham Study. *J Rheumatol* 2000;27:1032–7.
50. Bergink AP, Uitterlinden AG, Van Leeuwen JP, Hofman A, Verhaar JA, Pols HA. Bone mineral density and vertebral fracture history are associated with incident and progressive radiographic knee osteoarthritis in elderly men and women: the Rotterdam Study. *Bone* 2005;37:446–56.
51. Kumar NM, Hafezi-Nejad N, Guermazi A, Haj-Mirzaian A, Haugen IK, Roemer FW, *et al.* Association of Quantitative and Topographic Assessment of Heberden's Nodes With Knee Osteoarthritis: Data from the osteoarthritis initiative. *Arthritis Rheumatol* 2018;70(8):1234–9.
52. Engstrom G, Gerhardsson de Verdier M, Roloff J, Nilsson PM, Lohmander LS. C-reactive protein, metabolic syndrome and incidence of severe hip and knee osteoarthritis. A population-based cohort study. *Osteoarthritis Cartil* 2009;17:168–73.
53. Fan J, Zhu J, Sun L, Wang T, Li Y. Causal association of adipokines with osteoarthritis: A Mendelian randomization study. *Rheumatology (United Kingdom)* 2021;60(6):2808–15.
54. Meng H, Jiang L, Song Z, Wang F. Causal associations of circulating lipids with osteoarthritis: A bidirectional Mendelian randomization study. *Nutrients* 2022;14(7):1327.
55. Lo GH, Strayhorn MT, Driban JB, Price LL, Eaton CB, McAlindon TE. Subjective crepitus as a risk factor for incident symptomatic knee osteoarthritis: Data from the osteoarthritis initiative. *Arthritis Care Res* 2018;70:53–60.
56. Nordenvall R, Bahmanyar S, Adami J, Mattila VM, Fellander-Tsai L. Cruciate ligament reconstruction and risk of knee osteoarthritis: the association between cruciate ligament injury and post-traumatic osteoarthritis. a population based nationwide study in Sweden, 1987–2009. *PLoS ONE* 2014;9, e104681.
57. Dai Z, Niu J, Zhang Y, Jacques P, Felson DT. Dietary intake of fibre and risk of knee osteoarthritis in two US prospective cohorts. *Ann Rheum Dis* 2017;76(8):1411–9.
58. Konstari S, Sares-Jaske L, Heliovaara M, Rissanen H, Knekt P, Arokoski J, *et al.* Dietary magnesium intake, serum high sensitivity C-reactive protein and the risk of incident knee osteoarthritis leading to hospitalization-A cohort study of 4,953 Finns. *PLoS ONE* 2019;14, e0214064.
59. McAlindon TE, Jacques P, Zhang Y, Hannan MT, Aliabadi P, Weissman B, *et al.* Do antioxidant micronutrients protect against the development and progression of knee osteoarthritis? *Arthritis Rheum* 1996;39:648–56.
60. Johnson VL, Roe JP, Salmon LJ, Pinczewski LA, Hunter DJ. Does age influence the risk of incident knee osteoarthritis after a traumatic anterior cruciate ligament injury? *Am J Sports Med* 2016;44:2399–405.
61. Dahaghin S, Bierma-Zeinstra SMA, Reijman M, Pols HAP, Hazes JMW, Koes BW. Does hand osteoarthritis predict future hip or knee osteoarthritis? *Arthritis Rheum* 2005;52(11):3520–7.
62. Yoshimura N, Muraki S, Oka H, Kawaguchi H, Nakamura K, Tanaka S, *et al.* Does mild cognitive impairment affect the occurrence of radiographic knee osteoarthritis? A 3-year follow-up in the ROAD study. *BMJ Open* 2012;2, e001520.

63. Mork PJ, Holtermann A, Nilsen TI. Effect of body mass index and physical exercise on risk of knee and hip osteoarthritis: longitudinal data from the Norwegian HUNT Study. *J Epidemiol Commun Health* 2012;66:678–83.
64. Thorlund JB, Felson DT, Segal NA, Nevitt MC, Niu J, Neogi T, et al. Effect of knee extensor strength on incident radiographic and symptomatic knee osteoarthritis in individuals with meniscal pathology: Data from the multicenter osteoarthritis study. *Arthritis Care Res* 2016;68:1640–6.
65. Felson DT, Niu J, Clancy M, Sack B, Aliabadi P, Zhang Y. Effect of recreational physical activities on the development of knee osteoarthritis in older adults of different weights: the Framingham Study. *Arthritis Rheum* 2007;57:6–12.
66. Segal NA, Torner JC, Felson D, Niu J, Sharma L, Lewis CE, et al. Effect of thigh strength on incident radiographic and symptomatic knee osteoarthritis in a longitudinal cohort. *Arthritis Rheum* 2009;61:1210–7.
67. Sowers MR, McConnell D, Jannausch M, Buyuktur AG, Hochberg M, Jamadar DA. Estradiol and its metabolites and their association with knee osteoarthritis. *Arthritis Rheum* 2006;54:2481–7.
68. Zhang Y, McAlindon TE, Hannan MT, Chaisson CE, Klein R, Wilson PW, et al. Estrogen replacement therapy and worsening of radiographic knee osteoarthritis: the Framingham Study. *Arthritis Rheum* 1998;41:1867–73.
69. Liu T, Xu C, Driban JB, McAlindon T, Eaton CB, Lu B. Excessive alcohol consumption and the risk of knee osteoarthritis: a prospective study from the Osteoarthritis Initiative. *Osteoarthritis Cartil* 2022;30:697–701.
70. Runhaar J, Sanchez C, Taralla S, Henrotin Y, Bierma-Zeinstra SM. Fibulin-3 fragments are prognostic biomarkers of osteoarthritis incidence in overweight and obese women. *Osteoarthritis Cartil* 2016;24:672–8.
71. Campbell TM, McGonagle D. Flexion contracture is a risk factor for knee osteoarthritis incidence, progression and earlier arthroplasty: Data from the Osteoarthritis Initiative. *Ann Phys Rehabil Med* 2021;64, 101439.
72. Perry TA, Segal NA, Bowen C, Gates L, Arden N, Nevitt MC. Foot and ankle pain and risk of incident knee osteoarthritis and knee pain: Data from the Multicentre Osteoarthritis Study. *Osteoarthritis Cartil* 2021;3(4), 100210.
73. Lo GH, McAlindon TE, Kriska AM, Price LL, Rockette-Wagner BJ, Mandl LA, et al. Football increases future risk of symptomatic radiographic knee osteoarthritis. *Med Sci Sports Exerc* 2020;52:795–800.
74. Hannan MT, Felson DT, Anderson JJ, Naimark A. Habitual physical activity is not associated with knee osteoarthritis: the Framingham Study. *J Rheumatol* 1993;20:704–9.
75. Nevitt MC, Zhang Y, Javaid MK, Neogi T, Curtis JR, Niu J, et al. High systemic bone mineral density increases the risk of incident knee OA and joint space narrowing, but not radiographic progression of existing knee OA: The MOST study. *Ann Rheum Dis* 2010;69(1):163–8.
76. Zhang L, Lin C, Liu Q, Gao J, Hou Y, Lin J. Incidence and related risk factors of radiographic knee osteoarthritis: a population-based longitudinal study in China. *J Orthop Surg* 2021;16:474.
77. Prieto-Alhambra D, Judge A, Javaid MK, Cooper C, Diez-Perez A, Arden NK. Incidence and risk factors for clinically diagnosed knee, hip and hand osteoarthritis: influences of age, gender and osteoarthritis affecting other joints. *Ann Rheum Dis* 2014;73:1659–64.
78. Muraki S, Akune T, Oka H, Ishimoto Y, Nagata K, Yoshida M, et al. Incidence and risk factors for radiographic knee osteoarthritis and knee pain in Japanese men and women: a longitudinal population-based cohort study. *Arthritis Rheum* 2012;64:1447–56.
79. Ren Y, Hu J, Tan J, Tang X, Li Q, Yang H, et al. Incidence and risk factors of symptomatic knee osteoarthritis among the Chinese population: analysis from a nationwide longitudinal study. *BMC Public Health* 2020;20:1491.
80. Lo GH, Richard MJ, McAlindon TE, Park C, Strayhorn MT, Harkey MS, et al. Increased risk of incident knee osteoarthritis in those with greater work-related physical activity. *Occup Environ Med* 2022;79:543–9.
81. Livshits G, Zhai G, Hart DJ, Kato BS, Wang H, Williams FM, et al. Interleukin-6 is a significant predictor of radiographic knee osteoarthritis: The Chingford Study. *Arthritis Rheum* 2009;60:2037–45.
82. Niu J, Zhang YQ, Torner J, Nevitt M, Lewis CE, Aliabadi P, et al. Is obesity a risk factor for progressive radiographic knee osteoarthritis? *Arthritis Rheum* 2009;61:329–35.
83. Turkiewicz A, Timpka S, Thorlund JB, Ageberg E, Englund M. Knee extensor strength and body weight in adolescent men and the risk of knee osteoarthritis by middle age. *Ann Rheum Dis* 2017;76:1657–61.
84. Smith MV, Nepple JJ, Wright RW, Matava MJ, Brophy RH. Knee osteoarthritis is associated with previous meniscus and anterior cruciate ligament surgery among elite college American football athletes. *Sports Health* 2017;9:247–51.
85. Wills AK, Black S, Cooper R, Coppack RJ, Hardy R, Martin KR, et al. Life course body mass index and risk of knee osteoarthritis at the age of 53 years: Evidence from the 1946 British birth cohort study. *Ann Rheum Dis* 2012;71:655–60.
86. Staines KA, Hardy R, Samvelyan HJ, Ward KA, Cooper R. Life course longitudinal growth and risk of knee osteoarthritis at age 53 years: evidence from the 1946 British birth cohort study. *Osteoarthritis Cartil* 2021;29:335–40.
87. Veronese N, Stefanac S, Koyanagi A, Al-Daghri NM, Sabico S, Cooper C, et al. Lower limb muscle strength and muscle mass are associated with incident symptomatic knee osteoarthritis: A longitudinal cohort study. *Front Endocrinol* 2021;12, 804560.
88. Veronese N, Cooper C, Guglielmi G, Rizzoli R, Punzi L, Al-Daghri N, et al. Mediterranean diet and knee osteoarthritis outcomes: A longitudinal cohort study. *Clin Nutr* 2019;38:2735–9.
89. Barbour KE, Hootman JM, Helmick CG, Murphy LB, Theis KA, Schwartz TA, et al. Meeting physical activity guidelines and the risk of incident knee osteoarthritis: a population-based prospective cohort study. *Arthritis Care Res* 2014;66:139–46.
90. Kuusalo L, Felson DT, Wang N, Lewis CE, Torner J, Nevitt MC, et al. Metabolic osteoarthritis – relation of diabetes and cardiovascular disease with knee osteoarthritis. *Osteoarthritis Cartil* 2021;29(2):230–4.
91. Niu J, Clancy M, Aliabadi P, Vasan R, Felson DT. Metabolic syndrome, its components, and knee osteoarthritis: The Framingham Osteoarthritis Study. *Arthritis Rheumatol* 2017;69:1194–203.
92. Fernandez-Moreno M, Soto-Hermida A, Vazquez-Mosquera ME, Cortes-Pereira E, Relano S, Hermida-Gomez T, et al. Mitochondrial DNA haplogroups influence the risk of incident knee osteoarthritis in OAI and CHECK cohorts. A meta-analysis and functional study. *Ann Rheum Dis* 2017;76:1114–22.
93. Takiguchi R, Komatsu R, Kitamura K, Watanabe Y, Takahashi A, Kobayashi R, et al. Modifiable factors associated with symptomatic knee osteoarthritis: The Murakami cohort study. *Maturitas* 2019;128:53–9.
94. Felson DT, Anderson JJ, Naimark A, Walker AM, Meenan RF. Obesity and knee osteoarthritis. The Framingham Study. *Ann Intern Med* 1988;109:18–24.
95. Grotle M, Hagen KB, Natvig B, Dahl FA, Kvien TK. Obesity and osteoarthritis in knee, hip and/or hand: an epidemiological

- study in the general population with 10 years follow-up. *BMC Musculoskelet Disord* 2008;9:132.
96. Shiozaki H, Koga Y, Omori G, Tamaki M. Obesity and osteoarthritis of the knee in women: Results from the Matsudai Knee Osteoarthritis Survey. *Knee* 1999;6(3):189–92.
 97. Hart HF, van Middelkoop M, Stefanik JJ, Crossley KM, Bierma-Zeinstra S. Obesity is related to incidence of patellofemoral osteoarthritis: the Cohort Hip and Cohort Knee (CHECK) study. *Rheumatol Int* 2020;40:227–32.
 98. Toivanen AT, Heliovaara M, Impivaara O, Arokoski JPA, Knekt P, Lauren H, et al. Obesity, physically demanding work and traumatic knee injury are major risk factors for knee osteoarthritis—a population-based study with a follow-up of 22 years. *Rheumatology* 2010;49(2):308–14.
 99. Jin QIN, Barbour KE, Nevitt MC, Helmick CG, Hootman JM, Murphy LB, et al. Objectively measured physical activity and risk of knee osteoarthritis. *Med Sci Sports Exerc* 2018;50:277–83.
 100. Felson DT, Hannan MT, Naimark A, Berkeley J, Gordon G, Wilson PW, et al. Occupational physical demands, knee bending, and knee osteoarthritis: results from the Framingham Study. *J Rheumatol* 1991;18:1587–92.
 101. Kopec JA, Sayre EC, Schwartz TA, Renner JB, Helmick CG, Badley EM, et al. Occurrence of radiographic osteoarthritis of the knee and hip among African Americans and whites: a population-based prospective cohort study. *Arthritis Care Res* 2013;65:928–35.
 102. Paradowski PT, Lohmander LS, Englund M. Osteoarthritis of the knee after meniscal resection: long term radiographic evaluation of disease progression. *Osteoarthr Cartil* 2016;24:794–800.
 103. Driban JB, Eaton CB, Lo GH, Price LL, Lu B, Barbe MF, et al. Overweight older adults, particularly after an injury, are at high risk for accelerated knee osteoarthritis: data from the Osteoarthritis Initiative. *Clin Rheumatol* 2016;35:1071–6.
 104. Manninen P, Riihimäki H, Heliovaara M, Makela P. Overweight, gender and knee osteoarthritis. *Int J Obesity Related Metab Disord: J Int Assoc Study Obesity* 1996;20:595–7.
 105. Verweij LM, Van Schoor NM, Deeg DJH, Dekker J, Visser M. Physical activity and incident clinical knee osteoarthritis in older adults. *Arthritis Care Res* 2009;61(2):152–7.
 106. Felson DT, Niu J, Yang T, Torner J, Lewis CE, Aliabadi P, et al. Physical activity, alignment and knee osteoarthritis: data from MOST and the OAI. *Osteoarthr Cartil* 2013;21:789–95.
 107. Takagi S, Omori G, Koga H, Endo K, Koga Y, Nawata A, et al. Quadriceps muscle weakness is related to increased risk of radiographic knee OA but not its progression in both women and men: the Matsudai Knee Osteoarthritis Survey. *Knee Surg Sports Traumatol Arthrosc* 2018;26:2607–14.
 108. Shelbourne KD, Benner RW, Gray T. Results of anterior cruciate ligament reconstruction with patellar tendon autografts: objective factors associated with the development of osteoarthritis at 20 to 33 years after surgery. *Am J Sports Med* 2017;45:2730–8.
 109. Wei J, Gross D, Lane NE, Lu N, Wang M, Zeng C, et al. Risk factor heterogeneity for medial and lateral compartment knee osteoarthritis: analysis of two prospective cohorts. *Osteoarthr Cartil* 2019;27:603–10.
 110. Felson DT, Zhang Y, Hannan MT, Naimark A, Weissman B, Aliabadi P, et al. Risk factors for incident radiographic knee osteoarthritis in the elderly: the Framingham Study. *Arthritis Rheum* 1997;40:728–33.
 111. Cooper C, Snow S, McAlindon TE, Kellingray S, Stuart B, Coggon D, et al. Risk factors for the incidence and progression of radiographic knee osteoarthritis. *Arthritis Rheum* 2000;43:995–1000.
 112. Nishimura A, Hasegawa M, Kato K, Yamada T, Uchida A, Sudo A. Risk factors for the incidence and progression of radiographic osteoarthritis of the knee among Japanese. *Int Orthop* 2011;35:839–43.
 113. Snoeker B, Turkiewicz A, Magnusson K, Frobell R, Yu D, Peat G, et al. Risk of knee osteoarthritis after different types of knee injuries in young adults: a population-based cohort study. *Br J Sports Med* 2020;54:725–30.
 114. Misra D, Felson DT, Niu J, Brown C, Neogi T, Fielding RA, et al. Risk of knee osteoarthritis with obesity, sarcopenic obesity, and sarcopenia. *Arthritis Rheumatol* 2019;71:232–7.
 115. Rhon DI, Perez KG, Eskridge SL. Risk of post-traumatic knee osteoarthritis after knee injury in military service members. *Musculoskelet Care* 2019;17:113–9.
 116. Konstari S, Kaila-Kangas L, Jaaskelainen T, Heliovaara M, Rissanen H, Marniemi J, et al. Serum 25-hydroxyvitamin D and the risk of knee and hip osteoarthritis leading to hospitalization: a cohort study of 5274 Finns. *Rheumatology* 2014;53:1778–82.
 117. Kluzek S, Bay-Jensen AC, Judge A, Karsdal MA, Shorthose M, Spector T, et al. Serum cartilage oligomeric matrix protein and development of radiographic and painful knee osteoarthritis. A community-based cohort of middle-aged women. *Biomarkers* 2015;20:557–64.
 118. Nelson AE, Golightly YM, Kraus VB, Stabler T, Renner JB, Helmick CG, et al. Serum transforming growth factor-beta 1 is not a robust biomarker of incident and progressive radiographic osteoarthritis at the hip and knee: the Johnston County Osteoarthritis Project. *Osteoarthr Cartil* 2010;18:825–9.
 119. Zhou M, Yang S, Guo Y, Wang D, Qiu W, Wang B, et al. Shift work and the risk of knee osteoarthritis among Chinese workers: a retrospective cohort study. *Scand J Work Environ Health* 2020;1:152–60.
 120. Wilder FV, Hall BJ, Barrett JP. Smoking and osteoarthritis: is there an association? The Clearwater Osteoarthritis Study. *Osteoarthr Cartil* 2003;11:29–35.
 121. Veronese N, Koyanagi A, Stubbs B, Cooper C, Guglielmi G, Rizzoli R, et al. Statin use and knee osteoarthritis outcomes: a longitudinal cohort study. *Arthritis Care Res* 2019;71:1052–8.
 122. Lo GH, Strayhorn MT, Driban JB, Price LL, Eaton C, McAlindon TE. Crepitus as a risk factor for symptomatic knee osteoarthritis: Data from the osteoarthritis initiative. *Arthritis Rheumatol* 2016;68(ement 10):390–2.
 123. Lo GH, McAlindon TE, Katz JN, Driban JB, Price LL, Eaton CB, et al. Systolic and pulse pressure associate with incident knee osteoarthritis: data from the Osteoarthritis Initiative. *Clin Rheumatol* 2017;36:2121–8.
 124. Van Spij WE, Welsing PM, Bierma-Zeinstra SM, Bijlsma JW, Roorda LD, Cats HA, et al. The ability of systemic biochemical markers to reflect presence, incidence, and progression of early-stage radiographic knee and hip osteoarthritis: data from CHECK. *Osteoarthr Cartil* 2015;23:1388–97.
 125. Wise BL, Niu J, Zhang Y, Felson DT, Bradley LA, Segal N, et al. The association of parity with osteoarthritis and knee replacement in the multicenter osteoarthritis study. *Osteoarthr Cartil* 2013;21:1849–54.
 126. Takemoto G, Seki T, Takegami Y, Osawa Y, Makida K, Ochiai S, et al. The development of knee osteoarthritis and serum carotenoid levels among community-dwelling people in Japan. *Modern Rheumatol* 2022;32:205–12.
 127. Soutakbar H, Lamb SE, Silman AJ. The different influence of high levels of physical activity on the incidence of knee OA in overweight and obese men and women—a gender specific analysis. *Osteoarthr Cartil* 2019;27:1430–6.

128. Segal NA, Glass N, Felson DT, Hurley M, Yang M, Nevitt M, *et al.* The effect of quadriceps strength and proprioception on risk for knee osteoarthritis. *J Investig Med* 2010;58(4):680.
129. Carlesso LC, Niu J, Segal NA, Frey-Law LA, Lewis CE, Nevitt MC, *et al.* The effect of widespread pain on knee pain worsening, incident knee osteoarthritis (OA), and incident knee pain: The Multicenter OA (MOST) Study. *J Rheumatol* 2017;44:493–8.
130. Peregoy J, Wilder FV, Peregoy J, Wilder FV. The effects of vitamin C supplementation on incident and progressive knee osteoarthritis: a longitudinal study. *Public Health Nutr* 2011;14:709–15.
131. Felson DT, Zhang Y, Hannan MT, Naimark A, Weissman BN, Aliabadi P, *et al.* The incidence and natural history of knee osteoarthritis in the elderly. The Framingham Osteoarthritis Study. *Arthritis Rheum* 1995;38:1500–5.
132. Segal NA, Findlay C, Wang K, Torner JC, Nevitt MC. The longitudinal relationship between thigh muscle mass and the development of knee osteoarthritis. *Osteoarthr Cartil* 2012;20:1534–40.
133. Kim I, Kim HA, Seo YI, Song YW, Jeong JY, Kim DH. The prevalence of knee osteoarthritis in elderly community residents in Korea. *J Korean Med Sci* 2010;25:293–8.
134. Paterson KL, Kasza J, Hunter DJ, Hinman RS, Menz HB, Peat G, *et al.* The relationship between foot and ankle symptoms and risk of developing knee osteoarthritis: data from the osteoarthritis initiative. *Osteoarthr Cartil* 2017;25:639–46.
135. Sharma L, Song J, Dunlop D, Felson D, Lewis CE, Segal N, *et al.* Varus and valgus alignment and incident and progressive knee osteoarthritis. *Ann Rheum Dis* 2010;69:1940–5.
136. Sharma L, Chang AH, Jackson RD, Nevitt M, Moio KC, Hochberg M, *et al.* Varus thrust and incident and progressive knee osteoarthritis. *Arthritis Rheumatol* 2017;69:2136–43.
137. Li M, Zeng Y, Nie Y, Wu Y, Liu Y, Wu L, *et al.* Varus-valgus knee laxity is related to a higher risk of knee osteoarthritis incidence and structural progression: data from the osteoarthritis initiative. *Clin Rheumatol* 2022;41:1013–21.
138. Uitterlinden AG, Burger H, Huang Q, Odding E, Duijn CM, Hofman A, *et al.* Vitamin D receptor genotype is associated with radiographic osteoarthritis at the knee. *J Clin Investig* 1997;100:259–63.
139. Bergink AP, Uitterlinden AG, Van Leeuwen JPT, Buurman CJ, Hofman A, Verhaar JAN, *et al.* Vitamin D status, bone mineral density, and the development of radiographic osteoarthritis of the knee: The Rotterdam Study. *JRC: J Clin Rheumatol* 2009;15:230–7.
140. Wilder FV, Leaverton PE, Rogers MW, Lemrow NB. Vitamin supplements and radiographic knee osteoarthritis: The Clearwater Osteoarthritis Study. *J Musculoskelet Res* 2009;12(2):85–93.
141. Felson DT, Zhang Y, Anthony JM, Naimark A, Anderson JJ. Weight loss reduces the risk for symptomatic knee osteoarthritis in women. The Framingham Study. *Ann Internal Med* 1992;116:535–9.
142. Liu T, Xu C, Driban JB, Liang GY, Zhang XH, Hu FB, *et al.* Whole grain consumption and risk of radiographic knee osteoarthritis: a prospective study from the Osteoarthritis Initiative. *Rheumatology* 2022;21:21.
143. Nelson AE, Golightly YM, Renner JB, Schwartz TA, Liu F, Lynch JA, *et al.* Variations in hip shape are associated with radiographic knee osteoarthritis: Cross-sectional and longitudinal analyses of the Johnston county osteoarthritis project. *J Rheumatol* 2016;43(2):405–10.
144. Wilder F, Hall B, Barrett Jr J, Lemrow N. History of acute knee injury and osteoarthritis of the knee: a prospective epidemiological assessment: The Clearwater Osteoarthritis Study. *Osteoarthr Cartil* 2002;10:611–6.
145. Oliveria SA, Felson DT, Reed JI, Cirillo PA, Walker AM. Incidence of symptomatic hand, hip, and knee osteoarthritis among patients in a health maintenance organization. *Arthritis Rheum: Off J Am College Rheumatol* 1995;38:1134–41.
146. Perry TA, Wang X, Gates L, Parsons CM, Sanchez-Santos MT, Garriga C, *et al.* Occupation and risk of knee osteoarthritis and knee replacement: a longitudinal, multiple-cohort study. *Semin Arthritis Rheum* 2020;50:1006–14.
147. Jørgensen K, Pedersen B, Nielsen N, Hansen A, Jacobsen S, Frisch M. Socio-demographic factors, reproductive history and risk of osteoarthritis in a cohort of 4.6 million Danish women and men. *Osteoarthr Cartil* 2011;19:1176–82.
148. McAlindon TE, Wilson PW, Aliabadi P, Weissman B, Felson DT. Level of physical activity and the risk of radiographic and symptomatic knee osteoarthritis in the elderly: the Framingham study. *Am J Med* 1999;106:151–7.
149. Landsmeer ML, Runhaar J, Henrotin YE, Middelkoop van M, Oei EH, Vroegindewij D, *et al.* Association of urinary biomarker COLL2-1NO2 with incident clinical and radiographic knee OA in overweight and obese women. *Osteoarthr Cartil* 2015;23:1398–404.
150. Runhaar J, van Middelkoop M, Reijman M, Vroegindewij D, Oei EH, Bierma-Zeinstra SM. Malalignment: a possible target for prevention of incident knee osteoarthritis in overweight and obese women. *Rheumatology* 2014;53:1618–24.
151. Lee DH, Lee HS, Jang SH, Heu J-Y, Han K, Lee S-W. Decreased risk of knee osteoarthritis with taller height in an east asian population: A nationwide cohort study. *J Clin Med* 2023;13:92.
152. Wei Y, Zhang T, Liu Y, Liu H, Zhou Y, Su J, *et al.* Ultra-processed food consumption, genetic susceptibility, and the risk of hip/knee osteoarthritis. *Clin Nutr* 2024;43:1363–71.
153. Wu Y, Boer CG, Hofman A, Schiphof D, van Middelkoop M, Szilagyi IA, *et al.* Weight-bearing physical activity, lower-limb muscle mass, and risk of knee osteoarthritis. *JAMA Netw Open* 2024;7:e248968.
154. Zhou S, Wu L, Si H, Li M, Liu Y, Shen B. Association between nighttime sleep duration and quality with knee osteoarthritis in middle-aged and older Chinese: A longitudinal cohort study. *Arch Gerontol Geriatr* 2024;118, 105284.
155. Mahmoudian A, Lohmander LS, Mobasheri A, Englund M, Luyten FP. Early-stage symptomatic osteoarthritis of the knee—time for action. *Nat Rev Rheumatol* 2021;17:621–632.
156. Antony B, Jones G, Venn A, Cicuttini F, March L, Blizzard L, *et al.* Association between childhood overweight measures and adulthood knee pain, stiffness and dysfunction: a 25-year cohort study. *Ann Rheum Dis* 2015;74:711–7.
157. Meng T, Thayer S, Venn A, Wu F, Cicuttini F, March L, *et al.* Association of childhood adiposity measures with adulthood knee cartilage defects and bone marrow lesions: a 25-year cohort study. *Osteoarthr Cartil* 2018;26:1055–62.
158. Meng T, Venn A, Eckstein F, Wirth W, Cicuttini F, March L, *et al.* Association of adiposity measures in childhood and adulthood with knee cartilage thickness, volume and bone area in young adults. *Int J Obes* 2019;43:1411–21.
159. Briggs AM, Cross MJ, Hoy DG, Sánchez-Riera L, Blyth FM, Woolf AD, *et al.* Musculoskeletal Health Conditions Represent a Global Threat to Healthy Aging: A Report for the 2015 World Health Organization World Report on Ageing and Health. *Gerontologist* 2016;56(2):S243–55.
160. Hall KD, Ayuketah A, Brychta R, Cai H, Cassimatis T, Chen KY, *et al.* Ultra-processed diets cause excess calorie intake and

- weight gain: an inpatient randomized controlled trial of ad libitum food intake. *Cell Metab* 2019;30:67–77.e63.
161. Bülbül S. Exercise in the treatment of childhood obesity. *Türk Arch Pediatr/Türk Pediatri Arşivi* 2020;55:2.
 162. Barnes C, Hall A, Nathan N, Sutherland R, McCarthy N, Pettet M, et al. Efficacy of a school-based physical activity and nutrition intervention on child weight status: Findings from a cluster randomized controlled trial. *Prevent Med* 2021;153, 106822.
 163. Simmonds M, Llewellyn A, Owen CG, Woolacott N. Predicting adult obesity from childhood obesity: a systematic review and meta-analysis. *Obes Rev* 2016;17:95–107.
 164. Pronk N. Physical activity paradox: providing evidence-based guidance while closing research gaps. *Br J Sports Med* 2024;58:932–3.
 165. Sundstrup E, Seeberg KGV, Bengtsen E, Andersen LL. A systematic review of workplace interventions to rehabilitate musculoskeletal disorders among employees with physical demanding work. *J Occup Rehabil* 2020;30:588–612.
 166. Stephenson SD, Kocan JW, Vinod AV, Kluczynski MA, Bisson LJ. A comprehensive summary of systematic reviews on sports injury prevention strategies. *Orthop J Sports Med* 2021;9, 23259671211035776.