

Towards an understanding of neural responses to emotion in recent onset and chronic psoriasis: a feasibility brain imaging study

LADA, Georgia <<http://orcid.org/0000-0003-1926-2599>>, MCKIE, Shane <<http://orcid.org/0000-0002-4418-8944>>, KAFKAS, Alex <<http://orcid.org/0000-0001-5133-8827>>, MULLINGS, Emma <<http://orcid.org/0000-0001-5718-5112>>, GRIFFITHS, Christopher E.M. <<http://orcid.org/0000-0001-5371-4427>>, ELLIOT, Rebecca <<http://orcid.org/0000-0002-7602-010X>> and KLEYN, C. Elise <<http://orcid.org/0000-0003-4038-5833>>

Available from Sheffield Hallam University Research Archive (SHURA) at:
<https://shura.shu.ac.uk/37282/>

This document is the Published Version [VoR]

Citation:

LADA, Georgia, MCKIE, Shane, KAFKAS, Alex, MULLINGS, Emma, GRIFFITHS, Christopher E.M., ELLIOT, Rebecca and KLEYN, C. Elise (2026). Towards an understanding of neural responses to emotion in recent onset and chronic psoriasis: a feasibility brain imaging study. *JID Innovations*: 100476. [Article]

Copyright and re-use policy

See <http://shura.shu.ac.uk/information.html>

Journal Pre-proof

Towards an understanding of neural responses to emotion in recent onset and chronic psoriasis: a feasibility brain imaging study

Georgia Lada, Shane Mckie, Alex Kafkas, Emma Mullings, Christopher E.M. Griffiths, Rebecca Elliot, C. Elise Kleyn

PII: S2667-0267(26)00029-9

DOI: <https://doi.org/10.1016/j.xjidi.2026.100476>

Reference: XJIDI 100476

To appear in: *JID Innovations*

Received Date: 18 October 2025

Revised Date: 12 March 2026

Accepted Date: 19 March 2026



Please cite this article as: Lada G, Mckie S, Kafkas A, Mullings E, Griffiths CEM, Elliot R, Kleyn CE, Towards an understanding of neural responses to emotion in recent onset and chronic psoriasis: a feasibility brain imaging study, *JID Innovations* (2026), doi: <https://doi.org/10.1016/j.xjidi.2026.100476>.

This is a PDF of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability. This version will undergo additional copyediting, typesetting and review before it is published in its final form. As such, this version is no longer the Accepted Manuscript, but it is not yet the definitive Version of Record; we are providing this early version to give early visibility of the article. Please note that Elsevier's sharing policy for the Published Journal Article applies to this version, see: <https://www.elsevier.com/about/policies-and-standards/sharing#4-published-journal-article>. Please also note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2026 Published by Elsevier Inc. on behalf of Society for Investigative Dermatology.

Author Details

Manuscript Title:

Towards an understanding of neural responses to emotion in recent onset and chronic psoriasis: a feasibility brain imaging study

Short title:

Psoriasis and emotion: an fMRI study

Author names and affiliations:

Georgia Lada ^{1,2}, Shane Mckie ³, Alex Kafkas ⁴, Emma Mullings ⁵, Christopher E.M. Griffiths ^{6,7}, Rebecca Elliot ⁸, C. Elise Kleyn ¹

¹ Dermatology Centre, Northern Care Alliance NHS Foundation Trust, Manchester Academic Health Science Centre, Division of Musculoskeletal & Dermatological Sciences, The University of Manchester, Manchester, M6 8HD, United Kingdom

² Greater Manchester Mental Health NHS Foundation Trust, Manchester, M25 3BL, United Kingdom

³ Faculty of Biology, Medicine and Health, The University of Manchester, Manchester, UK

⁴ Division of Psychology, Communication and Human Neuroscience, School of Health Sciences, University of Manchester, Manchester, UK

⁵ Department of Psychology, Sociology & Politics, Sheffield Hallam University, Sheffield, UK

⁶ Centre for Dermatology Research, University of Manchester, Manchester, UK

⁷ Department of Dermatology, King's College Hospital, King's College London, London, UK

⁸ Division of Psychology and Mental Health, University of Manchester, Manchester, UK

City and country in which the work was done:

Manchester, United Kingdom

Corresponding Author name and contact information:

Georgia Lada, The Dermatopharmacology Unit, The University of Manchester, 1st Floor, Barnes Building, Salford Royal Hospital, Stott Lane, Greater Manchester M6 8HD, UK. Switchboard: 0161 789 7373. Email: georgia.lada@manchester.ac.uk.

Author ORCiDs:

Georgia Lada <https://orcid.org/0000-0003-1926-2599>

Shane Mckie <https://orcid.org/0000-0002-4418-8944>

Alex Kafkas <https://orcid.org/0000-0001-5133-8827>

Emma Mullings <https://orcid.org/0000-0001-5718-5112>

Christopher E.M. Griffiths <https://orcid.org/0000-0001-5371-4427>

Rebecca Elliot <https://orcid.org/0000-0002-7602-010X>

C. Elise Kleyn <https://orcid.org/0000-0003-4038-5833>

Author Contributions:

Conceptualization: RE, CEK; Methodology: GL, AK, SM, EM, RE, CEK; Data curation: SM, EM; Investigation: EM, CEK; Software: GL, AK, SM, EM; Formal analysis: GL, AK, SM, EM;

Visualization: GL, AK; Resources: CEM, RE, CEK; Supervision: RE, CEK; Writing- Original draft preparation: GL; Writing – review and editing: GL, AK, SM, EM, CEM, RE, CEK.

Data Availability Statement:

The data that support the findings of this study are available from the corresponding author upon reasonable request. The data are not publicly available due to privacy or ethical restrictions.

Ethics Statement:

The study was approved by the National Research Ethics Service NRES (Ref: 13/NW/0549).

Written, informed consent was obtained from all participants.

Clinical Trial Information (if applicable):

N/A.

Conflict of Interest:

GL has received speaker honoraria from Janssen, Lilly, Leo, and Novartis. CEMG has received honoraria and/or research grants from AbbVie, Almirall, Amgen, Anaptysbio, Artax, Boehringer Ingelheim, Bristol-Myers Squibb, Celltrion, Eli Lilly, Evelo Bioscience, Galderma, GSK, Imogene, Kyowa Kerin, Janssen, ONO Pharmaceuticals, Novartis, Pfizer, Sun Pharma and UCB Pharma. CEK conflicts of interest are (including consulting fees, research or institutional support, and educational grants): Almirall, Amgen, Celgene, Eli Lilly and Company, Janssen Pharmaceuticals, La Roche-Posay, LEO Pharma, Novartis, Pfizer, and UCB. CEK is also a

Visiting Professor at the University of Bolton. All other authors have no conflicts of interest to declare.

Declaration of Artificial Intelligence/Large Language Model use

The authors declare no use of Artificial Intelligence/Large Language Model in the preparation of this article.

Acknowledgments:

RE was supported by the NIHR Manchester Biomedical Research Centre (NIHR203308). The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care.

Funding statement:

A Pfizer I-CRP research grant contributed to funding the research project. AK is supported by the Medical Research Council (MR/Y004507/1). The funder had no involvement in the study design, data collection, analysis, interpretation or the writing of the manuscript.

Keywords:

psoriasis, fMRI, emotion, psychodermatology

Abbreviations:

Analysis of variance (ANOVA)

Blood Oxygenation Level Dependent (BOLD)

Chronic psoriasis (CD)

Estimated marginal means (EMMs)

Functional magnetic resonance imaging (fMRI)

General linear model (GLM)

International Affective Picture System (IAPS)

Psoriasis Area and Severity Index (PASI)


Recent onset psoriasis (ROD)

Region of interest (ROI)

Journal Pre-proof


Shorter duration of psoriasis is associated with an attenuated response to faces expressing disgust in the left fusiform cortex.

This may reflect early coping strategies of people experiencing skin-related aversive social cuing.



Duration of psoriasis

Age of onset modulates disgust processing in the hippocampus.



Right hippocampal response to disgust in people with chronic (CD) versus people with recent onset (ROD) psoriasis depending on age of onset

Age of psoriasis onset

TITLE

Towards an understanding of neural responses to emotion in recent onset and chronic psoriasis: a feasibility brain imaging study

SHORT TITLE

Psoriasis and emotion: an fMRI study

MANUSCRIPT WORD COUNT: 3,333

ABSTRACT WORD COUNT: 199

TABLE COUNT: 5

FIGURE COUNT: 2

¹ABBREVIATIONS

¹ **Abbreviations:** Analysis of variance (ANOVA), Blood Oxygenation Level Dependent (BOLD), Chronic psoriasis (CD), Estimated marginal means (EMMs), Functional magnetic resonance imaging (fMRI), General linear model (GLM), International Affective Picture System (IAPS), Psoriasis Area and Severity Index (PASI), Recent onset psoriasis (ROD), Region of interest (ROI)

ABSTRACT

Psoriasis is linked to significant stigmatization. Prior research suggests that people with psoriasis demonstrate altered neurobiological responses to disgust. However, chronically affected patients may develop coping mechanisms to disgust-related social cues.

We investigated whether duration of psoriasis is associated with more attenuated responses to disgust. We used brain functional magnetic resonance imaging while conducting a covert facial recognition task and a task involving emotional-stimulating pictures in patients with chronic psoriasis (CD), patients with recent onset psoriasis (ROD) and controls without skin disease.

We found no differences in disgust processing between patients and healthy controls. Shorter duration of psoriasis was marginally associated with an attenuated brain response to disgust in the left fusiform gyrus within the inferior temporal cortex. An inverse relationship was observed between the age of onset and the hippocampus response when comparing the CD and ROD patient groups.

Our findings suggest that both the duration and the age of psoriasis onset may modulate disgust processing in patients, possibly reflecting evolved learned strategies and disease coping mechanisms. Timely pharmacological and psychosocial interventions for psoriasis may be beneficial for people diagnosed during life stages that may increase vulnerability to neurocognitive changes. Further studies are needed to replicate these results.

INTRODUCTION

The psychosocial burden of psoriasis as well as the associated disability and life-changing impact are well recognized. Patients with psoriasis commonly experience stigmatization and overt public rejection because of the visibility of skin lesions (Germain et al., 2021) and have a

high burden of psychiatric morbidity, primarily depression (Parisi et al., 2019). Patients demonstrate an attentional bias to words signalling threat, including disgust, which may be caused by stigmatization fear and is followed by social avoidance patterns (Etty et al., 2024, Fortune et al., 2003). However, the neurocognitive processes underlying the internalized stigma, self-image and overall psychological consequences of psoriasis are not clear.

Kleyn et al. previously demonstrated differences in the neural processing of disgust in 13 male psoriasis patients compared to healthy controls, using a validated covert, face processing task implemented in functional Magnetic Resonance Imaging (fMRI) (Kleyn et al., 2009). In fMRI, local neural activity is associated with the BOLD (blood-oxygen-level-dependent) signal which reflects changes in blood oxygenation. For example, when neural activity increases in a brain region, local blood flow typically increases, producing a measurable change in the BOLD signal. In the study by Kleyn et al., patients showed a significantly reduced BOLD response to disgust in the insula cortex, which is considered a key brain region in disgust recognition and processing (Chapman and Anderson, 2012). Importantly, duration of psoriasis was proportional to attenuated insular signal.

Psoriasis duration has previously been linked to less distress and higher acceptance (Hill and Kennedy, 2002). However, it is not clear if the neural responses found in this previous study reflect an isolated protective coping mechanism in psoriasis or a fundamental change in patients' emotional processing in the form of a trait. Such changes may be related to self-perception and contribute to psychiatric pathology and functional impairment. Furthermore, disgust processing in the human brain is not restricted to the insula and it is not clear whether altered neural responses in psoriasis may extend to other areas, importantly the fusiform cortex, the amygdala and the orbitofrontal cortex, which have been linked to disgust and emotional face processing (Fusar-Poli et al., 2009, Habel et al., 2007, Harry et al., 2013, Schienle et al.,

2002). Understanding these processes in psoriasis is critical in order to develop appropriate interventions for the psychosocial impact of psoriasis and tailor them for patients.

We developed a feasibility study to investigate whether emotional processing of disgust in psoriasis is a core marker of its psychosocial effect and reflects long-standing psychological processes over time. We therefore hypothesised that patients with long-standing moderate-to-severe psoriasis (disease duration ≥ 10 years) would show decreased brain responses to disgust compared to newly diagnosed patients (disease duration ≤ 6 years) and healthy volunteers. Secondly, we aimed to explore synergistic effects of age of onset alongside disease duration on disgust processing given evidence suggesting a negative association between age and coping difficulties in psoriasis (Hill and Kennedy, 2002).

RESULTS

Participants

Twenty patients with long-standing psoriasis (50% women; mean age: 34.7 ± 8.4 years, range: 21–50; median disease duration 20 years, range 10–42), 16 patients with a recent onset of psoriasis (69% women; mean age: 27.5 ± 8.3 years, range: 19–44; median disease duration 2.5 years, range 0.2–6) and 19 controls (42% women; mean age: 32.9 ± 8.1 years, range: 21–48) with no history of skin disease participated in the study (Table 1).

The ROD patients were younger than the CD patients and had a significantly higher mean age of onset. The patient groups did not differ for psoriasis severity (PASI: mean score 5.6 ± 3.2 , range: 0.7–15.6 for the CD group and 5.1 ± 2.8 , range: 1.4–11.1 for the ROD group), psoriasis-related quality of life or psychometric scores for depression, disgust sensitivity or disgust propensity (Table 1).

FMRI Data

Whole brain analysis of BOLD responses to disgust-expressing faces compared with neutral faces across all three groups showed activations in regions within the bilateral middle occipital gyrus, the bilateral middle temporal and fusiform gyri and in the left insula. The opposite contrast showed activations in the anterior cingulate and medial prefrontal cortex and the bilateral calcarine cortex. Table 2 presents these activations, reflecting the regions where the group-level analysis in the total sample indicated a significant difference between the disgust and neutral conditions in the faces task along with the voxel counts representing the spatial extent of these significant activation clusters.

Whole-brain analysis of BOLD responses to disgust-inducing images, compared to neutral images, across all three groups revealed bilateral activations in the middle occipital cortex extending to the fusiform gyrus, the amygdala, the inferior frontal gyrus, the dorsal striatum, the thalamus, the right medial frontal gyrus, the cingulate gyrus, and the right supramarginal gyrus (Table 3). Conversely, the opposite contrast demonstrated activations in the right cuneus, insula, and superior motor cortex, as well as the left superior temporal gyrus (Table 3). Whole-brain analysis activations are summarised in Figure 1.

Overall, the findings of our whole-brain analysis across all three groups aligned with our a priori hypotheses and previous literature showing involvement of the insula, the fusiform gyrus, the amygdala and inferior frontal areas in processing disgust.

We then performed ROI analyses to investigate differences among the three groups within these areas of interest and the hippocampus, which was identified in our a priori hypothesis. To compare disgust processing across groups, we measured the difference in BOLD responses when viewing facial expressions of disgust and disgust-provoking images in the two respective experimental paradigms. For comparative purposes, we also measured the difference in BOLD

responses when viewing facial expressions of neutral and fearful/fear-provoking expressions and images.

During the Covert Emotion Processing Task, we found a group difference in the BOLD response to disgust in the left fusiform ($F(2, 49)=3.36, p=0.043$). Post-hoc analysis showed significantly higher signal values for the CD group than the ROD group (estimated marginal means (EMMs) 0.48 95% CIs 0.34-0.62 vs 0.18 95% CIs 0.006-0.36; p -value=0.036).

There were no further significant differences between the groups for any emotion or ROI in either task (Tables 4 and 5). Interestingly, however, there was a marginal group effect on the response to fearful faces in the left fusiform ($F(2, 49)=3.04, p=0.057$). Similarly to disgust, this effect reflected higher BOLD signal values for the CD group than the ROD group (EMMs 0.45 95% CIs 0.31-0.59 vs 0.18 95% CIs 0.004-0.36; p -value=0.053).

We did not find any differences of the BOLD response to neutral faces among the groups.

Furthermore, we explored whether the onset of psoriasis may moderate emotional processing of disgust in the two groups. During the Covert Emotion Processing Task, we found a significant interaction between the patient group and the age of onset in the right hippocampus ($F(1,29)=5.89, p=0.02$). This suggested an inverse trend in the relationship between the age of onset and the BOLD response for the CD and ROD groups. Following post-hoc tests, the association between the age of onset and the right hippocampus signal was not statistically significant for either of the groups (Figure 2). There were no significant interactions or main effects of the age of onset on the BOLD response in any other ROIs in this task. Finally, in the Picture Viewing (IAPS) Task, we did not find an interaction of age of onset with group and disgust processing in any ROI.

DISCUSSION

We found that shorter duration of psoriasis was associated with an attenuated BOLD response to disgust in the left fusiform. However, there were no significant differences in disgust processing in any ROI between patients and healthy controls. This is in contrast to the previous study by Kleyn et al., showing reduced insula response to disgust in patients with psoriasis compared to healthy controls (Kleyn et al., 2009). The reason for this discrepancy is not clear, but may be related to scanner differences between the two studies as well as different patient populations and sample sizes; for example, Kleyn et al. included a smaller male-only sample (thirteen men) limiting the extrapolation of its findings.

The fusiform cortex is involved in facial recognition, however growing evidence suggests an important role in emotional face processing (Harry et al., 2013). This is supported by our findings, where the activation trends in the left fusiform observed for disgust and, to lesser extent, fear processing were not replicated for neutral faces. Furthermore, our findings align with a large meta-analysis of emotional processing studies in humans, which reported that activation in the left fusiform gyrus supports disgust processing (Fusar-Poli et al., 2009). Expressed disgust is perceived as a cue for violating social norms and is often experienced as shame leading to social alienation (Giner-Sorolla and Espinosa, 2011). ROD patients' attenuated processing of expressed disgust in the left fusiform gyrus may, therefore, also reflect a coping strategy to this aversive social cuing. The lack of group differences for the picture-viewing task may indicate differential processing of personal experiences of disgust versus recognising others' disgust, further supporting the interpretation that psoriasis affects disgust processing in an interpersonal context.

Our secondary analysis indicated that age of onset might play a role in how people with psoriasis recognize and process others' expressed disgust. We found an inverse effect of the age of onset on the BOLD response in the right hippocampus for the CD and ROD groups, with a positive correlation trend in the CD group. It is not clear why this effect was only found for

the hippocampus. Although our findings implicate the hippocampus, one possibility requiring further investigation is that it regulates activity in the insula. Hippocampal connectivity with insula has been linked to internally generated disgust, supported by memory retrieval and imagery, when viewing disgust-provoking images (Fitzgerald et al., 2004). Additionally, hippocampus-insula connectivity has been found to increase with disgust-associated imagery (Jabbi et al., 2008). These changes may reflect the recall of disgust-associated autobiographical events and represent evolved strategies over time via learning and conditioning (Fitzgerald et al., 2004, Jabbi et al., 2008).

In psoriasis, neurocognitive restructuring may develop more effectively in individuals with early childhood disease onset, as well as in those who acquire psoriasis later in life, after completing their identity development and establishing their personal and social resources. Indeed, a number of studies have found an association between age and resilience in adults (Cohen et al., 2014, Portzky et al., 2010), with more frequent use of maladaptive mechanisms in adolescents and younger adults, including self-blame and hostile reaction (Irion and Blanchard-Fields, 1987). In adults with psoriasis, age has been negatively correlated with maladaptive coping and distress (Hill and Kennedy, 2002). People with adolescent and young adulthood onset of disease may, therefore, be less able to develop adequate emotional regulating defence mechanisms to cope with the consequences of psoriasis in the social context. Furthermore, recognition of others' facial expressions of disgust starts being shaped at around 9 years of age, which may also explain the differential processing we observed in the group with earlier childhood compared to late adolescence onset (Widen and Russell, 2013). People diagnosed in late adolescence and young adulthood could therefore benefit most from interventions offering psychosocial support and teaching adaptive coping skills such as Cognitive-Behavioural Therapy (CBT)-based programs to improve their emotional wellbeing and psychological functioning (Sansom-Daly et al, 2012).

Our study has some limitations. Although larger than the previous pilot sample of Kleyn et al., the present study sample remains relatively small and smaller effects of psoriasis on emotional processing may not be detectable. As a result, non-significant findings should be interpreted with caution and not as definitive evidence of no effect. Furthermore, given the exploratory nature of the ROI analysis, no correction for multiple comparisons was applied at the ROI analysis. This may increase the risk of Type I error and the reported p-values should be interpreted as exploratory rather than confirmatory. However, it is of note that both effects that were around the uncorrected significance level were in the fusiform gyrus. Replication in larger cohorts will be important to confirm these findings. Finally, we have tested for several demographic and disease characteristics among groups and controlled for age differences. However, it is possible that unmeasured confounding factors, such as other subclinical psychopathology, may have affected our findings.

In conclusion, we found no significant differences in the processing of expressed disgust between patients with psoriasis and controls. However, our results suggest that the duration of psoriasis and age of psoriasis onset may contribute to how patients process and cope with social reactions to their skin, which may consequently affect their experience of internalized stigma, feelings of shame and the severe psychosocial consequences of psoriasis. These effects warrant further research and could have implications on treatment decisions, particularly since stress from others' reactions and stress-perpetuating avoidance behaviours have been shown to be the most important markers of disability in psoriasis (Fortune et al., 1997). Early intervention and psychological support may be of most benefit for patients who experience psoriasis in their formative years (adolescence and early adulthood), particularly in the first years after diagnosis, as they may present with higher vulnerability, experience more severe distress and less resilience in their social interactions due to their skin.

MATERIALS AND METHODS

Recruitment

Patients were recruited from our weekly tertiary psoriasis clinic, local general practices and a database of patients with declared interest in research. Healthy volunteers responded to advertisement.

We included patients with a dermatologist-confirmed diagnosis of moderate-to-severe chronic plaque psoriasis. Healthy volunteers had no history of skin disease or significant disfigurement in their family. Inclusion criteria for all participants were: (i) aged between 18 and 50 years; (ii) right-handed; (iii) not taking psychotropic medication; (iv) no contraindications to fMRI scanning; (v) no history of severe, enduring mental or neurological illness; and (vi) not pregnant.

Clinical assessment

Following informed consent, all participants underwent clinical assessment on the day of the scan. This included full medical history, physical examination, Psoriasis Area and Severity Index (PASI), where appropriate, and, for women, a urine pregnancy test. PASI is a widely used physician-rated measure of psoriasis severity providing a composite score (range 0-72)(Fredriksson and Pettersson, 1978). The Dermatology Life Quality Index (DLQI) (Finlay and Khan, 1994) was used to assess the impact of psoriasis on quality of life.

Experimental design

Each participant was presented with a series of pictures, while brain scans were acquired. Participants were first presented with facial expressions of emotion, including disgusted, fearful and neutral expressions, as part of the covert recognition of faces task (Ekman and Friesen, 1976). We describe the protocol for this paradigm in detail elsewhere (Kleyn et al., 2009). In

summary, participants were shown face stimuli (expressing emotions and neutral faces with each face shown for 3 seconds) and were asked to determine the gender of the face. Covert brain responses to emotions could therefore be studied. In order to derive baseline measures for the Blood Oxygenation Level Dependent (BOLD) signal-based analysis, blocks of rest periods were also included.

In a separate task, each participant viewed a set of colour photographs, selected from the International Affective Picture System (IAPS) (Lang et al., 1997), which provides normative ratings for a set of photographs for investigations of emotion and attention. Participants were shown randomly 21 disgust-inducing pictures, 21 fear-inducing pictures and 21 pictures with neutral content (4 seconds for each picture).

Both tasks were included, because, although limited evidence suggests shared substrates between the experience of disgust and identification of expressions of disgust in others (Wicker et al., 2003), it is currently not clear whether facial expressions of disgust and disgust-inducing pictures are processed differently and engage distinct neural networks.

Neuroimaging parameters

Brain images were acquired in a Philips 3.0 Tesla MR scanner. T2*-weighted/echo-planar imaging (EPI) data were acquired separately from each task (for the faces task: 3x3x4mm voxels size; 80x80 matrix size; TR (repetition time)=2.3; TE (echo time)=dual echo of 12 and 30; total volumes=192; for the IAPS task: 3x3x4mm voxels size; 80x80 matrix size; TR=2.5; TE=dual echo of 12 and 30, total volumes=290). A T1-weighted structural image was also acquired for each participant (1mm voxel size isotropic) to exclude structural abnormalities and for use in spatial pre-processing of functional images. A healthy control with abnormal scan was excluded from the analysis.

Imaging analysis

Imaging data were analysed using Statistical Parametric Mapping (SPM5) (Penny et al., 2011), with a random effects model. We used the same imaging analysis methods for the two tasks (faces and IAPS) and analysed data from them separately.

EPI data were re-aligned to the first image using a standard six-parameter rigid body transformation and reslicing using sinc interpolation, coregistered to the individual T1-weighted images and spatially normalised to Montreal Neurological Institute (MNI) space. The data were then spatially smoothed using an 8 mm isotropic full width half maximum (FWHM) Gaussian kernel.

Following preprocessing, data were analysed using the general linear model (GLM) at the first-level analysis for each participant. The onset and duration of each block of interest (neutral, disgust, fear) were convolved with the canonical haemodynamic response function. The six movement parameters produced at realignment for each participant were used as regressors of no interest to account for residual effects of movement artefacts. A high-pass filter of 128 seconds was applied to the data.

To explore whole brain responses to disgust first-level t contrasts were created contrasting brain responses to disgust compared with neutral blocks (i.e., $\text{disgust} > \text{neutral}$ and $\text{neutral} > \text{disgust}$) across the whole sample. In more detail, in first-level (within-subject) analysis, a contrast map was created for each participant, representing the magnitude and direction of the difference in BOLD activation between disgust and neutral stimuli at each voxel. The BOLD signal is measured separately within each voxel, which is the basic three-dimensional unit of the brain image. Multiple neighbouring voxels showing a statistically significant BOLD response form an activation cluster and the number of voxels in the cluster reflects the spatial extent of the activation.

The individual contrast maps from the first-level analysis were then entered into the second-level (group-level) analysis. Second-level group analyses were implemented as one sample t-tests, exploring the group responses to disgust and neutral blocks and testing whether the BOLD signal is significantly greater for one condition relative to the other at each voxel across the sample. Significance in all analyses was determined at a cluster-corrected family-wise error (FWE) $p < 0.05$, from an initial voxel-based level of $p < 0.001$ (Flandin and Novak, 2020; Friston et al., 1994).

To explore group differences, we performed a region of interest (ROI) analysis in areas found to be activated in the whole sample that were also a priori hypothesised to play a role in disgust based on existing evidence. These include the bilateral anterior insula, fusiform gyrus, frontal inferior orbital cortex and amygdala (Fusar-Poli et al., 2009, Kleyn et al., 2009, Schienle et al., 2002). We also included the hippocampus in the analysis based on *a priori* hypotheses regarding its role in learning and retrieval (Kafkas et al., 2024). For the ROI analysis, activation data (parameter estimates) were extracted using 6mm spheres around the GLM local maxima from the whole-brain analysis.

Statistical analysis

Statistical analysis was carried out in Rstudio. Participant characteristics are reported as mean \pm standard deviation unless stated otherwise. We compared groups for baseline characteristics, including potentially confounding measured variables such as depression scores, using classical statistical tests as appropriate for the type of data, including chi-square tests, t-tests and Mann-Whitney U tests; one-way analysis of variance (ANOVA) or the Kruskal-Wallis test were used where means or medians, respectively, among all three groups were compared.

To compare the BOLD signal (parameter estimates) across the three groups, we performed ANOVAs for each ROI, adjusting for age to control for age effects on the hemodynamic response to emotional faces (Iidaka et al., 2002). For significant group effects in ANOVAs, we performed post-hoc comparisons using Tukey's method to adjust for multiple comparisons. To test effects of age of psoriasis onset on the BOLD response to disgust among patients, we performed ANOVAs using interaction terms between the age of onset and the patient group (dichotomous variable). All models were age-adjusted and checked for multicollinearity using variance inflation factor (VIF) <5 . The level of statistical significance was set at 0.05 for all analyses.

REFERENCES

- Chapman HA, Anderson AK. Understanding disgust. *Ann N Y Acad Sci* 2012;1251(1):62-76.
<https://doi.org/10.1111/j.1749-6632.2011.06369.x>.
- Chen A, Beck KM, Tan E, Koo J. Stigmatization in psoriasis. *Journal of Psoriasis and Psoriatic Arthritis* 2018;3(3):100-6. <https://doi.org/10.1177/2475530318781339>.
- Cohen M, Baziliansky S, Beny A. The association of resilience and age in individuals with colorectal cancer: an exploratory cross-sectional study. *J Geriatr Oncol* 2014;5(1):33-9. <https://doi.org/10.1016/j.jgo.2013.07.009>.
- Ekman P, Friesen WV. *Pictures of facial affect consulting psychologists press*. Palo Alto, CA 1976.
- Etty S, George DN, van Laarhoven AI, Kleyn CE, Walton S, Holle H. Attentional bias in psoriasis: The role of processing time and emotional valence. *Br J Health Psychol* 2024;29(3):533-50. <https://doi.org/10.1111/bjhp.12712>.
- Finlay AY, Khan G. Dermatology Life Quality Index (DLQI)—a simple practical measure for routine clinical use. *Clin Exp Dermatol* 1994;19(3):210-6.
<https://doi.org/10.1111/j.1365-2230.1994.tb01167.x>.
- Fitzgerald DA, Posse S, Moore GJ, Tancer ME, Nathan PJ, Phan KL. Neural correlates of internally-generated disgust via autobiographical recall: a functional magnetic resonance imaging investigation. *Neurosci Lett* 2004;370(2-3):91-6.
<https://doi.org/10.1016/j.neulet.2004.08.007>.
- Flandin G, Novak MJ. *fMRI data analysis using SPM. InfMRI: Basics and Clinical Applications*. Cham: Springer International Publishing; 2020. p. 89-116.
https://doi.org/10.1007/978-3-030-41874-8_8.

- Fortune D, Main C, O'sullivan T, Griffiths C. Quality of life in patients with psoriasis: the contribution of clinical variables and psoriasis-specific stress. *Br J Dermatol* 1997;137(5):755-60. <https://doi.org/10.1046/j.1365-2133.1997.19472062.x>.
- Fortune DG, Richards HL, Corrin A, Taylor RJ, Griffiths CE, Main CJ. Attentional bias for psoriasis-specific and psychosocial threat in patients with psoriasis. *J Behav Med* 2003;26:211-24. <https://doi.org/10.1023/A:1023408503807>.
- Fredriksson T, Pettersson U. Severe psoriasis—oral therapy with a new retinoid. *Dermatology* 1978;157(4):238-44. <https://doi.org/10.1159/000250839>.
- Friston KJ, Holmes AP, Worsley KJ, Poline JP, Frith CD, Frackowiak RS. Statistical parametric maps in functional imaging: a general linear approach. *Hum Brain Mapp* 1994;2(4):189-210.
- Fusar-Poli P, Placentino A, Carletti F, Landi P, Allen P, Surguladze S, et al. Functional atlas of emotional faces processing: a voxel-based meta-analysis of 105 functional magnetic resonance imaging studies. *J Psychiatry Neurosci* 2009;34(6):418-32. <https://doi.org/10.1139/jpn.0953>.
- Germain N, Augustin M, François C, Legau K, Bogoeva N, Desroches M, et al. Stigma in visible skin diseases—a literature review and development of a conceptual model. *J Eur Acad Dermatol Venereol* 2021;35(7):1493-504. <https://doi.org/10.1111/jdv.17110>.
- Giner-Sorolla R, Espinosa P. Social cuing of guilt by anger and of shame by disgust. *Psychol Sci* 2011;22(1):49-53. <https://doi.org/10.1177/0956797610392925>.
- Habel U, Windischberger C, Derntl B, Robinson S, Kryspin-Exner I, Gur RC, et al. Amygdala activation and facial expressions: explicit emotion discrimination versus implicit emotion processing. *Neuropsychologia* 2007;45(10):2369-77. <https://doi.org/10.1016/j.neuropsychologia.2007.01.023>.

- Harry B, Williams MA, Davis C, Kim J. Emotional expressions evoke a differential response in the fusiform face area. *Front Hum Neurosci* 2013;7:692.
<https://doi.org/10.3389/fnhum.2013.00692>.
- Hill L, Kennedy P. The role of coping strategies in mediating subjective disability in people who have psoriasis. *Psychol Health Med* 2002;7(3):261-9.
<https://doi.org/10.1080/13548500220139340>.
- Hrehorów E, Salomon J, Matusiak U, Reich A, Szepietowski JC. Patients with psoriasis feel stigmatized. *Acta Derm Venereol* 2012;92(1):67-72.
<https://doi.org/10.2340/00015555-1193>.
- Iidaka T, Okada T, Murata T, Omori M, Kosaka H, Sadato N, et al. Age-related differences in the medial temporal lobe responses to emotional faces as revealed by fMRI. *Hippocampus* 2002;12(3):352-62. <https://doi.org/10.1002/hipo.1113>.
- Irion JC, Blanchard-Fields F. A cross-sectional comparison of adaptive coping in adulthood. *J Gerontol* 1987;42(5):502-4. <https://doi.org/10.1093/geronj/42.502>.
- Jabbi M, Bastiaansen J, Keysers C. A common anterior insula representation of disgust observation, experience and imagination shows divergent functional connectivity pathways. *PLoS One* 2008;3(8):e2939. <https://doi.org/10.1371/journal.pone.0002939>.
- Kafkas A, Mayes AR, Montaldi D. The hippocampus supports the representation of abstract concepts: Implications for the study of recognition memory. *Neuropsychologia* 2024;199:108899. <https://doi.org/10.1016/j.neuropsychologia.2024.108899/>
- Kleyn CE, McKie S, Ross AR, Montaldi D, Gregory LJ, Elliott R, et al. Diminished neural and cognitive responses to facial expressions of disgust in patients with psoriasis: a functional magnetic resonance imaging study. *J Invest Dermatol* 2009;129(11):2613-9. <https://doi.org/10.1038/jid.2009.152>.

- Lang PJ, Bradley MM, Cuthbert BN. International affective picture system (IAPS): Technical manual and affective ratings. NIMH Center for the Study of Emotion and Attention 1997;1(39-58):3.
- Parisi R, Webb RT, Kleyn CE, Carr MJ, Kapur N, Griffiths CEM, et al. Psychiatric morbidity and suicidal behaviour in psoriasis: a primary care cohort study. *Br J Dermatol* 2019;180(1):108-15. <https://doi.org/10.1111/bjd.17004>.
- Penny WD, Friston KJ, Ashburner JT, Kiebel SJ, Nichols TE. *Statistical parametric mapping: the analysis of functional brain images*: Elsevier, 2011.
- Portzky M, Wagnild G, De Bacquer D, Audenaert K. Psychometric evaluation of the Dutch Resilience Scale RS-nl on 3265 healthy participants: a confirmation of the association between age and resilience found with the Swedish version. *Scand J Caring Sci* 2010;24:86-92. <https://doi.org/10.1111/bjd.17004>.
- Sansom-Daly UM, Peate M, Wakefield CE, Bryant RA, Cohn RJ. A systematic review of psychological interventions for adolescents and young adults living with chronic illness. *Health Psychol.* 2012;31(3):380. <https://doi.org/10.1037/a0025977>.
- Schienle A, Stark R, Walter B, Blecker C, Ott U, Kirsch P, et al. The insula is not specifically involved in disgust processing: an fMRI study. *Neuroreport* 2002;13(16):2023-6. <https://doi.org/10.1097/00001756-200211150-00006>.
- Wicker B, Keysers C, Plailly J, Royet J-P, Gallese V, Rizzolatti G. Both of us disgusted in My insula: the common neural basis of seeing and feeling disgust. *Neuron* 2003;40(3):655-64. [https://doi.org/10.1016/S0896-6273\(03\)00679-2](https://doi.org/10.1016/S0896-6273(03)00679-2).
- Widen SC, Russell JA. Children's recognition of disgust in others. *Psychol Bull* 2013;139(2):271. <https://doi.org/10.1037/a0031640>.

TABLES

Table 1. Baseline Characteristics

Baseline Characteristic	HC (n=19)	CD (n=20)	ROD (n=16)	p – value
Sex, female (%)	8 (42%)	10 (50%)	11 (69%)	0.277
Age years, mean (sd)	32.9 (8.1)	34.7 (8.4)	27.5 (8.3)	0.036
BMI, mean (sd)	23.1 (3.6)	27.3 (3.4)	25.0 (5.1)	0.009
Disgust Sensitivity, mean (sd)	9.5 (3.1)	8.9 (2.5)	10.3 (2.9)	0.338
Disgust Propensity, mean (sd)	12.9 (4.5)	15.2 (3.1)	15.6 (3.9)	0.088
BDI, median (IQR)	1 (1.5)	3.5 (7)	3.5 (11.3)	0.414
PASI, mean (sd)		5.6 (3.2)	5.1 (2.8)	0.598
DLQI, median (IQR)		5 (3.5)	5 (5.5)	0.469
Age of Onset, mean (sd)		12.6 (7.5)	24.7 (8.3)	<0.001
Disease Duration, median (IQR)		20 (11.25)	2.5 (3.1)	<0.001
Treatment				0.218

Topical		9 (45%)	8 (50%)	
Phototherapy		0 (0%)	2 (12.5%)	
Oral		6 (30%)	2 (12.5%)	
Biologics		2 (10%)	1 (6.2%)	
Other		2 (10%)	0 (0%)	
None		1 (5%)	3 (18.8%)	

Notes: sd=standard deviation, IQR= Interquartile Range, BMI=Body Mass Index, FNE=Fear of Negative Evaluation, BDI= Beck Depression Inventory, PASI= Psoriasis Area and Severity Index, DLQI= Dermatology Life Quality Index. Chi-square tests were used for sex and treatment data. T-tests and Mann-Whitney U tests were used where means and medians, respectively, among patient groups were compared. One-way analysis of variance (ANOVA) and the Kruskal-Wallis test were used where means and medians, respectively, among all three groups were compared.

Table 2. Brain responses during the recognition of facial expressions of disgust compared to neutral expressions

L/R	Region	Voxels	BA	T-value	MNI coordinates (x y z)
Disgust > Neutral					
L	Middle occipital gyrus	118	19	5.60	-36 -91 6
L	Middle temporal gyrus and fusiform gyrus	87	22, 37	4.90	-48 -61 -2
R	Middle temporal gyrus and fusiform gyrus	92	37	4.49	45 -61 -2
R	Middle occipital gyrus	54	19	4.34	42 -82 18
L	Insula	22	13	4.34	-36 -25 26
Neutral > Disgust					

R	Anterior cingulate and medial orbitofrontal cortex	283	32	5.59	18 50 -10
R & L	Calcarine cortex and lingual gyrus	215	17, 18	4.99	15 -85 -2
R	Middle cingulate & superior medial frontal cortex	88	32	4.32	9 32 30

Notes: Voxels refer to the volume elements of the three-dimensional brain image, which are the basic spatial units of measurement in fMRI. The voxel numbers in the table represent the total number of contiguous voxels within each significant activation cluster identified in the group-level analysis. The reported activations reflect only regions where the BOLD (blood-oxygen-level-dependent) response meets or exceeds the predefined statistical threshold, which was determined at an initial voxel-based level of $p < 0.001$ and cluster-corrected family-wise error (FWE) $p < 0.05$. BA refers to the respective Brodmann Area of the cerebral cortex. MNI coordinates refer to coordinates of each region according to the Montreal Neurological Institute (MNI) template, a commonly used standardised brain atlas. The t-value indicates the t-statistic for each region for the disgust>neutral and neutral>disgust contrasts.

Table 3. Brain responses to disgust-inducing images compared to neutral images

L/R	Region	Voxels	BA	T-value	MNI coordinates (x y z)
Disgust > Neutral					
L	Middle occipital and fusiform gyrus	1424	19, 37	11.71	-51 -73 6 & -42 -52 -10
R	Middle occipital and fusiform gyrus	1595	19, 37	10.15	30 -76 22 & 48 -67 6
R	Inferior frontal gyrus, insula and amygdala	811	45, 47, 13	8.83	48 11 18 & 39 29 -2
L & R	Thalamus and dorsal striatum	526		8.01	-3 -31 -2 & 9 5 2
R	Superior medial frontal and middle cingulate gyrus	522	9, 32	7.72	6 50 26 & 6 26 42
L	Amygdala, insula and inferior frontal gyrus	442	13, 47	6.98	-18 -4 -10 & -27 20 -10 & -48 2 26

L	Inferior parietal lobule (supramarginal gyrus)	69	40	4.72	-57 -28 30
Neutral > Disgust					
R	Insula	274	13	5.52	39 -19 14
R	Cuneus	97	18, 19	4.57	6 -82 26
R	Superior motor area	62	6	4.50	9 -19 54
L	Superior temporal gyrus	70	22	4.29	-57 -34 10
<p>Notes: Voxels refer to the volume elements of the three-dimensional brain image, which are the basic spatial units of measurement in fMRI. The voxel numbers in the table represent the total number of contiguous voxels within each significant activation cluster identified in the group-level analysis. The reported activations reflect only regions where the BOLD (blood-oxygen-level-dependent) response meets or exceeds the predefined statistical threshold, which was determined at an initial voxel-based level of $p < 0.001$ and cluster-corrected family-wise error (FWE) $p < 0.05$. BA refers to the respective Brodmann Area of the cerebral cortex. MNI coordinates refer to coordinates of each region according to the Montreal Neurological Institute (MNI) template, a commonly used standardised brain atlas. The t-value indicates the t-statistic for each region for the disgust>neutral and neutral>disgust contrasts.</p>					

Table 4. Faces Task – p-values from the statistical comparisons (ANOVAs) of BOLD responses in regions of interest across the three groups

Region Of Interest (ROI)	Disgust	Fear	Neutral
Amygdala left	0.610	0.481	0.633
Amygdala right	0.693	0.463	0.897
Frontal Inferior Orbital Cortex left	0.277	0.521	0.147
Frontal Inferior Orbital Cortex right	0.156	0.827	0.608
Fusiform left	0.043	0.057 ^a	0.299
Fusiform right	0.230	0.311	0.430
Hippocampus left	0.319	0.058	0.404
Hippocampus right	0.982	0.376	0.932
Insula anterior left	0.627	0.762	0.964
Insula anterior right	0.364	0.436	0.632
<p>Note: The BOLD response refers to the change in blood-oxygen-level–dependent signal measured during the faces fMRI task. ANOVA=Analysis Of Variance. All models were adjusted for age. Significant effects are presented in bold. ^a Trend to significance.</p>			

Table 5. IAPS task – p-values from the statistical comparisons (ANOVAs) of BOLD responses in regions of interest across the three groups

Region Of Interest (ROI)	Disgust	Fear	Neutral
Amygdala left	0.077	0.229	0.079
Amygdala right	0.259	0.515	0.289
Frontal Inferior Orbital Cortex right	0.745	0.571	0.239
Frontal Inferior Orbital Cortex left	0.613	0.682	0.151
Fusiform left	0.293	0.266	0.192
Fusiform right	0.183	0.163	0.176
Hippocampus left	0.298	0.333	0.193
Hippocampus right	0.469	0.654	0.422
Insula anterior left	0.103	0.468	0.458
Insula anterior right	0.624	0.858	0.834

Note: The BOLD response refers to the change in blood-oxygen-level–dependent signal measured during the IAPS fMRI task. ANOVA=Analysis Of Variance. All models were adjusted for age. Significant effects are presented in bold.

FIGURE LEGENDS

Figure 1. **Whole-brain activations for disgust.** (a) Whole-brain activations for disgust compared to neutral faces. (b) Whole-brain activations for disgust compared to neutral pictures.

Figure 2. **Blood Oxygenation Level Dependent (BOLD) signal effects of disgust in the right hippocampus in the Covert Emotion Processing Task in patients with chronic disease (CD) and patients with recent onset disease (ROD), depending on the age of psoriasis onset.** These effects are expressed as arbitrary units (a.u.) of GLM-estimated beta values. The shaded area around the estimated trend line represents confidence intervals. There was a significant interaction between the patient group and the age of onset ($F(1,29)=5.89$, $p=0.02$) suggesting an inverse trend in the relationship between the age of onset and the BOLD response for the CD and ROD groups. The association between the age of onset and the right hippocampus signal was not statistically significant for either of the groups.

