

**Is surgery more effective than non-surgical treatment for spinal stenosis and which non-surgical treatment is more effective? a systematic review**

MAY, Stephen and COMER, C

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/6976/>

---

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

**Published version**

MAY, Stephen and COMER, C (2013). Is surgery more effective than non-surgical treatment for spinal stenosis and which non-surgical treatment is more effective? a systematic review. *Physiotherapy*, 99 (1), 12-20.

---

**Copyright and re-use policy**

See <http://shura.shu.ac.uk/information.html>

**Table 2. Summary of included trials**

<b>Trial<sup>j</sup></b>	<b>Number</b>	<b>Interventions</b>	<b>Outcomes / follow-up<sup>a</sup></b>
Amundsen et al. [1]	31	1. Decompression surgery (13) 2. Orthosis, back school (18) Both groups general physical training	Clinician determined good or bad result at 6 months 1, 4 and 10 years. Good results: 1: 92%, 69%, 92%, 91%.2: 39%, 33%, 47%, 71%
<b>Comer et al. [7]<sup>j</sup></b>	40	1. Walking stick if not using one (20) 2. No walking stick (20)	ZCQ at 2 weeks: NS differences
<b>Cuckler et al. [9]</b>	37 <sup>d</sup>	1. ESI + procaine (20) 2. Saline + procaine (17) Both groups ESI if < 50% better	Success = 75% self-reported improvement at 24 hours and mean 20 months: NS differences
Eskola et al. [13]	40	1. Calcitonin then placebo 2. Placebo then calcitonin	VAS rest pain and jumping, walking distance 1, 3, 4, 6, 12 months. <b>Active stage V placebo: VAS rest (P=0.01); jump pain (P=0.001 / 0.019; walking distance (P=0.007 / 0.14) up to 3 months</b> No long term difference.
<b>Fukusaki et al. [16]</b>	53	1. Saline epidural injection 2. Mepivacaine epidural injection 3. ESI + mepivacaine	Walking distance improvement: 100m (excellent), 20-100m (good), <20m (poor) at 1 week, 1, 3 months <b>2 + 3 V 1 at 1 week (P&lt;0.01); NS at 1 and 3 months</b>
<b>Goren et al. [21]</b>	50	1. US + exercise (17) 2. Sham US + exercise (17) 3. Control (16)	VAS leg / back, Oswestry, treadmill test, medication after 3 weeks of treatment <b>1 + 2 V 3: Leg pain (P&lt;0.007); Oswestry (P&lt;0.014); 1 V 3: medication (P=0.016)</b> 1 V 2 = NS differences

<b>Koc et al. [31]</b>	29	<ol style="list-style-type: none"> <li>1. In-patient physical therapy (10)</li> <li>2. ESI (10)</li> <li>3. Controls (9)</li> </ol>	<p>VAS, flexion, treadmill test, sit-to-stand, Roland-Morris (RMD), NHP at 2 weeks, 1, 3, and 6 months  <b>2 v 3 at 2 weeks: VAS (P=0.008); RMD (P=0.007); NHP (P=0.004). SD in all groups. NS 1 V 2.</b></p>
<b>Kurihara et al. [32]</b>	146	<ol style="list-style-type: none"> <li>1. Opalman (15 mgd)<sup>h</sup> (69)</li> <li>2. Opalman (3 mgd)<sup>h</sup> (77)</li> </ol>	<p>Improvement in sensation, walking distance, leg pain standing pain at 6 months  <b>1 V 2: improvement (P=0.005); improvement in sensation (P=0.008); walking distance (P=0.019).</b></p>
<b>Lee et al. [33]</b>	99	<ol style="list-style-type: none"> <li>1. Interlaminar ESI (42)</li> <li>2. Bilateral transforaminal ESI (57)</li> </ol>	<p>NRS, Patient Satisfaction Index (PSI), 5-point pain score at 2 weeks, 2 and 4 months  <b>2 V 1 at 2w, 2 and 4m NPRS / pain score (P&lt;0.05)</b></p>
<b>Malmivaara et al. [35]</b>	94	<ol style="list-style-type: none"> <li>1. Decompression surgery (50)</li> <li>2. NSAID, back school, some individualised physical therapy (44)</li> </ol>	<p>Oswestry, NRS, treadmill test at 6, 12, 24 months  <b>1 V 2 entire follow-up period: Oswestry (P=0.01), leg pain walking (P=0.02), LBP walking (P=0.0003)</b></p>
<b>Manchikanti et al. [36]</b>	40	<ol style="list-style-type: none"> <li>1. Caudal ESI + anaesthetic (20)</li> <li>2. Caudal epidural anaesthetic (20)</li> </ol>	<p>NRS, Oswestry at 3, 6, 12 months  NS differences; SD over time</p>
<b>Manchikanti et al. [37]</b>	50	<ol style="list-style-type: none"> <li>1. Caudal ESI + anaesthetic (25)</li> <li>2. Percutaneous adhesiolysis (25)</li> </ol>	<p>NRS, Oswestry at 3, 6, 12 months  <b>2 V 1 entire follow-up period NRS and Oswestry (P&lt;0.0001)</b></p>
<b>Mariconda et al. [38]</b>	44	<ol style="list-style-type: none"> <li>1 Decompression surgery (22)</li> <li>2. Bed rest, orthosis, physical therapy (22)</li> </ol>	<p>Beaujon Scoring System<sup>9</sup> at 1, 2 years, and mean 47 months:  <b>1 V 2 at 2 years / long-term (P≤0.05 / ≤0.01)</b></p>
<b>Matsudaira et al. [39]</b>	79	<ol style="list-style-type: none"> <li>1. Prostaglandin (39)</li> <li>2. Etodolac (NSAID) (40)</li> </ol>	<p>SF 36, rating scale for back and leg pain and walking distance, improvement, satisfaction at 8 weeks  <b>1 V 2 SF 36 physical functioning, bodily pain,</b></p>

**mental health (P<0.01), role physical (P=0.03), walking distance, leg numbness, improvement, satisfaction (P<0.01)**

<b>Ng et al. [40]</b>	32 <sup>d</sup>	1. PRI bupivacaine (15) 2. PRI bupivacaine + steroid (17)	VAS back / leg pain, walking distance at 2, 4, 6, 12 weeks: NS difference in walking distance (only outcome separate data for spinal stenosis)
<b>Podichetty et al. [45]</b>	55	1. Nasal calcitonin (36) 2. Placebo (19)	VAS, Oswestry, walking distance, SF 36 at 6 weeks: NS differences
Porter & Miller [47]	42	1. Injected calcitonin (20) 2. Injected saline (placebo) (22)	Walking distance, pain, sleep, mobility, analgesics at 4, 8 weeks: NS differences
<b>Pua et al. [48]</b>	68	1. BWST <sup>b</sup> (33) 2. Cycling <sup>b</sup> (35)	Oswestry, Roland-Morris, VAS, patient-rated change at 3, 6 weeks: NS differences, SD over time
Sahin et al. [50]	45	1. Nasal calcitonin <sup>c</sup> (23) 2. Paracetamol <sup>c</sup> (22)	VAS, ROM, Roland-Morris, walking distance at 8 weeks: NS differences, SD over time
<b>Slatis et al. [53]</b>	83	1. Decompression surgery (45) 2. NSAID, back school, some individualised physical therapy (38)	Oswestry, NRS, treadmill test at mean 6 years NS difference; SD over time
<b>Tafazal et al. [56]</b>	40	1. Nasal calcitonin (20) 2. Placebo (20)	VAS back / leg pain, Oswestry, LBOS, walking distance at 4, 10, 16 weeks: NS differences
<b>Tafazal et al. [57]</b>	48 <sup>d</sup>	1. PRI bupivacaine (25) 2. PRI bupivacaine + steroid (23)	VAS back / leg pain, Oswestry, LBOS at 6, 12 weeks 1 year: <b>Oswestry at 3 months (P=0.04)</b>
<b>Uratsuji et al. [66]</b>	84	1. Opalman (30 mgd) <sup>h</sup> (29) 2. Opalman (15 mgd) <sup>h</sup> (32)	Self-reported improvement, functional tasks at 6 weeks. NS differences

		3. Opalman (6 mgd) <sup>h</sup> (23)	
Waikakul & Waikakul [67]	152	1. Methylcobalamin (70) 2. Control (82) Both groups – education, strengthening exercises, physical therapy, NSAID.	Pain on movement, ROM, SLR, Neurology, walking distance, medication at 6, 12, 18, 24 months: <b>1 v 2 at 6, 12, 18 months: walking distance (P&lt;0.05)</b>
Weinstein et al. [69]	304 <sup>e</sup>	1. Decompressive surgery (159) 2. Usual care <sup>f</sup> (145)	SF 36, Oswestry at 6 weeks, 3, 6, 12, 24 months: NS differences
Weinstein et al. [70]	289 <sup>e</sup>	1. Decompressive surgery (138) 2. Usual care <sup>f</sup> (151)	SF 36, Oswestry at 6 weeks, 3, 6, 12, 24 months: <b>1 V 2 SF 36 bodily pain at 2 years (P not stated)</b>
Weinstein et al. [71]	304 <sup>e</sup>	1. Decompressive surgery (159) 2. Usual care <sup>f</sup> (145)	SF 36, Oswestry at 3, 4 years: NS differences
<b>Whitman et al. [72]</b>	58	1. MT, BWST, flexion exercises (29) 2. Flexion exercises, walking, US (29)	Global rating of change (GRC), Oswestry, NPRS, SSS at 6 weeks, 1 year: <b>1 V 2 GRC at 6 weeks (P=0.0015)</b> . Other outcomes NS differences
Yaksi et al. [74]	55	1. Flexion / strengthening exercises traction, corset, NSAID (27) 2. As 1 + gabapentin (28)	Walking distance, VAS with movement, neurological deficit at 1, 2, 3, 4 months: <b>2 V 1 walking distance at 2, 3, 4 months (P=0.03, 0.04, 0.001); VAS at 3, 4 months (P=0.039, 0.006); improvement sensory loss at 4 months (P=0.04)</b>
Zucherman et al. [76]	191	1. Decompression surgery (100) 2. ESI (NSAID, physical therapy) (91)	SF 36, ZCQ, ZCS at 6 weeks, 6, 12 months: <b>1 V 2 at all time points ZCQ (P not stated), and SF 36 (P not stated)</b>
Zucherman et al. [77]	191	1. Decompression surgery (100)	ZCQ at 2 years:

2. ESI (NSAID, physical therapy) (91) **1 V 2 all domains of ZCQ (P<0.001)**

<sup>a</sup> = significant differences in bold (with more effective treatment given first)

<sup>b</sup> = in addition both groups received warm-up (heat, traction) and home flexion exercise programme

<sup>c</sup> = in addition both groups did exercise programme (heat, flexion and stabilisation exercises)

<sup>d</sup> = spinal stenosis patients only, trial also included patients with disc herniations

<sup>e</sup> = in randomised controlled trial, more patients in an observational study

<sup>f</sup> = physical therapy, ESI, education, home exercises, NSAID

<sup>g</sup> = combination: walking distance, leg pain rest / exertion, back pain, neurological deficit, medication, quality of life

<sup>h</sup> = mgd = micrograms per day; Japanese trademark name for prostaglandin E

<sup>j</sup> = high quality ( $\geq 6$  on PEDro scale) in bold

BWST = body-weight supported treadmill; ESI = epidural steroid injection; LBOS = Low Back Outcome Score; LBP = low back pain; MT = manual therapy; NPRS = Numeric Pain Rating Scale; NRS = Numeric (pain) Rating Scale; NS = not significant; NSAID = non-steroidal anti-inflammatory drugs; PRI = periradicular infiltration; ROM = range of movement; SD = significant difference; SSS = Spinal Stenosis Scale; US = ultrasound; VAS = visual analogue scale; ZCQ = Zurich Claudication Questionnaire; ZCS = Zurich Claudication Score.